

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Solid lipid nanoparticles and microemulsions for drug delivery The CNS.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/75935> since

Published version:

DOI:10.1016/S0079-6123(08)80010-6

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

H.S. Sharma (Ed.)

Progress in Brain Research, Vol. 180

ISSN: 0079-6123

Copyright © 2009 Elsevier B.V. All rights reserved.

CHAPTER 10

Solid lipid nanoparticles and microemulsions for drug delivery: the CNS

Maria Rosa Gasco¹, Lorenzo Priano^{2,3} and Gian Paolo Zara⁴

¹Nanovector s.r.l, Torino, Italy

²Department of Neurosciences, University of Turin, Torino, Italy

³IRCCS – Istituto Auxologico Italiano, Ospedale S. Giuseppe – Piancavallo, Oggebbio, Italy

⁴Department of Anatomy Pharmacology and Forensic Medicine, University of Turin, Torino, Italy

Abstract: The chapter examined solid lipid nanoparticles (SLN) and microemulsions, chosen as carriers of drugs, administered *in vivo* to be transported to the central nervous system. Drugs of different structures and for different therapies have been studied such as doxorubicin SLN stealth and nonstealth administered in rats by intravenous route, apomorphine SLN administered in rats by duodenal route, melatonin SLN administered by transdermal and oral routes in humans, and apomorphine microemulsion administered by transdermal route in Parkinson's patients. The pharmacokinetics of the drug, followed in most studies, put in evidence that the many important pharmacokinetic parameters were notably improved versus the drug alone or in a commercial formulation.

Keywords: solid lipid nanoparticles; microemulsions; drug delivery system; central nervous system

Introduction

The brain homeostasis is of primary importance for survival so that specific interfaces, also referred to as barriers, tightly regulate the exchange between the peripheral blood circulation and the cerebrospinal fluid (CSF) circulatory system. These barriers are represented by the choroid plexus epithelium, the arachnoid epithelium, and the blood–brain barrier (BBB). The concentration and clearance of endogenous and exogenous molecules, essential for the normal brain functions or dangerous because of their toxicity, are strictly regulated by the anatomic and physiologic features of each barrier (Abbott, 2002; Segal, 2000).

The presence of the BBB is certainly the most critical issue encountered in brain drug delivery. Among the possible strategies to deliver therapeutic molecules into the brain, namely, intracerebral, intraventricular, and intravascular delivery, the latest represents the most reliable one because of its potential efficacy, safety, and compliance (Silva, 2007).

Brain capillaries, differently from the peripheral capillaries, present no fenestrae, a low amount of pinocytosis vesicles and particular tight junctions also known zonula occludens. Tight junctions are structures that form a narrow and continuous seal surrounding each endothelial and epithelial cell at the apical border and are at strictly regulating the movements the molecules through the paracellular

01 pathway. These structures, together with the brain
02 endothelial cells, make an almost impermeable
03 barrier for drugs administered through the
04 peripheral circulation (Kniesel & Wolburg, 2000;
05 Lapierre, 2000).

06 A further contribution to the peculiar BBB
07 functions is given by the periendothelial structures
08 represented by astrocytes, pericytes, and the basal
09 membrane (Balahanov & Dore-Duffy, 1998; Lay
10 & Kuo, 2005).

11 The presence of BBB transport systems further
12 complicates the scenario. In fact, these transport-
13 ers may assist or hinder the drug delivery to the
14 brain. The carrier-mediated transport may be able
15 to shuttle drugs or prodrugs into the brain in
16 therapeutic concentrations, mimicking nutrients
17 or endogenous compounds (Conford & Hyman,
18 1999; Pardridge, 1998).

19 Unfortunately, the presence of active efflux
20 transporters to the BBB also limits the therapeutic
21 efficacy of drugs virtually able to access the brain.
22 The P-glycoprotein (P-gp) is an ATP-dependent
23 drug transport protein present at the apical mem-
24 branes of different epithelial cell types including
25 those forming the BBB.

26 Recently, it has been demonstrated, either *in*
27 *vitro* or *in vivo*, that BBB P-gp can prevent the
28 accumulation of many molecules including a vari-
29 ety of drugs in the brain (Stouch & Gudmundsson,
30 2002), and P-gp inhibition has been proposed as a
31 possible strategy to enhance the drug penetration
32 (Skinkel, 1999).

33 Different strategies have been studied for the
34 delivery of drugs to the brain. Indeed most part of
35 the small drug molecules and of large molecules
36 such as recombinant proteins or gene-based mole-
37 cules are not able to penetrate the BBB and many
38 efforts have been spent in the previous years
39 toward delivery and targeting of drugs to the
40 brain (de Boer & Gaillard, 2007). Many investiga-
41 tions have been carried out in the previous years
42 to improve brain tumors therapy with nanoparti-
43 culates; there are less number of studies regarding
44 colloidal carriers of drugs for neurological diseases
45 or of diagnostics. Liposomes, polymeric nanopar-
46 ticles, and solid lipid nanoparticles (SLN) have
47 been studied, with different approaches, and the
48 problems of overcoming the BBB.

In this chapter, we consider SLN and microemulsions as carriers for the delivery only of drugs active on the central nervous system (CNS). In particular, examining drugs used for therapy in neurological diseases, as many times their administration gives problems, such as high amount of drug administered by parenteral route, short half-life, high hydrophilicity, and poor transport through the BBB. The aim of all the researchers is to study if some improvements in pharmacokinetic parameters in laboratory animals and/or in humans could be achieved using colloidal formulations; the review considers studies on SLN and microemulsions carrying only drugs active on CNS.

Solid lipid nanoparticles

Different approaches are followed for the SLN preparation.

They can be prepared by high-pressure homogenization at elevated or low temperatures, via warm microemulsions, by solvent emulsification–evaporation–diffusion, by high-speed stirring, and/or sonication (Muller, Kader, & Gohla, 2000).

Here we refer only about SLN carrying drugs active on CNS (at brain level).

SLN carrying the lipophilic antipsychotic drug clozapine were prepared by hot homogenization followed by ultrasonication method. Clozapine has a very poor bioavailability (Manjunath & Venkateswarlu, 2005). The SLN were administered by intravenous (IV) and duodenal routes to Swiss albino mice. For the intravenous administration, stearylamine was entrapped with clozapine in SLN; the area under curve (AUC) in the brain increased up to 2.91-fold the one of clozapine suspension.

The same authors (Manjunath & Venkateswarlu, 2006) developed SLN as carriers of the highly lipophilic drug nitrendipine, using different triglycerides for the lipid matrix, soy lecithin, and Poloxamer 188. Positive and negative charged nitrendipine SLN were also produced and then examined to explore the influence of the charge on oral bioavailability. The different kinds of SLN were administered by IV and intraduodenal

01 routes to rats; pharmacokinetic parameters of
 02 nitrendipine SLN were examined, tissue distribu-
 03 tion studies were carried out in Swiss albino mice,
 04 against that of a nitrendipine suspension. Follow-
 05 ing IV administration nitrendipine-loaded SLN
 06 were found to be taken up to a greater extent in
 07 tested organs than nitrendipine suspension. The
 08 AUC and MRT of nitrendipine SLN were higher
 09 than those of nitrendipine suspension especially in
 10 brain and heart. Positively charged SLN were bet-
 11 ter taken up by the brain and moderately taken up
 12 by the heart. Reticuloendothelial system (RES)
 13 organs such as liver and spleen were compared
 14 with the ones after nitrendipine suspension admin-
 15 istration. The higher levels of the drug were main-
 16 tained for over 6 h in confront to only 3 h with
 17 nitrendipine suspension.

AU3

AU4

18 SLN were investigated for their ability to deli-
 19 ver quinine dihydrochloride for the management
 20 of cerebral malaria (Gupta, Jain, & Jain, 2007).
 21 Quinine was incorporated in SLN and successively
 22 coupling of SLN with transferrin (Tf) was
 23 achieved by a cross-linker. IV administration of
 24 Tf-conjugated SLN enhanced the brain uptake of
 25 quinine in confront to the SLN loaded of quinine
 26 alone.

27 In order to enhance the delivery of atazanavir, a
 28 HIV protease inhibitor, spherical SLN carrying
 29 the drug were tested at first using a well-charac-
 30 terized human brain microvessel endothelial cell
 31 line (hCMEC/D3). Cell viability experiments
 32 demonstrated that SLN exhibit no toxicity on
 33 hCMEC/D3 cells up to a concentration corre-
 34 sponding to 200 nM of the drug. Delivery of ³H-
 35 atazanavir by SLN led to a significantly higher
 36 accumulation by the endothelial cell monolayer
 37 as compared to the drug aqueous solution (Chat-
 38 topadhyay, Zastre, Wong, Wu, & Bendayan,
 39 2008).

40 The transport *in situ* of lipid nanoparticles to the
 41 brain was evaluated by Koziara, Lockman, Allen,
 42 and Mumper (2003); the lipidic nanoparticles were
 43 prepared by warm microemulsion precursors fol-
 44 lowed by hot homogenization technique. Their
 45 components were emulsified wax (E wax) or Brij
 46 72 as matrix, and water and Brij 78 as surfactant.
 47 The warm microemulsion was cooled upon stir-
 48 ring and the lipid SLN were obtained and

homogenized. The SLN were labelled with ³H
 cetyl alcohol. The transport of the nanoparticles
 was measured by an “*in situ*” rat brain perfusion
 method; significant uptake of SLN was obtained .
 suggesting CNS uptake. The same group studied
 also the effect that the addition of a thiamine
 ligand to NPs, obtained by microemulsion as pre-
 cursors, causes association with the BBB thiamine
 transporter (Lockman et al., 2003).

AU5

Muller and coworkers studied the preferential
 adsorption of blood protein onto intravenously
 injected particulate carriers from different origins
 (Luck, Paulke, Schroder, Blunk, & Muller, 1998);
 in particular, Apolipoprotein E (Apo E) on the
 surface of P80-coated SLN after their incubation
 in human plasma citrate. Delivery to the brain
 using nanoparticulate drug carriers in combination
 with the targeting principles of “differential pro-
 tein adsorption” has been proposed (Dehouck
 et al., 1997). The Pathfinder technology (Muller
 & Schmidt, 2002) exploits proteins present in the
 blood which absorb onto the surface of intrave-
 nously injected carriers for targeting nanoparticles
 to the brain. Apo E is one of such targeting mole-
 cules for the delivery of nanoparticles to the
 endothelial cells of the BBB. Apo E can play an
 important role in the transport of lipoprotein into
 brain via the low-density lipoprotein receptor pre-
 sent on the BBB. Atoquavone (Muller & Keck,
 2004; Scholler et al., 2001) is a drug poorly
 adsorbed after oral administration, showing poor
 therapeutic efficacy against toxoplasma encephal-
 itis (TE). Nanocrystals of the drug were pro-
 duced, their surface was modified with Tween 80
 leading to *in vivo* preferential absorption of Apo
 E; the nanosuspension was IV administered to a
 murine model of TE, obtaining the disappearance
 of parasites and of cysts at dose 10-fold smaller
 than the one of atoquavone administered by oral
 route.

AU6

Solid lipid nanoparticles from warm microemulsions

SLN can be achieved from warm microemulsions.
 Warm microemulsions are prepared at tem-
 perature ranging from 60°C to 80°C by using

AU7

01 melted lipids (such as triglycerides/fatty acids) as
02 oil, surfactants such as lecithin, and cosurfactants
03 (such as short-chain carboxylates, biliar salts); the
04 warm microemulsions are subsequently dispersed
05 in cold water. The nanodroplets of warm micro-
06 emulsion, using this procedure, become SLN; they
07 are successively washed by tangential flow filtra-
08 tion. SLN are spherical in shape and with a narrow
09 size distribution. The zeta potential is normally
10 high (30/40 mV) being positive or negative
11 depending on the starting formulation.

12 Hydrophilic and lipophilic molecules (drugs or
13 diagnostics) can be incorporated in SLN using
14 different methods.

15 SLN are able to carry drugs of different struc-
16 ture and lipophilicity, such as cyclosporine A
17 (Ugazio, Cavalli, & Gasco, 2002), paclitaxel
18 (Cavalli, Caputo, & Gasco, 2000), doxorubicin
19 (Fundaro, Cavalli, Bargoni, Vighetto, & Gasco,
20 2000), tobramycin (Cavalli et al., 2003), short-
21 chain fatty acids (Dianzani et al., 2006), peptides
22 (Morel, Cavalli, & Gasco, 1996), antisense oligo-
23 nucleotides (Brioschi et al., 2008), and melatonin
24 (MT) (Rezzani et al., 2009). Also diagnostic com-
25 pounds such as iron oxides (Pereira, 2003) have
26 been incorporated into SLN.

27 SLN can be internalized within 2–3 min into all
28 the tested cell lines (Miglietta, Cavalli, Bocca,
29 Gabriel, & Gasco, 2000; Serpe et al., 2006); admin-
30 istered by duodenal route and are targeted to
31 lymph (Bargoni et al., 1998). SLN stealth can
32 also be prepared to avoid their recognition by
33 the RES, thus prolonging their residence time
34 (Podio, 2001). SLN drug, unloaded or loaded,
35 stealth/or nonstealth, are transported through the
36 BBB (Podio, 2001; Zara et al., 2002).

39 *Drug-loaded solid lipid nanoparticles*

40
41 In the late 1990s SLN were proposed for brain
42 drug targeting by several groups (Yang, Zhu, Lu,
43 & Liang, 1999; Zara et al., 1999), which studied
44 the pharmacokinetics of two anticancer agents:
45 camptothecin and doxorubicin. After oral and IV
46 administration, they observed drug accumulation
47 into the brain.

Both stealth and nonstealth stearic acid
unloaded labelled SLN were found in rat CSF
20 min after IV administration even though low
amount of radioactivity was found in the CSF
samples collected from cysterna magna (Podio,
Zara, Carazzone, Cavalli, & Gasco, 2000).

When the same kind of SLN were loaded with
doxorubicin, significantly higher drug concentra-
tions were found in the brain of the animals treat-
ed with stealth SLN as compared to nonstealth
SLN and doxorubicin solution. The overall plasma
pharmacokinetics of stealth and nonstealth SLN
provided to be significantly different from that of
the doxorubicin solution (Fundaro et al., 2000).

R-apomorphine (10,11-dihydroxyapomorphine)
is a well-known potent short-acting dopamine ago-
nist at D1 and D2 dopamine receptors and it was
proposed as an antiparkinsonian drug more than a
century ago. It significantly reduces the severity
and duration of “off” periods and it is able to
reverse bradykinesia when administered alone.
Despite these favorable clinical effects, the drug’s
clinical use is somewhat limited by its pharmaco-
kinetic profile: short half-life (~30 min), rapid
clearance from the plasma, lack of storage and
retention in brain regions, poor oral bioavailabil-
ity (5%), and first-pass hepatic metabolism are
significant limitations to chronic oral administra-
tion. Our group evaluated a new formulation of
apomorphine in SLN (submitted data for publish-
ing); the study was designed to investigate the
pharmacokinetics and biodistribution of apomor-
phine incorporated in SLN, injected orally or
intravenously in rats.

In vitro the release over time of apomorphine
from the SLN dispersion was almost linear. After
IV administration the peak plasma concentration
was higher after apomorphine solution adminis-
tration than after apomorphine SLN. However,
the total area under curve (AUC_{tot}) was nonsigni-
ficantly different after SLN than apomorphine
solution. The terminal half-life was significantly
longer following apomorphine SLN.

Following intraduodenal administration we
found that the C_{max} and AUC_{tot} were significantly
higher with apomorphine SLN compared to apo-
morphine solution; on the contrary, the clearance

was shorter after apomorphine solution than after the SLN formulation.

In the brain the apomorphine concentration was significantly higher 30 min after apomorphine SLN IV administration versus solution; it was detected only at 4 h after apomorphine SLN injection.

After duodenal administration the drug was detectable in brain only at 30 min after apomorphine SLN administration. No drug was found neither at 4 h nor at 24 h after injection of either apomorphine SLN or the solution.

Furthermore, the free drug concentration was measured in human plasma and we showed that the release started after the absorption of the apomorphine SLN. We also measured the free apomorphine concentration in human blood over time. The amounts in question are relatively low, but may be sufficient to expect clinical effects when administered to parkinsonian patients. After apomorphine solution administration, the amounts of apomorphine determined in the plasma were by far lower than those from SLN, confirming previous studies on the duodenal administration of drug loaded and unloaded SLN (Fig. 1).

In order to furnish a general model for SLN-based delivery systems of drugs devoid of favorable pharmacokinetics, we have recently

incorporated MT in SLN (MT-SLN). MT has been chosen for our *in vivo* study because of its safeness in humans even at high dosages.

MT is a hormone produced by the pineal gland at night, involved in the regulation of circadian rhythms. For clinical purposes (mainly disorders of the sleep-wake cycle and insomnia in the elderly), exogenous MT administration should mimic the typical nocturnal endogenous MT levels, but its pharmacokinetics is not favorable due to its short half-life of elimination (DeMuro, Nafziger, Blask, Menhinick, & Bertino, 2000; Mallo et al., 1990). The pharmacokinetics of MT-SLN has been examined in humans after administration by oral and transdermal route (Priano et al., 2007). Three kinds of freeze-dried MT-SLN containing different amounts of MT were prepared and characterized: (a) MT-SLN: MT = 1.8% for *in vitro* experiments (average diameter: 85 nm, polydispersity index = 0.135); (b) MT-SLN: MT = 2% for transdermal application (average diameter = 91 nm, polydispersity index = 0.140); and (c) MT-SLN: MT = 4.13% for oral route (average diameter = 111 nm, polydispersity index = 0.189).

In vitro, MT-SLN produced a flux of MT of 1 $\mu\text{g}/\text{h}/\text{cm}^2$ through hairless mice skin, following a pseudo-zero-order kinetics (45). At the same time, *in vivo* study produced very interesting results,




Fig. 1. Plasma levels of free apomorphine and total apomorphine after duodenal administration of apomorphine solution or apomorphine SLN in rats.

confirming in humans that SLN can act as a reservoir that allows a constant and prolonged release of the included drugs (Peira et al., 2003). MT (3 mg) incorporated in SLN was orally administered at 8.30 a.m. to seven healthy subjects; for control purposes, 1 week later the same subjects received orally a standard formulation of MT at the same dose (3 mg) and again at 8.30 a.m. Compared to the MT standard solution, T_{max} observed after MT-SLN administration was delayed of about 20 min, while mean AUC and mean half-life of elimination were significantly higher (respectively $169,944.7 \pm 64,954.4$ pg/mL \times hour vs. $85,148.4 \pm 50,642.6$ pg/mL \times hour, $p = 0.018$; and 93.1 ± 37.1 min vs. 48.2 ± 8.9 min, $p = 0.009$). Even more, standard formulation and MT-SLN after oral administration produced similar peak plasma levels of MT, even if delayed of about half an hour in the case of MT-SLN. More interestingly, detectable and clinically significant MT plasma levels after MT-SLN oral administration were maintained for a longer period of time, suggesting that SLN orally administered to humans can yield a sustained release of the incorporated drug, a feature that could be particularly useful for molecules, such as MT, characterized by unfavorable kinetics (Priano et al., 2007). Previous studies

in laboratory animals indicated a probable targeting of SLN — either drug-loaded or unloaded — to lymph, after duodenal administration (Bargoni et al., 1998). Similarly, the significantly longer half-life of MT observed in the study of Priano et al. (2007) may suggest a targeting of MT-SLN to human lymph, even though the capsules used to administer SLN were not gastro-resistant. In fact, MT half-life of elimination has been calculated in about 40 min after an intravenous bolus and following oral administration low bioavailability and rapid clearance from plasma have been shown, primarily due to a marked first-pass hepatic metabolism. Moreover, pharmacokinetic analysis following transdermal administration of MT-SLN demonstrated that plasma levels of MT similar to those produced by oral administration may be achieved for more than 24 h (50). In 10 healthy subjects, SLN incorporating MT were administered transdermally by applying a patch at 8.30 a.m. and leaving it in place for 24 h. In this delivery system, MT absorption and elimination were slow (mean half-life of absorption = 5.3 ± 1.3 h; mean half-life of elimination = 24.6 ± 12.0 h) so that MT plasma levels above 50 pg/mL were maintained for at least 24 h (Figs. 2 and 3). Tolerability of MT-SLN administered transdermally or by oral




Fig. 2. MT plasma profile in humans after MT (♦) and MT-SLN (■) oral administration.




Fig. 3. MT plasma levels in humans at baseline and after MT-SLN transdermal administration (MT-SLN-TD).

route was good and no adverse effect occurred, apart from a predictable mild somnolence and transient erythema after gel application. This means that, at least at the doses used in that study (45), SLN administration via the oral or transdermal routes is safe.

In this context, we also tested transdermal MT-SLN for three consecutive nights in five patients suffering from delayed sleep phase syndrome (unpublished data), confirming the safeness of this formulation. Due to the small sample, however, the tendency of clinical benefits was present but statistical significance could not be reached, so that further investigations in larger samples are needed in order to evaluate the impact of this new formulation in clinical practice.

However, these very favorable results, obtained in humans administering MT-loaded SLN, clearly suggest that SLN can be considered effective *in vivo* delivery systems that could be suitably applied to different drugs, and in particular to those requiring prolonged high plasma levels but that have unfavorable pharmacokinetics. Finally, it must be stressed that, since doses and concentrations of drugs included in SLN can be varied, different plasma level profiles could be obtained, thus disclosing new chances for sustained delivery systems adaptable to a variety of clinical conditions (Priano et al., 2007).

Suitability of SLN to convey drugs into CNS is also confirmed by studies regarding baclofen included in SLN. Intrathecal baclofen administration represents the reference treatment for spasticity of spinal or cerebral origin. Nevertheless, surgical involvement together with risk of infection or catheter dysfunction may limit the number of potentially treatable patients (Dario & Tomei, 2004; Perot & Almeida-Silveira, 1994). In order to explore alternative and efficacious routes of administration, we studied a new pharmaceutical preparation characterized by SLN incorporating baclofen (baclofen-SLN) (submitted data for publishing). Baclofen concentration, after reconstitution with water of freeze-dried SLN, was 1.7 mg/mL. Groups of Wistar rats were injected intraperitoneally with physiological solution and unloaded SLN at 10 mL/kg (control groups), with baclofen-SLN (baclofen-SLN group), and baclofen solution (baclofen-sol group) at increasing dosages of 2.5, 5, 7.5, 8.5, and 10 mg/kg. At different times up to the fourth hour, efficacy testing was performed by means of H-reflex, while behavioral characterization was obtained using two scales validated for motor symptoms due to spinal lesions and sedation in rat models (Nemethy, Paroli, Williams-Russo, & Blanck, 2002; Tsunoda, Kuang, Tolley, Whitton, & Fujinami, 1998). Rats were sacrificed for detecting baclofen concentration in blood and tissue. Compared to baclofen-sol and control group, *H/M* amplitude curve after baclofen-SLN injection was characterized by a dose-dependent reduction at the first and second hours, so confirming efficacy, and a rebound increase at the fourth hour, indicating an unexpected belated spinal hyperexcitability (Fig. 4). Similarly, baclofen-SLN effect on behavioral scales was stronger compared to baclofen-sol group, with the maximum effects obtained at the first hour. Moreover, clinical effects were detectable after low dosages of baclofen-SLN (2.5 mg/kg) but only after higher dosages of baclofen-sol (7.5 mg/kg). After 4 h from the injection, only the rats treated with the higher dosages of baclofen-SLN still presented clinical signs consisting in sedation (8.5 mg/kg) or complete paralysis and piloerection (10 mg/kg). On the whole, these data suggest a dose-dependent modulation of spinal reflex excitability, which is

AU14




Fig. 4. H/M amplitude ratios after baclofen-solution (a) and after baclofen-SLN (b), at increasing doses, compared to control animal group. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

not so evident after administration of standard formulation of baclofen. Nevertheless, important cortical effects were also present. Clinical data were related with plasma and tissue concentrations. In fact, after 2 and 4 h only baclofen-SLN administration produced measurable baclofen plasma concentrations, with an almost linear decrease of baclofen appreciable for 4 h. On the contrary, undetectable amount of baclofen in plasma were noticed 2 h after administration of baclofen-sol. In brain, both the two formulations (baclofen in solution and in SLN) gave a maximum after 2 h but concentrations after SLN were almost twice the ones after solution. This last data might be due partly to the free drug already released and to baclofen-SLN overcoming the BBB. We realize that for clinical purposes this effect of baclofen-SLN is unwished, as it is responsible for sedation. However, baclofen-sol injections also produced sedation, even if weaker and corresponding to lower plasma concentrations, compared to baclofen-SLN. In conclusion, higher spinal and cortical effects of baclofen-SLN, compared to equivalent dosages of baclofen-sol, seem attributable to higher and more prolonged concentrations of drugs in plasma and brain.

As previously noted, unloaded SLN administered by duodenal route are targeted to lymph and the incorporated drug can be partly distributed in the brain; moreover, SLN can also be

prepared stealth for increasing their residence time (Bargoni et al., 1998; Fundaro et al., 2000; Podio, 2001; Zara et al., 2002). Other new studies will be directed toward a duodenal administration of baclofen-SLN stealth, not only for prolonging their residence time but also to target them to lymph, enhancing their bioavailability. Further research should also be directed toward the optimization of dosages and concentrations of baclofen included in SLN, in order to preserve the prolonged antispastic effect, peculiar of this new formulation, but devoid of clinically significant cortical effects.

Solid lipid nanoparticles as potential diagnostics

Superparamagnetic iron oxides are classified as contrast agents for magnetic resonance imaging (MRI). They are able to affect the water relaxation times T_1 and T_2 ; their ability in altering such properties is quantified by the parameter relaxivity. Iron oxides are able to affect preferentially the T_2 relaxation times of tissues (and are called T_2 -relaxing agents) while paramagnetic contrast agents such as Gd complexes affect mainly T_1 and are called T_1 -relaxing agents.

Iron oxides are insoluble in water; therefore, to be clinically used they must be transformed in modified colloids while their magnetic properties

AU15

01 should remain unchanged. The surface of the iron
02 oxide nanoparticles can be modified, covering
03 them by hydrophilic macromolecules; such as dex-
04 tran in the case of Endorem.

05 A research was performed in order to know
06 whether SLN can load iron oxides and whether
07 they are able to reach the brain. Two different
08 SLN, SLN-Fe^A and SLN^B containing iron oxides
09 were prepared from warm microemulsions and
10 studied at first *in vitro* (1). The comparison of
11 Fe-SLN was performed with Endorem. Both the
12 Fe-SLN preparations showed relaxometric prop-
13 erties similar to the ones of Endorem. The good
14 T_2 -relaxation-enhancing properties allow an *in*
15 *vivo* study of their distribution by MRI. Fe-
16 SLN^B, at the higher Fe concentration, were
17 administered IV to rats; the comparison was
18 performed with Endorem. Images obtained
19 after Endorem IV administration show early
20 modification, but soon return to baseline; these
21 findings are consistent with short Endorem
22 retention time in blood. Results from SLN-Fe^B
23 show a different behavior. For each part of the
24 brain, maximal SS is reached in the last images
25 (135 min after administration). SS increase from
26 the first to the last acquisition. This study shows
27 that after inclusion in SLN, Endorem becomes a
28 new type of contrast agent: Endorem is taken by
29 the liver and does not cross the BBB, while
30 Endorem containing SLN-Fe^B shows CNS
31 uptake. This means that SLN-Fe kinesis is
32 related to SLN and not to their iron oxide con-
33 tent as already seen.

36 Microemulsions

38 Microemulsions are transparent, thermodynami-
39 cally stable dispersions of water and oil, usually
40 stabilized by a surfactant and a cosurfactant. They
41 contain particles smaller than 0.1 μm . Microemul-
42 sions are often defined as thermodynamically
43 stable liquid solutions; the stability of microemul-
44 sions is a consequence of the ultralow interfacial
45 tension between the oil and water phases. A clear
46 distinction exists between microemulsion and
47 coarse emulsions. The latter are thermodynami-
48 cally unstable, droplets of their dispersed phase

are generally larger than 0.1 μm and consequently
their appearance is normally milky rather than
transparent.

The limits in the use of microemulsions in the
pharmaceutical field are chiefly from the need of
all the components to be acceptable, particularly
surfactants and cosurfactants — the amounts of
surfactants and cosurfactants required to form
microemulsions are usually higher than those
required for emulsions.

Recently, apomorphine was incorporated into
microemulsions to study whether they are a feasi-
ble vehicle for transdermal transport of this drug.
In the preparatory *in vitro* study (Peira, Scolari, &
Gasco, 2001), two different microemulsions whose
components were all biocompatible were studied:
the concentration of apomorphine was 3.9% in
each. Since apomorphine is highly hydrophilic, to
increase its lipophilicity, apomorphine–octanoic
acid ion pairs were formed. At pH 6.0, $\log P_{\text{app}}$
of apomorphine increased from 0.3 in the absence
of octanoic acid to $\log P_{\text{app}} = 2.77$ for a molar ratio
1:2.5 (apomorphine: octanoic acid). The flux of
drug from the two thickened microemulsions
through hairless mouse skin was, respectively,
100 and 88 $\mu\text{g}/\text{h}/\text{cm}^2$. The first formulation, having
the higher flux, was chosen for *in vivo* administra-
tion to Parkinson's patients.

For the *in vivo* study, 21 patients with idiopathic
Parkinson's disease who presented long-term
L-DOPA syndrome, motor fluctuations and pro-
longed “off” periods were selected (Priano et al.,
2004). Here, 10 g of apomorphine hydrochloride
(3.9%), included in microemulsion for transder-
mal delivery (Apo-MTD), was applied to a 100
 cm^2 skin area on the chest; the area was delimited
by 1-mm-thick biocompatible foam tape and cov-
ered with a polyester-based membrane and an
occlusive membrane to prevent evaporation. In
these conditions, a single layer of microemulsion
(1 mm thick) was directly in contact with the skin
surface and acted as a reservoir of apomorphine.
Apo-MTD was applied at 8.00 a.m. and left for 12
h. In all patients, except two, apomorphine was
detected in blood samples after a variable lag
time. Pharmacokinetic analysis revealed that epi-
cutaneous–transdermal apomorphine absorption
was rapid (mean half-life of absorption = 1.03 h)

01 with a variability among patients (half-life of
02 absorption, $SD = 1.39$ h). Mean C_{max} was above
03 the therapeutic range (mean $C_{max} = 42.81 \pm 11.67$
04 ng/mL), with a mean T_{max} of 5.1 ± 2.24 h. Ther-
05 apeutic concentrations of apomorphine were
06 reached after a mean latency of 45 min (range
07 18–125), and stable concentrations, above the
08 therapeutic range, continued for as long as Apo-
09 MTD was maintained in place. At the 12th hour,
10 Apo-MTD was removed, and the apomorphine
11 plasma concentration then decreased at a rate
12 comparable to that described for subcutaneous
13 administration (mean half-life of elimination
14 equal to 10.8 ± 1.93 h). C_{max} and AUC showed
15 good correlations with the reduction of “off” per-
16 iods duration and with the improvement of clinical
17 scores evaluating motor performances (r values
18 ranging from 0.49 to 0.56, with p values ranging
19 from 0.02 to 0.04). Apo-MTD overall tolerability
20 was good: systemic side effects were similar to
21 those caused by subcutaneous apomorphine injec-
22 tion (sleepiness, mild orthostatic hypotension, and
23 transient nausea), and in the case of nausea, they
24 were strictly related to the highest plasma level of
25 apomorphine. Moreover, regarding local side
26 effects, the large majority of patients (71.4%) pre-
27 sented a transient mild erythema at the site of
28 Apo-MTD application, with a complete regression
29 within 48 h, whereas only in two cases the
30 erythema lasted more than 3 days and required
31 local therapy. This study clearly demonstrated
32 that in most Parkinson’s patients Apo-MTD is
33 absorbed by the epicutaneous–transdermal route.
34 This result is in contrast with other reports, where
35 the transdermal route did not produce detectable
36 plasma levels of apomorphine, or in which no apo-
37 morphine was transported passively through the skin
38 (Gancher, Nutt, & Woodward, 1991; van der Geest,
39 van Laar, Gubbens-Stibbe, Boddé, & Danhof,
40 1997). Probably, this difference was mainly due to
41 the peculiar pharmaceutical preparation used. Even
42 if pharmacokinetic parameters are variable, Apo-
43 MTD demonstrated the feasibility of providing ther-
44 apeutic apomorphine plasma levels for much longer
45 periods of time than previously tested apomorphine
46 preparations (several hours), allowing a more con-
47 stant dopaminergic stimulation. These results are
48 encouraging and Apo-MTD might become of

clinical value in some parkinsonian patients suffering
from uncontrolled “wearing-off” and prolonged
“off” phenomena. On the contrary, because of the
lag time of about 1 h before therapeutic concentra-
tions are reached, Apo-MTD may not be the “ideal”
preparation for rapid relief of “off” periods.

Since Apo-MTD was found to provide constant
drug release over several hours, other studies have
been addressed to its use for the nocturnal sleep
disorders of Parkinson’s patients. Twelve parkin-
sonian patients underwent standard polysomno-
graphy on basal condition and during one night
treatment with Apo-MTD (applied to 100 cm^2
from 10 p.m. until 8 a.m.; Priano et al., 2003).
Sleep analysis during APO-MTD treatment in
comparison to basal condition showed very favor-
able findings: 16% increment of total sleep time,
12% increment of sleep efficiency, 16% increment
of stage 3 and 4 nonrapid eye movement
(NREM), 15% reduction of periodic limb move-
ments index, 22% reduction of arousal index, and
23% reduction of the “cycling alternating pattern”
rate, an objective measure of disruption and frag-
mentation of NREM sleep. Pharmacokinetic anal-
ysis confirmed the absorption of apomorphine
and the maintenance of therapeutic plasma levels
for several hours (mean $C_{max} = 31.8 \pm 9.7$ ng/mL;
mean $T_{max} = 3.1 \pm 1.6$ h; mean half-life of absorp-
tion = 1.2 ± 1.4 h; mean half-life of elimination =
 8.8 ± 1.9 h). On the whole, this study confirmed
that APO-MTD in Parkinson’s disease might be
able to reduce nocturnal anomalous movements,
akinesia, and rigidity, and might be efficacious for
reducing the instability of sleep maintenance typi-
cal of parkinsonian sleep.

References

- Abbott, N. J. (2002). Astrocyte-endothelial interactions and blood-brain-barrier permeability. *Journal of Anatomy*, 200, 629–638.
- Balahanov, R., & Dore-Duffy, P. (1998). Role of the CNS microvascular pericyte in the blood-brain barrier. *Journal of Neuroscience Research*, 53, 637–644.
- Bargoni, A., Cavalli, R., Caputo, O., Fundaro, A., Gasco, M. R., & Zara, G. P. (1998). Solid lipid nanoparticles in lymph and plasma after duodenal administration to rats. *Pharmaceutical Research*, 15, 745–750.
- Brioschi, A., Calderoni, S., Pradotto, L. G., Guido, M., Strada, A., Zenga, F., et al. (2008). Solid lipid nanoparticles

- 01 carrying oligonucleotides inhibit vascular endothelial grow
02 factor expression in rat glioma models. *Journal of Nanoneur-*
03 *oscience, 1*, 1–10.
- 04 Cavalli, R., Caputo, O., & Gasco, M. R. (2000). Preparation
05 and characterization of solid lipid nanospheres containing
06 paclitaxel. *European Journal of Pharmaceutical Science, 10*,
07 305–308.
- 08 Cavalli, R., Bargoni, A., Podio, V., Muntoni, E., Zara, G. P., &
09 Gasco, M. R. (2003). Duodenal administration of solid lipid
10 nanoparticles loaded with different percentages of tobramy-
11 cin. *Journal of Pharmaceutical Sciences, 92*, 1085–1094.
- 12 Chattopadhyay, N., Zastre, J., Wong, H. L., Wu, X. Y., &
13 Bendayan, R. (2008). Solid lipid nanoparticles enhance the
14 delivery of the HIV protease inhibitor, atazanavir, by a
15 human brain endothelial cell line. *Pharmaceutical Research,*
16 *25*, 2262–2271.
- 17 Conford, F. M., & Hyman, S. (1999). Blood-brain barrier per-
18 meability to small and large molecules. *Advanced Drug*
19 *Delivery Reviews, 36*, 145–163.
- 20 Dario, A., & Tomei, G. (2004). A benefit-risk assessment of
21 baclofen in severe spinal spasticity. *Drug Safety, 27*, 799–818.
- 22 de Boer, A. G., & Gaillard, P. J. (2007). Drug targeting to the
23 brain. *Annual Review of Pharmacology and Toxicology, 47*,
24 323–355.
- 25 Dehouck, R., Ferrari, L., Dehouck, M. P., Pierce, A.,
26 Torpier, G., & Cecchelli, R. (1997). A new function for the
27 LDL receptor. Transcytosis of LDL across the blood brain
28 barrier. *Journal of Cell Biology, 38*, 877–889.
- 29 DeMuro, R. L., Nafziger, A. N., Blask, D. E., Menhinick, A. M.,
30 & Bertino, J. S. (2000). The absolute bioavailability of oral
31 melatonin. *Journal of Clinical Pharmacology, 40*, 781–784.
- 32 Dianzani, C., Cavalli, R., Zara, G. P., Gallicchio, M., Lombardi, G.,
33 Gasco, M. R., et al. (2006). Cholesteryl butyrate solid lipid
34 nanoparticles inhibit adhesion of human neutrophils to endothe-
35 lial cells. *British Journal of Pharmacology, 148*, 648–656.
- 36 Fundaro, A., Cavalli, R., Bargoni, A., Vighetto, G. P., & Gasco,
37 M. R. (2000). Non-stealth and stealth solid lipid nanoparti-
38 cles (SLN) carrying doxorubicin: Pharmacokinetics and tis-
39 sue distribution after i.v. administration to rats.
40 *Pharmacological Research, 42*, 337–343.
- 41 Gancher, S. T., Nutt, J. G., & Woodward, W. R. (1991).
42 Absorption of apomorphine by various routes in parkinson-
43 ism. *Movement Disorders, 6*, 212–216.
- 44 Gupta, Y., Jain, A., & Jain, S. K. (2007). Transferrin conjugate
45 solid lipid nanoparticles for enhanced delivery of quinine
46 dihydrochloride to the brain. *Journal of Pharmacy and Phar-*
47 *macy, 59*, 935–940.
- 48 Kniesel, K., & Wolburg, H. (2000). Tight junctions of the blood
brain barrier. *Cellular and Molecular Neurobiology, 20*, 57–76.
- Koziara, J. M., Lockman, P. R., Allen, D. D., & Mumper, R. J.
(2003). In situ blood-brain barrier transport of nanoparticles.
Pharmaceutical Research, 20, 1772–1777.
- Lapierre, L. A. (2000). The molecular structure of the tight
junctions. *Advanced Drug Delivery Reviews, 41*, 255–264.
- Lay, C. H., & Kuo, K. H. (2005). The critical component to
establish in vitro BBB model: Pericyte. *Brain Research*
Reviews, 50, 258–265.
- Lockman, P. R., Oyewumi, M. O., Koziara, J. M., Roder, K. E.,
Mumper, R. J., & Allen D. D. (2003). Brain uptake of thiamine-
coated nanoparticles. *Journal of Controlled Release, 93*, 271–282.
- Luck, M., Paulke, B. R., Schroder, W., Blunk, T., & Muller,
R. H. (1998). Analysis of plasma protein adsorption on poly-
meric nanoparticles with different surface characteristics.
Journal of Biomedical Materials Research, 39, 478–485.
- Mallo, C., Zaidan, R., Galy, G., Vermeulen, E., Brun, J.,
Chazot, G., et al. (1990). Pharmacokinetics of melatonin in
man after intravenous infusion and bolus injection. *Eur-*
opean Journal of Clinical Pharmacology, 38, 297–301.
- Manjunath, K., & Venkateswarlu, V. (2005). Pharmacokinetics,
tissue distribution and bioavailability of clozapine solid lipid
nanoparticles after intravenous and intraduodenal adminis-
tration. *Journal of Controlled Release, 107*, 215–228.
- Manjunath, K., & Venkateswarlu, V. (2006). Pharmacokinetics,
tissue distribution and bioavailability of nitrendipine solid
nanoparticles after intravenous and intraduodenal adminis-
tration. *Journal of Drug Targeting, 14*, 632–645.
- Miglietta, A., Cavalli, R., Bocca, C., Gabriel, L., & Gasco,
M. R. (2000). Cellular uptake and cytotoxicity of solid lipid
nanospheres (SLN) incorporating doxorubicin or paclitaxel.
International Journal of Pharmaceutics, 210, 61–67.
- Morel, S., Cavalli, R., & Gasco, M. R. (1996). Thymopentin in
solid lipid nanoparticles. *International Journal of Pharmaceu-*
tics, 132, 259–262.
- Muller, R. H., Kader, K., & Gohla, S. (2000). Solid lipid nano-
particles (SLN) for controlled drug delivery — a review of
the state of the art. *European Journal of Pharmaceutics and*
Biopharmaceutics, 50, 161–177.
- Muller, R. H., & Schmidt, S. (2002). Path. Finder technol-
ogy for the delivery of drugs to the brain. *New Drugs, 2*,
38–42.
- Muller, R. H., & Keck, C. M. (2004). Challenges and solutions
for the delivery of biotech. Drugs — a review of drug nano-
crystal technology and lipid nanoparticle. *Journal of Biotech-*
nology, 113, 151–170.
- Nemethy, M., Paroli, L., Williams-Russo, P. G., & Blanck, T. J.
(2002). Assessing sedation with regional anesthesia: Inter-
rater agreement on a modified Wilson sedation scale.
Anesthesia and Analgesia, 94, 723–728.
- Pardrige, W. M. (1998). CNS drug design based on principles of
blood-brain barrier transport. *Journal of Neurochemistry, 70*,
1781–1792.
- Peira, E., Scolari, P., & Gasco, M. R. (2001). Transdermal
permeation of apomorphine through hairless mouse skin
from microemulsions. *International Journal of Pharmaceutics,*
226, 47–51.
- Peira, E., Marzola, P., Podio, V., Aime, S., Sbarbati, A., &
Gasco, M. R. (2003). In vitro and in vivo study of solid
lipid nanoparticles loaded with superparamagnetic iron
oxide. *Journal of Drug Targeting, 11*, 19–24.
- Perot, C., & Almeida-Silveira, M. I. (1994). The human H and
T reflex methodologies applied to the rat. *Journal of Neu-*
roscience Methods, 51, 71–76.
- Podio, V., Zara, G. P., Carazzone, R., Cavalli, R., & Gasco,
M. R. (2000). Biodistribution of stealth and non-stealth solid

- lipid nanoparticles after intravenous administration in rats. *Journal of Pharmacy and Pharmacology*, 52, 1057–1063.
- Priano, L., Albani, G., Brioschi, A., Guastamacchia, G., Calderoni, S., Lopiano, L., et al. (2003). Nocturnal anomalous movement reduction and sleep microstructure analysis in parkinsonian patients during 1-night transdermal apomorphine treatment. *Neurological Science*, 24, 207–208.
- Priano, L., Albani, G., Brioschi, A., Calderoni, S., Lopiano, L., Rizzone, M., et al. (2004). Transdermal apomorphine permeation from microemulsions: A new treatment in Parkinson's disease. *Movement Disorders*, 19, 937–942.
- Priano, L., Esposti, D., Esposti, R., Castagna, G., De Medici, C., Fraschini, F., et al. (2007). Solid lipid nanoparticles incorporating melatonin as new model for sustained oral and transdermal delivery systems. *Journal of Nanoscience and Nanotechnology*, 7, 3596–3601.
- Rezzani, R., Fabrizio Rodella, L., Fraschini, F., Gasco, M. R., Demartini, G., Musicanti, C., et al. (2009). Melatonin delivery in solid lipid nanoparticles: Prevention of cyclosporin A induced cardiac damage. *Journal of Pineal Research*, 46, 255–261.
- Segal, M. B. (2000). The choroid plexuses and the barriers between the blood and the cerebrospinal fluid. *Cellular and Molecular Neurobiology*, 20, 183–196.
- Scholler, N., Krause, K., Kayser, O., Muller, R. H., Borner, K., Hahn, H., et al. (2001). Atovaquone nanosuspensions show excellent therapeutic effect in a new murine model of reactivated toxoplasmosis. *Antimicrobial Agents and Chemotherapy*, 45, 1771–1779.
- Skinkel, A. H. (1999). P-glycoprotein, a gatekeeper in the blood-brain barrier. *Advanced Drug Delivery Reviews*, 36, 179–194.
- Serpe, L., Cavalli, R., Gasco, M. R., Muntoni, E., Cavalli, R., Panzanelli, P., et al. (2006). Intracellular accumulation and cytotoxicity of doxorubicin with different pharmaceutical formulations in human cancer cells. *Journal of Nanoscience and Nanotechnology*, 6, 3062–3069.
- Silva, A. S. (2007). Nanotechnology approaches for drug and small molecule delivery across the blood brain barrier. *Surgical Neurology*, 67, 113–116.
- Stouch, T. R., & Gudmundsson, O. (2002). Progress in understanding the structure-activity relationship of P-glycoprotein. *Advanced Drug Delivery Reviews*, 54, 315–328.
- Tsunoda, I., Kuang, L. Q., Tolley, N. D., Whitton, J. L., & Fujinami, R. S. (1998). Enhancement of experimental allergic encephalomyelitis (EAE) by DNA immunization with myelin proteolipid protein (PLP) plasmid DNA. *Journal of Neuropathology & Experimental Neurology*, 57, 758–767.
- Ugazio, E., Cavalli, R., & Gasco, M. R. (2002). Incorporation of cyclosporin A in solid lipid nanoparticles in solid lipid nanoparticles. *International Journal of Pharmaceutics*, 241, 341–344.
- van der Geest, R., van Laar, T., Gubbens-Stibbe, J. M., Boddé, H. E., & Danhof, M. (1997). Iontophoretic delivery of R- apomorphine — II: An in vivo study in patients with Parkinson's disease. *Pharmaceutical Research*, 14, 1804–1810.
- Yang, S. C., Zhu, J. B., Lu, Y., & Liang, C. Z. (1999). Body distribution of camptothecin solid lipid nanoparticles after oral administration. *Pharmaceutical Research*, 751–757.
- Zara, G. P., Cavalli, R., Fundaro, A., Bargonì, A., Caputo, O., & Gasco, M. R. (1999). Pharmacokinetics of doxorubicin incorporated in solid lipid nanospheres (SLN). *Pharmaceutical Research*, 40, 281–286.
- Zara, G. P., Cavalli, R., Bargonì, A., Fundaro, A., Vighitto, D., & Gasco, M. R. (2002). Intravenous administration to rabbits of non-stealth and stealth doxorubicin loaded solid lipid nanoparticles at increasing concentration of stealth agent: Pharmacokinetics and distribution of doxorubicin in brain and in other tissues. *Journal of Drug and Targeting*, 10, 327–335.

AU18

01 Chapter No: 10

02

03 Query No Contents

04

1 Please check the insertion of the colon after “delivery:” in the chapter title.

05

2 Please indicate the corresponding author.

06

3 Please provide the full form of ‘MRT’.

07

4 Please check throughout if the text “in confront” can be changed to “in contrast”.

08

5 Please provide the full form of ‘NP’.

09

6 Please check if the abbreviation of ‘apolipoprotein E’ is OK.

10

7 Please check if the unit for temperature is correct.

11

8 Please check if it is “biliar” or “bile”?

12

9 Please check the change of “short fatty acids” to “short-chain fatty acids”.

13

10 Reference “Pereira, 2003” is not listed in the reference list. Please provide.

14

11 Reference “Podio, 2001” is not listed in the reference list. Please provide.

15

12 Please provide the reference for “submitted data for publishing”.

16

13 Please check throughout the chapter the presence of numbers within the brackets, for example, “(45), (50), (1)”. Can they be deleted or are they references?

17

14 Please provide the reference for “submitted data for publishing”.

18

15 Please check if the word “unwished” can be changed to “undesirable/unwanted”.

19

16 The compound “SLN^B” is not clear. Please check throughout. Is it “SLN-Fe^B” or “Fe-SLN^B”.

21

17 Please provide the full form of ‘SS’.

22

18 Please provide volume details for Reference “Yang et al. 1999”.

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48