




# Solution Phase and Surface Photoisomerization of a Hydrazone Switch with a Long Thermal Half-Life

## Journal Article

### Author(s):

[Zheng, Liqing](#) ; [Yang, Sirun](#); [Lan, Jinggang](#); [Gyr, Luzia](#); [Goubert, Guillaume](#) ; [Qian, Hai](#); [Zenobi, Renato](#) 

### Publication date:

2019

### Permanent link:

<https://doi.org/10.3929/ethz-b-000375337>

### Rights / license:

[In Copyright - Non-Commercial Use Permitted](#)

### Originally published in:

Journal of the American Chemical Society 141(44), <https://doi.org/10.1021/jacs.9b07057>

### Funding acknowledgement:

741431 - Nanoscale Vibrational Spectroscopy of Sensitive 2D Molecular Materials (EC)

# Solution phase and surface photoisomerization of a hydrazone switch with long thermal half-life

Li-Qing Zheng<sup>a</sup>, Sirun Yang<sup>b</sup>, Jingtang Lan<sup>c</sup>, Luzia Gyr<sup>a</sup>, Guillaume Goubert<sup>a</sup>, Hai Qian<sup>b,d</sup>, Ivan Aprahamian<sup>b\*</sup> and Renato Zenobi<sup>a\*</sup>

a. Department of Chemistry and Applied Biosciences, ETH Zurich, Vladimir-Prelog-Weg 3, CH 8093, Switzerland

b. Department of Chemistry, Dartmouth College, Hanover, New Hampshire 03755, United States

c. Department of Chemistry, University of Zurich, Winterthurerstrasse 190, CH 8057, Switzerland

d. Department of Chemistry, University of Illinois at Urbana-Champaign, 505 S Mathews Avenue, Urbana, 61801, United States

---

**ABSTRACT:** Photoswitches can be employed for various purposes, with the half-life being a crucial parameter to optimize for the desired application. The switching of a photochromic hydrazone functionalized with a C6 alkyl thiolate spacer (**C6 HAT**) was characterized on a number of metal surfaces. **C6 HAT** exhibits a half-life of 789 years in solution. Tip-enhanced Raman spectroscopy (TERS) was used to study the photoisomerization of the **C6 HAT** self-assembled monolayers (SAM) on Au, Ag and Cu surfaces. The unique spectroscopic signature of the *E* isomer at 1580 and 1730  $\text{cm}^{-1}$  in TER spectra allowed for its discrimination from the *Z* isomer. It was found that **C6 HAT** switches on Au and Cu surfaces when irradiated with 415 nm, however it cannot isomerize on Ag surfaces, unless higher energy light is used. Based on this finding, and supported by density functional theory calculations, we propose a substrate-mediated photoisomerization mechanism to explain the behavior of **C6 HAT** on these different metal surfaces. This insight into the hydrazone's switching mechanism on metal surfaces will contribute to the further exploitation of this new family photochromic compounds on metal surfaces. Finally, although we found that the thermal isomerization rate of **C6 HAT** drastically increases on metal surfaces, the thermal half-life is still 6.9 days on gold, which is longer than the majority of azobenzene-based systems.

---

## INTRODUCTION

Molecular switches play an important role in various fields, including biology, chemistry, materials science and electronics, because<sup>[1]</sup> their structures and functions can be precisely controlled at the molecular level, using external stimuli, such as light, pH, or electric current.<sup>[2]</sup> These capabilities have made them very promising candidates for molecular electronics and high-density data storage applications.<sup>[3]</sup> Moreover, surfaces modified with molecular switches have been patterned and functionalized to act as regulators in organic field-effect transistors and organic light emitting diodes.<sup>[4]</sup> Various types of photo switches have been investigated over the years for such and other applications, including azobenzenes<sup>[7]</sup>, diarylethenes<sup>[8]</sup>, spiropyrans<sup>[9]</sup>, and more recently hydrazones.<sup>[10]</sup> The thermal isomerization half-life of these systems is a crucial characteristic that determines the type of application they can be used for.<sup>[5]</sup> While these switches have a wide range of thermal half-lives, ranging from microseconds to thousands of years in solution,<sup>[11,12]</sup> these values do not translate well when they are attached to metal surfaces, and in general a dramatic enhancement in isomerization rates is observed.<sup>[12a]</sup> For example, the thermal half-life of azobenzene derivatives decreases from 18.7h in solution to 45.3s when self-assembled on gold surfaces.<sup>[13]</sup> Consequently, photoswitches with very long thermal half-lives (e.g., bistable) are needed to build future devices. Investigating their molecular switching behav-

ior on metal surfaces at the nanoscale can provide crucial information for future implementations.

When exposed to an external stimulus, molecular switches typically isomerize, for example by undergoing a ring-closing/opening reaction,<sup>[14]</sup> or a *cis/trans* isomerization.<sup>[10,15]</sup> A great deal of work has been devoted to understanding the switching mechanism of molecules in solution,<sup>[17]</sup> and the influence of solvent, temperature and substituents on the outcome.<sup>[17a]</sup> However, when a molecular switch is attached to a metal surface, the new molecular environment and surface effects, further complicate the switching mechanism.<sup>[18]</sup> For example, switching in the solid state is inhibited to a certain degree as a result of steric hindrance. The metal surface can alter the switching mechanism of the adsorbed species. One reason for this is that excited adsorbates can relax via non-radiative decay, namely via electron transfer to the conduction band of the surface and energy transfer to the Fermi level of the surface,<sup>[18b, 19]</sup> leading to a decrease in the switching efficiency. For example, the observation that azobenzene fails to isomerize on Au (100) shows a strong surface effect.<sup>[20]</sup> Since molecular adsorbates can couple with the high density of states in the metal conduction band, the question arises whether there are other effects that interfere with the switching of molecules on metal surfaces upon light irradiation. For instance, Van Duyne and coworkers recently employed pump-probe surface-enhanced Raman spectroscopy (SERS) to demonstrate that photoinduced plasmons at a gold nanoparti-

cle surface can drive the *trans*-to-*cis* isomerization of *trans*-1,2-bis(4-pyridyl) ethylene (BPE).<sup>[21]</sup>

To reveal the isomerization mechanism of molecular switches on a metal surface down to the single molecular layer level, analytical tools with high surface sensitivity and a low detection limit are needed. Tip-enhanced Raman spectroscopy (TERS) combines plasmon-enhanced Raman spectroscopy with scanning probe microscopy, and therefore provides chemical fingerprint information with high sensitivity and morphological information on the sample with nanometer spatial resolution.<sup>[22]</sup> TERS has recently been used to probe the isomerization of adsorbed azobenzene thiol (ABT), which was triggered by the bias voltage of a scanning tunneling microscope (STM) in ultra-high vacuum.<sup>[23]</sup> Using ambient TERS, we reported that the photoisomerization yield of ABT molecules at Au grain boundaries is higher than on Au terraces, due to reduced *cis*-to-*trans* back-reaction rate at the grain boundaries. The reduced back-reaction rate is a result of the lower reaction enthalpy of *trans*-to-*cis* isomerization at Au steps.<sup>[24]</sup> However, the lifetime of *cis* ABT on Au surfaces was found to be only 5 h, which is comparable to TERS acquisition times making the measurements difficult.


To address this issue we decided to investigate the isomerization mechanism of recently developed bistable hydrazone photoswitches that exhibit thermal half-lives of up to 5000 years in solution.<sup>[5,12c]</sup> Herein, we first describe the molecular design of a hydrazone thiol (HAT) switch having a 6-carbon spacer (C6 HAT), allowing it to be conjugated to Au via thiol chemistry. Subsequently, we elaborate on the photoisomerization of HAT switches in solution, the solid-state, and assembled as a monolayer on a metal surface, using ultraviolet visible (UV-Vis) spectroscopy and TERS. We used characteristic peaks in the Raman spectrum of C6 HAT to demonstrate its photoisomerization, and quantitatively measured the extent of its switching using UV-Vis spectroscopy. Furthermore, to reveal the photoisomerization mechanism of C6 HAT in a monolayer, we studied the isomerization of HAT on different metals (i.e., Au, Ag and Cu). These studies, in corroboration with density functional theory (DFT) calculations, allowed us to propose a substrate-mediated isomerization mechanism to explain the different isomerization rates of C6 HAT on various metal surfaces.

## RESULTS AND DISCUSSION

**Molecular design, synthesis and solution behavior.** We chose a hydrazone skeleton<sup>[5]</sup> that has a half-life of 255 years in toluene in our design of C6 HAT. To minimize any electronic and steric effects on the hydrazone, the alkanethiol group was attached to the ester moiety of the switch. A 6 carbon chain linker (C6 HAT, Figure 1A) was selected to minimize the length of the molecule, but still allow it to form a well-organized SAM on the surface.<sup>[25]</sup> C6 HAT was synthesized using a Mitsunobu reaction between an appropriate hydrazone derivative (Scheme S1) and 6-bromohexan-1-ol, followed by reaction with potassium thioacetate and deprotection with hydrazine in MeCN (56 % yield). C6 HAT was fully characterized using NMR spectroscopy and high-resolution mass spectrometry (Figures S2-S5 in the Supporting Information).

UV/Vis spectroscopy was utilized to study the switching behavior of C6 HAT in toluene. The absorption spectrum of C6 HAT-Z shows a maximum ( $\lambda_{\max}$ ) at 368 nm (Figure 1b).


Irradiation the solution with 410 nm results in a shift of the  $\lambda_{\max}$  to 333 nm. The photoisomerization efficiency of C6 HAT was studied using <sup>1</sup>H NMR spectroscopy. Upon 410 nm light irradiation a sample of C6 HAT-Z yields a photostationary state (PSS) consisting of 98% C6 HAT-E isomer (Figure S6 in the Supporting Information). In addition to NMR spectroscopy, the ratio of *E* and *Z* isomers was calculated by fitting two Gaussian peaks to the UV-Vis absorption spectrum. PSS<sub>410</sub> of 96.5 % C6 HAT-E were obtained by this fit, which is in good agreement with the NMR results (Figure S7 in the Supporting Information). The quantum yield ( $\Phi$ ) of the process was determined to be  $4.8 \pm 0.1$  % (Figure S8 in the Supporting Information). The reverse switching can be induced by irradiation of the *E* rich sample with 340 nm light. The absorption maximum ( $\lambda_{\max}$ ) red-shifts to 364 nm with a PSS<sub>340</sub> consisting of 74% (NMR spectroscopy) C6 HAT-Z isomer, and a  $\Phi_{E \rightarrow Z}$  of  $9.4 \pm 0.4$  % (Figure S9 in the Supporting Information). Fitting the absorption spectrum (dotted line in Figure 1B) yields a PSS<sub>340</sub> containing 84 % of the *Z* isomer (Figure S10 in the Supporting Information). The switching process was repeated 10 times by alternation of the irradiation wavelength between 410 and 340 nm. No signs of sample degradation were observed (Figure S11 in the Supporting Information). The half-life ( $\tau_{1/2}$ ) of the *E* isomer was measured to be  $789 \pm 52$  years at 298 K using model compound **2** (Table S1 in the supporting information).



**Figure 1.** A. Light induced *Z/E* isomerization of C6 HAT; B. UV-vis spectra of C6 HAT in toluene ( $1 \times 10^{-5}$  M) before (solid line) and after irradiation with 410 nm light (dashed line), followed by irradiation with 340 nm light (dotted line).

**Raman spectroscopic markers and solid-state photoisomerization.** DFT calculations (B3LYP/6-31G\*) were performed to predict the differences in the Raman spectrum between the *Z* and *E* isomers of **C6 HAT** (Figure 2A).<sup>[26]</sup> Theoretical frequencies were scaled by a factor of 0.98 to correct for anharmonicity.<sup>[27]</sup> The intensity of the Raman spectrum of the *E* isomer is half as strong as that of the *Z* one, most likely as a result of the loss of extended conjugation in the *E* isomer, which leads to a smaller Raman cross-section.<sup>[26]</sup> The Raman bands at 1436 and 1475  $\text{cm}^{-1}$  ( $\nu_{\text{C=N}}$ ) of the *Z* isomer shift to 1576  $\text{cm}^{-1}$  upon isomerization. Moreover, the band at 1668  $\text{cm}^{-1}$  ( $\nu_{\text{C=O}}$ ) blue shifts to 1738  $\text{cm}^{-1}$  because of the disruption of the intramolecular H-bond (see Movie S1 and S2 in the Supporting Information). Upon *Z*→*E* isomerization, the C-N stretching mode slightly shifts from 1508 to 1522  $\text{cm}^{-1}$  and a small peak appears at 1027  $\text{cm}^{-1}$ . The bands at 1600 and 989  $\text{cm}^{-1}$  correspond to the C=C stretching mode and the ring breathing mode of the benzene, respectively. Upon isomerization, these two peaks do not shift. Therefore, these two peaks were chosen as references.

Confocal Raman measurements on solid **C6 HAT** (powder of pure material, Figure 2B) are in good agreement with the calculated spectra. After 415 nm light irradiation the intensity of the peak at 1431  $\text{cm}^{-1}$  decreases, while the signals at 1472 and 1677  $\text{cm}^{-1}$  disappear. Moreover, two new peaks appear at 1562 and 1704  $\text{cm}^{-1}$ , and a smaller one at 1027  $\text{cm}^{-1}$ . All these spectral changes indicate that the *Z* isomer underwent *Z*→*E* isomerization. The peaks at 994 and 1605  $\text{cm}^{-1}$  remain the same.




**Figure 2.** A. Simulated Raman spectra of the *Z* and *E* isomers of **C6 HAT**; B. Confocal Raman spectra of **C6 HAT** in the solid-state before and after irradiation at 415 nm, followed by irradiation at 340 nm. The light-blue bars indicate the Raman peak at 1431, 1472, and 1677  $\text{cm}^{-1}$ , which are characteristic for

the *Z* isomer. The yellow bars indicate the Raman peak at 1562 and 1704  $\text{cm}^{-1}$  which are characteristic for the *E* isomer.

Only the peak shift at 1522  $\text{cm}^{-1}$ , predicted by DFT, is not observed. Overall, the Raman spectrum of **C6 HAT**, after irradiation at 415 nm, is in very good agreement with the calculated spectrum of the *E* isomer. Based on the intensity decrease in the bands at 1431, 1472 and 1677  $\text{cm}^{-1}$ , almost 100% conversion from *Z* to *E* is achieved in the solid-state after 415 nm light irradiation. Upon further irradiation with 340 nm light, the intensities of the bands at 1431, 1472 and 1677  $\text{cm}^{-1}$  partially recover. Meanwhile, the intensities of the peaks at 1562 and 1704  $\text{cm}^{-1}$  decrease. Based on the peak area ratio between 1677 and 1704  $\text{cm}^{-1}$  in the Raman spectrum, we can calculate the *E*:*Z* and *E* ratio to be 66:34 (see Figure S12), which is smaller than that observed in solution. Previously, we reported on the photoisomerization of another hydrazone switch in solution and in the solid state, monitored by UV-Vis spectroscopy.<sup>[28]</sup> The switching efficiencies of **C6 HAT** obtained by UV-Vis spectroscopy and Raman spectroscopy are comparable to our earlier results.<sup>[28]</sup>

**Photo isomerization of a C6 HAT monolayer on Au surface.** TER spectra of a **C6 HAT** SAM on a template stripped (TS) Au substrate before and after 415 nm light irradiation were collected, as shown in Figure 3A (top panel). After 415 nm light irradiation, the intensities of the peaks at 1430, 1472 and 1508  $\text{cm}^{-1}$  decrease, while the intensities of the peaks at 1013 and 1050  $\text{cm}^{-1}$  increase, and new peaks appear at 1580 and 1730  $\text{cm}^{-1}$ . All these spectral changes indicate that *Z*→*E* isomerization occurred. To switch molecules back to the *Z* state, we heated the sample to 50 °C for 16 h and collected a TER spectrum afterwards. The bands at 1430 and 1472  $\text{cm}^{-1}$  recover, while the peak at 1580  $\text{cm}^{-1}$  still remains, indicating that part of **C6 HAT-E** molecules isomerized back to the *Z* form (see Figure S13). In the TER spectra, no Raman peak related to carbonaceous contaminants are observed, which indicates that neither UV irradiation nor laser irradiation induce any photodamage in **C6 HAT**. The assumption that there is no desorption of **C6 HAT** from the Au substrate upon UV irradiation was tested and proven by temperature-programmed desorption mass spectrometry (see Figure S14-S16).

We further collected TERS maps of **C6 HAT** SAMs on Au before and after 415 nm light irradiation ( $1 \times 1 \mu\text{m}^2$  size with a  $32 \times 32 \text{nm}^2$  pixel size) to estimate the isomerization ratio. All 1064 TERS spectra within a map are displayed as a waterfall plot (Figure 3B) to highlight the spectral differences. In the TERS image of a **C6 HAT** SAM on Au prior to 415nm light irradiation, the characteristic Raman peaks at 1430, 1472 and 1508  $\text{cm}^{-1}$  of *Z*-**C6 HAT** are observed. However, because of the limited enhancement of the used Ag tip, the Raman band at 1680  $\text{cm}^{-1}$ , which corresponds to the C=O stretching mode of the *Z* isomer, was invisible in this map (top panel of Figure 3B). We, therefore, recorded another TERS map, in which the peak at 1680  $\text{cm}^{-1}$  is clearly observed, using another Ag tip with higher enhancement (Figure S17 in Supporting Information). After irradiation at 415 nm, Raman peaks at 999, 1013, 1050, 1600 and 1680  $\text{cm}^{-1}$  were clearly observed in the image, as shown in Figure 3B (bottom panel). Moreover, we could clearly see that the peaks at 1430 and 1472  $\text{cm}^{-1}$  disappear and new signals arise at 1580 and 1730  $\text{cm}^{-1}$  simultaneously. As shown in the TERS image (Figure S18 in Supporting Information), the peak at 1730  $\text{cm}^{-1}$  appears in 79% of




**Figure 3.** (A) Baseline corrected TER spectra (top panel) of a **C6 HAT** monolayer on Au surface before (black) and after (red) 415 nm light irradiation, and average TER spectra (bottom panel) from the TERS maps of a **C6 HAT** monolayer on Au before (black) and after (red) 415 nm light irradiation. (B) Waterfall plot of all TER spectra from maps of a **C6 HAT** monolayer on Au before (top panel) and after (bottom panel) 415 nm light irradiation. The size of all maps is  $1 \times 1 \mu\text{m}^2$  with a 31 nm pixel size. All TER spectra were normalized using the Raman peak at  $1602 \text{ cm}^{-1}$ .

**Table 1.** Photochemical properties of **C6 HAT**


Parameter	Toluene	Solid-state <sup>a</sup>	Monolayer on	
			Au	Cu
$\lambda_{\text{max}}$ , <i>Z</i> -isomer (nm)	368	— <sup>b</sup>	378	364
$\lambda_{\text{max}}$ , <i>E</i> -isomer (nm)	333	— <sup>b</sup>	343	330
( <i>[E]</i> : <i>[Z]</i> ) <sub>PSS415</sub> (%)	98:2	99:1	82:18	62:38
( <i>[E]</i> : <i>[Z]</i> ) <sub>PSS340</sub> (%)	26:74	66:34	— <sup>b</sup>	— <sup>b</sup>
$\tau_{1/2}$ <sup>c</sup>	$789 \pm 52$ years	— <sup>b</sup>	$6.90 \pm 0.03$ days	— <sup>b</sup>

<sup>a</sup>Powder sample. <sup>b</sup>Not determined. <sup>c</sup>Half life of the thermal *E*-to-*Z* transformation.

the spectra, indicating that some of the *Z*-**C6 HAT** molecules have isomerized to the *E* state. The isomerization ratio of the **C6 HAT** SAM is smaller than that of **C6 HAT** bulk material (see Table 1), which could result from electron or energy transfer from the excited states of **C6 HAT** to the Au surface.



**Figure 4.** UV-Vis spectra of a **C6 HAT** SAM on a Au surface before (black line) and after 415 nm light irradiation (red line). The blue line is the sample after heating under dark for 16 h.



**Figure 5.** (A) Baseline corrected average TER spectra from the TERS maps of a **C6 HAT** SAM on a Ag surface before and after 415 nm light irradiation. (B) UV-Vis spectra of a **C6 HAT** SAM on a Ag surface before (black line) and after 415 nm light irradiation (red and blue lines). (C) Baseline corrected average TER spectra from the TERS maps of a **C6 HAT** SAM on a Cu surface before and after 415 nm light irradiation. (D) UV-Vis spectra of a **C6 HAT** SAM on a Cu surface before (black line) and after 415 nm light irradiation (red line). The sample was then kept in the dark and heated to 50 °C for 16 h (blue line). The size of all the maps is  $1 \times 1 \mu\text{m}^2$ .

All TER spectra within a map were accumulated and averaged to obtain statistical information (bottom panel of Figure 3A). The average TER spectra of **C6 HAT** SAM before and after irradiation at 415 nm show similar spectral features compared to the TER spectra shown in Figure 3A (top panel). Gaussian functions were fitted to the peaks at 1680 and 1730  $\text{cm}^{-1}$  corresponding to the *Z* and *E* isomers, respectively. The area ratio of these two peaks was used to determine that 82 % of the molecules in the SAM are in the *E* form (Table 1, Figure S19 in Supporting Information), which is consistent with the calculation based on the probability of the presence of the peak at 1580  $\text{cm}^{-1}$  in the TERS map (Figure S18 in Supporting Information). We also performed TERS line scans along Au grain edges and terraces with a 10 nm pixel size to measure the isomerization ratio at different sites (Figure S20 in Supporting Information). The presence of the peak at 1580  $\text{cm}^{-1}$  both on terraces and at grain edges indicates that photoisomerization can happen at both sites. Because of their low Raman cross-section, the peaks at 1680 or 1730  $\text{cm}^{-1}$  were not observed in line-scan TER spectra.

The photoisomerization of **C6 HAT** SAMs on Au was further investigated by UV-Vis spectroscopy. As shown in Figure 4, the maximum absorption peak of *Z*-**C6 HAT** SAM appears

at 378 nm. Upon 415 nm light irradiation it shifts to 343 nm, which indicates that *Z*→*E* isomerization has taken place. The sample was then heated to 50 °C for 16 h, which resulted in a shift of the  $\lambda_{\text{max}}$  back to 365 nm and a decrease of absorbance. This result indicates that most of the *E*-**C6 HAT** has isomerized back to the *Z* state. The absorbance decrease could be due to the desorption of **C6 HAT** molecules from the surface upon heating. The discrete steps at 370 and 420 nm are due to the change of grating. A control experiment was conducted to check whether irradiation by 340 nm can induce the *E*-to-*Z* isomerization of **C6 HAT** on Au surface. A **C6 HAT** SAM on Au was first irradiated with 415 nm light and then followed by irradiation at 340 nm. The absorption peak of **C6 HAT** shifts back to 346 nm, which means that irradiation at 340 nm barely induce *E*→*Z* isomerization on the Au surface (see Figure S21).


The thermal lifetime of the *E* isomers on the Au surface is an important characteristic of the surface bound switch. The absorbance change at 340 nm of a **C6 HAT** SAM on a Au surface after 415 nm light irradiation as a function of time is shown in Figure S22. Analysis of the data shows that the thermal back isomerization on the surface is dominated by first-order kinetics with a rate constant of  $1.16 \times 10^{-6} \text{ s}^{-1}$ . The half-life of the *E* isomer on the Au surface is thus around 6.90

$\pm 0.03$  days, which is the longest value obtained for such systems, and higher than a recently reported azobenzene derivative on Au surface.<sup>[13]</sup> As a thermodynamically unfavorable state, the long lifetime of the *E* form is attributed to the high energy barrier for the *E*→*Z* isomerization.<sup>[5]</sup> With such a relatively long thermal lifetime on the metal surface, it is a promising system towards the development of optical data storage devices.

**Photo isomerization of a C6 HAT monolayer on Ag or Cu surface.** To study the effect of the nature of the metal on the photoisomerization of a C6 HAT SAM, we also studied its isomerization on Ag or Cu surface. A waterfall plot of the spectra extracted from TERS images of C6 HAT SAM on Ag before and after 415 nm light irradiation is shown in Figure S23. The average TERS spectra are shown in Figure 5A, in which the Raman bands at 1430, 1473, 1506 and 1680  $\text{cm}^{-1}$ , characteristic peaks for *Z*-C6 HAT, are still observed, and no peaks appear at 1580 or 1730  $\text{cm}^{-1}$ . This suggests that no *Z*→*E* isomerization of C6 HAT occurs on the Ag surface. Moreover, there is no absorption peak shift observed in the UV-Vis spectrum of C6 HAT SAM on Ag after 415 nm light irradiation (Figure 5B), which is in agreement with the TERS results. Next we studied the isomerization of a C6 HAT SAM on a Cu surface. As shown in Figure 5C, after 415 nm light irradiation, we can clearly see that the intensities of the peaks at 1430, 1473 and 1680  $\text{cm}^{-1}$  decrease, with small peaks appearing at 1580 and 1730  $\text{cm}^{-1}$  in the average TERS spectrum of C6 HAT on Cu. The spectral change is seen more clearly in the waterfall plot of the TERS spectra extracted from images (see Figure S24). Gaussian functions were again fitted to the peaks at 1680  $\text{cm}^{-1}$  and 1730  $\text{cm}^{-1}$  corresponding to *Z* and *E* isomers, respectively. Based on the area ratio of these two peaks, 62 % of C6 HAT molecules in the SAM have isomerized (Table 1, see Figure S25 in Supporting Information). The lower isomerization efficiency is the main reason why the average spectral change of the TERS spectrum of C6 HAT on Cu is smaller than that on the Au surface. (see Table 1). Moreover, the molecular orientation of C6 HAT on Cu is different from that of C6 HAT on Au. In figure 5C, the new peaks that appear at 1576  $\text{cm}^{-1}$  and 1732  $\text{cm}^{-1}$  are less obvious than the peak intensity decrease at 1427 and 1470  $\text{cm}^{-1}$ , which is likely due to the surface selection rules of TERS. The UV-Vis spectroscopic results also support this observation. As shown in Figure 5D, the blue shift of the absorption maximum from 364 to 330 nm was seen after 415 nm light irradiation, which indicates that C6 HAT molecules on Cu surface underwent *Z*→*E* isomerization. The obtained sample was then heated to 50 °C for 16 h. The absorption maximum was found to shift to 339 nm, which indicates only a partially isomerization of *E*-C6 HAT to the *Z* form. The discrete step at 330 nm (blue spectrum) is due to the change from a tungsten lamp to a deuterium lamp during measurement. A control experiment was conducted to check whether irradiation by 340 nm can induce the *E*-to-*Z* isomerization of C6 HAT on a Cu surface. It is shown that irradiation by 340 nm light barely induce *E*→*Z* isomerization on the Cu surface, which is similar to the case of C6 HAT on Au (see Figure S26 in Supporting Information).


**Substrate-mediated photoisomerization mechanism.** The first possible isomerization mechanism that we evaluated (Figure 6A) is the direct photo-excitation of C6 HAT on the metal surfaces. Because of the quenching effects of metal surfaces, the isomerization of C6 HAT through direct molecular photoexcitation is suppressed, although to different de-

grees, by the metal surfaces. The failure of the isomerization of C6 HAT on Ag surface and the decrease in the isomerization efficiency of C6 HAT on the Au surface compared to the solid-state are indications of this phenomenon. A drop in isomerization efficiency could also result from steric hindrance. To exclude this effect, we investigated the isomerization of C6 HAT on Ag diluted in a 10-fold excess of hexanethiol using UV-Vis spectroscopy. As shown in Figure S27, the maximum absorbance of C6 HAT did not shift upon 415 nm light irradiation, indicating that C6 HAT molecules with a lower packing density on Ag cannot isomerize to the *E* form. Therefore, failure of the photoisomerization of C6 HAT on Ag must stem from electronic rather than steric effects. The fact that C6 HAT SAM can isomerize on Au and Cu with 415 nm light irradiation also supports this interpretation. The second possible mechanism is a substrate-mediated photoisomerization mechanism<sup>[29]</sup> in which the electronic structure of the molecules, and the density of states (DOS) of the substrate should be taken into account. Tegeder and co-workers proposed such a mechanism for the photoisomerization of TBA physically adsorbed on a Au surface.<sup>[29]</sup> UV light first generates hot holes in the Au d-band, which subsequently relax to the top of the d-band through Auger decay. These hot holes then undergo a charge transfer to the HOMO of TBA, leading to the formation of transient positive ions, which may subsequently result in the isomerization of TBA. Based on this theory, we calculated the orbital energy levels of C6 HAT bound to the different metal surfaces, and the DOS of the different metals using density functional theory (DFT). (see Figure S28 and Figure S29) As shown in Figure 6B, the top of the d band of Au is at 2.2 eV, which overlaps with the second HOMO (HOMO-2) of C6 HAT. We speculate that 415 nm light ( $E = 3$  eV) irradiation generates hot holes in the Au d-band, which first relax to the top of the d-band, and can then transfer to C6 HAT to form transient positive ions, leading to the isomerization of C6 HAT. In the case of Ag, because of



**Figure 6.** (A) Possible excitation and quenching mechanisms in the photoisomerization of C6 HAT SAM at different metal surfaces. (B-D) Proposed excitation mechanism for the photoisomerization of C6 HAT SAM at different metal surfaces.

the lower lying d-band (d-band edge of Ag is 3.9 eV), light-induced switching cannot be achieved with 415 nm light irradiation (Figure 6C). In contrast, the d-band edge of Cu is at 2.1 eV, so 415 nm light irradiation can induce the isomerization of **C6 HAT** SAM on Cu. (Figure 6D) Regarding the hot carrier transfer process from substrate to the C=N switchable group of **C6 HAT**, our hypothesis is that it occurs via direct electron tunneling. Our system is a typical donor-bridge-acceptor system<sup>[30]</sup>, in which the alkyl chain serves as the bridge between the metal substrate and the switchable group of **C6 HAT**. It is well known that in such a system, coherent electron tunneling from donor to acceptor can take place via virtual states in the bridge, referred to as superexchange mechanism.<sup>[30,31]</sup> Based on Marcus theory, the electron transfer rate decays exponentially with donor-acceptor distance.<sup>[30a, 30c, 32]</sup> In our case, the distance between donor and acceptor is only ca. 10 Å. For comparison, Murray et al. reported that the conductivity of Au clusters protected by an alkanethiol monolayer with a 6-carbon chain was ca.  $1.8 \times 10^{-4} \Omega^{-1}\text{cm}^{-1}$  and  $6.6 \times 10^{-4} \Omega^{-1}\text{cm}^{-1}$  depending on the size of the Au clusters.<sup>[30a]</sup> Therefore, direct electron tunneling from the Au surface to the C=N switch group of **C6 HAT** is clearly feasible.



**Figure 7.** (A) Baseline corrected average TERS spectra from the TERS maps of a **C6 HAT** SAM on a Au surface before and after 340 nm light irradiation. (B) Baseline corrected average TERS spectra from the TERS maps of a **C6 HAT** SAM on a Ag surface before and after 280 nm light irradiation. The size of all maps is  $1 \times 1 \mu\text{m}^2$  with a 31 nm pixel size.

To further verify the substrate-mediated photoisomerization mechanism, we irradiated the **C6 HAT** SAM on Au with 340 nm light ( $E = 3.65 \text{ eV}$ ), and the **C6 HAT** SAM on Ag with 280 nm light ( $E = 4.4 \text{ eV}$ ), respectively, to induce the  $Z \rightarrow E$  photoisomerization. As shown in Figure 7A, after 340 nm light irradiation, we can clearly see the intensity drop of the peaks at 1430, 1472, 1506 and 1680  $\text{cm}^{-1}$ , and new peaks

appearing at 1580 and 1730  $\text{cm}^{-1}$  in the average TER spectrum of **C6 HAT** on Au. These spectral changes indicate that  $Z \rightarrow E$  isomerization took place, which is in agreement with the substrate-mediated mechanism. In contrast, 340 nm light irradiation induces  $E \rightarrow Z$  isomerization in solution, showing that there is a large difference in the solution phase and monolayer switching behavior. According to the substrate-mediated mechanism, light irradiation with energy above 3.9 eV should induce  $Z \rightarrow E$  isomerization of **C6 HAT** on Ag surface. The average TER spectra of **C6 HAT** SAM on Ag before and after 280 nm light irradiation are shown in Figure 7B. The intensity of peaks at 1430, 1473 and 1506  $\text{cm}^{-1}$  drops, and new peaks come out at 1577 and 1730  $\text{cm}^{-1}$ , which is quite different from the TER spectrum when irradiating with 415 nm light (Figure 5A). This suggests that **C6 HAT** molecules isomerized to the  $E$  form with 280 nm light irradiation. In contrast, solid-phase **C6 HAT** cannot isomerize from the  $Z$  to the  $E$  form upon irradiation by either 340 or 280 nm light (Figure S30). Moreover, irradiating a **C6 HAT** solution with 280 nm light cannot induce  $Z \rightarrow E$  isomerization (Figure S31). Only a very tiny and negligible proportion of  $Z$ -**C6 HAT** isomerizes to the  $E$  form in solution upon irradiation with 340 nm light because of the small absorption of  $Z$ -**C6 HAT** at this wavelength (see Figure S32) These results support our proposed substrate-mediated photo-isomerization mechanism.

## CONCLUSIONS

A new bistable ( $\tau_{1/2} = 789 \text{ years}$ ) hydrazone photoswitch having a C6 alkyl thiolate linker (**C6 HAT**) was synthesized, and its switching behavior studied in solution, the solid-state and as a monolayer on various metal surfaces using confocal Raman TERS and UV-Vis spectroscopies. Interestingly, **C6 HAT** can isomerize on Au and Cu surfaces, but not on Ag upon irradiation by 415 nm light ( $E = 3 \text{ eV}$ ). Higher energy light is required to activate the switch on Ag. Based on these findings, we propose a substrate-mediated charge transfer mechanism for the isomerization of **C6 HAT** on metal surfaces. This mechanism is supported by DFT calculations and TERS results. In this process, the length of the carbon linker plays an important role. When increasing the carbon chain length, the proposed substrate-mediated photoisomerization mechanism would stop working at a certain distance, since the electron transfer rate decays exponentially with donor-acceptor distance. However, photoswitches are still able to isomerize at larger distances via direct photoexcitation, because at larger distances there is less quenching by the surface. Thus, the length of the carbon is a critical parameter to maximize the isomerization efficiency on metal surfaces. As for the  $E \rightarrow Z$  isomerization, it can be thermally activated, with a half-life of 6.9 days (at room temperature) on the Au surface. This is the longest thermal half-life ever measured for photoswitches on metal surfaces. The lifetime of the  $E$  molecules may depend on the nature of the metal, and on the linker length,<sup>[13]</sup> which suggests a thermal relaxation mechanism. The details of the relaxation mechanism are a topic that will require future studies to be clarified. This property, in addition to the ease of synthesis and assembly of the switches on metal surfaces, and very good separation of the Raman bands of the  $Z$  and  $E$  isomers, make this system highly valuable in the development of photo-controllable surfaces. One immediate application might be surface patterning using the photoswitch-



es and using TERS for the in-situ imaging of the UV light initiated surface isomerization.

## ASSOCIATED CONTENT

### Supporting Information

Experimental details, chemical synthesis, photoisomerization studies in toluene solution, desorption analysis by MS, TERS study of the isomerization of a C6 HAT SAM, thin-film UV-Vis spectroscopic study, theoretical calculations, confocal Raman spectrum analysis and the vibrational modes of the *Z* and *E* isomers.

The Supporting Information is available free of charge on the ACS Publications website.

All of the supported figures (PDF)

Vibrational modes of the *Z* and *E* isomers (movies)

## AUTHOR INFORMATION

### Corresponding Author

\* Email: ivan.aprahamian@dartmouth.edu

\* Email: renato.zenobi@org.chem.ethz.ch

### Author Contributions

The manuscript was written through contributions of all authors.

### Notes

The authors declare no competing financial interest.

The original data used in this publication are made available in a curated data archive at ETH Zurich (<https://www.research-collection.ethz.ch>) under the DOI: 10.3929/ethz-b-000353526.

## ACKNOWLEDGMENT

We thank Hao Yin (Xiamen University) and Jonas Bastian Metternich (ETH Zurich) for insightful discussions, Tian Liu (ETH Zurich), Dr. Alessandro Lauria (ETH Zurich), Madeleine Fellner (ETH Zurich) and Prof. Markus Niederberger (ETH Zurich) for UV-Vis spectroscopy measurements, Jacek Szczerbiński (ETH Zurich) for TPD-MS measurements. L.-Q. Z. is indebted to the Chinese Scholarship Council for a Ph.D. student fellowship. We thank the High Performance Computing Team at ETH Zurich for help with DFT calculations and the ERC program (grant # 741431 - 2DNanoSpec) for financial support. I. A. is thankful to the National Science Foundation (CHE-1807428) for the generous support.

## REFERENCES

- [1] (a) Szymanski, W.; Beierle, J. M. H.; Kistemaker, A. V.; Velema, W. A.; Feringa, B. L. Reversible Photocontrol of Biological Systems by the Incorporation of Molecular Photoswitches. *Chem. Rev.* **2013**, *113*, 6114-6178. (b) Russew, M.-M.; Hecht, S. Photoswitches: From Molecules to Materials. *Adv. Mater.* **2010**, *22*, 3348-3360.
- [2] (a) Fihey, A.; Perrier, A.; Browne, W. R.; Jacquemin, D. Multiphotochromic molecular systems. *Chem. Soc. Rev.* **2015**, *44*, 3719-3759. (b) Landge, S. M.; Aprahamian, I. A pH Activated Configurational Rotary Switch: Controlling the *E/Z* Isomerization in Hydrazones. *J. Am. Chem. Soc.*, **2009**, *131*, 18269-18271. (c) Choi, B. Y.; Kahng, S. J.

- Kim, S.; Kim, H.; Kim, H. W.; Song, Y. J.; Ihm, J.; Kuk, Y. Conformational Molecular Switch of the Azobenzene Molecule: A Scanning Tunneling Microscopy Study. *Phys. Rev. Lett.* **2006**, *96*, 1-4.
- [3] (a) Robertus, J.; Browne, W. R.; and Feringa, B. L. Dynamic control over cell adhesive properties using molecular-based surface engineering strategies. *Chem. Soc. Rev.* **2010**, *39*, 354-378. (b) Zheng, Y. B.; Kiraly, B.; Cheunkar, S.; Huang, T. J.; Weiss, P. S. Incident-angle-modulated molecular plasmonic switches: a case of weak exciton-plasmon coupling. *Nano Lett.*, **2011**, *11*, 2061-2065. (c) Balzani, V.; Credi, A.; Raymo, F. M.; Stoddart, J. F. Artificial molecular machines. *Angew. Chem. Int. Ed.*, **2000**, *39*, 3349-3391.
  - [4] (a) Orgiu, E.; Samori, P. 25th anniversary article: organic electronics marries photochromism: generation of multifunctional interfaces, materials, and devices. *Adv. Mater.*, **2014**, *26*, 1827-1845. (b) Jerca, V. V.; Hoogenboom, R. Photocontrol in Complex Polymeric Materials: Fact or Illusion? *Angew. Chem. Int. Ed.* **2018**, *57*, 7945-7947; Steuerung komplexer Polymermaterialien mit Licht: Wirklichkeit oder Illusion? *Angew. Chem.*, **2018**, *130*, 8073-8075. (c) Ratner, M. Molecular electronics: Pushing electrons around. *Nature*, **2000**, *404*, 137-138. (d) Cuniberti, G.; Fagas, F. and Richter, K. Introducing Molecular Electronics, ed. Springer, Berlin, Heidelberg, **2005**. (e) Gahl, C.; Brete, D.; Leyssner, F.; Koch, M.; McNellis, E. R.; Mielke, J.; Carley, R.; Grill, L.; Reuter, K.; Tegeder, P.; Weinelt, M. Coverage- and temperature-controlled isomerization of an imine derivative on Au (111). *J. Am. Chem. Soc.*, **2013**, *135*, 4273-4281.
  - [5] Qian, H.; Pramanik, S.; Aprahamian, I. Photochromic hydrazone switches with extremely long thermal half-lives. *J. Am. Chem. Soc.*, **2017**, *139*, 9140-9143.
  - [6] Hirshberg, Y. Reversible formation and eradication of colors by irradiation at low temperatures. A photochemical memory model. *J. Am. Chem. Soc.* **1956**, *78*, 2304-2312.
  - [7] Nacci, C.; Baroncini, M.; Credi, A.; Grill, L. Reversible Photoswitching and Isomer - Dependent Diffusion of Single Azobenzene Tetramers on a Metal Surface. *Angew. Chem. Int. Ed.* **2018**, *57*, 15034-15039.
  - [8] (a) Pijper, A.; T. C.; Kudernac, T.; Katsonis, N.; van der Maas, M.; Feringa, B. L.; van Wees, B. J. Reversible light induced conductance switching of asymmetric diarylethenes on gold: surface and electronic studies. *Nanoscale*, **2013**, *5*, 9277-9282. (b) Irie, M. Diarylethenes for memories and switches. *Chem. Rev.* **2000**, *100*, 1685-1716.
  - [9] Klajn, R. Spiropyran-based dynamic materials. *Chem. Soc. Rev.* **2014**, *43*, 148-184.
  - [10] (a) Aprahamian, I. Hydrazone switches and things in between. *Chem. Commun.*, **2017**, *53*, 6674-6684. (b) van Dijken, D. J.; Kovaříček, P.; Ihrig, S. P.; Hecht, S. Acylhydrazones as Widely Tunable Photoswitches. *J. Am. Chem. Soc.* **2015**, *137*, 14982-14991.
  - [11] (a) Irie, M.; Mohri, M. Thermally irreversible photochromic systems. Reversible photocyclization of diarylethene derivatives. *J. Org. Chem.* **1988**, *53*, 803-808. (b) Nakamura, S.; Irie, M. Thermally irreversible photochromic systems. A theoretical study. *J. Org. Chem.* **1988**, *53*, 6136-6138. (c) Irie, M. Diarylethenes for memories and switches. *Chem. Rev.* **2000**, *100*, 1685-1716. (d) Tian, H.; Yang, S. Recent progresses on diarylethene based photochromic switches. *Chem. Soc. Rev.* **2004**, *33*, 85-97. (e) Zhang, J. J.; Zou, Q.; Tian, H. Photochromic materials: more than meets the eye. *Adv. Mater.* **2013**, *25*, 378-399.
  - [12] (a) Pace, G.; Ferri, V.; Grave, C.; Elbing, M.; Hänisch, C. V.; Zharnikov, M.; Mayor, M.; Rampi, M. A.; Samori, P. Cooperative light-induced molecular movements of highly ordered azobenzene self-assembled monolayers. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 9937-9942. (b) Schweighauser, L.; Strauss, M. A.; Bellotto, S.; Wegner, H.

- A. Attraction or repulsion? London dispersion forces control azobenzene switches. *Angew. Chem. Int. Ed.* **2015**, *54*, 13436-13439; Anziehung oder Abstoßung? London - Dispersionswechselwirkungen kontrollieren Azobenzol-basierte molekulare Schalter. *Angew. Chem.* **2015**, *127*, 13636-13639. (c) Li, Q.; Qian, H.; Shao, B.; Hughes, R. P.; Aprahamian, I. Building strain with large macrocycles and using it to tune the thermal half-lives of hydrazone photochromes. *J. Am. Chem. Soc.* **2018**, *140*, 11829-11835.
- [13] Schlimm, A.; Löw, R.; Rusch, T.; Röhricht, F.; Strunskus, T.; Tellkamp, T.; Sönnichsen, F.; Manthe, U.; Magnussen, O.; Tuczek, F.; Herges, R. Long - Distance Rate Acceleration by Bulk Gold. *Angew. Chem. Int. Ed.* **2019**, *58*, 6574-6578.
- [14] Katsonis, N.; Kudernac, T.; Walko, M.; van der Molen, S. J.; van Wees, B. J.; Feringa, B. L. Reversible conductance switching of single diarylethenes on a gold surface. *Adv. Mater.*, **2006**, *18*, 1397-1400.
- [15] Kumar, A. S.; Ye, T.; Takami, T.; Yu, B.-C.; Flatt, A. K.; Tour, J. M.; Weiss, P. S. Reversible photo-switching of single azobenzene molecules in controlled nanoscale environments. *Nano Lett.*, **2008**, *8*, 1644-1648.
- [16] Durr, H. Perspectives in Photochromism: A Novel System Based on the 1, 5 - Electrocyclization of Heteroanalogous Pentadienyl Anions. *Angew. Chem. Int. Ed.*, **1989**, *28*, 413-431.
- [17] (a) Bandara, H. M. D.; Burdette, S. C.; Photoisomerization in different classes of azobenzene. *Chem. Soc. Rev.* **2012**, *41*, 1809-1825. (b) Landge, S. M.; Tkatchouk, K.; Benitez, D.; Elhabiri, D. A. L. M.; Goddard III, W. A.; Aprahamian, I. Isomerization mechanism in hydrazone-based rotary switches: lateral shift, rotation, or tautomerization? *J. Am. Chem. Soc.*, **2011**, *133*, 9812-9823.
- [18] (a) Napper, A. M.; Liu H.; Waldeck, D. H. The Nature of Electronic Coupling between Ferrocene and Gold through Alkanethiolate Monolayers on Electrodes: The Importance of Chain Composition, Interchain Coupling, and Quantum Interference. *J. Phys. Chem. B*, **2001**, *105*, 7699-7707. (b) Comstock, M. J.; Levy, N.; Kirakosian, A.; Cho, J.; Lauterwasser, F.; Harvey, J. H.; Strubbe, D. A.; Frechet, J. M. J.; Trauner, D.; Louis, S. G.; Crommie, M. F. Reversible photomechanical switching of individual engineered molecules at a metallic surface. *Phys. Rev. Lett.*, **2007**, *99*, 38301.
- [19] (a) Zhou, X.-L.; Zhu X.-Y.; White, J. M. Photochemistry at adsorbate/metal interfaces. *Surf. Sci. Rep.*, **1991**, *13*, 73-220. (b) Wen, J.; Li, W.; Chen, S.; Ma, J. Simulations of molecular self-assembled monolayers on surfaces: packing structures, formation processes and functions tuned by intermolecular and interfacial interactions. *Phys. Chem. Chem. Phys.*, **2016**, *18*, 22757-22771. (c) Pourghaz, Y.; Dongare, P.; Thompson D. W.; Zhao, Y. Click functionalized poly(p-phenylene ethynylene)s as highly selective and sensitive fluorescence turn-on chemosensors for Zn<sup>2+</sup> and Cd<sup>2+</sup> ions. *Chem. Commun.*, **2011**, *47*, 11014-11016.
- [20] Alemani, M.; Selvanathan, S.; Ample, F.; Peters, M. V.; Rieder, K. H.; Moresco, F.; Joachim, C.; Hecht, S.; Grill, L. Adsorption and switching properties of azobenzene derivatives on different noble metal surfaces: Au (111), Cu (111), and Au (100). *J. Phys. Chem. C* **2008**, *112*, 10509-10514.
- [21] Sprague-Klein, E. A.; Negru, B.; Madison, L. R.; Coste, S. C.; Rugg, B. K.; Felts, A. M.; McAnally, M. O.; Banik, M.; Apkarian, V. A.; Wasielewski, M. R.; Ratner, M. A.; Seideman, T.; Schatz, G. C.; Van Duyne, R. P. Photoinduced Plasmon-Driven Chemistry in trans-1,2-Bis(4-pyridyl)ethylene Gold Nanosphere Oligomers. *J. Am. Chem. Soc.*, **2018**, *140*, 10583-10592.
- [22] (a) Wang, X.; Huang, S.-C.; Huang, T.-X.; Su, H.-S.; Zhong, J.-H.; Zeng, Z.-C.; Li, M.-H.; Ren, B. Tip-enhanced Raman spectroscopy for surfaces and interfaces. *Chem. Soc. Rev.* **2017**, *46*, 4020-4041. (b) Anderson, M. S. Locally enhanced Raman spectroscopy with an atomic force microscope. *Appl. Phys. Lett.* **2000**, *76*, 3130-3132. (c) Hayazawa, N.; Inouye, Y.; Sekkat, Z.; Kawata, S. Metalized tip amplification of near-field Raman scattering. *Opt. Commun.* **2000**, *183*, 333-336. (d) Pettinger, B.; Schambach, P.; Villagómez, C. J.; N. Scott, Tip-enhanced Raman spectroscopy: near-fields acting on a few molecules. *Annu. Rev. Phys. Chem.* **2012**, *63*, 379-399. (e) Wickramasinghe, H. K.; Chaigneau, M.; Yasukuni, R.; Picardi, G.; Ossikowski, R. Billion-fold increase in tip-enhanced Raman signal. (f) Zhang, R.; Zhang, Y.; Dong, Z. C.; Jiang, S.; Zhang, C.; Chen, L. G.; Zhang, L.; Liao, Y.; Aizpurua, J.; Luo, Y.; Yang, J. L.; Hou, J. G. Chemical mapping of a single molecule by plasmon-enhanced Raman scattering. *Nature* **2013**, *498*, 82-86. (g) Zhong, J.-H.; Jin, X.; Meng, L.; Wang, X.; Su, H.-S.; Yang, Z.-L.; Williams, C. T.; Ren, B. Probing the electronic and catalytic properties of a bimetallic surface with 3 nm resolution. *Nat. Nanotechnol.* **2017**, *12*, 132-136.
- [23] Tallarida, N.; Rios, L.; Apkarian, V. A.; Lee, J. Isomerization of one molecule observed through tip-enhanced Raman spectroscopy. *Nano Lett.* **2015**, *15*, 6386-6394.
- [24] Zheng, L.-Q.; Wang, X.; Shao, F.; Hegner, M.; Zenobi, R. Nanoscale Chemical Imaging of Reversible Photoisomerization of an Azobenzene - Thiol Self - Assembled Monolayer by Tip - Enhanced Raman Spectroscopy. *Angew. Chem. Int. Ed.* **2017**, *56*, 1025-1029.
- [25] (a) Love, J. C.; Estroff, L. A.; Kriebel, J. K.; Nuzzo, R. G.; Whitesides, G. M. Self-Assembled Monolayers of Thiolates on Metals as a Form of Nanotechnology. *Chem. Rev.* **2005**, *105*, 1103-1169; (b) Claridge, S. A.; Liao, W.-S.; Thomas, J. C.; Zhao, Y.; Cao, H. H.; Cheunkar, S.; Serino, A. C.; Andrews, A. M.; Weiss, P. S. From the bottom up: dimensional control and characterization in molecular monolayers. *Chem. Soc. Rev.*, **2013**, *42*, 2725-2745.
- [26] Zheng, Y. B.; Payton, J. L.; Weiss, P. S. Surface-enhanced Raman spectroscopy to probe reversibly photoswitchable azobenzene in controlled nanoscale environments. *Nano Lett.* **2011**, *11*, 3447-3452.
- [27] Merrick, J. P.; Moran, D.; Radom, L. An evaluation of harmonic vibrational frequency scale factors. *J. Phys. Chem. A*, **2007**, *111*, 11683-11700.
- [28] Shao, B.; Baroncini, M.; Qian, H.; Bussotti, L.; Donato, M. D.; Credi, A.; Aprahamian, I. Solution and Solid-State Emission Toggling of a Photochromic Hydrazone. *J. Am. Chem. Soc.* **2018**, *140*, 12323-12327.
- [29] Wolf, M.; Tegeder, P. Reversible molecular switching at a metal surface: A case study of tetra-*tert*-butyl-azobenzene on Au (111). *Surface Science*, **2009**, *603*, 1506-1517.
- [30] (a) Wuelfing, W. P.; Green, S. J.; Pietron, J. J.; Cliffel, D. E.; Murray, R. W. Electronic Conductivity of Solid-State, Mixed-Valent, Monolayer-Protected Au Clusters. *J. Am. Chem. Soc.* **2000**, *122*, 11465-11472. (b) Wang, G. R.; Wang, L.; Rendeng, Q.; Wang, J.; Luo, J.; Zhong, C.-J. Correlation between nanostructural parameters and conductivity properties for molecularly-mediated thin film assemblies of gold nanoparticles. *J. Mater. Chem.* **2007**, *17*, 457-462. (c) Avila, A.; Gregory, B. W.; Niki, K.; Cotton, T. M. An Electrochemical Approach to Investigate Gated Electron Transfer Using a Physiological Model System: Cytochrome c Immobilized on Carboxylic Acid-Terminated Alkanethiol Self-Assembled Monolayers on Gold Electrodes. *J. Phys. Chem. B* **2000**, *104*, 2759-2766.
- [31] (a) Guo, L.-H.; Facci, J. S.; Mclendon, G. Distance dependence of electron transfer rates in bilayers of a ferrocene Langmuir-Blodgett monolayer and a self-assembled monolayer on gold. *J. Phys. Chem.* **1995**, *99*, 8458-8461. (b) Khoshtariya, D. E.; Dolidze, T. D.; Sarauli, D.; van Eldik, R. High - Pressure Probing of a Changeover in the

- Charge - Transfer Mechanism for Intact Cytochrome c at Gold/Self - Assembled Monolayer Junctions. *Angew. Chem. Int. Ed.* **2006**, *45*, 277-281. (c) Closs, G. L.; Miller, J. R. Intramolecular long-distance electron transfers in organic molecules. *Science* **1988**, *240*, 440-447.
- [32] Kim, Y.; Wilson, A. J.; Jain, P. K. The Nature of Plasmonically Assisted Hot-Electron Transfer in a Donor-Bridge-Acceptor Complex. *ACS Catal.* **2017**, *7*, 4360-4365.

