

**SOME CRITERIA FOR THE ABSOLUTE SUMMABILITY OF
A FOURIER SERIES**

GEN-ICHIRO $\widehat{\text{SUNOUCHI}}$

(Received May 8, 1967)

Let

$$(1) \quad f(x) \sim a_0/2 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx) = \sum_{n=0}^{\infty} A_n(x)$$

and $\sigma_n^k(x)$ ($k > -1$) denote the n -th (C, k) mean of Fourier series (1). If the series

$$\sum_{n=0}^{\infty} |\sigma_n^k(x) - \sigma_{n-1}^k(x)|$$

is convergent, we say that the series (1) is absolutely summable (C, k) or summable $|C, k|$. We denote the integral modulus of continuity of f by

$$\omega_p(t, f) = \sup_{0 < h < t} \left\{ \int_{-\pi}^{\pi} |f(x+h) - f(x-h)|^p dx \right\}^{1/p} \quad (1 \leq p < \infty).$$

H.C. Chow [1] proved the following theorem.

THEOREM A. *Let $1 \leq p \leq 2$. If*

$$\omega_p(t, f) = O \left\{ \left(\log \frac{1}{t} \right)^{-1-\delta} \right\} \quad (\delta > 0),$$

then the series (1) is summable $|C, \alpha|$ almost everywhere for $\alpha > 1/p$.

On the other hand, P.L. Ul'yanov [5] proved the following theorem.

THEOREM B. *If*

$$\omega_2(t, f) = O \left\{ \left(\log \frac{1}{t} \right)^{-1/2-\delta} \right\} \quad (\delta > 0),$$

then the series (1) is summable $|C, \alpha|$ almost everywhere for $\alpha > 1/2$.

We will sharpen Theorem A into the form of Theorem B.

THEOREM 1. Let $1 < p \leq 2$. If

$$(2) \quad \omega_p(t, f) = O \left\{ \left(\log \frac{1}{t} \right)^{-1/2-\delta} \right\} \quad (\delta > 0),$$

then the series (1) is summable $|C, \alpha|$ almost everywhere for $\alpha > 1/p$.

For proof of Theorem, we need three lemmas.

LEMMA 1. If (2) is true, then the n -th partial sum of series (1), $s_n(f)$ has the approximation such as

$$\left\{ \int_{-\pi}^{\pi} |f(x) - s_n(x)|^p dx \right\}^{1/p} = O \{ (\log n)^{-1/2-\delta} \}$$

PROOF. If $T_n(x)$ is an arbitrary trigonometric polynomial of degree n , then

$$\|f - s_n(f)\|_p \leq \|f - T_n\|_p + \|s_n(f - T_n)\|_p \leq A_p \|f - T_n\|_p$$

by the M. Riesz theorem and the order of best approximation is $\omega_p(1/n) = (\log n)^{-1/2-\delta}$ in this case.

LEMMA 2. Under the condition of Theorem,

$$(3) \quad \sum_{n=0}^{\infty} \lambda(n) A_n(x)$$

is a Fourier series of a function of class L^p , where

$$\begin{aligned} \lambda(0) &= \lambda(1) = 1, \\ \lambda(n) &= (\log n)^{1/2+\varepsilon} \quad (\delta > \varepsilon > 0) \quad (n = 2, 3, \dots). \end{aligned}$$

PROOF. Let denote by $t_n(x)$ the partial sum of (3), then

$$\begin{aligned} t_n(x) - f(x) &= \lambda(0) \{A_0(x) - f(x)\} + \sum_{k=1}^n \lambda(k) A_k(x) \end{aligned}$$

$$= \sum_{k=0}^{n-1} \{s_k(x) - f(x)\} \Delta\lambda(k) + \{s_n(x) - f(x)\} \lambda(n).$$

Applying Minkowski's inequality,

$$\begin{aligned} & \|t_n - \lambda(0)f\|_p \\ & \leq \sum_{k=0}^{n-1} \|s_k - f\|_p \Delta\lambda(k) + \|s_n - f\|_p \lambda(n) \\ & \leq C_1 + C_2 \sum_{k=2}^{n-1} (\log k)^{-1/2-\delta} \frac{1}{k} (\log k)^{-1/2+\varepsilon} + C_3 (\log n)^{-1/2-\delta} (\log n)^{1/2+\varepsilon} \\ & \leq C_1 + C_2 \sum_{k=2}^{n-1} \frac{1}{k(\log k)^{1+\delta-\varepsilon}} + C_3 \frac{1}{(\log n)^{\delta-\varepsilon}} \leq C, \end{aligned}$$

which is an absolute constant by Lemma 1. Hence

$$\|t_n(x)\|_p = O(1)$$

and (1) is a Fourier series of a function of the class L^p .

LEMMA 3. *If $f(x)$ belongs to the class $L^p(1 < p \leq 2)$, then the series $\sum \mu(n)A_n(x)$ is summable $|C, \alpha|(\alpha > 1/p)$ almost everywhere, provided that*

$$\mu(0) = \mu(1) = 1, \quad \mu(n) = (\log n)^{-1/2-\varepsilon} (\varepsilon > 0), \quad (n = 2, 3, \dots).$$

This is known, [Chow, 2].

The proof of Theorem is almost completed. That is to say, from Lemma 2 and Lemma 3

$$\sum_{n=0}^{\infty} A_n(x)$$

is $|C, \alpha|$ summable $(\alpha > 1/p)$ almost everywhere.

We can prove the following theorem with the same method also.

THEOREM 2. *If $1 < p \leq 2$ and*

$$\omega_p(t, f) = O \left\{ \left(\log \frac{1}{t} \right)^{-\left(1-\frac{1}{p}+\frac{1}{2}+\delta\right)} \right\}$$

then (1) is $|C, 1/p|$ summable almost everywhere.

PROOF. We apply a result of Kojima [3] instead of Lemma 3.

THEOREM 3. *If $f(e^{i\theta})$ belongs to the class H , and the integral modulus of continuity of complex $f(e^{i\theta})$ be*

$$\omega_1(t, f) = O \left\{ \left(\log \frac{1}{t} \right) \right\}^{-(1/2+\delta)}$$

then the complex Fourier series of $f(e^{i\theta})$ is $|C, 1|$ summable almost everywhere.

PROOF. We use a result of the author [4] and the fact that the power series of bounded variation is absolutely continuous.

Literatures

- [1] H. C. CHOW, Some new criteria for the absolute summability of a Fourier series and its conjugate series, Journ. London Math. Soc., 30(1955), 133-142.
- [2] H. C. CHOW, An extension of a theorem of Zygmund and its application, Journ. London Math. Soc., 29(1954), 189-198.
- [3] M. KOJIMA, On the absolute summability factors of power series, Tôhoku Math. Journ., 18(1966), 60-70.
- [4] G. SUNOUCHI, On the absolute summability of Fourier series, Journ. Math. Soc. Japan, 1(1949), 122-129.
- [5] P. L. UL'YANOV, Solved and unsolved problem in the theory of trigonometric and orthogonal series, Uspehi Mat. Nauk, 19(1964), 3-69.

MATHEMATICAL INSTITUTE
TÔHOKU UNIVERSITY
SENDAI, JAPAN.