

Some factors affecting the growth of prosobranch veligers

M. C. PILKINGTON and V. FRETTER

University of Reading; Reading, England

EXTRAIT: Sur quelques facteurs intervenant dans la croissance des véligères de Proso-branches. Le vélum des véligères de Prosobranches recueille toutes les particules que ses cils composés peuvent maîtriser. Chacune des dix espèces d'algues unicellulaires offertes aux larves de *Crepidula fornicata* et de *Nassarius reticulatus* subit l'action mécanique de l'estomac. Sauf pour *Chlamydomonas parkeae*, que les larves ne réussissent pas à détruire, et pour *Phaeodactylum tricorutum* dont les frustules sont perforées, les parois cellulaires sont ramollies ou fragmentées. Les cellules de la glande digestive ingèrent le contenu cellulaire y compris le pigment, mais non les débris des parois. Les pigments sont rejetés ultérieurement; ils retardent le cycle de l'activité glandulaire. La croissance est relativement bonne avec des cellules de *Monochrysis lutheri* et de *Pyramimonas grossii*, qui se rompent facilement, et avec *Phaeodactylum tricorutum*; mais il se forme des boules fécales volumineuses qui peuvent obturer la cavité palléale; les grandes frustules de *Phaeodactylum tricorutum* entraînent un effet purgatif. *Chlamydomonas parkeae*, *Brachiomonas submarina*, *Exuviaella pusilla* et *Olisthodiscus* sp. sont de mauvais aliments; les deux dernières sont toxiques pour les larves, spécialement *Olisthodiscus* sp. Nourries de *Cricosphaera* sp. *carterae* les deux espèces présentent une bonne croissance et un développement jusqu'à la métamorphose; il en est de même pour *Crepidula fornicata* nourrie de *Exuviaella baltica* et de *Nassarius reticulatus* alimenté avec *Dunaliella primolecta*. Un aliment convenable détermine un aspect particulier du comportement alimentaire. Il ne semble pas que les larves sélectionnent leur aliment lorsqu'elles reçoivent simultanément deux ou trois espèces d'algues, mais de petites cellules facilement maîtrisées par le vélum sont absorbées plus fréquemment que des cellules volumineuses. L'apport occasionnel d'aliments bactériens ne semble causer aucun dommage. Un aliment artificiel (par exemple la farine de blé) permet pendant 5 à 6 jours de conserver les larves et d'obtenir leur croissance. Dans toutes les expériences la température et l'éclairage, dont les fluctuations retentissent sur la croissance des larves, furent maintenus constants. La manipulation des larves se fit à l'aide de pipettes. Dans la mer les détritiques organiques doivent constituer un apport alimentaire important.

INTRODUCTION

The majority of marine prosobranch gastropods in temperate waters have a planktotrophic veliger. Observations on their distribution at inshore stations off Plymouth reveal that during daylight hours they are feeding in the surface waters and, in considerable numbers, well below the compensation depth. The extended velum edged with compound cilia and a food groove leading to the mouth propels the larva

forwards and simultaneously collects food; the velum and foot are uppermost, the visceral mass below. Organic and inorganic particles are collected. If organic particles are available the stomach is filled rapidly and further intake is regulated while the meal is being digested (FRETTER & MONTGOMERY 1968). When feeding stops the velum may be partly withdrawn and the larva sinks or it may swim actively to a lower level where it is maintained by the languid beat of the velar cilia (FRETTER 1967).

Food particles collected by the velum are manipulated in the gut by cilia and muscles. Cilia pass them straight along the oesophagus to the ventral chamber of the stomach where they are retained and rotated vigorously against the gastric shield, mixed with digestive juices and subjected to mechanical breakdown. A muscular network surrounding each of the two lobes of the digestive gland effects a rhythmical pumping action, drawing the stomach contents into the lobes, so that the food is brought into direct contact with the ingesting cells of the epithelium, and then expelling them, together with secretion and waste. The food is retained in the ventral chamber for a length of time which may be correlated with its apparent food value to the larva and its accessibility to digestive enzymes. Unwanted remains are later directed by cilia and muscles into the style sac where the rotary action of the cilia aggregates them and mixes them with a viscous secretion. These remains include plant pigments once intimately linked with food and passed with them into the cells of the digestive gland; the pigments are soon excreted to the stomach. The faecal mass is sucked from the style sac into the initial part of the intestine and its passage to the anus is effected mainly by muscles. The faeces are compacted into rods which are usually slow to disintegrate.

In high concentrations of a good food a hungry larva will fill the stomach in a few minutes, and then stop feeding while digestion of the meal is underway. If only inorganic particles are available on which there is no organic scum they are collected rapidly and passed directly to the intestine for egestion. Feeding is continuous. Thus the presence of digestible food in the stomach initiates the digestive processes and clean inorganic matter fails to do this. A similar mechanism is found in *Amphioxus lanceolatus* in which feeding stops when the gut is filled with utilizable food and is continuous if colloidal graphite is collected (BONE 1961). If sand grains rather than organic particles are collected by *Artemia salina* the rate of collection is ten times that of plant cells of equivalent size and they are speeded through the gut for egestion (REEVE 1963). Differences in the rate of collection of organic and inorganic particles have not been estimated for veligers.

If food is abundant, particles collected by the velum are rejected from the region of the mouth when the stomach is adequately full. If it is scarce and the rate of collection does not exceed the rate of digestion, feeding is more or less continuous.

In order to study the fate of food particles in the gut and to compare the growth rates and efficiency of different foods it was necessary to formulate experimental conditions under which the larvae would remain healthy and metamorphose. Some indication of the sensitivity of the veligers to external conditions will be given when the standardization of the experimental conditions are discussed.

Studies on the growth of molluscan larvae fed on unicellular algae and bacteria have been concentrated on oysters and clams which are of economic importance. The

present study on prosobranchs, a group relatively neglected except for the work of SCHELTEMA (1962, 1967) and PAULSON & SCHELTEMA (1968), differs from others in that it incorporates such observations on gut contents and faecal waste as may help with an understanding of the relative value of different foods. The larvae of *Crepidula fornicata* (L.) and *Nassarius reticulatus* (L.) were used since their egg capsules are readily obtainable and provide an abundance of larvae of known age. The initial average size of healthy larvae was surprisingly uniform in terms of shell length (*C. fornicata* 0.37–0.40 mm; *N. reticulatus* 0.32–0.34 mm). Shell length was used to estimate growth. In order to obtain healthy *C. fornicata* only the older capsules from which larvae were about to hatch were taken from the care of the parent. The conditions under which larvae of these two prosobranchs were kept proved successful for other species taken from the plankton.

STANDARDIZATION OF EXPERIMENTAL CONDITIONS

As soon as the larvae hatched they were placed in filtered sea water in acid-cleaned glass containers. Polythene vessels proved harmful. It is essential to keep the bacterial population low otherwise growth is impaired (WALNE 1956a, 1958). The water used for all experiments was collected from outside the Plymouth Breakwater to avoid sewage contamination, and passed through glass filters (pore size $3\ \mu$) under pressure. Filtering through Oxoid membranes (pore size 0.5–1.0 μ), a slower process necessitating frequent change of filter, was of no apparent advantage. Fresh membrane-filtered water contained 4.77×10^6 bacteria per 10 ml sample compared with 6.02×10^6 in a glass-filtered sample, but there was no significant difference in their bacterial counts after larvae had been kept in each type of water without food for 3.5 days at 12° C. There were disadvantages in using water treated with antibiotics. A mixture of sodium penicillin G and streptomycin sulphate (50 I.U. & 0.05 mg/ml) has been used successfully to control bacteria in cultures of oyster larvae (WALNE 1958), and this was used in preliminary experiments with prosobranch veligers. The treated water contained 4.77×10^8 per 10 ml bacteria if the antibiotic solutions were fresh, but 1.26×10^6 per 10 ml if they were a week old and had been stored under refrigeration to retard deterioration. For effective bacterial control the antibiotic solution must be fresh and renewed with daily water change, especially in view of the fact that deterioration is more rapid once the solution is contaminated with bacteria (MARTINDALE 1967).

The method of water change generally used for the culture of bivalve larvae and used by SCHELTEMA (1962) for larvae of *Nassarius obsoletus* and *N. vibex*, involves pouring the water through a coarse filter which retains the larvae. This treatment proved detrimental to growth, and larvae of *Crepidula fornicata* died after two filtrations. The larvae were therefore transferred to fresh water by means of a pipette. This method, more reliable for growth experiments, is time consuming and imposes a limit on the number of animals used. With large numbers shell measurements can be taken on a limited number of veligers during the course of the experiments and these larvae then cast away. With small numbers this may not be practic-

able and the larvae must be returned to the experiment. The results of twice weekly measurements on the growth of 2 day old larvae of *Nassarius reticulatus* over a period of 3 weeks were compared with those of larvae measured only at the beginning and end of the experiment. Although the latter group consistently grew more than the former the difference was not always significant ($P = 0.2$ and 0.1).

The volume of sea water available to the larva under experimental conditions may affect growth. To test this the growth of 20 veligers of *Crepidula fornicata* in a boiling tube with 30 ml sea water was compared with that of 333 veligers in 500 ml sea water in a beaker; the depth of the water was 8 cm and 9 cm respectively. A column of water is essential for the successful functioning of the gut (FRETTER & MONTGOMERY 1968). Either *Cricosphaera* sp. or *Exuviaella baltica*, at concentrations of 40×10^3 , 20×10^3 or 2×10^3 cells/ml, were provided for food, the illumination was constant at 155 lux and the temperature was maintained at 12°C ; the water was renewed twice weekly over the 3 week period. A similar experiment was set up using *Nassarius reticulatus* veligers. Growth in the larger vessels was better, but in all cases the differences in growth were not significant at the 10% level. *Crepidula fornicata* larvae fed with *Cricosphaera* sp. grew significantly better ($P = 0.001$) in both boiling tube and beaker than with *E. baltica*. This agrees with later results. In experiments which were set up to compare the food values of various algal cultures the smaller volumes of water and numbers of larvae were used.

Veligers grew better in the constant illumination used in the last experiment (155 lux) than in the dark. Moreover, in the dark the mortality of algal cells stimulated the growth of bacterial populations and water needed to be changed at least daily. Larvae under continuous illumination showed better growth than when subjected to alternating 12 h periods of light and dark. The experiments were therefore carried out under constant illumination from a Mazda daylight strip giving an intensity of 155 lux at the water surface. Care was taken to ensure that all vessels in an experiment were subjected to the same light intensity since growth rate is influenced by intensity (Fig. 1).




Fig. 1: *Nassarius reticulatus*. Results of experiments with 2 batches of 20 larvae fed *Cricosphaera* sp. *carterae* (20×10^3 cells/ml) for 3 weeks, to show relationship between growth of shell and illumination

The surface water temperatures off Plymouth have been recorded for Station E₁ and ranged from 8.95–10.07° C in early spring to 16.15–18.30° C in late summer (ARMSTRONG & BUTLER 1960a, b, 1962, 1963) over 4 years. *Nassarius reticulatus* starts to breed in March and *Crepidula fornicata* in May so 12° C was considered a suitable constant temperature for the experiments. Low temperatures reduced larval activity and high temperatures favoured the increase of bacteria. At 8° C *Crepidula fornicata* larvae in both large and small volumes of water collected cells of *Cricosphaera* sp. and *Exuviaella baltica* and appeared to digest them, though slowly, but did not grow. At 4° C *Nassarius reticulatus* larvae did not collect plant cells. For both species a sudden lowering of water temperature from 12° C to below 8° C caused healthy larvae, actively feeding, to remain stationary with the velum extended, though not fully, and the ciliary beat reduced. They regained full activity when, after 6 h, the temperature was raised.

The water in which the larvae were kept was not agitated or aerated since it was difficult to ensure equality of treatment in the series of experimental vessels and any localized excessive movement of water had a retarding effect on growth.

ALGAL FOODS

The algal foods (Table 1) were from cultures maintained at Plymouth Laboratory; they were not bacteria free. Old cultures were not used. To compare relative food values each food was given at three concentrations 40×10^3 , 20×10^3 and 2×10^3 cells/ml. The last occurs frequently in surface waters of Plymouth Sound; the others may be associated with high density blooms (BAINBRIDGE 1957). The volume of algal culture added to each experimental vessel to give the required concentration was calculated from haemocytometer counts of the stock culture. The value obtained could be only approximate and suffers from the severe limitation that it does not allow for differences in cell size and shape. Differences in cell size have been allowed for (DAVIS & GUILLARD 1958, WALNE 1963) by centrifugation for a standard time in graduated haematocrit tubes. However these determinations are dependent on cell density as well as volume. Table 2 shows that cell density varies greatly in the same culture within a few hours. Such changes in density must affect the distribution of the food in the experimental vessels.

In nature veligers are likely to encounter the higher concentrations of algal cells used in the experiments on growth only briefly. Prolonged access to high concentrations might be harmful. To investigate this *Monochrysis lutheri* and *Pyramimonas grossii* were used up to 12×10^4 cells/ml and *Cricosphaera* sp. *carterae* 60×10^3 cells/ml.

KORRINGA (1951), working on larvae of *Ostrea edulis*, suggested that toxic concentrations of metabolites are manifest if there are more than 5,000 flagellates/ml. IMAI & HATANAKA (1949) working with larvae of *Crassostrea gigas* kept the concentration of *Monas* sp. below 1×10^3 – 2×10^3 cells/ml. These authors had a maximal concentration of only 2×10^2 larvae/litre. The concentration used in the present experiments is of the order of 7×10^2 litre. WALNE (1956b) on the other

Table 1

Cultures used for food. Culture numbers refer to the Plymouth collection.
The approximate size of the cells is in brackets

Order	Culture number	Species
Dinophyceae	28	<i>Exuviaella baltica</i> LOHM (9–15 μ)
	184	<i>Exuviaella pusilla</i> J. SCHILLER (8–10 μ)
Haptophyceae	156	<i>Cricosphaera</i> ap. <i>carterae</i> (BRAARUD et FAGERL.) BRAARUD (10–18 μ)
Chrysophyceae	75	<i>Monochrysis lutheri</i> DROOP (6–10 \times 2–3 μ)
Xanthophyceae	239	<i>Olisthodiscus</i> sp. (10–15 μ)
Prasinophyceae	78	<i>Pyramimonas grossii</i> PARKE (5.5–8 \times 4.5–5.5 μ)
Chlorophyceae	81	<i>Dunaliella primolecta</i> BUTCH. (5–10 μ)
	404	<i>Brachiomonas submarina</i> BOHLIN var. <i>pulsifera</i> DROOP (8 \times 20 μ)
	285	<i>Chlamydomonas parkeae</i> Ettl (3–8 μ)
Bacillariophyceae	100	<i>Phaeodactylum tricornutum</i> BOHLIN (8–35 μ)

Table 2

Volume of cells deposited by centrifugation for 15 min at 250 R.P.M.
at different times of day

Alga	Volume of cells deposited in ml at			
	11.00	12.30	14.30	17.30
<i>Exuviaella baltica</i>	0.0	0.01	0.0	0.0
<i>Olisthodiscus</i> sp.	0.0		0.006	0.01
<i>Cricosphaera</i> ap. <i>carterae</i>	0.008		0.008	0.014
<i>Brachiomonas submarina</i>	0.005		0.008	0.005

hand, had a concentration of 5×10^8 larvae/litre and found that while the optimal concentration varied with the species of flagellate, none of the utilizable species gave toxic effects up to 15×10^8 cells/ml and for certain species up to 25×10^8 /ml. This may be correlated with the larger number of larvae utilizing the food.

GROWTH WITH ALGAL FOODS

Increase in shell length of *Crepidula fornicata* veligers over 3–4.5 week periods was used to demonstrate the relative food values of the 10 algal species. The most vigorous larvae were selected 24 h after hatching and 20 were placed in each of a series of boiling tubes containing 30 ml filtered sea water. The tubes were covered with perforated parafilm, to reduce aerial bacterial contamination, as soon as the algal cells had been added. Similar experiments were set up with *Nassarius reticulatus* veligers and a selected number of algal cultures, and mixed cultures were given to

both species of veliger. The larvae fed vigorously from the beginning of the experiments and all species of algae were ingested. Shell measurements were recorded twice weekly when water and food were renewed and an average of these measurements was used to estimate the relative growth with the different foods. Experiments conducted over 3 breeding seasons showed considerable uniformity of results except for *Crepidula fornicata* collected from the Plymouth area in 1967. Larvae which hatched over the breeding period (May–September) took up to 4 days to deplete the yolk store as compared with 24 h in larvae from the same areas in 1966 and 1968. Although all larvae were kept under identical conditions growth in the 1967 brood was generally poor and this, together with indications of a softening shell in an exceptionally large number of individuals 14 days old, was a sign of an unhealthy stock. This season's results for *Crepidula fornicata* were discarded.

Mean growth in shell length over the whole period of the experiment are given as histograms (Fig. 2). These histograms express the results of single experiments with *Crepidula fornicata* (a, 1966; c, 1968) and *Nassarius reticulatus* (b and d, 1967). Important differences in results from other experiments are referred to later. The larval life of *C. fornicata* is about 5 weeks (CHIPPERFIELD 1951), that is shorter than that of *N. reticulatus* which under laboratory conditions is up to 2 months (LEBOUR 1931). The larvae of *C. fornicata* which grew well were kept under the same conditions after the experimental period to see whether with such a limited diet, and no substratum other than the glass surface, metamorphosis occurred.

Single algal cultures

With each experiment a batch of 20 unfed veligers served as controls. They lived an average of 3 weeks. Their increase in shell length was probably due to the utilization of remaining food reserves and organic matter, including bacteria in the water. Unfed larvae kept in water treated with antibiotics grew less. The histograms show that certain cultures promoted little growth and were either lethal or of no food value so that the larvae starved. *Olisthodiscus* sp. inhibited growth of veligers of *Nassarius reticulatus* at the 2 highest concentrations and they died within 10 days. In the lowest concentration they survived another week, but had grown only as much as the controls. This was the only concentration at which *Crepidula fornicata* survived to the end of the experiment, though the larvae in the other two grew to twice the size of the controls. These algal cells are toxic to the veligers and although the larvae of *Crepidula fornicata* continued to eat and grow at the lowest concentration they died 10 days after the end of the experimental period. Larvae also died before the end of the experiment when given *Exuviaella pusilla*. Observations on stomach contents showed that whereas *Olisthodiscus* sp. fragmented moderately easily, the cells of *E. pusilla* needed a more violent mechanical action before their contents were available for digestion. The food promoted more growth in *N. reticulatus* than in *C. fornicata*. All larvae of *N. reticulatus*, and those of *C. fornicata* in the highest concentration of the alga, died in 3 weeks; the rest of the larvae were unhealthy and died a few days after the end of the experiment. *Chlamydomonas parkeae* and *Brachiomonas submarina*

were also poor foods, but had no toxic effects, and larvae taken from them after a week and given better foods thrived; this did not hold for *Olisthodiscus* sp.

Cricosphaera sp. *carterae* was the best food for both species especially at the higher concentrations. The poor growth of *Crepidula fornicata* at the lowest did not appear to be due to periods of food depletion due to only twice weekly renewal, for larvae given daily renewal of algal cells did not show appreciable increase in growth. There is a tendency for these algal cells, and the large cells of *Brachiomonas submarina* to settle on the bottom of the tube and become less accessible to the veliger, which,




Fig. 2: Mean growth in shell length of veligers of *Crepidula fornicata* and *Nassarius reticulatus* fed on single or mixed algal cultures. Broken line indicates shell length of unfed larvae. The numbers refer to Plymouth cultures, and the 3 columns above each indicate growth at 40×10^3 (left), 20×10^3 (middle) and 2×10^3 (right) cells/ml. Where there are only 2 columns the lowest of the three concentrations was not used, and where only one the highest was used. (a) *C. fornicata*, fed on single algal cultures 4 weeks; (b) *N. reticulatus*, fed on single algal cultures 4 weeks; (c) *C. fornicata*, fed on single and mixed algal cultures 3 weeks; (d) *N. reticulatus*, fed on single and mixed algal cultures $4\frac{1}{2}$ weeks

however, will attempt to sweep them up with the velum. *Exuviaella baltica* was good for *Crepidula fornicata* and in some experiments the larvae grew as well with this food at a concentration of 20×10^3 cells/ml as with *Cricosphaera* sp. *carterae*. This was always the optimal concentration for *E. baltica*. The consistently poor value of

this food to *Nassarius reticulatus* and the greater value of *Dunaliella primolecta* may be indicative of the differences in food requirements of the two species of veliger.

Crepidula fornicata reached the swimming-crawling stage after 40 days feeding on *Exuviaella baltica* at the optimal concentrations, and also at approximately the same time in the two higher concentrations of *Cricosphaera* ap. *carterae*. Larvae eventually metamorphosed, though the swimming-crawling stage was unusually prolonged, presumably through lack of a suitable substratum.

In other experiments the two small-celled algae *Monochrysis lutheri* and *Pyramimonas grossii* were given to *Crepidula fornicata* larvae at the two higher concentrations and 2 others, 80×10^3 and 12×10^4 cells/ml, to test the effect of high density on growth. The results (Tab. 3) were compared with those from larvae fed with *Cricosphaera* ap. *carterae*, the maximum concentration of which was 60×10^3 cells/ml. All the larvae remained healthy.

Table 3

Mean growth increments in shell length of 10 veligers of *Crepidula fornicata* in high concentrations of 3 algal foods over a period of 3 weeks

Species	Concentration (cells/ml)	Growth (μ)
<i>Pyramimonas grossii</i>	20×10^3	80
	40×10^3	80
	80×10^3	70
	12×10^4	50
<i>Monochrysis lutheri</i>	20×10^3	80
	40×10^3	90
	80×10^3	120
	12×10^4	170
<i>Cricosphaera</i> ap. <i>carterae</i>	20×10^3	120
	40×10^3	140
	60×10^3	180

Increase in the number of available cells of *Monochrysis lutheri* and *Cricosphaera* ap. *carterae* resulted in increase in growth. At the highest concentration of *M. lutheri* growth approached that at the highest concentration of *C. ap. carterae*. In contrast the two lower concentrations of *Pyramimonas grossii* were the best and growth decreased in the two higher. These experiments were carried out at the end of the breeding season and this may account for the fact that growth increase in the best food, *C. ap. carterae*, was considerably less than in previous experiments.

The food value of *Monochrysis lutheri* and *Pyramimonas grossii* varied from experiment to experiment at lower concentrations. They are both moderately good foods but sometimes one is better and sometimes the other; a constant factor is that growth is increased with the concentration in the former and decreased in the latter. Variations in their food value may be due to some differences in the cultures.

Mixed algal cultures

Cells of more than one species of alga were mixed in equal proportions and the combined concentrations were 40×10^3 , 20×10^3 or 2×10^3 cells/ml. The numbers of food cells available to the larvae were thus comparable to those in previous experiments; in some experiments only the two higher concentrations were used. The results of feeding larvae with *Cricosphaera* sp. *carterae* and either of the 2 algae which had some lethal effect, *Olisthodiscus* sp. and *Exuviaella pusilla*, showed that the retardation of growth and toxicity caused by *Olisthodiscus* sp. was still evident in both species of veliger, but there was less evidence of *E. pusilla* being detrimental. In the mixture with *E. pusilla* the growth of *Nassarius reticulatus* was improved and the larvae died only in the highest of the 3 concentrations. Results of other experiments are shown in Figure 2, c and d.

Crepidula fornicata larvae which were fed on *Cricosphaera* sp. *carterae* mixed with a poorer food, either *Monochrysis lutheri* or *Pyramimonas grossii*, at a concentration of 40×10^3 cells/ml, grew as well as with *Cricosphaera* sp. *carterae* alone at this concentration. This suggests that they were feeding selectively, a phenomenon described by PAULSON & SCHELTEMA (1968) for the larvae of *Nassarius obsoletus*. However direct observation showed that cells of each alga were ingested and broken down in the stomach. The smaller cells of *Monochrysis lutheri* and *Pyramimonas grossii* were ingested in greater numbers than *Cricosphaera* sp. *carterae* in accordance with the ease with which the velar cilia manipulated them. The cells of *C. ap. carterae* and *Exuviaella baltica* are more closely related in shape and size and about the same number of these were ingested when they were given together. *E. baltica* is a poor food for *Nassarius reticulatus*, but in this mixture growth was considerably improved at the three concentrations. A mixture of these two algae and *Dunaliella primolecta* produced the best growth in *N. reticulatus*.

The results of feeding with mixture of 2 algae not including *Cricosphaera* sp. *carterae* – *Pyramimonas grossii* and *Monochrysis lutheri*, or *Dunaliella primolecta* and *Exuviaella baltica* – show that growth is better than when the poorer food of each pair is given singly.

BACTERIAL FOODS

Experiments with bivalves have shown the deleterious effects of bacteria as foods when larvae are exposed to them for a few days (DAVIS 1953, GUILLARD 1959). The fate of bacteria in the gut of a veliger has not been studied, nor the effect of exposing the larva to bacterial suspensions for brief periods of time. To carry out observations on these, experiments were set up as for algal cultures.

Agar cultures of 8 strains of Gram-negative bacteria from the inshore surface water of Cardigan Bay were supplied by Dr. M. RHODES and Mr. D. WYNN-WILLIAMS and 5 were provisionally identified. According to the test methods given by SHEWAN (1963) 3 were *Vibrio* spp. (K4B, K12B, K27A), all motile, and 2 were *Flavobacterium* spp. (K4A, K28). Another strain was very similar to *Pseudobacterium lateri-*

ceum (KRIS 1963). The maximal size of the bacteria was $1.48 \times 0.92 \mu$; the minimal size $0.46 \times 0.46 \mu$. These measurements were taken with a calibrated shearing ocular using hanging drops of inoculum, but they were not from the actual cultures used for the experiments; they thus give some indication of the particle size collected by the velum. A suspension of cells for feeding to the larvae was obtained by washing 3–5 ml glass-filtered sea water over the surface of a young slope culture. The concentration in the experiments was 10^6 – 10^7 ml. Veligers of *Nassarius reticulatus* were exposed to each of the eight strains of bacteria for periods of up to 1 h, and during this time the gut contents of some were examined under the phase-contrast microscope. Some of the remaining larvae were transferred to a suspension of *Cricosphaera* sp. *carterae* and the rest left in the bacterial suspension. For observations on growth 2 batches of 20 *Nassarius reticulatus* larvae were exposed to motile bacteria for 30 min, transferred to fresh filtered sea water for 5 min and then into another experimental vessel with *C. ap. carterae* at a concentration of 4×10^4 cells/ml. They were left with the algal food 2.5 days when the feeding cycle was repeated. The growth of these larvae over a period of 10 days was compared with that of others fed only on the same concentration of *C. ap. carterae* during this period and with food and water renewed at 2.5 day intervals. All larvae for these experiments were from the same batch of eggs. After hatching they had been kept in glass-filtered sea water 48 h at 12°C .

All strains of bacteria were eaten by the veligers and after 5 min a moderately dense collection was mixed with secretion in the stomach. They were sucked into the lobes of the digestive gland, especially the larger lobe, and taken by the ingesting cells. No bacterium was seen to be motile in the gut: they were passed along rapidly by the oesophageal cilia and into the whirling stomach contents. As compared with algal foods which must be broken mechanically before the digestive gland can deal with them, the ingestion of bacteria by the gland was rapid. It was not until the larvae were removed from the bacterial foods and fed on *Cricosphaera* sp. *carterae* that the ingesting cells were seen to be emitting bacterial waste; they soon became yellow with the algal pigments. After feeding on the plant cells 1.25 h there was faecal waste containing residues from both foods. Whole bacteria appeared to be in it, but changes in their structure revealed by the phase-contrast microscope suggested that something other than the mucous coat which envelops each cell may have been utilized by the larvae. The activity of the larvae was not impaired, but larvae not removed from bacterial food were unhealthy after 1.5 days. An examination of their digestive glands showed the ingesting cells packed with bacteria and apparently incapable of voiding them as though injured by their toxic metabolites.

The average growth in shell length of 40 veligers, which had been exposed to bacteria for 4 half-h periods during 10 days and otherwise fed on *Cricosphaera* sp. *carterae*, was 80μ . For veligers fed on *C. ap. carterae* only it was 90μ . Both batches of larvae remained healthy after the experiment. The results show that the occasional high intake of these bacteria is not injurious to the healthy veligers provided that an abundance of good food, moderately free from bacteria, is otherwise available.

DISCUSSION

The factors influencing the value of algal species as food for gastropod larvae include the composition and texture of the cell wall as well as cell size and contents. The walls of algal cells are of varied composition and texture. There is no evidence that they are utilized by any prosobranch larva and their shattered remains can be found in the faeces; no fragment of wall has been seen in the cells of the digestive gland. The mechanical treatment the cells receive from the moment they enter the stomach weakens or disintegrates the walls so that cell contents are made accessible to the gastric juices. The resistance of cells with complete cellulose walls suggests the absence of a cellulase (FRETTER & MONTGOMERY 1968). Thus *Chlamydomonas parkeae* is retained in the stomach of *Crepidula fornicata* and *Nassarius reticulatus*, but the digestive gland is not coloured with the algal pigment and cells pass into the intestine apparently unharmed. It was used only in growth experiments with *C. fornicata*. The slight growth which resulted may have been due to the digestion of the mucous coat surrounding each cell and bacteria and organic matter from the culture medium introduced with the algal cells. The cell contents of another chlorophycean, *Brachiomonas submarina*, were obtainable after considerable battering against the gastric shield. Utilization of this food was slow and poor growth resulted. On the contrary, the cellulose walls of *Exuviaella baltica* offer much less resistance since each is composed of 2 watch-glass shaped thecae which can be forced apart in about 10 min and soon after this the cell empties. Cells of the second species of this genus, *E. pusilla*, are emptied more slowly.

Cells of *Monochrysis lutheri* and *Pyramimonas grossii* fragment rapidly in the stomach, but because of their size and shape they provide a higher proportion of undigestible cell covering to utilizable inclusions. A workable stomachful is about 60 cells and from these the faecal waste is bulky. In healthy larvae this waste is flushed away by the flow of water through the mantle cavity which is maintained by the osphradial cilia. But, if larvae are less vigorous, excessive waste from high concentrations of food will accumulate near the anus. These faeces and also those resulting from a meal of the diatom *Phaeodactylum tricorutum* are frequently uncompacted and disintegrate as they are dropped. The spiny frustules of the diatom are discarded intact and some still have protoplasmic contents. The larvae are wasteful with this food: the cells are easily ingested and their presence in the stomach stimulates the muscular pulsations of the lobes of the digestive gland, but, perhaps because they irritate the gut, an undiluted meal of these cells, especially if they are large, acts as a purgative.

The best food, *Cricosphaera* sp. *carterae*, comprises large, almost spherical cells, their diameter being 10–18 μ , and about 30 are found in a stomachful. They are battered vigorously, but the walls do not fragment. The flagella are soon lost, the calcite plates of the walls are forced apart as the protoplasm swells and within 15 min the plant pigments colour the digestive gland and some cells are empty. The emptied remains are retained in the stomach while other cells are digested. Undigested cells are rare in the faeces.

The ease with which the contents of algal cells are made available for digestion

is important in a mixed diet. If cells of *Cricosphaera* sp. *carterae* are eaten with the easily fragmented cells of *Monochrysis lutheri* or *Pyramimonas grossii*, or even with cells of *Olisthodiscus* sp. or *Exuviaella baltica* which need a moderate mechanical force to weaken them, some *Cricosphaera* sp. *carterae* cells will leave the gut undigested. This is especially noticeable with the smaller cells which are so easily ingested. Their fragments buffer the walls of *C. ap. carterae* against the hurling action of the gastric cilia and the stomach is emptied before all the cells are digested. The stomach is emptied in response to a bulky accumulation of waste or the availability of more food. When *Olisthodiscus* sp. is eaten with *C. ap. carterae* its cell contents are the more easily available so that its toxic properties appear virtually unreduced.

Plant pigments as well as cell walls are waste to the veliger. All pigments are taken into the ingesting cells and later leave them, so they delay the cycle of events in the digestive gland. PARSONS et al. (1961) have estimated indigestible matter of some algal species with respect to crude fibre, as % total carbohydrate, and pigment content as % dry weight. For *Cricosphaera* sp. *carterae* these are 1.7 % and 1.1 % respectively, for *Monochrysis lutheri* 3.6 % and 0.8 % and for *Phaeodactylum tricornutum* 2.5 % and 2.9 %. *C. ap. carterae* is moderately low in crude fibre and pigment, in agreement with the observation that little waste is produced, and *Monochrysis lutheri* is high in crude fibre. However an unspecified species of *Exuviaella baltica* had 37.0 % total carbohydrate as crude fibre. It would be of interest to know more about the 2 species of *E. baltica* which give such contrasting results in growth experiments with prosobranch veligers.

When different species of alga are grown under approximately similar conditions they tend to resemble each other in the relative amount of crude protein, fats and hydrolysable polysaccharide (COLLYER & FOGG 1955). Some differences do exist between classes, but these are usually small compared with the differences exhibited by a single species during the course of its growth in culture (FOGG 1955).

COWEY & CORNER (1966) and CHAU et al. (1967) have estimated the amino-acids for several species of unicellular algae including Plymouth cultures used in these experiments (100, 75, 404, 156, 81, 285 and 239). COWEY & CORNER point out the similarity between the amino-acid spectra of the algal species and suggest that differences in food value cannot be explained in terms of amino-acid composition. This is further supported by DROOP (1966) working on the response of amoebae to different algal foods. Earlier work by COWEY & CORNER (1962) demonstrates that the amino-acid composition of particulate matter in sea water, algal cells, *Calanus* and fish muscle are very similar, suggesting that protein of a certain amino-acid composition may be typical of food chains in the sea. In contrast to amino-acids the component fatty-acid composition of marine phytoplankton lacks uniformity. CHUECAS & RILEY (1969) have studied 27 species of marine phytoplankton and their results indicate that the assemblages of unsaturated acids vary widely from one organism to another.

The concentration of algal food producing the best growth of veligers in the laboratory is much higher than they would normally be exposed to in Plymouth Sound. Similarly WALNE (1965) quotes a considerable amount of field data which points to the fact that oyster larvae are often living at lower densities of algal cells than

those giving best growth in the laboratory, and experiments by CUSHING (1959) suggest such a discrepancy for *Calanus*. WALNE concludes that the organic detritus in the sea (CORNER 1961) comprises part of the oysters' food. COWEY & CORNER (1963) have shown that as far as amino-acids are concerned *Calanus* would gain no advantage nutritionally from a selection of algal cells in preference to the amino-acid containing fraction of particulate matter as a whole. Obviously the same must hold for prosobranch veligers which ingest any particle not too big for the velar cilia to manipulate and retain the organic ones in the stomach for digestion.

Species of algae differ considerably in food value, even those belonging to the same genus as has been shown in the present experiment for *Exuviaella baltica* and *E. pusilla*. Previously it has been demonstrated that *Dunaliella euchlora* is a very poor food for clam larvae and another species of this genus good (DAVIS & GUILLARD 1958); *Cosmarium impressulum* and *Scenedesmus spinosus* are good for *Daphnia magna* whilst *C. tetraophthalmum*, *S. oahuensis* and *S. quadricula* are poor or mediocre (LEFÈVRE 1942, PROVASOLI et al. 1959). Some of these differences may be associated with differences in micronutrients and vitamins. Deficiencies in some algal foods have been made good under experimental conditions. The life span of *Daphnia magna* fed on a diet of *Chlamydomonas parkeae* was tripled by the addition of 200 mg/l pantothenic acid and egg production increased tenfold (FRITSCH 1953). Similarly riboflavin (alone or in combination with calcium pantothenate), thiamine hydrochloride and pyridoxine hydrochloride significantly increased the rate of growth of veligers of *Crassostrea virginica* and *Ostrea lurida* (DAVIS & CHANLEY 1956). However, these vitamins had no effect on the growth rate of the larvae of *Mercenaria mercenaria*. The requirements of the different species of bivalve larvae might be expected to vary and those of the veligers of *Crepidula fornicata* and *Nassarius reticulatus* even more so. It has been shown for *C. fornicata* that *Exuviaella baltica* approached *Cricosphaera* sp. *carterae* in nutritive value, but was a poor food for *N. reticulatus*. The adult *C. fornicata* is a microphagous feeder and the adult *N. reticulatus* essentially a carrion feeder and it is surprising that *C. ap. carterae* proved a good food for both, even up to the time of metamorphosis. *C. fornicata* veligers given only this food were ready to metamorphose after 4–6 weeks and *N. reticulatus* after 8 weeks.

A study of *Olisthodiscus* sp. which is toxic to these and other prosobranch veligers reveals a low lipid and carbohydrate content (RICKETTS 1966) and a high concentration of acid-soluble phosphorus compounds which may reflect an unidentified storage product. Only 64% of its total dry weight has been accounted for in terms of known substances. BIDWELL (1957) has found that the main storage product of photosynthesis is *d*-mannitol. Although even young cultures of this food are lethal to both young and old veligers of prosobranchs, larvae of *Sabellaria alveolata* are apparently unaffected by the alga. The largest larvae of *S. alveolata* reared in some experiments were those which in their older stages were given *Olisthodiscus* sp. alone or with another flagellate (WILSON 1968).

SUMMARY

1. Larvae of *Crepidula fornicata* (L.) and *Nassarius reticulatus* (L.) were used for experiments. The conditions under which they thrived proved successful for veligers of other species.
2. They were kept in glass-filtered sea water (pore size 3.0μ) in acid-clean glass containers, provided with algal foods and handled carefully by means of a pipette. Trapping larvae in a coarse filter as a means of transferring them from one vessel to another was injurious.
3. Shell length was used to estimate growth.
4. The growth of 20 veligers in 30 ml sea water (depth 8 cm) was compared with that of 333 veligers in 500 ml (depth 9 cm). Growth in the larger volume was better, but in all cases the differences were not significant at the 10 % level.
5. Growth rate is influenced by light intensity. For comparing the value of different foods experiments were carried out under a constant intensity of 155 lux at the water surface. In the dark, mortality of algal cells stimulates growth of bacteria.
6. The water temperature was maintained at 12° C. Low temperatures, even 8° C, reduce the activity of veligers of *Crepidula fornicata* and *Nassarius reticulatus*; high temperatures favour bacterial growth.
7. The growth of recently hatched veligers feeding on one of 10 species of unicellular algae and on some mixtures of these was recorded for 2 (*C. fornicata*) or 3 (*N. reticulatus*) breeding seasons. Food was given at different concentrations (2×10^3 , 20×10^3 , 40×10^3 cells/ml) which were calculated from haemocytometer counts of the stock cultures, though this gives only an approximate value. Experiments lasted up to 4.5 weeks and a few for a longer period.
8. *Cricosphaera* ap. *carterae* and *Exuviaella baltica* were the best foods for *Crepidula fornicata*, especially at higher concentrations, and larvae were ready to metamorphose in 40 days or less. *C.* ap. *carterae* and *Dunaliella primolecta* were good for *Nassarius reticulatus*, especially the former, and *E. baltica* consistently poorer.
9. *Monochrysis lutheri* and *Pyramimonas grossii* were moderately good foods, but with these none of the larvae metamorphosed. When fed to *Crepidula fornicata* at high concentrations (80×10^3 , 120×10^3 cells/ml) growth of the former approached that with *Cricosphaera* ap. *carterae* whilst with the latter growth decreased with increased concentration. The food value of *Phaeodactylum tricornutum* is lower; the large frustules irritate the gut and act as a purgative.
10. *Chlamydomonas parkeae*, *Brachiomonas submarina*, *Exuviaella pusilla* and *Olisthodiscus* sp. are poor foods; the last two are toxic.
11. When the food was 2 species of alga mixed in equal proportions the good value of *Cricosphaera* ap. *carterae* was still evident. Examination of the stomach contents showed that the larvae were not feeding selectively on this alga. When the second alga was one with smaller cells (*Monochrysis lutheri* or *Pyramimonas grossii*) these were ingested in greater numbers in accordance with the ease with which the velar cilia manipulated them.
12. Algal cells are subjected to mechanical treatment in the stomach; their walls may

- be shattered but no fragment has been seen in the cells of the digestive gland. The resistance of cells with complete cellulose walls suggests the absence of a cellulase.
13. With some algal foods (species of *Monochrysis*, *Pyramimonas*, *Phaeodactylum*) the walls produce a high proportion of faecal waste which, in less vigorous larvae, may clog the exhalant passage of the mantle cavity.
 14. Plant pigments are egested by the veliger. They are intimately linked with food, taken into ingesting cells of the digestive gland and later excreted. They thus delay the cycle of events in the gland.
 15. Differences in food value of the various algae may be due to differences in micronutrients and vitamins essential to growth. The fact that one species of alga may produce good growth in one species of veliger and not another must reflect either differences in requirements or in assimilation of the food.
 16. Prosobranch veligers are found in numbers well below the compensation depth as well as in other areas where the density of algal cells is low. It is suggested that organic detritus in the sea is an important item of food.

Acknowledgements. This work was carried out at the Plymouth Laboratory and was supported by a grant from the Science Research Council. We thank Dr. M. PARKE who supplied the algal cultures and gave valuable advice, the crews and staff of the Marine Biological Association, Plymouth, and, in particular, the Director, Dr. J. E. SMITH, F.R.S., for providing space and facilities. One of us (M.C.P.) wishes to thank the British Association for the use of their table at the Laboratory. We are also indebted to Dr. M. RHODES and Mr. D. WYNN-WILLIAMS for supplying the cultures of bacteria and the information concerning them.

LITERATURE CITED

- ARMSTRONG, F. A. J. & BUTLER, E. I., 1960a. Chemical changes in sea water off Plymouth during 1958. *J. mar. biol. Ass. U.K.* **39**, 299-302.
- — 1960b. Chemical changes in sea water off Plymouth during 1959. *J. mar. biol. Ass. U.K.* **39**, 525-528.
- — 1962. Chemical changes in sea water off Plymouth during 1960. *J. mar. biol. Ass. U.K.* **42**, 253-258.
- — 1963. Chemical changes in sea water off Plymouth in 1961. *J. mar. biol. Ass. U.K.* **43**, 75-78.
- BAINBRIDGE, R., 1957. The size, shape and density of marine phytoplankton concentrations. *Biol. Rev.* **32**, 91-115.
- BIDWELL, R. G. S., 1957. Photosynthesis and metabolism of marine algae. I. Photosynthesis of two marine flagellates compared with *Chlorella*. *Can. J. Bot.* **55**, 945-950.
- BONE, Q., 1961. The organization of the atrial nervous system of *Amphioxus* (*Branchiostoma lanceolatum* PALLAS). *Phil. Trans. R. Soc. (B)* **243**, 241-269.
- CHAU, Y. K., CHUECAS, L. & RILEY, J. P., 1967. The component combined amino acids of some marine phytoplankton species. *J. mar. biol. Ass. U.K.* **47**, 543-554.
- CHIPPERFIELD, P. N. J., 1951. The breeding of *Crepidula fornicata* (L.) in the river Blackwater, Essex. *J. mar. biol. Ass. U.K.* **30**, 49-71.
- CHUECAS, L. & RILEY, J. P., 1969. Component fatty acids of the total lipids of some marine phytoplankton. *J. mar. biol. Ass. U.K.* **49**, 97-116.
- COLLYER, D. M. & FOGG, G. E., 1955. Studies on fat accumulation by algae. *J. exp. Bot.* **6**, 256-275.

- CORNER, E. D. S., 1961. On the nutrition and metabolism of zooplankton. I. Preliminary observations on the feeding of the marine copepod, *Calanus helgolandicus* (CLAUS). *J. mar. biol. Ass. U.K.* **41**, 5-16.
- COWEY, C. B. & CORNER, E. D. S., 1962. The amino-acid composition of *Calanus finmarchicus* (CLAUS) in relation to that of its food. *Rapp. P.-v. Réun. Cons. perm. int. Explor. Mer* **153**, 124-128.
- — 1963. On the nutrition and metabolism of zooplankton. II. The relationship between the marine copepod *Calanus helgolandicus* and particulate material in Plymouth sea water, in terms of amino-acid composition. *J. mar. biol. Ass. U.K.* **43**, 495-511.
- — 1966. The amino-acid composition of certain unicellular algae and of the faecal pellets produced by *Calanus finmarchicus* when feeding on them. *In: Some contemporary studies in marine science*. Ed. by H. Barnes. Allen & Unwin, London, 225-231.
- CUSHING, D. H., 1959. On the nature of production in the sea. *Fishery Invest., Lond. (Ser. 2)* **22**, 1-40.
- DAVIS, H. C., 1953. On food and feeding of larvae of the American oyster, *C. virginica*. *Biol. Bull. mar. biol. Lab., Woods Hole* **104**, 334-350.
- & CHANLEY, P. E., 1956. Effects of some dissolved substances on bivalve larvae. *Proc. natn. Shellfish. Ass.* **46**, 59-74.
- & GUILLARD, R. R., 1958. Relative value of ten genera of micro-organisms as food for oyster and clam larvae. *Fishery Bull. Fish Wildl. Serv. U.S.* **58**, 293-304.
- DROOP, M. R., 1966. The role of algae in the nutrition of *Heteramoeba clara* DROOP, with notes on *Oxyrrhis marina* DUJARDIN and *Philodina roseola* EHRENBERG. *In: Some contemporary studies in marine science*. Ed. by H. Barnes. Allen & Unwin, London, 269-282.
- FOGG, G. E., 1965. Algal cultures and phytoplankton ecology. Athlone press, London, 126 pp.
- FRETTER, V., 1967. The prosobranch veliger. *Proc. malac. Soc. Lond.* **37**, 357-366.
- & MONTGOMERY, M. C., 1968. The treatment of food by prosobranch veligers. *J. mar. biol. Ass. U.K.* **48**, 499-520.
- FRITSCH, R. H., 1953. Die Lebensdauer von *Daphnia* spec. bei verschiedener Ernährung, besonders bei Zugabe von Pantothenensäure. *Z. wiss. Zool.* **157**, 35-56.
- GUILLARD, R. R. L., 1959. Further evidence of the destruction of bivalve larvae by bacteria. *Biol. Bull. mar. biol. Lab., Woods Hole* **117**, 258-266.
- IMAI, T. & HATANAKA, M., 1949. On the artificial propagation of the Japanese common oyster, *Ostrea gigas* THUN. by non-coloured naked flagellates. *Bull. Inst. agric. Res. Tohoku Univ.* **1**, 33-46.
- KORRINGA, P., 1951. Difficulties encountered in tank-breeding of oysters (*Ostrea edulis*). *Rapp. P. v. Réun. Cons. perm. int. Explor. Mer* **128**, 35-38.
- KRISS, A. E., 1963. Marine microbiology. Transl. from the Russian. Oliver & Boyd, Edinburgh, 536 pp.
- LEBOUR, M. V., 1931. The larval stages of *Nassarius reticulatus* and *N. incrassatus*. *J. mar. biol. Ass. U.K.* **17**, 797-816.
- LEFÈVRE, M., 1942. L'utilisation des algues d'eau douce par les cladocères. *Bull. biol. Fr. Belg.* **76**, 250-276.
- MARTINDALE, W., 1967. Penicillins and other antibiotics. *In: Todd, R. G.: Extra Pharmacopoeia*. The Pharmaceutical Press, London, 1804 pp.
- PARSONS, T. R., STEPHENS, K. & STRICKLAND, J. D. H., 1961. On the chemical composition of eleven species of marine phytoplankters. *J. Fish. Res. Bd Can.* **18**, 1001-1016.
- PAULSON, T. C. & SCHELTEMA, R. S., 1968. Selective feeding on algal cells by the veliger larvae of *Nassarius obsoletus* (Gastropoda, Prosobranchia). *Biol. Bull. mar. biol. Lab., Woods Hole* **134**, 481-489.
- PROVASOLI, L., SHIVAIKI, K. & LANCE, J. R., 1959. Nutritional idiosyncrasies of *Artemia* and *Tigriopus* in monoxenic culture. *Ann. N. Y. Acad. Sci.* **77**, 250-261.
- REEVE, M. R., 1963. The filter-feeding of *Artemia*, I, II, III. *J. exp. Biol.* **40**, 195-221.
- RICKETTS, T. R., 1966. On the chemical composition of some unicellular algae. *Phytochemistry* **5**, 67-76.

- SCHELTEMA, R. S., 1962. Pelagic larvae of New England intertidal gastropods. I. *Nassarius obsoletus* SAY and *Nassarius vibex* SAY. *Trans. Am. microsc. Soc.* **81**, 1-11.
- 1967. The relationship of temperature to the larval development of *Nassarius obsoletus* (Gastropoda). *Biol. Bull. mar. biol. Lab., Woods Hole* **132**, 253-265.
- SHEWAN, J. M., 1963. The differentiation of certain genera of gram negative bacteria frequently encountered in marine environments. In: Symposium on marine microbiology. Ed. by C. H. Oppenheimer. C. C. Thomas, Springfield, Ill., 499-521.
- WALNE, P. R., 1956a. Bacteria in experiments on rearing oyster larvae. *Nature, Lond.* **178**, 91.
- 1956b. Experimental rearing of larvae of *Ostrea edulis* L. in the laboratory. *Fishery Invest., Lond. (Ser. 2)* **20** (9), 1-23.
- 1958. The importance of bacteria in laboratory experiments on rearing the larvae of *Ostrea edulis* (L.) *J. mar. biol. Ass. U.K.* **37**, 415-425.
- 1963. Observations on the food value of seven species of algae to the larvae of *Ostrea edulis*. 1. Feeding experiments. *J. mar. biol. Ass. U.K.* **43**, 767-784.
- 1965. Observations on the influence of food supply and temperature on the feeding and growth of the larvae of *Ostrea edulis* L. *Fishery Invest., Lond. (Ser. 2)* **24** (1), 1-45.
- WILSON, D. P., 1968. Some aspects of the development of eggs and larvae of *Sabellaria alveolata* (L.). *J. mar. biol. Ass. U.K.* **48**, 367-386.

First author's address: MARGARET C. PILKINGTON
Portobello Biological Station
New Zealand