

SOME NEW RESULTS ABOUT HAMMERSTEIN EQUATIONS¹

BY HAIM BRÉZIS AND FELIX E. BROWDER

Communicated by S. S. Chern, August 1, 1973

Let Ω be a σ -finite measure space. Let K be a (nonlinear) monotone operator and let $(Fu)(x) = f(x, u(x))$ be a Niemytski operator. We consider the Hammerstein type equation

$$(1) \quad u + KF u = g.$$

A detailed discussion and a complete bibliography about equation (1) can be found in [3]. The new feature about the results we present here is the fact that we do not assume any coercivity for F . When F is monotone and K maps $L^1(\Omega)$ into $L^\infty(\Omega)$, there is no growth restriction on F either (cf. Theorem 1). The monotonicity of F can be weakened when K is compact (cf. Theorem 4). Also some of these results are valid for systems in the case where F is the gradient of a convex function (cf. Theorem 5).

Assume

(2) K is a monotone hemicontinuous mapping from $L^1(\Omega)$ into $L^\infty(\Omega)$ which maps bounded sets into bounded sets,

(3) $f(x, r): \Omega \times \mathbf{R} \rightarrow \mathbf{R}$ is continuous and nondecreasing in r for a.e. $x \in \Omega$, and is integrable in x for all $r \in \mathbf{R}$.

THEOREM 1. *Under the assumptions (2) and (3), equation (1) has one and only one solution $u \in L^\infty(\Omega)$ for every $g \in L^\infty(\Omega)$.*

Uniqueness. Let u_1 and u_2 be two solutions of (1). By the monotonicity of K we get

$$\int_{\Omega} (u_1(x) - u_2(x)) \cdot (f(x, u_1(x)) - f(x, u_2(x))) dx \leq 0$$

which implies that $f(x, u_1(x)) = f(x, u_2(x))$ a.e. on Ω and therefore by (1), $u_1 = u_2$.

In proving existence of u we shall use the following

LEMMA 1. *Let X be a Banach space and let $K: X \rightarrow X^*$ and $F: X^* \rightarrow X$ be two monotone hemicontinuous operators. Let $\{u_n\} \subset X^*$, $\{v_n\} \subset X$ and*

AMS (MOS) subject classifications (1970). Primary 47H05, 47H15, 45G05.

¹ Partially supported by NSF GP-28148.

$\{w_n\} \subset X^*$ be three sequences such that

- (4) u_n converges to u in X^* for the weak* topology,
 - (5) $F(u_n)$ converges to v in X for the weak topology,
 - (6) v_n converges to v in X for the weak topology,
 - (7) Kv_n converges to $g-u$ in X^* for the weak* topology,
 - (8) $\langle w_n, F(u_n) \rangle - \langle Kv_n, v_n \rangle \rightarrow 0$,
 - (9) $\langle g_n, F(u_n) \rangle \rightarrow \langle g, v \rangle$ where $g_n = u_n + w_n$.
- Then $u + KF u = g$.

PROOF OF LEMMA 1. We have

$$\langle u_n - u, F(u_n) \rangle = \langle g_n - w_n - u, F(u_n) \rangle.$$

By the monotonicity of K we get

$$\langle Kv_n, v_n \rangle \geq \langle Kv_n, v \rangle + \langle Kv, v_n - v \rangle$$

and thus

$$\liminf \langle Kv_n, v_n \rangle \geq \langle g - u, v \rangle.$$

By (8) we have

$$\liminf \langle w_n, Fu_n \rangle \geq \langle g - u, v \rangle.$$

Consequently, $\limsup \langle u_n - u, F(u_n) \rangle \leq 0$. Since F is pseudomonotone (cf. [1]), we conclude that $v = Fu$ and $\langle u_n, F(u_n) \rangle \rightarrow \langle u, v \rangle$. Also $\langle Kv_n, v_n \rangle \rightarrow \langle g - u, v \rangle$ since $\langle w_n, F(u_n) \rangle = \langle g_n - u_n, F(u_n) \rangle \rightarrow \langle g, v \rangle - \langle u, v \rangle$. Thus

$$\lim \langle Kv_n, v_n - v \rangle = 0,$$

and again, since K is pseudomonotone, we conclude that $g - u = Kv = KF u$.

PROOF OF THEOREM 1. By a shift we can always assume that $f(x, 0) = 0$ and that $K0 = 0$ (note that (1) can be written as $u + \tilde{K}Fu = \tilde{g}$, where $\tilde{F}v = Fv - F0$, $\tilde{K}v = K(v + F(0)) - KF0$ and $\tilde{g} = g - KF0$). Let Ω_n be an increasing sequence of finite measure subsets of Ω such that $\bigcup_n \Omega_n = \Omega$. Let χ_n be the characteristic function of Ω_n . Let F_n be F truncated by n , i.e.,

$$\begin{aligned} f_n(x, r) &= f(x, r) && \text{whenever } |f(x, r)| < n, \\ &= nf(x, r)/|f(x, r)| && \text{whenever } |f(x, r)| \geq n. \end{aligned}$$

The equation

$$(10) \quad u_n + \chi_n K \chi_n F_n(u_n) = \chi_n g$$

has a solution.

Indeed the mapping $K_n: v \mapsto \chi_n K \chi_n v$ is monotone hemicontinuous from $L^2(\Omega)$ into itself.

On the other hand, the (multivalued) operator A defined on $L^2(\Omega)$ by

$$Av = \{w \in L^2(\Omega); v(x) = \chi_n(x) f_n(x, w(x)) \text{ a.e. on } \Omega\}$$

is maximal monotone in $L^2(\Omega)$ and $D(A)$ is bounded in $L^2(\Omega)$ ($|v|_{L^2} \leq n(\text{meas } \Omega_n)^{1/2}$, $v \in D(A)$). Consequently, $R(A+K_n)=L^2(\Omega)$ (cf. [2]) and (10) has a solution.

Multiplying (10) through by $F_n(u_n)$ and using the monotonicity of K we get

$$(11) \quad \int_{\Omega} u_n \cdot F_n(u_n) \, dx \leq \int_{\Omega} \chi_n g F_n(u_n) \, dx.$$

Let $C=2\|g\|_{L^\infty}$; we have

$$\begin{aligned} \int_{\Omega} u_n F_n(u_n) \, dx &= \int_{|u_n| \geq C} u_n F_n(u_n) \, dx + \int_{|u_n| < C} u_n F_n(u_n) \, dx \\ &\geq C \int_{|u_n| \geq C} |F_n(u_n)| \, dx - C \int_{|u_n| < C} |F_n(u_n)| \, dx \\ &\geq C \int_{\Omega} |F_n(u_n)| \, dx - 2C \int_{|u_n| < C} |F_n(u_n)| \, dx. \end{aligned}$$

Using (11) we obtain

$$\int_{\Omega} |F_n(u_n)| \, dx \leq 4 \int_{|u_n| \leq C} |F_n(u_n)| \, dx \leq 4 \int_{|u_n| \leq C} |f(x, u_n(x))| \, dx \leq C'$$

by assumption (3).

Going back to (10), we conclude that $\{u_n\}$ remains bounded in $L^\infty(\Omega)$. Therefore, by assumption (3), there is some function $h \in L^1(\Omega)$ such that

$$(12) \quad |F_n(u_n)(x)| \leq |f(x, u_n(x))| \leq h(x) \quad \text{a.e. on } \Omega.$$

We apply now Lemma 1 with $v_n = \chi_n F_n(u_n)$, $w_n = \chi_n K v_n$, $g_n = \chi_n g$. By extracting a subsequence, we can always assume that

- u_n converges to u weak* in $L^\infty(\Omega)$,
- $F(u_n)$ converges to v weakly in $L^1(\Omega)$,
- v_n converges to v weakly in $L^1(\Omega)$,
- g_n converges to g weak* in $L^\infty(\Omega)$.

Hence

- w_n converges to $g-u$ weak* in $L^\infty(\Omega)$,
- Kv_n converges to $g-u$ weak* in $L^\infty(\Omega)$.

It remains to verify (8) and (9). We have

$$\begin{aligned} \langle w_n, F(u_n) \rangle &= \int_{\Omega} \chi_n K v_n \cdot F(u_n) \, dx = \int_{\Omega} K v_n \chi_n F(u_n) \, dx \\ &= \int_{\Omega} K v_n \cdot v_n \, dx + \int_{\Omega} \chi_n K v_n (F(u_n) - F_n(u_n)) \, dx. \end{aligned}$$

The last term can be bounded by

$$C \int_{|F(u_n)| > n} |Fu_n| \, dx \leq C \int_{|h| > n} |h(x)| \, dx$$

which tends to zero as $n \rightarrow +\infty$ and (8) follows.

Finally (9) holds since

$$\langle g_n, F(u_n) \rangle = \int_{\Omega} \chi_n g F(u_n) \, dx = \int_{\Omega} g F(u_n) \, dx + \int_{\Omega} (\chi_n - 1) g F(u_n) \, dx,$$

and the last term goes to zero by Lebesgue's theorem.

THEOREM 2 (CONTINUOUS DEPENDENCE). *Under the assumptions (2) and (3), $F(I+KF)^{-1}$ is strongly continuous from $L^\infty(\Omega)$ into $L^1(\Omega)$ and $(I+KF)^{-1}$ is demicontinuous (from $L^\infty(\Omega)$ strong into $L^\infty(\Omega)$ weak*). If in addition K is strongly continuous from $L^1(\Omega)$ into $L^\infty(\Omega)$, then $(I+KF)^{-1}$ is strongly continuous from $L^\infty(\Omega)$ into $L^\infty(\Omega)$.*

PROOF. We shall prove a slightly stronger result. Let g_n be a bounded sequence in $L^\infty(\Omega)$ such that $g_n \rightarrow g$ a.e. on Ω . Let $u_n = (I+KF)^{-1}g_n$ and let $u = (I+KF)^{-1}g$. We are going to show that $F(u_n) \rightarrow F(u)$ in $L^1(\Omega)$.

We know, from the proof of Theorem 1, that $\{u_n\}$ is bounded in $L^\infty(\Omega)$ and there is some $h \in L^1(\Omega)$ such that $|F(u_n)| \leq h$ a.e. on Ω . Since

$$\int_{\Omega} (u_n - u)(F(u_n) - F(u)) \, dx \leq \int_{\Omega} (g_n - g)(F(u_n) - F(u)) \, dx$$

and the right hand side goes to zero by Lebesgue's theorem, we can extract a subsequence such that

$$(u_{n_k} - u)(F(u_{n_k}) - F(u)) \rightarrow 0 \quad \text{a.e. on } \Omega.$$

Consequently, $F(u_{n_k}) \rightarrow F(u)$ a.e. on Ω and hence $F(u_{n_k}) \rightarrow F(u)$ in $L^1(\Omega)$. By the uniqueness of the limit we conclude that $F(u_n) \rightarrow F(u)$ in $L^1(\Omega)$.

Using similar arguments, we can prove some variants of Theorem 1.

THEOREM 3. *Assume K is monotone, hemicontinuous and bounded from $L^{p'}(\Omega)$ into $L^p(\Omega)$. Assume $f(x, r): \Omega \times \mathbf{R} \rightarrow \mathbf{R}$ is continuous and nonincreasing in r for a.e. $x \in \Omega$ and is measurable in x for all $x \in \mathbf{R}$, and satisfies*

$$|f(x, r)| \leq c(x) + c_0 |r|^{p-1} \quad \text{a.e. } x \in \Omega, \text{ for all } r \in \mathbf{R}$$

where $c \in L^{p'}(\Omega)$.

Then (1) has a unique solution $u \in L^p(\Omega)$ for every $g \in L^p(\Omega)$.

THEOREM 4. *Assume K is monotone, hemicontinuous from $L^1(\Omega)$ into $L^\infty(\Omega)$ and maps bounded sets of $L^1(\Omega)$ into compact sets of $L^\infty(\Omega)$.*

Assume $f(x, r)$ is continuous in r for a.e. $x \in \Omega$ and there exists M such that

$$(f(x, r) - f(x, 0))r \geq 0 \text{ for a.e. } x \in \Omega \text{ and for all } |r| \geq M.$$

Suppose $f(x, r)$ is measurable in x for all $r \in \mathbf{R}$ and for every constant C ,

$$\int_{|r| \leq C} |f(x, r)| \text{ is integrable.}$$

Then (1) has a solution $u \in L^\infty(\Omega)$ for every $g \in L^\infty(\Omega)$.

The case of systems. Assume

(13) K is monotone hemicontinuous and bounded from $L^1(\Omega; \mathbf{R}^n)$ into $L^\infty(\Omega; \mathbf{R}^n)$.

(14) $f(x, r): \Omega \times \mathbf{R}^n \rightarrow \mathbf{R}^n$ is continuous in r for a.e. $x \in \Omega$ and trimonotone in r , i.e., for a.e. $x \in \Omega$ and for any sequence $r_0, r_1, r_2, r_3 = r_0$ we have

$$\sum_{i=1}^3 (f(x, r_i), r_i - r_{i-1}) \geq 0$$

(for example, the gradient of a convex function is trimonotone, see [4]).

(15) $f(x, r)$ is measurable in x for all $r \in \mathbf{R}$ and for every constant C

$$\int_{|r| \leq C} |f(x, r)| \text{ is integrable.}$$

THEOREM 5. Under the assumptions (13), (14), (15), equation (1) has a unique solution $u \in L^\infty(\Omega; \mathbf{R}^n)$ for every $g \in L^\infty(\Omega; \mathbf{R}^n)$.

In order to bound Fu in L^1 , we use the following

LEMMA 2. Assume (14) and (15) hold. Then for any constant $\rho > 0$, there exists $h_\rho \in L^1(\Omega)$ such that

$$\rho |f(x, r)| \leq (f(x, r) - f(x, 0), r) + h_\rho(x) \text{ for a.e. } x \in \Omega, \text{ all } r \in \mathbf{R}^n.$$

Uniqueness follows from the following

LEMMA 3. Assume B is continuous and trimonotone from a Hilbert space H into itself. Let $u, v \in H$ be such that

$$(Bu - Bv, u - v) = 0.$$

Then $Bu = Bv$.

Along the same lines one can prove the following lemma which leads to stability results.

LEMMA 4. Assume B is trimonotone and Hölder continuous with exponent $\alpha \leq 1$ (i.e., $|Bu - Bv| \leq L|u - v|^\alpha$ for all $u, v \in H$).

Then there exists a constant $k > 0$ such that

$$(Bu - Bv, u - v) \geq k |Bu - Bv|^{1+1/\alpha} \quad \text{for all } u, v \in H.$$

REFERENCES

1. H. Brézis, *Équations et inéquations non linéaires dans les espaces vectoriels en dualité*, Ann. Inst. Fourier (Grenoble) **18** (1968), fasc. 1, 115–175. MR **42** #5113.
2. F. E. Browder, *Nonlinear maximal monotone operators in Banach spaces*, Math. Ann. **175** (1968), 89–113. MR **36** #6989.
3. ———, *Nonlinear functional analysis and nonlinear integral equations of Hammerstein and Urysohn type*, Contributions to Nonlinear Functional Analysis, E. Zarantonello ed. Academic Press, New York.
4. R. T. Rockafellar, *Characterization of subdifferentials of convex functions*, Pacific J. Math. **17** (1966), 497–510. MR **33** #1769.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, ILLINOIS 60637