

This article was downloaded by: [Cochin University of Science & Technology]

On: 24 July 2014, At: 22:12

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Communications in Statistics - Theory and Methods

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Ista20>

Some Properties of Weighted Distributions in the Context of Repairable Systems

S. M. Sunoj^a & S. S. Maya^a

^a Department of Statistics, Cochin University of Science and Technology, Cochin, Kerala, India

Published online: 19 Aug 2006.

To cite this article: S. M. Sunoj & S. S. Maya (2006) Some Properties of Weighted Distributions in the Context of Repairable Systems, Communications in Statistics - Theory and Methods, 35:2, 223-228

To link to this article: <http://dx.doi.org/10.1080/03610920500439968>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Distribution Theory

Some Properties of Weighted Distributions in the Context of Repairable Systems

S. M. SUNOJ AND S. S. MAYA

Department of Statistics, Cochin University of Science and Technology,
Cochin, Kerala, India

In this article we introduce some structural relationships between weighted and original variables in the context of maintainability function and reversed repair rate. Furthermore, we prove some characterization theorems for specific models such as power, exponential, Pareto II, beta, and Pearson system of distributions using the relationships between the original and weighted random variables.

Keywords Maintainability function; Reversed hazard rate; Weighted distributions.

Mathematics Subject Classification 62E10; 62N05.

1. Introduction

The concept of weighted distributions was introduced by Rao (1965) in connection with modeling statistical data, in situations where the usual practice of employing standard distributions for the purpose was not found appropriate. A formal definition of a weighted distribution is obtained by considering a probability space $(\Omega, \mathfrak{F}, P)$ and a random variable (rv) $X : \Omega \rightarrow H$, where $H = (a, b)$ is an interval of the real line. In the continuous case, if $f(x)$ is the probability density function (pdf) of X and $w(\cdot)$ a non negative function satisfying $\mu_w = E(w(X)) < \infty$, then the rv Y with pdf

$$f_w(x) = \frac{w(x)f(x)}{\mu_w}, \quad a < x < b \quad (1.1)$$

is said to have weighted distribution, corresponding to the distribution of X . The definition in the discrete case is analogs. The concept of weighted distribution has been employed in various practical problems such as analysis of family size, study of

Received September 15, 2004; Revised and Accepted July 7, 2005

Address correspondence to S. M. Sunoj, Department of Statistics, Cochin University of Science and Technology, Cochin, 682 022 Kerala, India; E-mail: smsunoj@cusat.ac.in

albinism, human heredity, aerial survey and visibility bias, line transcend sampling, renewal theory, cell cycle analysis and pulse labeling, efficacy of early screening for disease, etiological studies, statistical ecology, and reliability modeling. For detailed investigation of research in this area, we refer to Patil and Rao (1977), Gupta and Kirmani (1990), Navarro et al. (2001), Nair and Sunoj (2003), and Pakes et al. (2003).

Motivated by the applicability of weighted distributions in various fields, Gupta and Keating (1986) found it worthwhile to investigate the structural relationships between the distributions of X and Y in the context of reliability. Later, Gupta and Kirmani (1990) gave a detailed account of the survey of literature and extended the study of weighted distributions in the context of reliability and life testing. They considered the reliability measures useful in modeling life length studies such as reliability function, failure rate, and mean residual life function (MRLF) of the rv Y for comparison. The major relationships established by them are

$$S_w(x) = \frac{E(w(X) | X > x)}{\mu_w} S(x) \quad (1.2)$$

$$h_w(x) = \frac{w(x)}{E(w(X) | X > x)} h(x) \quad (1.3)$$

$$r_w(x) = \frac{r(x)}{w(x) + A(x)} \int_x^b \frac{w(t) + A(t)}{r(t)} \exp\left(-\int_x^t \frac{du}{r(u)}\right) dt \quad (1.4)$$

where $S(x)$, $h(x)$, and $r(x)$ are survival function, failure rate, and MRLF of X , respectively, defined by $S(x) = P(X > x)$, $h(x) = \frac{f(x)}{S(x)}$, and $r(x) = E(X - x | X > x) = \frac{1}{S(x)} \int_x^\infty S(t) dt$, and $S_w(x)$, $h_w(x)$, and $r_w(x)$ are the corresponding functions for Y with $A(x) = E(w(X) - w(x) | X > x)$, the mean residual weight. When $w(x) = x$ and $a > 0$, the weighted distribution (1.1) then termed as the length biased distribution denoted by $f^*(x)$.

Even if an extensive work has been carried out on weighted distributions using various reliability measures such as failure rate, MRLF, and vitality function in univariate and multivariate set up, very little has been explored in the context of repair time models. Accordingly, in the present article, we obtain some structural relationships between the original and weighted random variables and prove characterization theorems arising out of it for models such as power, exponential, Pareto II, beta, and Pearson system of distributions.

2. Maintainability and Reversed Repair Rate

Maintainability of a system provides a measure of the reparability of the system, when it fails or it is the probability of repairing a failed component/system in a specified period of time (see Rao, 1992). Various probability distributions may be used to present an item's repair time data. Once the repair time distributions are identified, the corresponding maintainability function may be obtained. The maintainability functions are used to predict the probability that a repair, beginning at time $x = 0$, will be accomplished in a time x . Therefore the maintainability function, $F(x)$, for any distribution is given by $F(x) = P(X \leq x)$, where X is the repair time. When X represents the repair time, the usual distribution function is termed as the maintainability function as it gives the probability that required

maintenance will be successfully completed in a given time period. Then the reversed repair rate is defined as

$$\lambda(x) = \frac{f(x)}{F(x)} \tag{2.1}$$

where $f(x)$ is the pdf of the repair time and $F(x)$ is the maintainability function, which is absolutely continuous. The interval of support of F being (a, b) with $a = \inf\{x : F(x) > 0\}$ and $b = \sup\{x : F(x) < 1\}$. When X represents the repair time of a component/system, the probability that it is repaired during the time $(x - \varepsilon, x)$ (ε is a small positive number) is approximately equal to $\varepsilon\lambda(x)$, whereas when X represents the lifetime, $\lambda(x)$ then termed as reversed hazard rate (RHR) (see for example, Block et al., 1998; Shaked and Santhikumar, 1994). In survival analysis, RHR has been found to be important for the estimation of survival function in the presence of left censored observations (see Anderson et al., 1993; Kalbfleish and Lawless, 1989). Let

$$m(x) = E(w(X) | X \leq x) = \frac{1}{F(x)} \int_a^x w(t)f(t)dt. \tag{2.2}$$

It is well known that $\lambda(x)$ and $m(x)$ uniquely determine $F(x)$ by

$$F(x) = \exp\left(-\int_x^b \lambda(t)dt\right) = \exp\left(-\int_x^b \frac{dm(t)}{w(t) - m(t)}\right). \tag{2.3}$$

See expression (1.2) in Navarro and Ruiz (1996) and Remark 2.2 in Navarro et al. (1998), respectively.

Recently, Nair et al. (2005) proved some characterization theorems using a relationship between RHR and conditional expectation (2.3). Further, Nair and Asha (2004) gave a review of the literature on RHR and developed certain identities connecting the distribution function, density function, and survival function in terms of RHR and the hazard rate.

We now study the structure of the weighted random variable Y in comparison to the original variable X .

$$F_w(x) = P(Y \leq x) = \int_a^x f_w(t)dt = \frac{m(x)}{\mu_w} F(x) \tag{2.4}$$

where $\mu_w = Ew(X) < \infty$. The corresponding reversed repair rate becomes

$$\lambda_w(x) = \frac{f_w(x)}{F_w(x)} = \frac{w(x)}{m(x)} \lambda(x). \tag{2.5}$$

3. Characterizations

In this section we prove some theorems useful to compare the original and weighted rv and also prove characterization theorems for specific models such as power, exponential, Pareto II, beta, and Pearson system of distributions using the relationships between the original and weighted rv.

- Theorem 3.1.** (a) $X \underset{lr}{\leq} Y(\geq)$ if, and only if, w is non decreasing (non increasing).
 (b) $X \underset{rfr}{\leq} Y(\geq)$ if, and only if, m is non decreasing (non increasing).
 (c) $X \underset{st}{\leq} Y(\geq)$ if, and only if, $\mu_w \leq m(x)(\geq)$.

Here $\underset{lr}{\leq}$, $\underset{rfr}{\leq}$, and $\underset{st}{\leq}$ denotes the likelihood ratio, the reversed failure rate, and the stochastic orders. It is well known (see p. 402 in Navarro et al., 1997) that

$$X \underset{lr}{\leq} Y \Rightarrow X \underset{rfr}{\leq} Y \Rightarrow X \underset{st}{\leq} Y.$$

Results (a) and (c) were given by Gupta and Kirmani (1990). Result (b) can be obtained immediately from (2.4) and the definition of the reversed hazard rate order.

Theorem 3.2. If $\alpha(x) = \frac{F_w(x)}{F(x)}$ and $\mu_w = Ew(X)$, then

$$F(x) = \exp\left(-\int_x^b \frac{\alpha'(t)}{\frac{w(t)}{\mu_w} - \alpha(t)} dt\right). \quad (3.1)$$

Theorem 3.3. If $\beta(x) = \frac{\lambda(x)}{\lambda_w(x)}$, then

$$F(x) = \exp\left(-\int_x^b \frac{\beta'(t)w(t) + \beta(t)w'(t)}{(1 - \beta(t))w(t)} dt\right). \quad (3.2)$$

The proofs of Theorems 3.2 and 3.3 are similar to that of Theorems 1 and 2 in Navarro et al. (2001).

Theorem 3.4. Let $A^*(x) = E(x - X | X \leq x)$. If X is increasing reversed repair rate and $\frac{A^*(x)}{x}$ is non decreasing, then Y is also increasing reversed repair rate.

Proof. X is increasing reversed repair rate implies

$$\lambda(x_1) \leq \lambda(x_2) \quad \text{for all } x_1 \leq x_2,$$

$$\text{i.e., } \left(\frac{x_1 - A^*(x_1)}{x_1}\right)\lambda^*(x_1) \leq \left(\frac{x_2 - A^*(x_2)}{x_2}\right)\lambda^*(x_2) \quad \text{for all } x_1 \leq x_2$$

$$\Rightarrow \lambda^*(x_1) \leq \frac{\left(1 - \frac{A^*(x_2)}{x_2}\right)}{\left(1 - \frac{A^*(x_1)}{x_1}\right)} \lambda^*(x_2)$$

$$\Rightarrow \lambda^*(x_1) \leq \lambda^*(x_2) \quad \text{for all } x_1 \leq x_2. \quad \square$$

Theorem 3.5. The ratio of the relationship

$$\frac{F_w(x)}{F(x)} = 1 - x(1 + Cx)\lambda(x) \quad (3.3)$$

if and only if X has Pareto II distribution with

$$F(x) = 1 - (1 + ax)^{-c}; \quad x > 0, \quad a, c > 0 \quad (3.4)$$

for $C > 0$, exponential distribution with

$$F(x) = 1 - e^{-ax}; \quad x > 0, \quad a > 0 \tag{3.5}$$

for $C = 0$, or beta distribution with

$$F(x) = 1 - (1 - ax)^c; \quad 0 < x < \frac{1}{a}, \quad c > 0 \tag{3.6}$$

for $C < 0$.

The proof can be obtained from Theorem 3 in Ruiz and Navarro (1994).

Further, Table 1 provides some characterization theorems for power and beta distributions based on their functional forms of $\alpha(x)$ and $\beta(x)$.

We now prove a characterization theorem that provide the relationships between reversed repair rate and right truncated moments of the original and weighted rv for the Pearson system of distributions.

Let X be an absolutely continuous rv with pdf $f(x)$, the distribution of X is said to be a member of the Pearson System of distributions if $f(x)$ satisfies the differential equation

$$\frac{f'(x)}{f(x)} = \frac{-(x + a)}{b_0 + b_1x + b_2x^2} \tag{3.7}$$

where a, b_0, b_1 , and b_2 are real valued. The shape of the distribution depends on the values of the parameters a, b_0, b_1 , and b_2 .

In a study on Pearson system of distributions, Sankaran and Nair (1993) derived the conditions under which Pearson and Ord families are form-invariant with respect to the length-biased sampling. According to them, the members of the Pearson system satisfying the differential equation (3.7) with $b_2 \neq 1$, have the same type of distributions for Y if and only if $b_0 = 0$. Later, Asadi (1998) extended this result to size-biased sampling of order α (i.e., $w(x) = x^\alpha$).

Theorem 3.6. *Let X be a non negative, non degenerate rv with density function $f(x)$ and suppose that $w(x) = x^\alpha$. Then X is a member of Pearson system of distributions of the form (3.7) with $b_0 = 0$ and $\lim_{x \rightarrow a} (b_1x + b_2x^2)f(x) = 0$ if and only if*

$$\frac{\lambda_w(x)}{\lambda(x)} = K \frac{(\mu_\alpha - m_\alpha(x))}{(\mu^* - m^*(x))} \tag{3.8}$$

where $K = \frac{1-(\alpha+2)b_2}{1-2b_2}$, $1 - 2b_2 \neq 0$, $\mu_\alpha = E(X^\alpha)$, and $m_\alpha(x) = E(X^\alpha | X \leq x)$.

Table 1

Different functional forms of $\alpha(x)$ and $\beta(x)$			
$w(x) = x$	$F(x)$	$\alpha(x)$	$\beta(x)$
power	$(\frac{x}{p})^d$	Cx	C
beta	$(px - q)^d$	$A + Bx$	$\frac{Ax+B}{x}$

The proof of Theorem 3.6 is similar to the analogous reversed result given in Asadi (1998).

Acknowledgments

The authors wish to thank the referees and the editor for the very useful and constructive comments on the earlier version of this article, which led to a substantial improvement.

References

- Anderson, P. K., Borgan, O., Gill, R. D., Keiding, N. (1993). *Statistical Methods Based on Counting Processes*. New York: Springer Verlag.
- Asadi, M. (1998). Characterization of the Pearson system of distributions based on reliability measures. *Statist. Pap.* 39:347–360.
- Block, H. W., Savits, T. H., Singh, H. (1998). The reversed hazard rate function. *Probab. Eng. Inform. Sci.* 12:69–90.
- Gupta, R. C., Keating, J. P. (1986). Relations for reliability measures under length biased sampling. *Scand. J. Statist.* 13:49–56.
- Gupta, R. C., Kirmani, S. N. U. A. (1990). The role of weighted distributions in stochastic modeling. *Commun. Statist. Theor. Meth.* 19(9):3147–3162.
- Kalbfleish, J. D., Lawless, J. F. (1989). Inference based on retrospective ascertainment: an analysis of the data on transfusion-related AIDS. *J. Amer. Statist. Assoc.* 84:360–372.
- Nair, N. U., Sunoj, S. M. (2003). Form-invariant bivariate weighted models. *Statistics* 37(3):259–269.
- Nair, N. U., Asha, G. (2004). Characterizations using failure and reversed failure rates. *J. Ind. Soc. Prob. Statist.* 8:45–56.
- Nair, N. U., Sankaran, P. G., Asha, G. (2005). Characterizations of distributions using reliability concepts. *J. Appl. Statist. Sci.* (to appear).
- Navarro, J., Ruiz, J. M. (1996). Failure rate functions for doubly truncated random variables. *IEEE Trans. Reliabil.* 45:685–690.
- Navarro, J., Belzunce, F., Ruiz, J. M. (1997). New stochastic orders based on double truncation. *Probab. Eng. Inform. Sci.* 11:395–402.
- Navarro, J., Ruiz, J. M., Zoroa, N. (1998). A unified approach to characterization problem using conditional expectations. *J. Statist. Plann. Infer.* 69(2):193–207.
- Navarro, J., Del Aguila, Y., Ruiz, J. M. (2001). Characterizations through reliability measures from weighted distributions. *Statistical Pap.* 42:395–402.
- Pakes, A. G., Navarro, J., Ruiz, J. M., Del Aguila, Y. (2003). Characterizations using weighted distributions. *J. Statist. Plann. Infer.* 116:389–420.
- Patil, G. P., Rao, C. R. (1977). The weighted distribution: a survey of their applications. In: Krishnaiah, P. R., ed. *Applications of Statistics*. Amsterdam: North Holland Publishing Company, pp. 383–405.
- Rao, C. R. (1965). On discrete distributions arising out of methods of ascertainment. In: Patil, G. P., ed. *Classical and Contagious Discrete Distributions*. Calcutta: Pergunon Press and Statistical Publishing Society. Also reprinted in *Sankhya A* 27:311–324.
- Rao, S. S. (1992). *Reliability-Based Design*. New York: McGraw Hill, Inc.
- Ruiz, J. M., Navarro, J. (1994). Characterizations of distributions by relationships between failure rate and mean residual life. *IEEE Trans. Reliab.* 43(4):640–644.
- Sankaran, P. G., Nair, N. U. (1993). On form invariant length biased models from Pearson family. *J. Ind. Statist. Assoc.* 31:85–89.
- Shaked, M., Santhikumar, J. G. (1994). *Stochastic Orders and Their Applications*. New York: Academic Press.