

Some Sharp Restriction Theorems for Homogeneous Manifolds

*Laura De Carli and Alex Iosevich**

Communicated by John J. Benedetto

1. Introduction

Let M denote a submanifold of \mathbb{R}^{n+l} of codimension l . Let \mathcal{R} denote a restriction operator

$$\mathcal{R}f(\eta) = \int e^{-i\langle x, \eta \rangle} f(x) dx, \quad \eta \in M, \quad f \in \mathcal{S}(\mathbb{R}^{n+l}). \quad (1.1)$$

We wish to find an optimal range of exponents p such that

$$\|\mathcal{R}f\|_{L^2(M, d\sigma)} \leq C_p \|f\|_{L^p(\mathbb{R}^{n+l})}, \quad (1.2)$$

where $d\sigma$ is a compactly supported measure on M .

When M is a codimension one surface in \mathbb{R}^{n+1} with non-vanishing Gaussian curvature, the estimate (1.2) is well understood. A celebrated result due to Stein and Thomas says that if M has non-vanishing Gaussian curvature, then the estimate (1.2) holds with $p = \frac{2(n+2)}{n+4}$. In higher dimensions, the situation is more complicated. When M is a codimension two surface in \mathbb{R}^{n+2} satisfying a non-degeneracy assumption, the estimate (1.2) holds with $p = \frac{2(n+4)}{n+8}$. (See [1] and Theorem 7 in Section 3 below). However, a sharp necessary and sufficient condition is not currently available.

It should also be noted that even in codimension one, the more general (L^p, L^q) estimates for the restriction operator are not fully understood, except in the dimension two. (See [14] for a detailed discussion). We shall not address this issue here.

The purpose of this article is to establish the estimate (1.2) in the case when

$$M = \{(x, x_{k+1}, \dots, x_{k+l}) : x_{k+1} = \phi_1(x), \dots, x_{k+l} = \phi_l(x)\},$$

where $\phi_j \in C^\infty(\mathbb{R}^n \setminus \{0\})$ is homogeneous of degree $m_j \geq 1$.

Acknowledgements and Notes. *This work was supported in part by NSF Grant DMS97 0682S.

Let $\mathcal{F}[d\sigma]$ denote the Fourier transform of $d\sigma$. By a theorem of Greenleaf (see [3]), the inequality (1.2) holds for $p = \frac{2(l+\gamma)}{2l+\gamma}$ if

$$|\mathcal{F}[d\sigma](R\zeta)| \leq C(1 + R)^{-\gamma}, \quad \zeta \in S^{n+1}. \tag{1.3}$$

We shall see below that isotropic Fourier transform estimates do not yield the sharp restriction theorem in codimension two or higher (see, e.g., [1]). It should be noted that even in codimension one, it is not known whether the exponent given by Greenleaf’s theorem is sharp.

The best possible isotropic rate of decay of $\mathcal{F}[d\sigma]$ for the homogeneous manifold M defined above is $-\frac{n}{\max_j\{m_j\}}$. An application of Greenleaf’s theorem yields the estimate (1.2) with

$$p_0 = \frac{2(n + lm)}{n + 2lm}, \tag{1.4}$$

where $m = \max_j\{m_j\}$.

However, the following homogeneity argument due to Knapp suggests that the optimal exponent for the estimate (1.2) is

$$p_0 = \frac{2(m_1 + \dots + m_l + n)}{n + 2(m_1 + \dots + m_l)}. \tag{1.5}$$

Indeed, let \mathcal{R} denote the restriction operator defined above. Let $\hat{f}_\delta = h$, where h is the characteristic function of a rectangle in \mathbb{R}^{n+l} with sides of lengths $(1, 1, \dots, 1, C, \dots, C)$, C large.

Then

$$\|f_\delta\|_p \approx \delta^{(1-1/p)(n+m_1+\dots+m_l)} \quad \text{and} \quad \|\mathcal{R}f_\delta\|_2 \approx \delta^{n/2}. \tag{1.6}$$

Hence, (1.2) can only hold if $p \leq \frac{2(n+m_1+\dots+m_l)}{n+2(m_1+\dots+m_l)}$.

We will establish the estimate (1.2) for a homogeneous manifold M , with the exponent p_0 given by Knapp’s homogeneity argument, under a variety of conditions on the level sets of the graphing functions $\phi_1, \phi_2, \dots, \phi_l$.

Our main results are the following. The first result gives us a good description of L^2 restriction theorems for two-dimensional submanifolds in codimension 2 given as graphs of homogeneous polynomials. (Please see Definition 1 and 2 below for the precise description of finite type. Please see Definition 3 below for the description of the order of vanishing along a line.)

Theorem 1.

(See Theorem 12 in Section 3). Let

$$S = \{(x_1, x_2, x_3, x_4) : x_3 = \phi_1(x_1, x_2), x_4 = \phi_2(x_1, x_2),\} \tag{1.7}$$

where $\phi_1, \phi_2 \in C^\infty(\mathbb{R}^2)$ are homogeneous polynomials of degree m_1 and m_2 , respectively ($m_1 \geq m_2 \geq 2$). Suppose that there exists a non-zero constant c such that $\phi_1(x)|_{\{x:\phi_2(x)=1\}} = c$. Let $Z_0 = \{(x_1, x_2) : \phi_2(x_1, x_2) = 0\}$, $Z_1 = \{(x_1, x_2) : \nabla\phi_2(x_1, x_2) = (0, 0)\}$, and $Z_2 = \{(x_1, x_2) : H\phi_2(x_1, x_2) = 0\}$, where $H\phi_2$ denotes the determinant of the Hessian matrix of ϕ_2 . Suppose that the curve $\{x : \phi_2(x) = 1\}$ is of finite type m at each point of $Z_2 \cap \{x : \phi_2(x) = 1\}$ and that ϕ_2 vanishes of order $\leq M$ along the lines contained in $Z_0 \cup Z_1$. Then (1.2) holds for every $p < p_0$, where p_0 is the sharp exponent

$$\frac{m_1 + m_2 + 2}{m_1 + m_2 + 1}, \tag{1.8}$$

if $m_1 + m_2 \geq \max\left\{2M\left(1 + \frac{m_1}{m_2}\right), 2m\right\}$. Furthermore (1.2) holds with

$$(1) \quad p < \max\left\{\frac{2(1+M(1+\frac{m_1}{m_2}))}{1+2M(1+\frac{m_1}{m_2})}, \frac{2(m+1)}{2m+1}\right\} \text{ if } \min\{2M(1 + \frac{m_1}{m_2}), 2m\} \leq m_1 + m_2 < \max\{2M(1 + \frac{m_1}{m_2}), 2m\},$$

$$(2) p < \min \left\{ \frac{2(1+M(1+\frac{m_1}{m_2}))}{1+2M(1+\frac{m_1}{m_2})}, \frac{2(m+1)}{2m+1} \right\} \text{ if } 4 \leq m_1 + m_2 < \min\{2M(1 + \frac{m_1}{m_2}), 2m\}.$$

The following result answers a question posed by Fulvio Ricci about the restriction theorems for manifolds given as graphs of quadratic monomials. The proof relies on the precise asymptotics of the Fourier transforms of certain distributions obtained by Shintani (see [12], and Lemma 7 below). It has been brought to our attention that this result is implied by a theorem announced in the Bulletin of the AMS by Gerd Mockenhaupt. We enclose the proof for the reader's convenience, and to motivate the related results proved in this article. (See, e.g., Theorem 3 below).

Theorem 2.

(See Theorem 13 below). Let $S = \{(x, x_{n+1}, \dots, x_{n+l}) \in \mathbb{R}^{n+l} : x_{n+1} = \phi_1(x), \dots, x_{n+l} = \phi_l(x)\}$, where $l = \frac{n(n+1)}{2}$, and the ϕ_j denote the distinct monomials of degree 2. Then the estimate (1.2) holds with the sharp exponent $p_0 = \frac{2(n+2)}{2n+3}$.

The following result generalizes Theorem 2 to manifolds given as joint graphs of smooth functions of higher order of homogeneity. The proof relies on the non-isotropic decay estimates for the associated Fourier transform of the surface carried measure. The observation that the non-isotropic decay estimates are useful to obtain sharp restriction theorems in codimension > 1 is not new. See, for example, the work of Christ [1] and Prestini [11].

Theorem 3.

(See Theorem 14 below). Let S denote a compact piece of the manifold $\{(x, x_{n+1}, \dots, x_{n+l}) \in \mathbb{R}^{n+l} : x_{n+1} = \phi_1(x), \dots, x_{n+l} = \phi_l(x)\}$, where $\phi_j \in C^\infty(\mathbb{R}^n/0)$ is homogeneous of degree $m \geq 2n$. Suppose that no linear combination of the ϕ_j s vanishes on a subset of positive measure of S^{n-1} . Let $\Phi(x) = (\phi_1(x), \dots, \phi_l(x))$. Suppose that $\Phi(\omega) \neq (0, \dots, 0)$, $\omega \in S^{n-1}$. Then the estimate (1.2) holds with the sharp exponent

$$p_0 = \frac{2(n + lm)}{n + 2lm} . \tag{1.9}$$

Many of our results are based on the non-isotropic decay estimates for the associated Fourier transform of the surface carried measure. A sample result is the following.

Theorem 4.

(See Theorem 6 below) Let $S = \{(x, x_{n+1}, \dots, x_{n+l}) \in \mathbb{R}^{n+l} : x_{n+1} = \phi_1(x), \dots, x_{n+l} = \phi_l(x)\}$, where each $\phi_j \in C^\infty(\mathbb{R}^n/0)$ is homogeneous of degree $m \geq 2n$. Let $\Phi(x) = (\phi_1(x), \dots, \phi_l(x))$. Let $\Phi_\lambda(x) = \langle \Phi(x), \lambda \rangle$. Let

$$F(\xi, \lambda) = \int e^{i(x,\xi)+\lambda_1\phi_1(x)+\dots+\lambda_l\phi_l(x)} \psi(x) dx , \tag{1.10}$$

where ψ is a smooth cutoff function. Then

$$|F(\xi, \lambda)| \leq C \int_{S^{n-1}} |\Phi_\lambda(\omega)|^{-\frac{n}{m}} d\omega . \tag{1.11}$$

This article is organized as follows. In Section 2 we will prove some estimates related to the decay of the Fourier transform of the surface carried measure. In particular, we will estimate the decay of the Fourier transform of the surface carried measure, in any codimension, in terms of the integrability of the multiplicative inverses of the graphing functions ϕ_j restricted to the unit sphere. Using this technique we shall also obtain an accurate non-isotropic estimate for the Fourier transform of the surface carried measure in the case when every graphing function has the same homogeneity.

In Section 3 we will apply the results of Section 2 along with the results of Christ [1], Prestini [11], and a variety of scaling arguments to obtain a sharp estimate (1.2) with the exponent p_0 given by (1.5).

In Section 4 we will use the non-isotropic estimates from Section 2 to study restriction theorems in the case when every graphing function is homogeneous of the same degree $m \geq 2$.

2. The Decay of the Fourier Transform of the Surface Carried Measure

We will need the following definitions. (See, e.g., [1, 3]).

Nonvanishing Gaussian curvature: Let Σ be a submanifold of \mathbb{R}^{N+1} of codimension 1 equipped with a smooth compactly supported measure $d\mu$. Let $J : \Sigma \rightarrow S^N$ be the usual Gauss map taking each point on Σ to the outward unit normal at that point. We say that Σ has everywhere nonvanishing Gaussian curvature if the differential of the Gauss map dJ is always nonsingular.

Strong curvature condition: Let S be a submanifold of \mathbb{R}^{n+l} of codimension l equipped with a smooth compactly supported measure $d\mu$. Suppose that S is a joint graph of smooth functions g_1, g_2, \dots, g_l , where $g_j : \mathbb{R}^n \rightarrow \mathbb{R}$. Let $\mathcal{N}_{x_0}(S)$ denote the l -dimensional space of normals to S at a point x_0 . We say that S satisfies the strong curvature condition (SCC) if for all $x_0 \in S$ in some neighborhood of the support of $d\mu$,

$$\det D^2(v_1g_1(x) + v_2g_2(x) + \dots + v_lg_l(x)) \neq 0, \quad \forall v \in \mathcal{N}_{x_0}, \quad (2.1)$$

where D^2 denotes the Hessian matrix.

N-curvature condition: Let $S \in \mathbb{R}^{n+l}$ be defined as above. We say that S satisfies the N -curvature condition if the rank of the Hessian matrix in (2.1) is greater than or equal to N everywhere.

Our main results are the following.

Theorem 5.

Let $S = \{(x, x_{n+1}, \dots, x_{n+l}) \in \mathbb{R}^{n+l} : x_{n+1} = \phi_1(x), \dots, x_{n+l} = \phi_l(x)\}$, where $\phi_j \in C^\infty(\mathbb{R}^n \setminus 0)$ is homogeneous of degree $m_j \geq 2$. Suppose that no ϕ_i is a constant multiple of any ϕ_j for $i \neq j$. Suppose that $(\phi_j)^{-1} \in L^{\rho_j}(S^{n-1})$, $0 < \rho_j \leq \frac{n}{m_j}$. Let μ denote the number of distinct m_j s. Suppose that $\frac{n}{m_j} \leq \frac{1}{\mu+1}$ for each j . Let $\lambda = (\lambda_1, \dots, \lambda_l)$, and define

$$F(\xi, \lambda) = \int e^{i((x, \xi) + \lambda_1\phi_1(x) + \dots + \lambda_l\phi_l(x))} \psi(x) dx, \quad (2.2)$$

where ψ is a smooth cutoff function.

Then

$$|F(\xi, \lambda)| \leq C(1 + |\xi| + |\lambda|)^{-\rho}, \quad (2.3)$$

where $\rho = \min_j \{\rho_j\}$.

Theorem 6.

Let $S = \{(x, x_{n+1}, \dots, x_{n+l}) \in \mathbb{R}^{n+l} : x_{n+1} = \phi_1(x), \dots, x_{n+l} = \phi_l(x)\}$, where each $\phi_j \in C^\infty(\mathbb{R}^n \setminus 0)$ is homogeneous of degree $m \geq 2$. Let $\Phi(x) = (\phi_1(x), \dots, \phi_l(x))$. Let $\Phi_\lambda(x) = \langle \Phi(x), \lambda \rangle$. Let $F(\xi, \lambda)$ be defined as in (2.2). Then

$$|F(\xi, \lambda)| \leq C \int_{S^{n-1}} |\Phi_\lambda(\omega)|^{-\frac{n}{m}} d\omega. \quad (2.4)$$

Lemma 1.

Let $S \in \mathbb{R}^{n+l}$ and $F(\xi, \lambda)$ be defined as above. Let

$$F_0(\xi, \lambda) = \int e^{i((x, \xi) + \lambda_1\phi_1(x) + \dots + \lambda_l\phi_l(x))} \psi_0(x) dx, \quad (2.5)$$

where ψ_0 is a smooth cutoff function supported away from the origin, and each $\phi_j \in C^\infty(\mathbb{R}^n \setminus \{0\})$ is homogeneous of degree $m_j \geq 2$. Suppose that

$$|F_0(\xi, \lambda)| \leq C(1 + |\xi| + |\lambda|)^{-\gamma}, \quad \gamma > 0. \tag{2.6}$$

Then

$$|F(\xi, \lambda)| \leq C(1 + |\xi| + |\lambda|)^{-\frac{n}{\max_j \{m_j\}}}, \tag{2.7}$$

if $\gamma \max_j \{m_j\} \geq n$.

Lemma 2.

Let $S \in \mathbb{R}^{n+l}$ be defined as above. Suppose that $S|_{\{x_{n+j_1}=1, x_{n+j_2}=1, \dots, x_{n+j_s}=1\}}$, $1 \leq s < l$, is an $(n - s)$ -dimensional submanifold of codimension l , of the hyperplane

$$\{(x, x_{n+1}, \dots, x_{n+l}) : x_{n+j_1} = 1, \dots, x_{n+j_s} = 1\},$$

satisfying the N -curvature condition. Let $F_0(\xi, \lambda)$ be defined as in (2.5). Then

$$|F_0(\xi, \lambda)| \leq C(1 + |\xi| + |\lambda|)^{-\frac{N}{2}}. \tag{2.8}$$

Lemma 3.

Let $S \in \mathbb{R}^{n+l}$ be defined as above with $l = 2$. Suppose that there exists a constant c , such that $\phi_1|_{\{x:\phi_2(x)=1\}} \equiv c$. Suppose that $m_1 \neq m_2$. Suppose that $H\phi_1$, the Hessian matrix of ϕ_1 , has rank ≥ 2 away from the origin. Suppose that $\phi_1(\omega) \neq 0$, $\omega \in S^{n-1}$. Let $F_0(\xi, \lambda)$ be defined as in (2.5). Then

$$|F_0(\xi, \lambda)| \leq C(1 + |\xi| + |\lambda|)^{-\frac{1}{2}}. \tag{2.9}$$

Lemma 4.

Let $S \in \mathbb{R}^{n+l}$ be defined as in Lemma 3, with $n = l = 2$. Then the assumptions of Theorem 8 (see Section 3) are satisfied at every $x \neq (0, 0)$ iff $\phi_2(\omega) \neq 0$, $\omega \in S^1$, and the level set $\{x : \phi_2 = 1\}$ has non-vanishing Gaussian curvature.

Proof of Theorem 5. We shall prove the theorem under the assumption that all the m_j s are distinct. The general statement follows by combining the terms with the same homogeneity. Recall that

$$F(\xi, \lambda) = \int e^{i((x,\xi)+\lambda_1\phi_1(x)+\dots+\lambda_l\phi_l(x))} \psi(x) dx, \tag{2.10}$$

where without loss of generality ψ is radial, and each $\phi_j \in C^\infty(\mathbb{R}^n)$ is homogeneous of degree $m_j \geq 2$.

$$\text{Let } \Omega = \left\{ \omega \in S^{n-1} : |\phi_1(\omega)| > \frac{1}{|\lambda_1|}, \dots, |\phi_l(\omega)| > \frac{1}{|\lambda_l|} \right\}.$$

Let

$$F(\xi, \lambda) = \int_{\frac{x}{|x|} \in S^{n-1} \setminus \Omega} + \int_{\frac{x}{|x|} \in \Omega} = F_{S^{n-1} \setminus \Omega} + F_\Omega. \tag{2.11}$$

Taking absolute values inside the integral we see that

$$|F_{S^{n-1} \setminus \Omega}(\xi, \lambda)| \leq C \left| S^{n-1} \setminus \Omega \right|, \tag{2.12}$$

which by Chebyshev's inequality is bounded by

$$C \max_j \left\{ |\lambda_j|^{-\rho_j} \int_{S^{n-1}} |\phi_j(\omega)|^{-\rho_j} d\omega \right\} \leq C |\lambda|^{-\rho}, \tag{2.13}$$

for $|\lambda|$ large.

Let $\psi_0 \in C_0^\infty(\mathbb{R})$ be supported in the interval $[\frac{1}{2}, 4]$, such that $\psi_0 \equiv 1$ inside $[1, 2]$, and

$$\sum_{k=0}^{\infty} \psi_0(2^k s) \equiv 1. \quad (2.14)$$

Let

$$F_\Omega^k(\xi, \lambda) = \int \int e^{i(r(\omega, \xi) + \lambda_1 r^{m_1} \phi_1(\omega) + \dots + \lambda_l r^{m_l} \phi_l(\omega))} \psi_0(2^k r) r^{n-1} dr d\omega. \quad (2.15)$$

A change of variables sending $r \rightarrow 2^{-k} r$ shows that

$$F_\Omega^k(\xi, \lambda) = 2^{-nk} F_\Omega^0(2^{-k} \xi, 2^{-m_1 k} \lambda_1, \dots, 2^{-m_l k} \lambda_l). \quad (2.16)$$

Let $\mathcal{A} = \{1, 2, \dots, l\}$.

We must estimate

$$\sum_{k=0}^{\infty} 2^{-nk} \left| F_\Omega^0(2^{-k} \xi, 2^{-m_1 k} \lambda_1, \dots, 2^{-m_l k} \lambda_l) \right| = \sum_{\mathcal{B} \subset \mathcal{A}} I_{\mathcal{B}}, \quad (2.17)$$

where

$$\begin{aligned} I_{\mathcal{B}} &= \sum_{\{\lambda_j \phi_j(\omega) \leq 2^{m_j k}, j \in \mathcal{B}, \lambda_j \phi_j(\omega) > 2^{m_j k}, j \notin \mathcal{B}\}} 2^{-nk} \left| F_\Omega^0(2^{-k} \xi, 2^{-m_1 k} \lambda_1, \dots, 2^{-m_l k} \lambda_l) \right| \\ &= \sum_{S_{\{\mathcal{B}, \leq\}} \cap S_{\{\mathcal{B}, >\}}} 2^{-nk} \left| F_\Omega^0(2^{-k} \xi, 2^{-m_1 k} \lambda_1, \dots, 2^{-m_l k} \lambda_l) \right|. \end{aligned} \quad (2.18)$$

To estimate each $I_{\mathcal{B}}$ we shall need the fact that the curve $(r, r^{m_1}, \dots, r^{m_l})$ has non-vanishing curvature and torsion away from the origin, so long as all the m_j s are distinct. An elementary van der Corput type estimate shows that the Fourier transform of the measure carried by this curve decays of order $-\frac{1}{l+1}$. We shall also use the fact that F_Ω^0 is bounded. More precisely,

$$\begin{aligned} |I_{\mathcal{B}}| &\leq \int_{\Omega} \left(C \sum_{S_{\{\mathcal{B}, \leq\}}} 2^{-nk} \right) + \left(C \sum_{S_{\{\mathcal{B}, >\}}} 2^{-nk} \left(\sum_{j \in \mathcal{B}} 2^{-m_j k} |\phi_j(\omega) \lambda_j| \right)^{-\frac{1}{l+1}} \right) d\omega \\ &\leq \max_{j \in \mathcal{A}} |\lambda_j|^{-\frac{n}{m_j}} \int_{\Omega} |\phi_j(\omega)|^{-\frac{n}{m_j}} d\omega, \end{aligned} \quad (2.19)$$

provided that $\frac{n}{m_j} \leq \frac{1}{l+1}$, for each $j \in \mathcal{A}$.

Moreover,

$$\begin{aligned} |\lambda_j|^{-\frac{n}{m_j}} \int_{\Omega} |\phi_j(\omega)|^{-\frac{n}{m_j}} d\omega &= |\lambda_j|^{-\rho_j} \int_{\Omega} |\phi_j(\omega)|^{-\rho_j} |\phi_j(\omega) \lambda_j|^{\rho_j - \frac{n}{m_j}} d\omega \\ &\leq |\lambda_j|^{-\rho_j} \int_{S^{n-1}} |\phi_j(\omega)|^{-\rho_j} d\omega, \end{aligned} \quad (2.20)$$

since $\rho_j \leq \frac{n}{m_j}$.

Hence, the expression (2.19) is bounded by

$$\max_{j \in \mathcal{A}} |\lambda_j|^{-\rho_j} \int_{S^{n-1}} |\phi_j(\omega)|^{-\rho_j} d\omega \leq C(|\lambda|)^{-\rho}, \quad (2.21)$$

for $|\lambda|$ large.

This completes the proof if $|\xi| \leq C|\lambda|$. However, if this is not the case, the gradient of the phase function $\langle x, \xi \rangle + \lambda_1 \phi_1(x) + \dots + \lambda_l \phi_l(x)$ is bounded away from zero, and an integration by parts argument (see [14, p.364]) shows that $|F(\xi, \lambda)| \leq C_N 1 + |\xi| + |\lambda|^{-N}$, for any $N > 0$. \square

Proof of Theorem 6. Let $\lambda = t\nu = t(\nu_1, \dots, \nu_l)$. Let $\Phi_\nu(x) = \langle \Phi(x), \nu \rangle$. We rewrite $F(\xi, \lambda)$ in the form

$$\int e^{i(\langle x, \xi \rangle + t\Phi_\nu(x))} \psi(x) dx, \tag{2.22}$$

where ψ is a smooth cutoff function. Let $\Omega = \{\omega \in S^{n-1} : |\Phi_\nu(\omega)| > \frac{1}{t}\}$. Let

$$F(\xi, \lambda) = \int_{\frac{x}{|x|} \in S^{n-1} \setminus \Omega} + \int_{\frac{x}{|x|} \in \Omega} = F_{S^{n-1} \setminus \Omega} + F_\Omega. \tag{2.23}$$

Taking the absolute values inside the integral, we see that

$$|F_{S^{n-1} \setminus \Omega}(\xi, \lambda)| \leq C |S^{n-1} \setminus \Omega| = C \left| \left\{ \omega \in S^{n-1} : \left| \Phi_\nu^{-1}(\omega) \right| > t \right\} \right|, \tag{2.24}$$

which by Chebyshev's inequality is bounded by

$$t^{-\frac{n}{m}} \int |\Phi_\nu(\omega)|^{-\frac{n}{m}} d\omega. \tag{2.25}$$

Let $\psi_0 \in C_0^\infty(\mathbb{R})$ be supported in the interval $[\frac{1}{2}, 4]$, such that $\psi_0 \equiv 1$ inside $[1, 2]$, and

$$\sum_{k=0}^\infty \psi_0(2^k s) \equiv 1. \tag{2.26}$$

Let

$$F_\Omega^k(\xi, \lambda) = \int \int e^{i(r\langle \omega, \xi \rangle + tr^m \Phi_\nu(\omega))} r^{n-1} \psi_0(2^k r) d\omega dr. \tag{2.27}$$

A change of variables sending $r \rightarrow 2^{-k}r$ shows that $F_\Omega^k(\xi, \lambda) = 2^{-nk} F_\Omega^0(2^{-k}\xi, 2^{-mk}\lambda)$. We must estimate

$$\sum_{k=0}^\infty 2^{-nk} \left| F_\Omega^0(2^{-k}\xi, 2^{-mk}\lambda) \right| = \sum_{\{t\Phi_\nu(\omega) \leq 2^{mk}\}} + \sum_{\{t\Phi_\nu(\omega) > 2^{mk}\}} = I + II. \tag{2.28}$$

To estimate II we shall use the fact that away from zero the Fourier transform of the measure supported on the curve (r, r^m) decays of order $-\frac{1}{2}$. To estimate I we shall just use the fact that F_Ω^0 is bounded. More precisely,

$$|I| \leq C \int_\Omega \sum_{\{t\Phi_\nu(\omega) \leq 2^{mk}\}} 2^{-nk} d\omega \leq Ct^{-\frac{n}{m}} \int |\Phi_\nu(\omega)|^{-\frac{n}{m}} d\omega, \tag{2.29}$$

and

$$|II| \leq C \int_\Omega \sum_{\{t\Phi_\nu(\omega) > 2^{mk}\}} 2^{-nk} \left| 2^{-mk} \Phi_\nu(\omega) t \right|^{-\frac{1}{2}} d\omega \leq Ct^{-\frac{n}{m}} \int_\Omega |\Phi_\nu(\omega)|^{-\frac{n}{m}} d\omega, \tag{2.30}$$

as long as $m \geq 2n$.

This completes the proof if $|\xi| \leq C|\lambda|$. However, if this is not the case, the gradient of the phase function $\langle x, \xi \rangle + \lambda_1 \phi_1(x) + \dots + \lambda_l \phi_l(x)$ is bounded away from the origin, and an integration

by parts argument (see [14, p.364]) shows that $|F(\xi, \lambda)| \leq C_N(1 + |\xi| + |\lambda|)^{-N}$, for any $N > 0$. \square

Proof of Lemma 1. Let

$$F_k(\xi, \lambda) = \int e^{i((x,\xi)+\lambda_1\phi_1(x)+\dots+\lambda_l\phi_l(x))} \psi_0(2^k x) dx . \tag{2.31}$$

A change of variables sending $x \rightarrow 2^{-k}x$ shows that

$$F_k(\xi, \lambda) = 2^{-nk} F_0\left(2^{-k}\xi, 2^{-m_1k}\lambda_1, \dots, 2^{-m_lk}\lambda_l\right) . \tag{2.32}$$

Let $\mathcal{A} = \{1, 2, \dots, l\}$. We must estimate

$$\sum_{k=0}^{\infty} 2^{-nk} \left| F_0\left(2^{-k}\xi, 2^{-m_1k}\lambda_1, \dots, 2^{-m_lk}\lambda_l\right) \right| = \sum_{B \subset \mathcal{A}} I_B , \tag{2.33}$$

where

$$\begin{aligned} I_B &= \sum_{\{|\lambda_j| \leq 2^{m_j k}, j \in B, |\lambda_j| > 2^{m_j k}, j \notin B\}} 2^{-nk} \left| F_0\left(2^{-k}\xi, 2^{-m_1k}\lambda_1, \dots, 2^{-m_lk}\lambda_l\right) \right| \\ &= \sum_{S_{\{B, \leq\}} \cap S_{\{B, >\}}} 2^{-nk} \left| F_0\left(2^{-k}\xi, 2^{-m_1k}\lambda_1, \dots, 2^{-m_lk}\lambda_l\right) \right| . \end{aligned} \tag{2.34}$$

Using the assumed decay of F_0 , and the fact that, in particular, F_0 is bounded, we get

$$|I_B| \leq C_1 \sum_{S_{\{B, \leq\}}} 2^{-nk} + C_2 \sum_{S_{\{B, >\}}} 2^{-nk} \left(\sum_{j \notin B} 2^{-m_j k} |\lambda_j| \right)^{-\gamma} \tag{2.35}$$

$$\leq C \max_{j \in \mathcal{A}} |\lambda_j|^{-\frac{n}{m_j}} \leq C |\lambda|^{-\frac{n}{\max_j \{m_j\}}} , \tag{2.36}$$

as long as $\max_j \{m_j\} \leq n\gamma$.

This completes the proof if $|\xi| \leq C|\lambda|$. However, if this is not the case, the gradient of the phase function $(x, \xi) + \lambda_1\phi_1(x) + \dots + \lambda_l\phi_l(x)$ is bounded away from the origin. An integration by parts argument (see [14, p.364]) shows that $|F(\xi, \lambda)| \leq C_N(1 + |\xi| + |\lambda|)^{-N}$, for any $N > 0$. \square

Proof of Lemma 2. Assume that $x_{n+j_1} = x_{n+1}, \dots, x_{n+j_s} = x_{n+s}$. In what follows we will denote $x \in \mathbb{R}^n$ by (x', x'') , with $x' \in \mathbb{R}^s, x'' \in \mathbb{R}^{n-s}$ and by $J_{x'}(f)$ [resp. $J_{x''}(f)$] and $D_{x'}^2 g$ (resp. $D_{x''}^2 g$) the Jacobian of a function $f(x', x'')$ and the Hessian matrix of a function $g(x', x'')$ computed with respect to the x' (resp. x'') variables. Let $\phi = (\phi_1, \dots, \phi_s)$.

Take P on the support of $d\sigma$ such that $\phi(P) = (1, 1, \dots, 1)$. Since $S_{\{x_{n+1}=1, \dots, x_{n+s}=1\}}$ is by assumption a submanifold of codimension l of the hyperplane $\{(x, x_{n+1}, \dots, x_{n+l}) : x_{n+1} = 1, \dots, x_{n+s} = 1\}$, the Jacobian of the function $\phi, J(\phi)$, has rank s at P .

There is no loss of generality if we assume that $J_{x'}\phi(P)$ is the identity in the space of $s \times s$ matrices, and that $J_{x''}\phi(P) = 0$

By the implicit function theorem there exists a smooth function $\psi(x'') : \mathbb{R}^{n-s} \rightarrow \mathbb{R}^s$ such that $\phi(\psi(x''), x'') \equiv 1$ in a neighborhood of P'' .

An application of the chain rule yields:

$$(1) \quad J\phi(\psi(x''), x'') = J_{x'}\phi(\psi(x''), x'') \times J\psi(x'') + J_{x''}\phi(\psi(x''), x'') \equiv 0, \text{ which implies that } J\psi(P'') = 0,$$

- (2) $\frac{\partial}{\partial x_{s+j}} J\psi(P'') + \frac{\partial}{\partial x_{s+j}} J_{x''}\phi(P) = 0$ for every $j \leq n + s$, and
 (3) $D^2(\phi_{s+j})(\psi(x''), x'')|_{x''=P''} = \sum_{k=1}^s \frac{\partial}{\partial x_k} \phi_{s+j}(\psi(x''), x'')|_{x''=P''} D^2(\psi_k)(P'')$
 $+ D_{x''}^2 \phi_{s+j}(\psi(x''), x'')$ for every $j \leq n - s$.

From (2) we have that, for every $k \leq s$, the Hessian matrix of ψ_k at P'' , $D^2\psi_k(P'')$, is $-D_{x''}^2\phi_k(P'')$, and from (3) that $D_{x''}^2\phi_{s+j}(P)$ can be written as a linear combination of the Hessian matrices of the functions ψ_k and of the Hessian matrix of $\phi_{s+j}(\psi(x''), x'')$ at P'' .

Let $\Phi = \langle x, \xi \rangle + \lambda_1\phi_1 + \dots + \lambda_l\phi_l$ denote the phase function of F as in (2.2). By the above remark, $D_{x''}^2\Phi(P)$ can be written as a linear combination of the Hessian matrices of the functions $\psi_k(x'')$ and the function $\phi_{s+j}(\psi(x''), x'')$ at P'' . Since $S_{\{|x_{n+j}=1, \dots, x_{n+s}=1\}}$ satisfies the N -curvature condition, the rank of every linear combination of the above matrices is N for every $\lambda \neq 0$. This shows that the rank of the Hessian matrix of Φ is $\geq N$ and hence that (2.7) holds. \square

Proof of Lemma 3. A theorem of Littman [9] says that if a surface in codimension one has at least k non-vanishing principal curvatures, then the Fourier transform of the surface carried measure decays of order $-\frac{k}{2}$. The proof of that theorem shows that F_0 has the required decay if the rank of the Hessian matrix of the phase function $\Phi(x) = \langle x, \xi \rangle + \lambda_1\phi_1(x) + \lambda_2\phi_2(x)$ is ≥ 1 for every x on the support of χ and for every $(\lambda_1, \lambda_2) \neq (0, 0)$.

We observe now that $\phi_1(x) \equiv c\phi_2^{\frac{m_1}{m_2}}(x)$. Indeed, if $\phi_2(x_0) \neq 0$, then $\phi_2\left(\frac{x_0}{\phi_2(x_0)^{\frac{1}{m_2}}}\right) = 1$, and since $\phi_1|_{\{\phi_2(x)=1\}} \equiv c$, then $\phi_1\left(\frac{x_0}{\phi_2(x_0)^{\frac{1}{m_2}}}\right) = \frac{\phi_1(x_0)}{\phi_2(x_0)^{\frac{m_1}{m_2}}} = c$. Then $\phi_1(x_0) = c\phi_2(x_0)^{\frac{m_1}{m_2}}$ for every $x_0 \in \mathbb{R}^n$ such that $\phi_2(x_0) \neq 0$. But the argument that we have just used shows that $\phi_2(x_0) \neq 0$ iff $\phi_1(x_0) \neq 0$. Hence, $\phi_1(x) = c\phi_2(x)^{\frac{m_1}{m_2}}$ for every $x \in \mathbb{R}^n$.

By the chain rule,

$$D_{i,j}\phi_1(x) = c \frac{m_1}{m_2} D_{i,j}\phi_2(x)\phi_2(x)^{\frac{m_1}{m_2}-1} + c \frac{m_1}{m_2} \left(\frac{m_1}{m_2} - 1\right) D_i(\phi_2) D_j(\phi_2)\phi_2(x)^{\frac{m_1}{m_2}-2}, \tag{2.37}$$

for $i, j, \leq n$. Then

$$D^2(\Phi)(x) = \left(\lambda_2 + c\lambda_1 \frac{m_1}{m_2} \phi_2(x)^{\frac{m_1}{m_2}-1}\right) D^2(\phi_2)(x) + c\lambda_1 \frac{m_1}{m_2} \left(\frac{m_1}{m_2} - 1\right) \phi_2(x)^{\frac{m_1}{m_2}-2} (D_1(\phi_2)(x) \nabla(\phi_2)(x) \cdots D_n(\phi_2)(x) \nabla(\phi_2)(x)). \tag{2.38}$$

Let P be a point on the support of χ . If we set $\lambda_2 + c\lambda_1 \frac{m_1}{m_2} \phi_2(P)^{\frac{m_1}{m_2}-1} = 0$, we observe that $D^2(\Phi)(P)$ equals a matrix whose rank is 1 if and only if $\nabla(\phi_2)(P) \neq (0, \dots, 0)$, $\phi_2(P)^{\frac{m_1}{m_2}-2} \neq 0$, and zero elsewhere. Since we assumed that ϕ_2 does not vanish away from the origin, Euler's homogeneity relations guarantee that $\nabla(\phi_2)(P)$ does not vanish. Consequently, the rank of $D^2(\Phi)(P)$ is at most 1.

To show that $D^2(\Phi)(P)$ cannot be zero, we observe that if this were the case we would have

$$- \frac{\left(\lambda_2 + c\lambda_1 \frac{m_1}{m_2} \phi_2(P)^{\frac{m_1}{m_2}-1}\right)}{c\lambda_1 \frac{m_1}{m_2} \left(\frac{m_1}{m_2} - 1\right) \phi_2(P)^{\frac{m_1}{m_2}-2}} D^2\phi_2(P)$$

$$= (D_1(\phi_2)(P) \nabla(\phi_2)(P) \dots D_n(\phi_2)(P) \nabla(\phi_2)(P)) . \tag{2.39}$$

The coefficient that multiplies $D^2\phi_2(P)$ cannot be zero because the matrix on the right-hand side has rank one. On the other hand, the matrix on the left-hand side has rank ≥ 2 by assumption, hence the equality in (2.39) can never hold.

This concludes the proof of the theorem. \square

Proof of Lemma 4. After perhaps rotating and dilating the coordinates, we can work in a neighborhood of the point $(0,1)$.

There exists a constant c such that $\phi_1|_{\{x:\phi_2=1\}} \equiv c, \phi_1 \equiv c\phi_2^{\frac{m_2}{m_1}}$. (See proof of Lemma 3.)

By the chain rule $D_{i,j}\phi_1(x_1, x_2) =$

$$\begin{aligned} & \frac{m_1}{m_2} D_{i,j}\phi_2(x_1, x_2) \phi_2(x_1, x_2)^{\frac{m_1}{m_2}-1} \\ & + \frac{m_1}{m_2} \left(\frac{m_1}{m_2} - 1 \right) D_i(\phi_2) D_j(\phi_2) \phi_2(x_1, x_2)^{\frac{m_1}{m_2}-2} , \end{aligned} \tag{2.40}$$

for $i, j, \leq 2$.

Observe that by Euler's homogeneity relations

- (1) $D_1(\phi_2)(0, 1) = m_2\phi_2(0, 1)$,
- (2) $D_2(\phi_2)(0, 1) = \frac{1}{m_2-1}D_{12}\phi_2(0, 1)$, and
- (3) $D_{11}(\phi_2)(0, 1) = m_2(m_2 - 1)\phi_2(0, 1)$.

In order to show that the sufficient condition of Theorem 6 is verified, we must show that the determinant of the matrix $J(x, y)$, whose rows are $\nabla(\frac{x_1^2}{2}D_{11}\phi_1(0, 1) + \frac{x_2^2}{2}D_{22}\phi_1(0, 1) + x_1x_2D_{12}\phi_1(0, 1))$ and $\nabla(\frac{x_1^2}{2}D_{11}\phi_2(0, 1) + \frac{x_2^2}{2}D_{22}\phi_2(0, 1) + x_1x_2D_{12}\phi_2(0, 1))$, is not a square. A direct computation shows that the discriminant of the determinant of $J(x_1, x_2)$ is

$$\frac{m_1^2 \left(\frac{m_1}{m_2} - 1 \right)^2}{(m_2 - 1)^2} (\phi_2(0, 1))^{2(\frac{m_1}{m_2}-1)} [\det(H(\phi_2))(0, 1)]^2 , \tag{2.41}$$

which does not vanish by the assumptions on ϕ_2 . Hence, J^{-a} is integrable for $a < 1$. This concludes the proof of the lemma. \square

3. Restriction Theorems-Scaling

We will need the following results.

Theorem 7. [1]

Let

$$S = \left\{ (x, x_{n+1}, x_{n+2}) \in \mathbb{R}^{n+2} : x_{n+1} = g_1(x), \quad x_{n+2} = g_2(x) \right\} , \tag{3.1}$$

where $g_j \in C^\infty(\mathbb{R}^n)$.

Suppose that (1.2) holds with $p_0 = \frac{2(n+4)}{n+8}$. Then for every $k \leq \frac{n}{2}$ and every θ , the expression

$$\left(\frac{d}{d\theta} \right)^k \left(\det D^2(\cos(\theta)g_1(x) + \sin(\theta)g_2(x)) \right) \tag{3.2}$$

does not vanish.

Theorem 8. [11]

Let

$$S = \{(x, x_{n+1}, \dots, x_{2n}) \in \mathbb{R}^{n+n} : x_{n+1} = g_1(x), \dots, x_{2n} = g_n(x)\}, \tag{3.3}$$

where $g_j \in C^\infty(\mathbb{R}^n)$. Let G_j denote the quadratic part of the Taylor expansion of g_j . Suppose that the vectors $\{\nabla(\frac{\partial}{\partial x_i} G_j)(0)\}_{\substack{i \leq n \\ j \leq n}}$ span \mathbb{R}^n . Let $J(x)$ denote the determinant of the matrix $(\nabla_x G_1, \dots, \nabla_x G_n)$. Suppose that $J^{-a} \in L^1(S^{n-1})$, for any $a < 1$. Then (1.2) holds with $p_0 = \frac{6}{5}$.

Note that when $n = 2$, the assumptions of Theorem 8 are equivalent to the necessary condition (3.2) in Theorem 7. In particular, the conditions of Theorem 8 are necessary and sufficient in that case.

Before stating our main results, we need to introduce the following definitions.

Definition 1.

Let $\Gamma : I \rightarrow \mathbb{R}^2$, where I is a compact interval in \mathbb{R} , and Γ is smooth. We say that Γ is finite type if $\langle (\Gamma(x) - \Gamma(x_0)), \mu \rangle$ does not vanish of infinite order for any $x_0 \in I$, and any unit vector μ .

We will also need a more precise definition to specify the order of vanishing at each point. Let a_0 denote a point in the compact interval I . We can always find a smooth function γ , such that in a small neighborhood of a_0 , $\Gamma(s) = (s, \gamma(s))$, where $s \in I$.

Definition 2.

Let Γ be defined as before. Let $\Gamma(s) = (s, \gamma(s))$ in a small neighborhood of a_0 . We say that Γ is finite type m at a_0 if $\gamma^{(k)}(a_0) = 0$ for $1 \leq k < m$, and $\gamma^{(m)}(a_0) \neq 0$.

Our main results are the following.

Lemma 5.

Let $S = \{(x, x_{n+1}, \dots, x_{n+l}) \in \mathbb{R}^{n+l} : x_{n+1} = \phi_1(x), \dots, x_{n+l} = \phi_l(x)\}$, where each $\phi_j \in C^\infty(\mathbb{R}^n \setminus \{0\})$, homogeneous of degree $m_j \geq 1$. Let $d\sigma$ denote a compactly supported smooth measure on S , and let $d\sigma_0 = \chi(x)d\sigma$, where $\chi(x)$ is a smooth cutoff function supported away from the origin. Let $Tf(x) = f * \widehat{d\sigma}$ and let $T_0f(x) = f * \widehat{d\sigma_0}$. Suppose that $T_0 : L^{q_0}(\mathbb{R}^{n+l}) \rightarrow L^{q'_0}(\mathbb{R}^{n+l})$ is a bounded operator. Then $T : L^{p_0(m_1, \dots, m_l)}(\mathbb{R}^{n+l}) \rightarrow L^{p'_0(m_1, \dots, m_l)}(\mathbb{R}^{n+l})$ is a bounded operator, where

$$p_0(m_1, \dots, m_l) = \frac{2(n + m_1 + \dots + m_l)}{n + 2(m_1 + \dots + m_l)}, \tag{3.4}$$

as long as $p_0(m_1, \dots, m_l) \leq q_0$.

Theorem 9.

Let

$$S = \{(x, x_{n+1}, \dots, x_{n+l}) : x_{n+1} = \phi_1(x), \dots, x_{n+l} = \phi_l(x)\}, \tag{3.5}$$

where $\phi_j \in C^\infty(\mathbb{R}^n \setminus \{0\})$ is homogeneous of degree $m_j \geq 2$. Suppose that S satisfies the assumptions of Theorem 5. Then the estimate (1.2) holds with

$$p_0 = \frac{2(l + \rho)}{2l + \rho}. \tag{3.6}$$

Theorem 10.

Let

$$S = \{(x, x_{n+1}, \dots, x_{n+l}) : x_{n+1} = \phi_1(x), \dots, x_{n+l} = \phi_l(x)\}, \tag{3.7}$$

where $\phi_j \in C^\infty(\mathbb{R}^n \setminus \{0\})$ is homogeneous of degree $m_j \geq 2$. Suppose that S satisfies the assumptions of Lemma 2. Then the estimate (1.2) holds with the sharp exponent

$$p_0 = \frac{2(n + m_1 + \dots + m_l)}{n + 2(m_1 + \dots + m_l)}, \tag{3.8}$$

provided that $(m_1 + \dots + m_l) \geq \frac{2ln}{N}$.

Theorem 11.

Let

$$S = \{(x, x_{n+1}, x_{n+2}) : x_{n+1} = \phi_1(x), x_{n+2} = \phi_2(x)\}, \quad n > 2, \tag{3.9}$$

where $\phi_j \in C^\infty(\mathbb{R}^n \setminus \{0\})$ is homogeneous of degree $m_j \geq 2$. Suppose that there exists a nonzero constant c , such that

$$\phi_1 \Big|_{\{x:\phi_2(x)=1\}} \equiv c.$$

Suppose that $\phi_1(\omega) \neq 0, \omega \in S^{n-1}$. If the rank of the Hessian matrix of ϕ_1 is ≥ 2 , then (1.2) holds with $p_0 = \frac{2(m_1+m_2+n)}{n+2(m_1+m_2)}$, provided that $m_1 + m_2 \geq 4n$.

In order to introduce Theorem 12, we need the following result which was stated and proved by the second author in [7].

Lemma 6.

Let $Z_0 = \{(x_1, x_2) : P(x_1, x_2) = 0\}$. Let $Z_1 = \{(x_1, x_2) : \nabla P(x_1, x_2) = (0, 0)\}$. Let $Z_2 = \{(x_1, x_2) : HP(x_1, x_2) = 0\}$, where $HP(x_1, x_2)$ denotes the determinant of the Hessian matrix of P . Then for each $j = 0, 1, 2, Z_j = \{(0, 0)\} \cup \bigcup_{k=1}^{N_j} L_k$, where each L_k is a line through the origin, and $N_j < \infty$. Moreover, $Z_1 = Z_0 \cap Z_2$.

Proof of Lemma 6. Let P_j denote the partial derivative of P with respect to x_j . Since P is homogeneous of degree m , P_j is homogeneous of degree $m - 1$, and $HP(x_1, x_2)$ is homogeneous of degree $2(m - 2)$. By homogeneity, if Z_j contains a point (x_1, x_2) , it also contains a line through the origin containing that point. Since P is a polynomial, there can be at most a finite number of such lines. This proves the first assertion of the lemma.

By the Euler homogeneity relations,

$$\begin{aligned} mP(x_1, x_2) &= x_1P_1(x_1, x_2) + x_2P_2(x_1, x_2), \\ (m - 1)P_1(x_1, x_2) &= x_1P_{11}(x_1, x_2) + x_2P_{12}(x_1, x_2), \end{aligned} \tag{3.10}$$

and

$$(m - 1)P_2(x_1, x_2) = x_1P_{21}(x_1, x_2) + x_2P_{22}(x_1, x_2),$$

where the $\{P_{jk}\}$ denote the second partial derivatives. Hence, $Z_0 \subset Z_1$. If we write the equations for P_1 and P_2 in matrix form, we see that $(m - 1) \Delta P(x_1, x_2)$ is obtained by applying the Hessian matrix of P to the vector (x_1, x_2) . Hence, $Z_2 \subset Z_1$. Putting these observations together we see that $Z_0 \cap Z_2 \subset Z_1$.

Suppose that both P and HP vanish along a line through the origin, which without loss of generality we take to be the x_1 -axis. Then $mP(x_1, 0) = x_1P_1(x_1, 0)$. This implies that $P_1(x_1, 0) = 0$. Also, $(m - 1)P_1(x_1, 0) = x_1P_{11}(x_1, 0)$. This implies that $P_{11}(x_1, 0) = 0$. $P_2(x_1, 0) = x_1P_{12}(x_1, 0)$. By assumption,

$$HP(x_1, 0) = P_{11}(x_1, 0)P_{22}(x_1, 0) - P_{12}^2(x_1, 0) = 0. \tag{3.11}$$

Since $P_{11}(x_1, 0) = 0$, we must conclude that $P_{12}(x_1, 0) = 0$, which implies that $P_2(x_1, 0) = 0$. This proves that $\nabla P(x_1, 0) = (0, 0)$ and hence that $Z_1 \subset Z_0 \cap Z_2$. This completes the proof of the lemma. \square

We shall need the following definition:

Definition 3.

Let $f \in C^\infty(\mathbb{R}^2)$. We say that f vanishes of order M along the line $L = \{(x_1, x_2) : x_1 = s_1t, x_2 = s_2t, t \in \mathbb{R}\}$ if M is the largest positive integer so that $f(s_1t, s_2t) = t^M g(t)$, where $g \in C^\infty(\mathbb{R}^2)$ is allowed to vanish only at the origin.

Theorem 12.

Let $S = \{(x_1, x_2, x_3, x_4) : x_3 = \phi_1(x_1, x_2), x_4 = \phi_2(x_1, x_2), \}$ where $\phi_1, \phi_2 \in C^\infty(\mathbb{R}^2)$ are homogeneous polynomials of degree m_1 and m_2 , respectively, ($m_1 \geq m_2 \geq 2$). Suppose that there exists a non-zero constant c such that $\phi_1(x)|_{\{x:\phi_2(x)=1\}} = c$. Let Z_0, Z_1, Z_2 be defined as in Lemma 6 with respect to ϕ_2 . Suppose that the curve $\{x : \phi_2(x) = 1\}$ is of finite type m at each point of $Z_2 \cap \{x : \phi_2(x) = 1\}$ and that ϕ_2 vanishes of order $\leq M$ along the lines contained in $Z_0 \cup Z_1$. Then (1.2) holds every $p < p_0$, where p_0 is the sharp exponent

$$\frac{m_1 + m_2 + 2}{m_1 + m_2 + 1}, \quad (3.12)$$

if $m_1 + m_2 \geq \max\left\{2M\left(1 + \frac{m_1}{m_2}\right), 2m\right\}$. Furthermore (1.2) holds with

$$(1) p < \max\left\{\frac{2(1+M(1+\frac{m_1}{m_2}))}{1+2M(1+\frac{m_1}{m_2})}, \frac{2(m+1)}{2m+1}\right\} \text{ if } \min\{2M(1 + \frac{m_1}{m_2}), 2m\} \leq m_1 + m_2 <$$

$$\max\{2M(1 + \frac{m_1}{m_2}), 2m\},$$

$$(2) p < \min\left\{\frac{2(1+M(1+\frac{m_1}{m_2}))}{1+2M(1+\frac{m_1}{m_2})}, \frac{2(m+1)}{2m+1}\right\} \text{ if } 4 \leq m_1 + m_2 < \min\{2M(1 + \frac{m_1}{m_2}), 2m\}.$$

Proof of Lemma 5. Let

$$\widehat{d\sigma}(\xi, \lambda) = \int_{\mathbb{R}^n} e^{i((x,\xi)+\lambda_1\phi_1(x)+\dots+\lambda_l\phi_l(x))} \psi(x) dx, \quad (3.13)$$

where $\psi(x)$ is a cutoff function. Let ρ be a cutoff function supported in the interval (1,2) such that $\sum_{j=0}^{+\infty} \rho(2^j t) = 1$ for every t , and let

$$\widehat{d\sigma}_j(\xi, \lambda) = \int_{\mathbb{R}^n} e^{i((x,\xi)+\lambda_1\phi_1(x)+\dots+\lambda_l\phi_l(x))} \rho(2^j t) dx. \quad (3.14)$$

If we make the change of variables sending $x \rightarrow 2^{-j}x$ we can write:

$$\widehat{d\sigma}_j(\xi, \lambda) = 2^{-jn} \widehat{d\sigma}_0(2^{-j}\xi, 2^{-m_1j}\lambda_1, \dots, 2^{m_lj}\lambda_l). \quad (3.15)$$

Let τ_j denote the nonisotropic dilation

$$\tau_j f(\xi, \lambda_1, \dots, \lambda_l) = f(2^j\xi, 2^{m_1j}\lambda_1, \dots, 2^{m_lj}\lambda_l). \quad (3.16)$$

Then

$$\widehat{d\sigma}_j * f = 2^{-nj} (\tau_{-j} \widehat{d\sigma}_0 * f)(\xi, \lambda) = 2^{-nj} (\tau_{-j} \widehat{d\sigma}_0 * (\tau_{-j} \tau_j f))(\xi, \lambda). \quad (3.17)$$

A change of variables shows that $\tau_{-j} \widehat{d\sigma}_0 * (\tau_{-j} \tau_j f) = 2^{j(n+m_1+\dots+m_l)} \tau_{-j}(\widehat{d\sigma}_0 * \tau_j f)$.

It follows that

$$\begin{aligned} \|\widehat{d\sigma}_j * f\|_{q'_0} &= 2^{-nj+j(n+m_1+\dots+m_l)} \|\tau_{-j}(\widehat{d\sigma}_0 * \tau_j f)\|_{q'_0} \\ &= 2^{j(m_1+\dots+m_l)} 2^{\frac{j}{q'_0}(n+m_1+\dots+m_l)} \|\widehat{d\sigma}_0 * (\tau_j f)\|_{q'_0} \\ &\leq C 2^{j(m_1+\dots+m_l)} 2^{\frac{j}{q'_0}(n+m_1+\dots+m_l)} \|\tau_j f\|_{q_0} \\ &= C 2^{j(m_1+\dots+m_l)} 2^{j\left(\frac{1}{q'_0} - \frac{1}{q_0}\right)(n+m_1+\dots+m_l)} \|f\|_{q_0}. \end{aligned} \quad (3.18)$$

The series $\sum_{j=1}^{\infty} \|\widehat{d\sigma}_j * f\|_{q'_0}$ converges, provided that

$$m_1 + \cdots + m_l + \left(\frac{1}{q'_0} - \frac{1}{p_0}\right)(n + m_1 + \cdots + m_l) < 0, \quad (3.19)$$

which yields

$$q_0 < \frac{2(n + m_1 + \cdots + m_l)}{n + 2(m_1 + \cdots + m_l)}. \quad (3.20)$$

This concludes the proof of the lemma. \square

Proof of Theorem 9. The application of Greenleaf's theorem (see (1.3) above) yields the result. \square

Proof of Theorem 10. Let F_0 be defined as in (2.5). By Lemma 2

$$|F_0(\xi, \lambda)| \leq C(1 + |\xi| + |\lambda|)^{-\frac{N}{2}}. \quad (3.21)$$

By a theorem of Greenleaf (see (1.3) above) the inequality (1.2) holds with $q_0 = \frac{2(2l+N)}{4l+N}$. An application of Lemma 5 completes the proof. \square

Proof of Theorem 11. Let F_0 be defined as in (2.5). By Lemma 3

$$|F_0(\xi, \lambda)| \leq C(1 + |\xi| + |\lambda|)^{-\frac{1}{2}}. \quad (3.22)$$

Applying Greenleaf's theorem as above we get $q_0 = \frac{10}{9}$. An application of Lemma 5 completes the proof. \square

Proof of Theorem 12. Let Σ be the level set $\{x : \phi_2(x) = 1\}$, and let

$$\widehat{d\sigma}(\xi, \lambda_1, \lambda_2) = \int_{\mathbb{R}^2} e^{i((x, \xi) + \lambda_1 \phi_1(x) + \lambda_2 \phi_2(x))} \chi(x) dx, \quad (3.23)$$

where χ is a smooth cutoff function. Let Z_0 , Z_1 , and Z_2 be defined as in Lemma 6 with respect to ϕ_2 . Recall that $Z_0 \cup Z_1 \cup Z_2$ is the union of a finite number of lines through the origin.

Let $\{\Gamma_j(x)\}_{j \leq N_1}$ and $\{T_j(x)\}_{j \leq N_2}$ be two finite families of cones in \mathbb{R}^2 with the following properties:

- i) Each $\Gamma_j(x)$ contains exactly one line of $Z_0 \cup Z_1$, and each $T_j(x)$ contains exactly one line of Z_2
- ii) $\Gamma_j \cap \Gamma_i = \{0, 0\}$ if $i \neq j$, $T_j \cap T_i = \{0, 0\}$ if $i \neq j$, and $\Gamma_j \cap T_i = \{0, 0\}$.

Let α_j be the characteristic function of Γ_j and let β_j be the characteristic function of T_j . Then

$$\begin{aligned} \widehat{d\sigma}(\xi, \lambda_1, \lambda_2) &= \int_{\mathbb{R}^2} e^{i((x, \xi) + \lambda_1 \phi_1(x) + \lambda_2 \phi_2(x))} \chi(x) \left(\sum_{j=1}^{N_1} \alpha_j(x) \right) dx \\ &\quad + \int_{\mathbb{R}^2} e^{i((x, \xi) + \lambda_1 \phi_1(x) + \lambda_2 \phi_2(x))} \chi(x) \left(\sum_{j=1}^{N_2} \beta_j(x) \right) dx \\ &\quad + \int_{\mathbb{R}^2} e^{i((x, \xi) + \lambda_1 \phi_1(x) + \lambda_2 \phi_2(x))} \chi(x) \left(1 - \sum_{j=1}^{N_1} \alpha_j(x) - \sum_{j=1}^{N_2} \beta_j(x) \right) dx \\ &= \widehat{d\sigma}_1 + \widehat{d\sigma}_2 + \widehat{d\sigma}_3. \end{aligned} \quad (3.24)$$

We first consider

$$\widehat{d\sigma}_3(\xi, \lambda_1, \lambda_2) = \int_{\mathbb{R}^2} e^{i((x,\xi)+\lambda_1\phi_1(x)+\lambda_2\phi_2(x))} \bar{\eta}(x) \chi(x) dx, \quad (3.25)$$

where we have set $\bar{\eta}(x) = 1 - \sum_{j=1}^{N_1} \alpha_j(x) - \sum_{j=1}^{N_2} \beta_j(x)$. On the support of $\bar{\eta}$ the curvature of $\Sigma = \{x : \phi_2(x) = 1\}$ never vanishes and ϕ_2 vanishes only at zero.

We recall that by the Stein-Thomas observation, (1.2) is equivalent to the inequality

$$\|\widehat{d\sigma} * f\|_{L^{p'}(\mathbb{R}^4)} \leq C_j \|f\|_{L^p(\mathbb{R}^4)}. \quad (3.26)$$

Let ρ be a smooth cutoff function supported in the interval (1,2), such that $\sum_{j=0}^{+\infty} \rho(2^j t) \equiv 1$.

Let

$$\widehat{d\sigma}_{3,j}(\xi, \lambda_1, \lambda_2) \int_{\mathbb{R}^2} e^{i((x,\xi)+\lambda_1\phi_1(x)+\lambda_2\phi_2(x))} \rho(2^j x) \bar{\eta}(x) dx. \quad (3.27)$$

The assumptions of Lemma 4 are satisfied on the support of $\rho(2^j x)\eta(x)$, hence the inequality (3.26) holds for the measure $d\sigma_{3,j}$ with $p = \frac{6}{5}$. Since the sharp exponent p_0 cannot exceed $\frac{6}{5}$, the estimate (3.26) holds for $p \leq p_0$, provided that $m_1 + m_2 \geq 4$.

If we make the change of variables sending $x \rightarrow 2^{-j}x$, and if we observe that $\bar{\eta}$ is invariant with respect to dilations, we see that

$$\begin{aligned} \widehat{d\sigma}_{3,j}(\xi, \lambda) &= 2^{-2j} \int_{\mathbb{R}^2} e^{i((2^{-j}x,\xi)+\lambda_1 2^{-m_1 j} \phi_1(x)+\lambda_2 2^{-m_2 j} \phi_2(x))} \rho(x) \bar{\eta}(x) dx \\ &= 2^{-2j} \widehat{d\sigma}_{3,0}(2^{-j}\xi, 2^{-m_1 j} \lambda_1, 2^{-m_2 j} \lambda_2). \end{aligned} \quad (3.28)$$

Without loss of generality we can replace $\bar{\eta}$ by a function $\eta \in C^\infty(\mathbb{R}^n/\{0\})$, homogeneous of degree zero, whose support coincides with the support of $\bar{\eta}$.

Let τ_j denote the nonisotropic dilation

$$\tau_j f(\xi, \lambda_1, \lambda_2) = f(2^j \xi, 2^{m_1 j} \lambda_1, 2^{m_2 j} \lambda_2). \quad (3.29)$$

Then

$$\widehat{d\sigma}_{3,j} * f = 2^{-2j} (\tau_{-j} \widehat{d\sigma}_{3,0} * f)(\xi, \lambda_1, \lambda_2) = 2^{-2j} (\tau_{-j} \widehat{d\sigma}_{3,0} * (\tau_j f))(\xi, \lambda_1, \lambda_2). \quad (3.30)$$

A change of variables shows that $\tau_{-j} \widehat{d\sigma}_{3,0} * (\tau_j f) = 2^{j(2+m_1+m_2)} \tau_{-j}(\widehat{d\sigma}_{3,0} * \tau_j f)$. It follows that

$$\begin{aligned} \|\widehat{d\sigma}_{3,j} * f\|_{p'} &= 2^{-2j+j(2+m_1+m_2)} \|\tau_{-j}(\widehat{d\sigma}_{3,0} * \tau_j f)\|_{p'} \\ &= 2^{j(m_1+m_2)} 2^{\frac{j}{p'}(2+m_1+m_2)} \|\widehat{d\sigma}_{3,0} * (\tau_j f)\|_{p'} \\ &\leq C 2^{j(m_1+m_2)} 2^{\frac{j}{p'}(2+m_1+m_2)} \|\tau_j f\|_p \\ &= C 2^{j(m_1+m_2)} 2^{j(\frac{1}{p'}-\frac{1}{p})(2+m_1+m_2)} \|f\|_p. \end{aligned} \quad (3.31)$$

The series $\sum_{j=1}^{\infty} \|\widehat{d\sigma}_{3,j} * f\|_{p'}$ converges, provided that

$$m_1 + m_2 + \left(\frac{1}{p'} - \frac{1}{p}\right)(n + m_1 + m_2) < 0, \quad (3.32)$$

which yields $p < p_0$. Hence, the measure $d\sigma_3$ satisfies the inequality (3.26) with $p < p_0$.

We consider now $\widehat{d\sigma}_2(\xi, \lambda_1, \lambda_2) = \int_{\mathbb{R}^2} e^{i((x,\xi)+\lambda_1\phi_1(x)+\lambda_2\phi_2(x))} \chi(x) \left(\sum_{j=1}^{N_2} \beta_j(x) \right) dx$.

Let

$$\widehat{d\mu}(\xi, \lambda_1, \lambda_2) = \int_{\mathbb{R}^2} e^{i((x,\xi)+\lambda_1\phi_1(x)+\lambda_2\phi_2(x))} \chi(x) \beta_j(x) dx, \quad (3.33)$$

where $j \leq N_2$ is fixed.

In the proof of Lemma 3 we observed that $\phi_1(x)|_{\Sigma} = c$ implies that $\phi_1(x) = c\phi_2(x)^{\frac{m_1}{m_2}}$ for every $x \in \mathbb{R}^2$. Observe that Σ is star-shaped with respect to the origin, because if $x_0 \in \Sigma$, then for every $t \leq 1$, $\phi_1(tx_0) = t^{m_1} \leq 1$. In the polar coordinates associated to Σ ,

$$\widehat{d\mu}(\xi, \lambda_1, \lambda_2) = \int_{\Sigma} \int_0^{+\infty} e^{i(r(\omega,\xi)+c\lambda_1 r^{m_1}+\lambda_2 r^{m_2})} r \chi(r\omega) \beta_j(\omega) dr d\omega. \quad (3.34)$$

Without loss generality χ is radial. Consider the (unique) point of $Z_2 \cap \Sigma \cap T_j$, which can be taken to be $(0,0)$. Suppose that $\Sigma \cap T_j$ is supported in a sufficiently small neighborhood of $(0,0)$. Since Σ is finite type m , it can be written as the graph of a smooth function $\psi(t) = t^m g(t)$, where $g(0) \neq 0$, and

$$\widehat{d\mu}(\xi, \lambda_1, \lambda_2) = \int_{\mathbb{R}} \int_0^{+\infty} e^{i(rt\xi_1+rt^m g(t)\xi_2+c\lambda_1 r^{m_1}+\lambda_2 r^{m_2})} r \chi(r) \beta_j(t, t^m g(t)) dr dt. \quad (3.35)$$

Let $\rho \in C^\infty(\mathbb{R}^2)$ be supported in $(\frac{1}{4}, 4)$, such that $\rho \equiv 1$ in $(1,2)$, and $\sum_{j=1}^{+\infty} \rho(2^j t) \equiv 1$. Let

$$\widehat{d\mu}_j(\xi, \lambda_1, \lambda_2) = \int_{\mathbb{R}} \int_0^{+\infty} e^{i(rt\xi_1+rt^m g(t)\xi_2+c\lambda_1 r^{m_1}+\lambda_2 r^{m_2})} \rho(2^j t) \chi(r) r dr dt. \quad (3.36)$$

The integral with respect to t is supported over a dyadic piece of Σ where the Gaussian curvature does not vanish. By Lemma 4 and the Stein-Thomas observation, the estimate (3.26) holds for the measure $d\mu_j$, for $p \leq p_0$, with a constant C_j .

In order to estimate C_j we make a change of variables in the expression for $\widehat{d\mu}_j$ setting $s = 2^j t$. We have

$$\widehat{d\mu}_j(\xi, \lambda_1, \lambda_2) = 2^{-j} \int_{\mathbb{R}} \int_0^{+\infty} e^{i(r2^{-j}s\xi_1+r2^{-mj} s^m g(2^{-j}t)\xi_2+c\lambda_1 r^{m_1}+\lambda_2 r^{m_2})} \chi(r) \rho(s) r dr ds. \quad (3.37)$$

Let τ_j be the nonisotropic dilation

$$\tau_j(f)(x_1, x_2, x_3, x_4) = f(2^{-j}x_1, 2^{-mj}x_2, x_3, x_4) \text{ and let}$$

$$\widehat{d\widetilde{\mu}}_j = \int_{\mathbb{R}} \int_0^{+\infty} e^{i(rs\xi_1+rs^m g(2^{-j}t)\xi_2+c\lambda_1 r^{m_1}+\lambda_2 r^{m_2})} \chi(r) \rho(s) r dr ds. \quad (3.38)$$

Then

$$\widehat{d\mu}_j * f = 2^{-j} (\tau_j(\widehat{d\widetilde{\mu}}_j) * f) = 2^{-j} (\tau_j(\widehat{d\widetilde{\mu}}_j) * \tau_j(\tau_{-j}f)). \quad (3.39)$$

A change of variables shows that $\tau_j(\widehat{d\widetilde{\mu}}_j) * \tau_j \tau_{-j}f = 2^{j(m+1)} \tau_j(\widehat{d\widetilde{\mu}}_j * \tau_{-j}f)$, and that $\|\tau_j \psi\|_{L^q(\mathbb{R}^4)} = 2^{j\frac{(m+1)}{q}} \|\psi\|_{L^q(\mathbb{R}^4)}$ for every $\psi \in L^q(\mathbb{R}^4)$ and $q \geq 1$. Then we can write the following string of inequalities:

$$\begin{aligned} \|\widehat{d\mu}_j * f\|_{L^{p'}(\mathbb{R}^4)} &= 2^{jm} \|\tau_j(\widehat{d\widetilde{\mu}}_j * \tau_{-j}f)\|_{L^{p'}(\mathbb{R}^4)} \\ &= 2^{jm+j\frac{(m+1)}{p'}} \|\widehat{d\widetilde{\mu}}_j * \tau_j f\|_{L^{p'}(\mathbb{R}^4)} \\ &\leq 2^{jm+j\frac{(m+1)}{p'}} C_j \|\tau_j f\|_{L^p(\mathbb{R}^4)} \\ &= 2^{jm+j(m+1)(\frac{1}{p}-\frac{1}{p'})} C_j \|f\|_{L^p(\mathbb{R}^4)}. \end{aligned} \quad (3.40)$$

We must prove that the constants C_j in the above expression are uniformly bounded. In fact the sum $\sum_{j=1}^{+\infty} \|\widehat{d\mu}_j * f\|_{L^{p'}(\mathbb{R}^4)}$ converges if $m + (m + 1)(\frac{1}{p'} - \frac{1}{p}) < 0$, hence if $p < \frac{2(m+1)}{2m+1}$. Since $\frac{2(m+1)}{2m+1} \geq p_0$ when $m_1 + m_2 \geq 2m$, the estimate (3.26) holds for the measure $d\mu$, and consequently for the measure $d\sigma_2$, for $p \leq p_0$, provided that $m_1 + m_2 \geq 2m$, and with $p = \frac{2(m+1)}{2m+1}$ if $m_1 + m_2 < 2m$.

From the proof of the theorem of Greenleaf it follows that the bounds for the constants C_j depend on a finite number of derivatives of the phase function of $\widehat{d\mu}_j$, $\Phi_j(r, s) = rs\xi_1 + rs^m g(2^{-j}t)\xi_2 + c\lambda_1 r^{m_1} + \lambda_2 r^{m_2}$. Since Φ_j is a smooth function, then, for j large, $D^\beta \Phi_j(r, s) \approx D^\beta (rs\xi_1 + rs^m g(0)\xi_2 + c\lambda_1 r^{m_1} + \lambda_2 r^{m_2})$. This shows that the constant C_j s are uniformly bounded.

We now consider $\widehat{d\sigma}_1(\xi, \lambda_1, \lambda_2) = \int_{\mathbb{R}^2} e^{i((x,\xi) + \lambda_1 \phi_1(x) + \lambda_2 \phi_2(x))} \chi(x) \left(\sum_{j=1}^{N_1} \alpha_j(x) \right) dx$.

Fix $j \leq N_1$. After perhaps a rotation of coordinates we may assume that $\phi_{2|\Gamma_j}$ vanishes along the x_2 axis. Then $\phi_2(x_1, x_2)$ can be written as $x_1^M g(x_1, x_2)$, where g does not vanish on the x_2 axis (except perhaps at the origin) if $\Gamma_j \cap S^1$ is small enough.

Let

$$\widehat{d\mu}(\xi, \lambda_1, \lambda_2) = \int_{\mathbb{R}^2} e^{i((x,\xi) + c\lambda_1 x_1^M g(x) + \lambda_2 x_1^M g(x))} \chi(x) \alpha_j(x) dx, \tag{3.41}$$

where we have set $\gamma = \frac{m_1}{m_2}$. Let $\rho \in C^\infty(\mathbb{R}^2)$ be a cutoff supported in $(\frac{1}{4}, 4)$ such that $\rho \equiv 1$ in $(1, 2)$ and $\sum_{j=1}^{+\infty} \rho(t) \equiv 1$. Let

$$\widehat{d\mu}_j(\xi, \lambda) = \int_{\mathbb{R}^2} e^{i((x,\xi) + c\lambda_1 x_1^{\gamma M} g^\gamma(x) + \lambda_2 x_1^M g(x))} \rho\left(2^j \frac{x_1}{x_2}\right) \chi(x) dx. \tag{3.42}$$

The above integral is defined over a cone of \mathbb{R}^2 where the curvature of Σ never vanishes, and ϕ_2 vanishes only at the origin.

Let $\rho_0 \in C^\infty(\mathbb{R}^2)$ be supported in $(\frac{1}{4}, 4)$, such that $\rho \equiv 1$ in $(1, 2)$, and $\sum_{k=0}^{+\infty} \rho_0(2^k t) \equiv 1$.

Let

$$\widehat{d\mu}_{j,k}(\xi, \lambda) = \int_{\mathbb{R}^2} e^{i((x,\xi) + c\lambda_1 x_1^{\gamma M} g^\gamma(x) + \lambda_2 x_1^M g(x))} \rho\left(2^j \frac{x_1}{x_2}\right) \rho_0(2^k x) dx. \tag{3.43}$$

The assumptions of Lemma 4 are satisfied on the support of $\rho_0(2^k x) \rho(2^j \frac{x_1}{x_2})$, and hence the estimate (3.26) holds for the measure $d\mu_{j,k}$ with $p \leq p_0$.

If we make the change of variables sending $x \rightarrow 2^{-k}x$,

$$\begin{aligned} \widehat{d\mu}_{j,k}(\xi, \lambda) &= 2^{-2k} \int_{\mathbb{R}^2} e^{i((2^{-k}x,\xi) + \lambda_1 2^{-m_1 k} \phi_1(x) + \lambda_2 2^{-m_2 k} \phi_2(x))} \rho\left(2^j \frac{x_1}{x_2}\right) \rho_0(x) dx \\ &= 2^{-2k} \widehat{d\mu}_{j,0}(2^{-k}\xi, 2^{-m_1 k} \lambda_1, 2^{-m_2 k} \lambda_2). \end{aligned} \tag{3.44}$$

Let τ_k denote the nonisotropic dilation

$$\tau_k f(\xi, \lambda_1, \lambda_2) = f(2^k \xi, 2^{m_1 k} \lambda_1, 2^{m_2 k} \lambda_2). \tag{3.45}$$

Then

$$\begin{aligned} \widehat{d\mu}_{j,k} * f &= 2^{-2k} (\tau_{-k} \widehat{d\mu}_{0,k} * f)(\xi, \lambda_1, \lambda_2) \\ &= 2^{-2k} (\tau_{-k} \widehat{d\mu}_{0,k} * (\tau_{-k} \tau_k f))(\xi, \lambda_1, \lambda_2). \end{aligned} \tag{3.46}$$

A change of variables shows that $\tau_{-k} \widehat{d\mu}_{j,k} * (\tau_{-k} \tau_k f) = 2^{k(2+m_1+m_2)} \tau_{-k}(\widehat{d\mu}_{j,k} * \tau_k f)$. It follows that

$$\|\widehat{d\mu}_{j,k} * f\|_{p'} = 2^{-2k+k(2+m_1+m_2)} \|\tau_{-k}(\widehat{d\mu}_{j,k} * \tau_k f)\|_{p'}$$

$$\begin{aligned}
&= 2^{k(m_1+m_2)} 2^{\frac{k}{p'}(2+m_1+m_2)} \|\widehat{d\mu_{j,k}} * (\tau_k f)\|_{p'} \\
&\leq C 2^{k(m_1+m_2)} 2^{\frac{k}{p'}(2+m_1+m_2)} \|\tau_k f\|_p \\
&= C 2^{k(m_1+m_2)} 2^{k(\frac{1}{p'}-\frac{1}{p})(2+m_1+m_2)} \|f\|_p.
\end{aligned} \tag{3.47}$$

The series $\sum_{k=1}^{\infty} \|\widehat{d\mu_{j,k}} * f\|_{p'}$, converges provided that

$$m_1 + m_2 + \left(\frac{1}{p'} - \frac{1}{p}\right) (n + m_1 + m_2) < 0, \tag{3.48}$$

which yields $p < p_0$. The above argument shows that we can assume that the measure $d\mu_j$ is supported away from zero. Hence, by Lemma 4 and the above observation, the inequality (3.26) holds for the measure $d\mu_j$, for $p \leq p_0$, with a constant C_j . In order to estimate C_j we perform the change of variables in (3.42) sending $x_1 \rightarrow 2^{-j}x_1, x_2 \rightarrow x_2$. We obtain

$$\widehat{d\mu_j}(\xi_1, \xi_2, \lambda_1, \lambda_2) = 2^{-j} \widehat{d\widetilde{\mu}_j}(2^{-j}\xi_1, \xi_2, 2^{-M\gamma j}\lambda_1, 2^{-Mj}\lambda_2), \tag{3.49}$$

where we have set

$$\begin{aligned}
\widehat{d\widetilde{\mu}_j}(\xi, \lambda) &= \int_{\mathbb{R}^2} e^{i((x,\xi)+c\lambda_1 x_1^{\gamma M} g^\gamma(2^{-j}x_1, x_2) + \lambda_2 x_1^M g(2^{-j}x_1, x_2))} \\
&\quad \rho\left(\frac{x_1}{x_2}\right) \chi(2^{-j}x_1, x_2) dx.
\end{aligned} \tag{3.50}$$

Let τ_j be the nonisotropic dilation $\tau_j f(x_1, x_2, x_3, x_4) = f(2^{-j}x_1, x_2, 2^{-M\gamma j}x_3, 2^{-Mj}x_4)$. Then

$$\widehat{d\mu_j} * f = 2^{-j} (\tau_j (\widehat{d\widetilde{\mu}_j}) * f) = 2^{-j} (\tau_j (\widehat{d\widetilde{\mu}_j}) * \tau_j (\tau_{-j} f)). \tag{3.51}$$

A change of variables shows that

$$\tau_j (\widehat{d\widetilde{\mu}_j}) * \tau_j (\tau_{-j} f) = 2^{j(1+M(1+\gamma))} \tau_j (\widehat{d\widetilde{\mu}_j} * \tau_{-j} f), \tag{3.52}$$

and that $\|\tau_j g\|_{L^q(\mathbb{R}^4)} = 2^{j\frac{(1+M(1+\gamma))}{q}} \|g\|_{L^q(\mathbb{R}^4)}$ for every $q \geq 1$. Then we can write the following string of inequalities:

$$\begin{aligned}
\|\widehat{d\mu_j} * f\|_{L^{p'}(\mathbb{R}^4)} &= 2^{j(M(1+\gamma))} \|\tau_j (\widehat{d\widetilde{\mu}_j} * \tau_{-j} f)\|_{L^{p'}(\mathbb{R}^4)} \\
&= 2^{jM(1+\gamma)+j\frac{(1+M(1+\gamma))}{p'}} \|\widehat{d\widetilde{\mu}_j} * \tau_j f\|_{L^{p'}(\mathbb{R}^4)} \\
&\leq 2^{j(M(1+\gamma))+j\frac{M(1+\gamma)+1}{p'}} C^j \|\tau_j f\|_{L^p(\mathbb{R}^4)} \\
&= 2^{jM(1+\gamma)+j(M(1+\gamma)+1)(\frac{1}{p'}-\frac{1}{p})} C^j \|f\|_{L^p(\mathbb{R}^4)}.
\end{aligned} \tag{3.53}$$

If we show that the constants C_j in the above expression are uniformly bounded, then we are done.

In fact the sum $\sum_{j=1}^{+\infty} \|\widehat{d\mu_j} * f\|_{L^{p'}(\mathbb{R}^4)}$ converges if

$$M(1+\gamma) + (M(1+\gamma)+1)\left(\frac{1}{p'} - \frac{1}{p}\right) < 0 \text{ hence if}$$

$$p < \frac{2(1+M(\gamma+1))}{1+2M(\gamma+1)}. \tag{3.54}$$

Since $\frac{2(1+M(\gamma+1))}{1+2M(\gamma+1)} \geq p_0$ when $m_1 + m_2 \geq 2M(1+\gamma)$, the estimate (3.26) holds for the measure $d\mu$, and consequently for the measure $d\sigma_1$, with $p < p_0$, provided that $m_1 + m_2 \geq 2M(1+\gamma)$, and with $p < \frac{2(1+M(\gamma+1))}{1+2M(\gamma+1)}$ if $4 < m_1 + m_2 < 2M(1+\gamma)$.

By a theorem of Greenleaf, the bounds for the constants C_j in (3.53) depend only on a finite number of derivatives of the phase function of $\widehat{d\mu}_j$, $\Phi_j(x) = (c\lambda_1 x_1^{\gamma M} g^\gamma(2^{-j}x_1, x_2) + \lambda_2 x_1^M g(2^{-j}x_1, x_2))$. Since the above function is smooth, then, for j large, $D^\beta \Phi_j(x) \approx D^\beta (c\lambda_1 x_1^{\gamma M} g^\gamma(0, x_2) + \lambda_2 x_1^M g(0, x_2))$. This shows that the constants C_j are uniformly bounded, thus concluding the proof of the theorem. \square

4. Restriction Theorems - Non-Isotropic Estimates

Theorem 13.

Let $S = \{(x, x_{n+1}, \dots, x_{n+l}) \in \mathbb{R}^{n+l} : x_{n+1} = \phi_1(x), \dots, x_{n+l} = \phi_l(x)\}$, where $l = \frac{n(n+1)}{2}$, and the ϕ_j denote the distinct monomials of degree 2. Then the estimate (1.2) holds with the sharp exponent $p_0 = \frac{2(n+2)}{2n+3}$.

Theorem 14.

Let S denote a compact piece of the manifold $\{(x, x_{n+1}, \dots, x_{n+l}) \in \mathbb{R}^{n+l} : x_{n+1} = \phi_1(x), \dots, x_{n+l} = \phi_l(x)\}$, where $\phi_j \in C^\infty(\mathbb{R}^n/0)$ is homogeneous of degree $m \geq 2n$. Suppose that no linear combination of the ϕ_j s vanishes on a subset of positive measure of S^{n-1} . Let $\Phi(x) = (\phi_1(x), \dots, \phi_l(x))$. Suppose that $\Phi(\omega) \neq (0, \dots, 0)$, $\omega \in S^{n-1}$. Then the estimate (1.2) holds with the sharp exponent p_0 given by (1.4).

Remark 1.

The restriction $m \geq 2n$ in Theorem 14 is not necessary. In fact, using the techniques in [8], one can prove Theorem 14 under the weaker restriction $m \geq n$.

Remark 2.

Theorem 14 implies the natural generalization of Theorem 13 to the case where $l = C_m^{m+n-1}$, ($C_b^a = \frac{a!}{b!(a-b)!}$), and the ϕ_j are the distinct monomials of degree m .

Proof of Theorem 13. Let $\lambda = (\lambda_1, \dots, \lambda_l)$, and let

$$\widehat{d\sigma}(\xi, \lambda) = \int_{\mathbb{R}^n} e^{i((x, \xi) + \lambda_1 \phi_1 + \dots + \lambda_l \phi_l(x))} dx .$$

Let A_λ be the matrix associated to the quadratic form $\lambda_1 \phi_1(x) + \dots + \lambda_l \phi_l(x)$, and

$$\widehat{d\sigma}(\xi, \lambda) = \int_{\mathbb{R}^n} e^{i((x, \xi) + (x, A_\lambda x))} dx . \tag{4.1}$$

Thus, $\widehat{d\sigma}(\xi, \lambda)$ is the Fourier transform of $e^{i(x, A_\lambda x)}$, and an easy generalization of the well-known formula for the Fourier transform of the Gaussian functions, (see, e.g., [10, p.186]), yields

$$\widehat{d\sigma}(\xi, \lambda) = \frac{(2\pi)^{\frac{n}{2}}}{|\det A_\lambda|^{1/2}} e^{-\frac{i}{2}(\xi, A_\lambda^{-1} \xi) + \frac{\pi i}{4} \text{sign}(A_\lambda)} . \tag{4.2}$$

Let

$$\begin{aligned} K_z(\xi, \lambda) &= \psi(z) \det(A_\lambda)^z \widehat{d\sigma}(\xi, \lambda) \\ &= \psi(z) (2\pi)^{\frac{n}{2}} |\det A_\lambda|^{z-1/2} e^{-\frac{i}{2}(\xi, A_\lambda^{-1} \xi) + \frac{\pi i}{4} \text{sign}(A_\lambda)} , \end{aligned} \tag{4.3}$$

with

$$\psi(z) = (2\pi)^{nz + \frac{n(n+1)}{2}} \prod_{j=0}^{n-1} \Gamma^{-1} \left(z + 1 + \frac{j}{2} \right) e^{-i\pi \frac{n}{2} (z + \frac{n+1}{2})} , \tag{4.4}$$

where Γ is the standard Gamma function. Let $T_z(f)(\xi, \lambda) = (f * K_z)(\xi, \lambda)$.

We will prove that T_z is a continuous family of operators when $\text{Re}(z) \in [-\frac{n+1}{2}, \frac{1}{2}]$, is analytic when $\text{Re}(z) \in (-\frac{n+1}{2}, \frac{1}{2})$, and that

- i) $\|T_z(f)\|_{L^\infty(\mathbb{R}^n)} \leq C_1(z)\|f\|_{L^1(\mathbb{R}^n)}$, when $\text{Re}(z) = \frac{1}{2}$,
- ii) $\|T_z(f)\|_{L^2(\mathbb{R}^n)} \leq C_2(z)\|f\|_{L^2(\mathbb{R}^n)}$, when $\text{Re}(z) = -\frac{n+1}{2}$,
- iii) $C_1(z)$ and $C_2(z)$ have at most exponential growth with respect to $\text{Im}(z)$.

Stein’s analytic interpolation theorem (see, e.g., [13]) will then imply that $T_z : L^{p_0}(\mathbb{R}^{n+l}) \rightarrow L^{p_1}(\mathbb{R}^{n+l})$ is a bounded operator when $z = 0$. Since $T_z(f) = f * \widehat{d\sigma}$, the Stein–Thomas observation (see the proof of Theorem 12) implies the conclusion of our theorem.

To prove (i), we observe that when $z = \frac{1}{2} + iy$, with $y \in \mathbb{R}$, then $K_z(\xi, \lambda) = \widehat{d\sigma}(\xi, \lambda) \det(A_\lambda)^{\frac{1}{2}+iy} \psi(\frac{1}{2} + iy)$ is bounded by $C_1(y) = \pi^{\frac{n}{2}} |\psi(\frac{1}{2} + iy)|$. By the Hausdorff–Young inequality we have that $\|f * K_{1+iy}\|_\infty \leq C_1(y)\|f\|_1$. (i) is then satisfied, and one can check, using Stirling’s formula, that $C_1(y)$ has at most exponential growth.

To prove (ii), it is enough to show that $|\widehat{K}_z|$ is a bounded function when $\text{Re}(z) = -\frac{n+1}{2}$. To compute the Fourier transform of $K_z(\xi, \lambda)$ with respect to ξ , $\mathcal{F}_\xi(K_z)(x, \lambda)$, we use again the formula (4.3) obtaining

$$\mathcal{F}_\xi(K_z)(x, \lambda) = \psi(z) |\det A_\lambda|^z e^{i(x, A_\lambda x)}. \tag{4.5}$$

Hence

$$\widehat{K}_z(x, x') = \psi(z) \int_{\mathbb{R}^l} e^{i((x', \lambda) + (x, A_\lambda x))} \det(A_\lambda)^z d\lambda, \tag{4.6}$$

where we have set $x' = (x_{n+1}, \dots, x_{n+l})$. We recall that the above identities hold in the distribution sense.

Since the phase of the above integral is a linear function of λ , we reduce to computing the Fourier transform of $\det(A_\lambda)^z$. We need the following lemma (see [12, p.48]).

Lemma 7.

Let $V_{\mathbb{R}}$ be the space of the real and symmetric matrices and let $V_i \subset V_{\mathbb{R}}$ be the subset of the matrices with i positive and $n - i$ negative eigenvalues. Let $G_i(z)$ be the distribution

$$G_i(z)(f) = \int_{V_i} f(Y) |\det(Y)|^2 dY \tag{4.7}$$

where dY is the standard Euclidean measure on $V_{\mathbb{R}}$. Then the distribution $G_i(z)$, viewed as a function of z , has analytic continuation to a meromorphic function in the whole complex plane satisfying

$$\widehat{G}_i(z) = \psi^{-1}(z) \sum_{j=0}^n c_{i,j}(z) G_j \left(-z - \frac{n+1}{2} \right), \tag{4.8}$$

where ψ is as in (4.4) and the $C_{i,j}(z)$ are bounded coefficients.

From the above formula we deduce that up to bounded constants,

$$\mathcal{F}(\det(A_\lambda)^z)(\eta) = \psi^{-1}(z) \det(A_\eta)^{-z - \frac{n+1}{2}}, \tag{4.9}$$

where the above formula holds in the distribution sense. Since $\text{Re}(z) = \frac{n+1}{2}$, the above is a bounded function of η . This shows that \widehat{K}_z is a bounded function of (x, x') , and completes the proof of the theorem. \square

Proof of Theorem 14. Let

$$\widehat{d\sigma}(\xi, \lambda) = \int_{\mathbb{R}^n} e^{i((x,\xi)+\lambda_1\phi_1(x)+\lambda_l\phi_l(x))} \chi(x) dx, \tag{4.10}$$

where χ is a smooth cutoff function. By Theorem 6 we can write

$$\widehat{d\sigma}(\xi, \lambda) = \widetilde{B}(\xi, \lambda) \tag{4.11}$$

where $|\widetilde{B}(\xi, \lambda)| \leq \int_{S^{n-1}} \frac{d\omega}{\left| \sum_{j=1}^l \phi_j(\omega)\lambda_j \right|^{\frac{n}{m}}}$.

Let

$$B(\lambda) = \int_{S^{n-1}} \frac{d\omega}{\left| \sum_{j=1}^l \phi_j(\omega)\lambda_j \right|^{\frac{n}{m}}}, \tag{4.12}$$

$$K_z(\xi, \lambda) = \widehat{d\sigma}(\xi, \lambda) B^z(\lambda) \psi(z),$$

with

$$\psi(z) = \Gamma^{-1}\left(-\frac{zn}{2m} + \frac{1}{2}\right) \Gamma^{-1}\left(-\frac{nz}{m} + l\right),$$

where Γ is the standard Gamma function. Let $T_z(f)(\xi, \lambda) = (f * K_z)(\xi, \lambda)$.

We will prove that T_z is a continuous family of operators when $\text{Re}(z) \in [-1, \frac{lm}{n}]$, is analytic when $\text{Re}(z) \in (-1, \frac{lm}{n})$, and that

- i) $\|T_z(f)\|_{L^\infty(\mathbb{R}^n)} \leq C_1(z) \|f\|_{L^1(\mathbb{R}^n)}$, when $\text{Re}(z) = -1$,
- ii) $\|T_z(f)\|_{L^2(\mathbb{R}^n)} \leq C_2(z) \|f\|_{L^2(\mathbb{R}^n)}$, when $\text{Re}(z) = \frac{ml}{n}$,
- iii) $C_1(z)$ and $C_2(z)$ have at most exponential growth with respect to $\text{Im}(z)$.

Stein’s analytic interpolation theorem (see, e.g., [13]) will then imply that $T_z : L^{p_0}(\mathbb{R}^{n+l}) \rightarrow L^{p_0'}(\mathbb{R}^{n+l})$ is a bounded operator when $z = 0$. Since $T_z(f) = f * \widehat{d\sigma}$, the Stein–Thomas observation (see the proof of Theorem 12) implies the conclusion of our theorem.

To prove the estimate (i) we observe that when $z = -1 + iy$, with $y \in \mathbb{R}$, then $K_z(\xi, \lambda) = \widehat{d\sigma}(\xi, \lambda) B^{iy-1}(\lambda) \psi(-1 + iy)$ is bounded by $C_1(y) = C|\psi_1(-1 + iy)|$. By the Hausdorff–Young inequality we have that $\|f * K_{-1+iy}\|_\infty \leq C_1(y) \|f\|_1$. The estimate (i) is then satisfied, and one can check using Stirling’s formula that $C_1(y)$ has at most exponential growth.

To prove the estimate (ii), it is enough to show that \widehat{K}_z is a bounded function when $\text{Re}(z) = \frac{ml}{n}$. Since $d\sigma$ is a finite measure, it is enough to prove that $\psi(z) \widehat{B}^z(\lambda)$ is bounded.

Let $X(\omega)$ be the vector defined by the equation $\sum_{j=1}^l \lambda_j \phi_j(\omega) = \langle \lambda, X(\omega) \rangle$. Then

$$\widehat{B}_z(y) = \psi(z) \int_{\mathbb{R}^l} \left(\int_{S^{n-1}} \frac{d\omega}{|\langle \lambda, X(\omega) \rangle|^{\frac{n}{m}}} \right)^z e^{i\langle \lambda, y \rangle} d\lambda. \tag{4.13}$$

Since $\widehat{B}_z(y)$ is homogeneous of degree zero with respect to y , we can assume $|y| = 1$.

In polar coordinates with respect to λ , with $\lambda = r\eta$, we have

$$\widehat{B}_z(y) = \psi(z) \int_0^{+\infty} r^{-\frac{n}{m}z+l-1} \int_{S^{l-1}} \left(\int_{S^{n-1}} \frac{e^{\frac{i}{z}r\langle \eta, y \rangle} d\omega}{|\langle \eta, X(\omega) \rangle|^{\frac{n}{m}}} \right)^z d\eta dr. \tag{4.14}$$

Let $\chi(r) \in C^\infty(\mathbb{R})$ be such that $\chi(r) \equiv 1$, when $r \in (0, 1)$, and $\chi(r) \equiv 0$ when $r \in (2, +\infty)$. Let

$$\widehat{B}_{z,M}(y) = \psi(z) \int_0^{+\infty} r^{-\frac{n}{m}z+l-1} \chi\left(\frac{r}{M}\right) \int_{S^{l-1}} \left(\int_{S^{n-1}} \frac{e^{\frac{i}{z}r\langle \eta, y \rangle} d\omega}{|\langle \eta, X(\omega) \rangle|^{\frac{n}{m}}} \right)^z d\eta dr. \tag{4.15}$$

We will prove that $\widehat{B_{z,M}}$ is bounded by a constant C independent of M and that $\widehat{B_{z,M}} \rightarrow \widehat{B_z}$ in distribution sense as $M \rightarrow \infty$. From the above it follows that $\widehat{B_z}$ is bounded. Indeed, since the balls are sequentially compact in the weak $*$ topology in $L^\infty(\mathbb{R}^l)$, there exists a sequence $\{\widehat{B_{z,M_j}}\}_{j \in \mathbb{N}} \subset \{\widehat{B_{z,M}}\}$ which converges to a bounded function in the weak $*$ topology of L^∞ , and hence converges also in a distribution sense. Consequently, $\widehat{B_z} = \lim_{j \rightarrow +\infty} \widehat{B_{z,M_j}}$, which is a bounded function.

Recalling that by assumption the vector $X(\omega)$ is never zero on S^{n-1} , we can construct an orthogonal matrix A_ω with the property that $A_\omega \frac{X(\omega)}{|X(\omega)|} = (1, 0, \dots, 0)$. The first row of A_ω is $\frac{X(\omega)}{|X(\omega)|}$ and the other rows are a set of $l - 1$ vectors which, together with $\frac{X(\omega)}{|X(\omega)|}$, determine an orthonormal basis of \mathbb{R}^l for every $\omega \in S^{n-1}$.

We make the change of variables $\eta \rightarrow A_\omega^t \eta$ in the expression for $\widehat{B_{z,M}}$. Since A_ω is orthogonal, the change of variables maps S^{l-1} into itself and the determinant of the Jacobian matrix of the transformation is 1. We obtain

$$\begin{aligned} \widehat{B_{z,M}}(y) &= \psi(z) \int_0^{+\infty} r^{-\frac{n}{m}z+l-1} \chi\left(\frac{r}{M}\right) \int_{S^{l-1}} \left(\int_{S^{n-1}} \frac{|X(\omega)|^{\frac{n}{m}}}{|\langle \eta, A_\omega X(\omega) \rangle|^{\frac{n}{m}}} e^{\frac{i}{z}r\langle \eta, A_\omega y \rangle} d\omega \right)^z d\eta dr \\ &= \psi(z) \int_0^{+\infty} r^{-\frac{n}{m}z+l-1} \chi\left(\frac{r}{M}\right) \int_{S^{l-1}} \frac{1}{|\eta_1|^{\frac{nz}{m}}} \left(\int_{S^{n-1}} |X(\omega)|^{\frac{n}{m}} e^{\frac{i}{z}r\langle \eta, A_\omega y \rangle} d\omega \right)^z d\eta dr. \end{aligned} \tag{4.16}$$

The integral with respect to ω is a continuous function of r, η, y , say $F(r, \eta, y)$. In particular $F(0, \eta, y) = \int_{S^{n-1}} |X(\omega)|^{\frac{n}{m}} d\omega < \infty$.

Since $\text{Re}(z) \geq l$ by assumption, we can check that we can compute at least $l - 1$ derivatives of $F(r, \eta, y)^z$ with respect to r and η . Then

$$\begin{aligned} \widehat{B_{z,M}}(y) &= \psi(z) \int_0^{+\infty} r^{-\frac{n}{m}z+l-1} \chi\left(\frac{r}{M}\right) \int_{S^{l-1}} \frac{1}{|\eta_1|^{\frac{nz}{m}}} F^z(r, \eta, y) dr d\eta \\ &= \psi(z) \int_0^{+\infty} r^{-\frac{n}{m}z+l-1} \chi\left(\frac{r}{M}\right) \int_{-1}^1 \int_{\{|\eta'|=(1-|\eta_1|^2)^{\frac{1}{2}}\}} \frac{F^z(r, \eta, y)}{|\eta_1|^{\frac{nz}{m}}} dr d\eta_1 d\eta'. \end{aligned} \tag{4.17}$$

Here we set $\eta = (\eta_1, \eta')$, and we let $d\eta'$ be the measure on the $(l - 2)$ -dimensional sphere $\{|\eta'| = (1 - |\eta_1|^2)^{\frac{1}{2}}\}$. If we make a change of variables in the above integral letting $\eta' \rightarrow (1 - |\eta_1|^2)^{\frac{1}{2}} \eta'$, we get

$$\begin{aligned} \widehat{B_{z,M}}(y) &= \psi(z) \int_0^{+\infty} r^{-\frac{n}{m}z+l-1} \chi\left(\frac{r}{M}\right) \int_{-1}^1 \frac{(1 - |\eta_1|^2)^{\frac{l-2}{2}}}{|\eta_1|^{\frac{nz}{m}}} \\ &\quad \times \int_{S^{l-2}} F^z\left(r, \eta_1, \eta' (1 - |\eta_1|^2)^{\frac{1}{2}}, y\right) d\eta_1 d\eta'. \end{aligned} \tag{4.18}$$

We recall that the distribution $\frac{r^{-\frac{n}{m}z+l-1}}{\Gamma(l-\frac{nz}{m})}$ is an entire function of z which coincides with the Dirac distribution δ_0 when $z = \frac{ml}{n}$, and that the distribution $\frac{|\eta_1|^{\frac{-nz}{m}}}{\Gamma(-\frac{zn}{m}+\frac{1}{2})}$ is an entire function of z which coincides with the $(k - 1)$ th derivative of the Dirac distribution δ_0 when $l = 2k + 1$ and $z = \frac{ml}{n}$. With that in mind we consider a smooth function $\rho(t)$ which is $\equiv 1$ when $t \in (0, \frac{1}{4})$ and is $\equiv 0$ when

$t \in (\frac{3}{4}, 1)$, and we write

$$\begin{aligned} \widehat{B_{z,M}}(y) &= \psi(z) \left(\int_{-1}^1 \rho(\eta_1^2) \cdots + \int_{-1}^1 (1 - \rho(\eta_1^2)) \cdots \right) \\ &= \psi(z) I_1(z, M, y) + \psi(z) I_2(z, M, y). \end{aligned} \tag{4.19}$$

Since

$$(1 - \rho(\eta_1^2)) (1 - \eta_1^2)^{\frac{l-2}{2}} \int_{S^{l-2}} F^z \left(r, \eta_1, \eta' (1 - |\eta_1|^2)^{\frac{1}{2}}, y \right) d\eta' d\eta_1 = F_2(r, \eta_1) \tag{4.20}$$

is bounded and continuous with respect to r and η_1 , we can write

$$\psi(z) I_2(z, M, y) = \Gamma^{-1} \left(-\frac{zn}{2m} + \frac{1}{2} \right) \chi \left(\frac{r}{M} \right) \int_{-1}^1 F_2(r, \eta_1) d\eta_1 \Big|_{r=0}. \tag{4.21}$$

Thus $|\psi(z) I_2(z, M, y)|$ is bounded by a constant which does not depend on M and has at most exponential growth with respect to $\text{Im}(z)$.

We shall now estimate $\psi(z) I_1(z, M, y)$. By our previous observations, the function

$$\rho(\eta_1^2) (1 - \eta_1^2)^{\frac{l-2}{2}} \int_{S^{l-2}} F^z \left(r, \eta_1, \eta' (1 - |\eta_1|^2)^{\frac{1}{2}}, y \right) d\eta' d\eta_1 = F_1(r, \eta_1) \tag{4.22}$$

can be differentiated $l - 1$ times with respect to η_1 and its derivatives are continuous functions of r . Then, if $l = 2k + 1$,

$$\psi(z) I_1(z, M, y) = \frac{d^{k-1}}{d\eta_1} \left(\chi \left(\frac{r}{M} \right) F_1(r, \eta_1) \right) \Big|_{\eta_1=r=0}. \tag{4.23}$$

If $l \neq 2k + 1$, and if $h = \lceil \frac{\text{Re}(l)}{2} \rceil$ we use the formula (3) in [4, pg.51], obtaining:

$$\begin{aligned} I_1(z, M, y) &= \chi \left(\frac{r}{M} \right) \int_0^{+\infty} \eta_1^{-\frac{nz}{m}} \left(F_1(r, \eta_1) + F_1(r, -\eta_1) \right. \\ &\quad \left. - 2 \left(F_1(r, 0) + \frac{\eta_1^2}{2!} \frac{\partial^2}{\partial \eta_1} F_1(r, 0) + \cdots + \frac{\eta_1^{2h}}{(2h)!} \frac{\partial^{2h}}{\partial \eta_1} F_1(r, 0) \right) \right) d\eta_1 \Big|_{r=0} \end{aligned} \tag{4.24}$$

Thus, $|\psi(z) I_1(z, M, y)|$ is bounded by a constant which does not depend on M . This shows that $\widehat{K_{z,M}}$ is bounded by a uniform constant. An easy adaptation of the above argument shows that $\widehat{K_{z,M}}$ converges to $\widehat{K_z}$ in the distribution sense as $M \rightarrow \infty$.

This completes the proof of the theorem. \square

Acknowledgments

The authors wish to thank Eric Sawyer for financial support provided through his NSERC grant. The second author wishes to thank Professor Sawyer for numerous helpful conversations over the last two years, which had a profound influence on this work.

The authors wish to thank Fulvio Ricci for suggesting Theorem 13, and for related helpful suggestions.

References

- [1] Christ, M. Restriction of the Fourier transform to submanifolds of low codimension, Ph.D Thesis, University of Chicago, Illinois.
- [2] De Carli, L. and Iosevich, A. (1995). A restriction theorem for flat manifolds of codimension two, *Illinois M.J.* (to appear).
- [3] Greenleaf, A. (1982). Principal curvature and harmonic analysis, *Indiana Math. J.*, **30**, 519–537.
- [4] Gel'fand, I.M. and Shilov, G.E. (1964). Generalized Functions Vol. 1, Academic Press.
- [5] Hironaka, H. (1964). Resolution of singularities of an algebraic variety over a field of characteristic 0, *Annal. Math.*, **79**, 109–203, 205–326.
- [6] Iosevich, A. (1994). Maximal operators associated to families of flat curves in the plane, *Duke Math. J.*, **76**.
- [7] Iosevich, A. (1995). Maximal averages over homogeneous hypersurfaces in \mathbb{R}^3 , *Forum Mathematicum* (to appear).
- [8] Iosevich, A. and Sawyer, E. (1995). Oscillatory integrals and maximal averaging operators associated to homogeneous hypersurfaces, *Duke Math. J.* (to appear).
- [9] Littman, W. (1963). Fourier transforms of surface carried measures and differentiability of surface averages, *Bull. Am. Math. Soc.*, **69**, 766–770.
- [10] Rudin, W. (1973). *Functional analysis*, McGraw-Hill, New York.
- [11] Prestini, E. (1979). Restriction theorems for the Fourier transform to some manifolds in \mathbb{R}^n , *Proc. Symp. Pure Math.*, XXXV Part 1.
- [12] Shintani, T. (1975). On zeta function associated with the vector space of quadratic forms, *J. F. Sc. U. Tokyo*, 25–65.
- [13] Sogge, C.D. (1991). *Fourier Integrals in Classical Analysis*, Oxford University Press.
- [14] Stein, E.M. (1993). *Harmonic Analysis*, Princeton University Press, **43**, Princeton, NJ.

Received May 3, 1996

University of Naples, Naples, Italy

Wright State University, Dayton, Ohio

e-mail: iosevich@zara.math.wright.edu