

## SOME THEOREMS ON CONVEX POLYGONS

BY  
E. ALTMAN

**ABSTRACT.** In this paper diagonals of various orders in a (strict) convex polygon  $P_n$  are considered. The sums of lengths of diagonals of the same order are studied. A relationship between the number of consecutive diagonals which do not intersect a given maximal diagonal and lie on one side of it and the order of the smallest diagonal among them is established. Finally a new proof of a conjecture of P. Erdos, considered already in [1], is given.

**I. Notation and nomenclature.** (1) A plane convex  $n$ -sided polygon will be denoted by

$$P_n = A_1A_2 \cdots A_n$$

( $A_i$  are the vertices). Let  $j \leq \left\lfloor \frac{n}{2} \right\rfloor$ . A diagonal  $A_iA_{i+j}$  (where  $i+j$  is taken mod  $n$ ), i.e. a diagonal cutting off  $j$  sides of the polygon, is said to be a diagonal of order  $j$ ; the sides of the polygon are diagonals of the 1-st order. Clearly,  $P_n$  contains diagonals of  $\left\lfloor \frac{n}{2} \right\rfloor$  distinct orders.

(2) The sum of lengths of the diagonals of the  $j$ -th order will be denoted by

$$u_j = \sum_{i=1}^n A_iA_{i+j}.$$

For  $n=2N$ , the corresponding sum

$$u_N = \sum_{i=1}^N A_iA_{i+N}$$

includes every diagonal of the  $N$ -th order twice.

(3) The various lengths of the diagonals of  $P_n$  will be denoted by

$$d_1 > d_2 > \cdots$$

A diagonal of length  $d_x$  is said to be of the  $x$ -th degree.

**II. THEOREM 1.** If  $0 < q < p \leq \left\lfloor \frac{n}{2} \right\rfloor$ , then

$$u_q < u_p.$$

---

Received by the editors March 9, 1970 and, in revised form, January 25, 1971.




Figure 1.

**Proof.** (1) Let  $N = \lfloor \frac{n}{2} \rfloor$ . We have to prove that

$$u_{k+1} > u_k$$

for  $k = 1, 2, \dots, N-1$ .

(2) We first prove that  $u_N > u_{N-1}$ . Two cases will be distinguished:

(a)  $n$  is even:  $n = 2N$ ,

Consider a convex quadrilateral (Fig. 1)

$$A_i A_{i+1} A_{i+N} A_{i+N+1}$$

In every convex quadrilateral, the sum of two opposite sides is smaller than the sum of the two diagonals. Hence:

$$(A) \quad A_i A_{i+N} + A_{i+1} A_{i+N+1} > A_{i+N+1} A_i + A_{i+1} A_{i+N}$$

The diagonals of this quadrilateral are diagonals of the  $N$ -th order in  $P_n$ , while the sides appearing in the inequality (A) are diagonals of the  $(N-1)$ -th order in  $P_n$ . Summation of (A) for  $i = 1, \dots, n$  yields

$$(\bar{A}) \quad \begin{aligned} 2u_N &> 2u_{N-1} \\ u_N &> u_{N-1} \end{aligned}$$

(b)  $n$  is odd:  $n = 2N + 1$ .

Consider a convex quadrilateral (Fig. 2)

$$A_i A_{i+1} A_{i+N} A_{i+N+1}$$

Here we have

$$(B) \quad A_i A_{i+N} + A_{i+1} A_{i+N+1} > A_{i+N+1} A_i + A_{i+1} A_{i+N}$$




Figure 2.

$A_{i+N+1}A_i$ ,  $A_iA_{i+N}$  and  $A_{i+1}A_{i+N+1}$  are diagonals of the  $N$ -th order in  $P_n$ , while  $A_{i+1}A_{i+N}$  is a diagonal of the  $(N-1)$ -th order in  $P_n$ . Summation of (B) for  $i=1, \dots, n$  yields

$$\begin{aligned} \text{(B)} \quad & 2u_N > u_N + u_{N-1} \\ & u_N > u_{N-1} \end{aligned}$$

(3) We make now an induction assumption that the theorem holds for  $k$ , i.e.

$$u_{k+1} > u_k$$

and prove that it holds for  $k-1$ , i.e.

$$u_k > u_{k-1}$$

To this end consider the convex quadrilateral (Fig. 3)  $A_iA_{i+1}A_{i+k}A_{i+k+1}$ .

We have

$$\text{(C)} \quad A_iA_{i+k} + A_{i+1}A_{i+k+1} > A_iA_{i+k+1} + A_{i+1}A_{i+k}$$

$A_{i+1}A_{i+k+1}$  and  $A_iA_{i+k}$  are diagonals of the  $k$ -th order in  $P_n$ ,  $A_iA_{i+k+1}$  is a diagonal of the  $(k+1)$ -th order, and  $A_{i+1}A_{i+k}$  a diagonal of the  $(k-1)$ -th order. Summation




Figure 3.

of (C) for  $i=1, \dots, n$  yields

$$(C) \quad 2u_k > u_{k+1} + u_{k-1}.$$

By our assumption

$$u_{k+1} > u_k$$

hence

$$2u_k > u_k + u_{k-1}$$

$$u_k > u_{k-1}$$

and Theorem 1 is hereby proved.

(4) REMARK 1. By ( $\bar{C}$ ):

$$2u_k > u_{k+1} + u_{k-1}$$

hence

$$u_k > \frac{u_{k+1} + u_{k-1}}{2} \quad \left( 1 < k < \left\lfloor \frac{n}{2} \right\rfloor \right),$$

i.e., the sequence  $u_1, u_2, \dots, u_N$  of the sums of the diagonals of consecutive orders increases at a slower rate than an arithmetic progression.

REMARK 2. By inscribing a convex polygon in  $P_n$ , several relations are obtainable between the sums  $u_j$ , in the same way as in the preceding proof. For example, by inscribing a triangle  $A_i A_{i+k} A_{i+k+l}$  ( $k+l \leq \left\lfloor \frac{n}{2} \right\rfloor$ ) we obtain

$$A_i A_{i+k} + A_{i+k} A_{i+k+l} > A_i A_{i+k+l}$$

and summation of this inequality for  $i=1, 2, \dots, n$  yields

$$u_k + u_l > u_{k+l} \quad \left( k+l \leq \left\lfloor \frac{n}{2} \right\rfloor \right)$$

REMARK 3. Let  $B_1 B_2 \dots B_n B_{n+1}$  be a  $n$ -sided polygonal line, consisting of segments which are parallel and equal to the consecutive diagonals of the  $k$ -th order in  $P_n$

$$B_i B_{i+1} \uparrow \uparrow A_i A_{i+k}$$

$$B_i B_{i+1} = A_i A_{i+k}$$

It is easily seen, by using vectors, that  $B_1 B_2 \dots B_n B_{n+1}$  is a closed polygon, i.e.  $B_{n+1} = B_1$ . It is called the  $k$ -th derivative of the polygon  $P_n$  and is denoted by  $P_n^{(k)}$ . This polygon is convex when  $P_n$  is convex. The  $l$ -th derivative of the polygon  $P_n^{(k)}$  will be denoted by  $P_n^{(kl)}$ . The commutativity  $P_n^{(kl)} = P_n^{(lk)}$  can be proven by using vectors. Examples of derivatives are shown in Fig. 4<sub>1,2,3</sub>.




Figure 4\_1




Figure 4\_2




Figure 4\_3

III. (1) Let  $P_n = A_1A_2 \cdots A_n$  be a plane convex  $n$ -sided polygon, with  $A_1A_n$  as maximal diagonal (i.e. not smaller than any diagonal in  $P_n$ ), in other words  $\overline{A_1A_n} = d_1$  (see §I (3)).

Any diagonal  $A_kA_l$ ,  $1 < k < l < n$ , is said to be parallel to  $A_1A_n$ , or briefly a parallel.




Figure 5.

Two parallels  $A_sA_t$  and  $A_uA_v$  are said to be consecutive if

$$s < t < v$$

$$s < u < v$$

(2) LEMMA 1<sup>(1)</sup>. *If  $A_sA_t$  and  $A_uA_v$ , are two consecutive parallels, at least one of them is smaller than  $A_sA_v$ .*

**Proof**<sup>(2)</sup>. Suppose  $l$  a support line of the convex hull of  $A_1, A_2, \dots, A_n$  (Fig. 5) which is parallel to  $A_sA_v$  and contains  $A_1$ . There is, then, a point  $P$ , on the segment  $A_1A_t$  such that line  $PA_n$  is parallel to  $l$ .

Let  $A_1A_t \cap A_sA_v = Q$  and  $A_tA_n \cap A_sA_v = R$ . Since  $\|A_sA_t\| \geq \|A_sA_v\|$  it follows from the triangle inequality that  $\|A_tQ\| > \|QR\|$ . From linearity we have  $\|PA_t\| > \|PA_n\|$  and again the triangle inequality implies that  $\|A_1A_t\| > \|A_1A_n\|$ . This contradicts the fact that  $A_1A_n$  is a diameter of  $P_n$ . Clearly, there is no assertion to make about the relation of  $A_uA_v$  to  $A_sA_v$ .

The obtained contradiction proves the lemma.

A sequence of parallels

$$A_{s_1}A_{t_1}, A_{s_2}A_{t_2}, \dots, A_{s_k}A_{t_k}$$

where for  $i=1, \dots, k-1$  the parallels  $A_{s_i}A_{t_i}$  and  $A_{s_{i+1}}A_{t_{i+1}}$  are consecutive, will be called a chain of consecutive parallels (Fig. 6).

(3) THEOREM 2. *Given a chain of  $f$  consecutive parallels in a strict convex polygon  $P_n$ . Let  $x$  be the degree of the smallest diagonal in the chain (see [§I (3)]). Then*

<sup>(1)</sup><sup>(2)</sup> The author's original proof of Lemma 1 was based on separate case arguments ( $u \not\leq t$ ). The proof below proposed by the referee, makes these case arguments superfluous.




Figure 6.

(a) If  $A_1A_n$  is the only diagonal of the 1-st degree in  $P_n$ , then

$$f \leq x-2$$

(b) If  $A_1A_n$  is not the only diagonal of the 1-st degree in  $P_n$ , then

$$f \leq x-1$$

Thus there are no parallels of the 1-st degree, no two consecutive parallels of the 2-nd degree, no three consecutive parallels of the 3-rd degree, etc. In case (a), there are no parallels of the 2-nd degree, no two consecutive parallels of the 3-rd degree, etc.

Proof by induction on  $f$ .

(4) **Proof for  $f=1$ .** We have to prove that there is no parallel of the 1-st degree, and that in case (a) there is even no parallel of the 2-nd degree.

Let  $A_iA_j$  be a parallel (Fig. 7) and let  $x$  be its degree. Consider the convex quadrilateral

$$A_1A_iA_jA_n.$$




Figure 7.




Figure 8.

The sum of two opposite sides is smaller than that of the diagonals, i.e.

$$A_1A_j + A_iA_n > A_1A_n + A_iA_j = d_1 + d_x$$

so that either  $A_1A_j$  or  $A_iA_n$  must have a length  $d_y$  exceeding  $d_x$ . Hence  $x \geq 2$ . In case (a),  $A_1A_n$  is the only diagonal of length  $d_1$ , hence

$$d_x < d_y < d_1$$

i.e.,

$$x > y > 1$$

or  $x \geq 3$ . The theorem is hereby proved for  $f=1$ .

(5) Now assume that the theorem holds for a chain of  $k$  consecutive parallels. Let a chain  $C$  of  $k+1$  parallels:

$$A_{s_1}A_{t_1}, A_{s_2}A_{t_2}, \dots, A_{s_{k+1}}A_{t_{k+1}}$$

be given, and let  $x$  be the degree of the smallest diagonal in  $C$ . We inscribe in  $C$  a chain  $C'$  of  $k$  consecutive parallels (Fig. 8) by connecting the origin of every diagonal of  $C$  (except the last), to the end of the next one. The chain  $C'$  will thus consist of the parallels

$$A_{s_1}A_{t_2}, A_{s_2}A_{t_3}, \dots, A_{s_k}A_{t_{k+1}}$$

which are consecutive, as is easily shown.

By Lemma 1, the diagonal  $A_{s_i}A_{t_{i+1}}$  of  $C'$  exceeds one of the diagonals  $A_{s_i}A_{t_i}$ ,  $A_{s_{i+1}}A_{t_{i+1}}$  of  $C$ . The length of any diagonal of  $C'$  thus exceeds  $d_x$ , hence the degree of the smallest diagonal in  $C'$  is at most  $x-1$ . By the assumption that the theorem holds for  $f=k$ , we have: In case (a):  $k \leq (x-1)-2 = x-3$ . Hence

$$k+1 \leq x-2$$

In case (b):

$$k \leq (x-1)-1 = x-2$$






Figure 9.

hence

$$k+1 \leq x-1.$$

So the theorem holds for  $f=k+1$  as well. Theorem 2 is hereby proved.

(6) REMARK. Existence of a parallel  $A_kA_i$  of the 2-nd degree implies that (Fig. 9):

$$A_1A_i + A_kA_n > A_kA_i + A_1A_n = d_1 + d_2.$$

This is possible only if

$$A_1A_i = A_kA_n = d_1.$$

Thus existence of a parallel of the 2-nd degree is possible only if another diagonal of length  $d_1$  originates from each end point of  $A_1A_n$ .

By the same induction as in (5), we conclude that the existence of a chain of consecutive parallels, satisfying

$$f = x-1$$

is possible only if another diagonal of length  $d_1$  originates from each end point of  $A_1A_n$ .

(7) COROLLARY TO THEOREM 2. *If a chain  $A_iA_{i+1} \cdots A_{i+f}$  of  $f$  consecutive sides of a plane convex  $n$ -sided polygon  $P_n = A_1A_2 \cdots A_n$  is not cut by a maximal diagonal  $A_kA_l$  of  $P_n$  (Fig. 10), and the degree of the smallest side of the chain is  $x$ , then by Theorem 2 it follows that:*

(a) *If the maximal diagonal is the only diagonal of length  $d_1$ , then*

$$f \leq x-2.$$

(b) *If there are other diagonals of length  $d_1$ , then*

$$f \leq x-1.$$

Moreover,  $f=x-1$  is possible only if another diagonal of length  $d_1$  originates from each end point of the maximal diagonal.




Figure 10.

**IV. A new solution to a problem by P. Erdos.** In [1], the author proved the following conjecture by P. Erdos:

**THEOREM 3.** *In every plane strictly convex  $n$ -sided polygon  $P_n$  there are at least  $\left\lfloor \frac{n}{2} \right\rfloor$  different distances between various pairs of vertices.<sup>(3)</sup>*

Here a proof of this theorem will be given based on Corollary (7).

**Proof.** Two cases will be distinguished:

(a) There are no two maximal diagonals with a common end point (Fig. 11):

Let  $A_i A_m$  be a maximal diagonal. It must cut off at least  $\left\lfloor \frac{n}{2} \right\rfloor$  sides of  $P_n$ ; hence there is a chain of  $\left\lfloor \frac{n}{2} \right\rfloor - 2$  consecutive sides of  $P_n$ , which is not cut by  $A_i A_m$ .

Let  $x$  be the degree of the smallest side of this chain. There is no other maximal diagonal originating from either end point of  $A_i A_m$ . Hence, by Corollary (7).

$$x - 2 \geq \left\lfloor \frac{n}{2} \right\rfloor - 2$$

$$x \geq \left\lfloor \frac{n}{2} \right\rfloor.$$

<sup>(3)</sup> Define the distance set of the vertex set  $\{P_1, \dots, P_n\}$  of points in a real normed linear space by:  $\{\|P_i P_j\| \mid 1 \leq i < j \leq n\}$ .

With the referee's proof of Lemma 2, Theorem 3 can read: The distance set of the vertex set of a plane strictly convex polygon of  $n$  sides in a strictly convex real normed linear space consists of at least  $\left\lfloor \frac{n}{2} \right\rfloor$  positive numbers.




Figure 11.

The polygon has, therefore, a diagonal whose degree is not less than  $\lceil \frac{n}{2} \rceil$ .

(b) There are two maximal diagonals with a common end point (Fig. 12). Clearly, one of them (denote by  $A_i A_m$ ) must cut off at least  $\lceil \frac{n}{2} \rceil + 1$  sides; hence there is a chain of  $\lceil \frac{n}{2} \rceil - 1$  sides of  $P_n$ , which is not cut by  $A_i A_m$ . For this chain




Figure 12.

we have, by Corollary (7),

$$x-1 \geq \left[ \frac{n}{2} \right] - 1$$

$$x \geq \left[ \frac{n}{2} \right].$$

Hence the polygon comprises at least  $\left[ \frac{n}{2} \right]$  different distances. The conjecture is hereby proved.

ACKNOWLEDGMENT. The author wishes to thank the referee for his elegant proof of the key Lemma 1 and for other important remarks.

#### REFERENCE

1. F. Altman, *On a Problem by P. Erdos*, Amer. Math. Monthly, **70** (1963), 148-157.

TECHNION-ISRAEL INSTITUTE OF TECHNOLOGY,  
HAIFA, ISRAEL.