

<https://helda.helsinki.fi>

Characterization of *Leuconostoc gasicomitatum* sp. nov.,
Associated with Spoiled Raw Tomato-Marinaded Broiler Meat
Strips Packaged under Modified-Atmosphere Conditions

Björkroth, Johanna

American Society for Microbiology (ASM)
2000

A p p l i e d a n d E n v i r o n m e n t a l M i c r o b i o l o g y . 2 0

<http://hdl.handle.net/1975/525>

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Characterization of *Leuconostoc gasicomitatum* sp. nov., Associated with Spoiled Raw Tomato-Marinaded Broiler Meat Strips Packaged under Modified-Atmosphere Conditions

K. JOHANNA BJORKROTH,^{1*} ROLF GEISEN,² ULRICH SCHILLINGER,² NORBERT WEISS,³ PAUL DE VOS,⁴ WILHELM H. HOLZAPFEL,² HANNU J. KORKEALA,¹ AND PETER VANDAMME⁴

Department of Food and Environmental Hygiene, University of Helsinki,¹ Helsinki, Finland and Toxicology, Federal Research Centre for Nutrition and Consumer Protection,² Karlsruhe, Germany; Collection of Microorganisms and Cell Cultures, Braunschweig,³ Germany; and Laboratory of Microbiology, University of Ghent,⁴ Ghent, Belgium

Received 15 February 2000/Accepted 7 June 2000

Lactic acid bacteria (LAB) associated with gaseous spoilage of modified-atmosphere-packaged, raw, marinated broiler meat strips were identified on the basis of a restriction fragment length polymorphism (RFLP) (ribotyping) database containing DNAs coding for 16S and 23S rRNAs (rDNAs). A mixed LAB population dominated by *Leuconostoc* species resembling *Leuconostoc gelidum* caused the spoilage of the product. *Lactobacillus*, *Lactobacillus curvatus* gram-positive rod phenotypically similar to heterofermentative *Lactobacillus* species were the other main organisms detected. An increase in pH together with extreme bulging of packages suggested a rare LAB spoilage type called "protein swell." This spoilage characterized by excessive production of gas due to amino acid decarboxylation, and the rise in pH is associated with the subsequent deamination of amino acids. Protein swell has not previously been associated with meat product. A polyphasic approach, including classical phenotyping, whole-cell protein electrophoresis, and 23S rDNA RFLP, 16S rDNA sequence analysis, and DNA-DNA reassociation analysis, was used for the identification of the dominant *Leuconostoc* species. In addition to the RFLP analysis, phenotyping, whole-cell protein analysis, and 16S rDNA sequence homology indicated that the most similar to the spoilage-associated species. The two spoilage strains studied possessed 98.8 and 99.0% 16S rDNA sequence homology with the *gelidum* type strain. DNA-DNA reassociation, however, clearly distinguished the two species. The same strains showed only 22 and 34% hybridization with the *gelidum* strain. These results warrant a separate species status, and we propose the name *Leuconostoc gasicomitatum* sp. nov. for this spoilage-associated *Leuconostoc* species.

Lactic acid bacteria (LAB) are the dominant spoilage organisms, which have also been set as the retail shelf life. In vacuum or modified-atmosphere (MA)-packaged meat products (1, 2, 8, 10, 23, 31). Spoilage is mainly caused by bulging due to gas formation 5 days after packaging (3, 4, 26, 28; W. H. Holzappel and E. S. Geisen, Abstr. 32nd Eur. Meet. Meat Res. Workers, p. 26, 1986) that time, only this tomato-marinaded product was spoiled by *Leuconostoc* (7, 14, 28, 43, 49) species. The activities of these organisms at stationary phase produce the compounds associated with sensory spoilage (22). Depending on the product, this quality deterioration usually starts after packaging, and it is manifested mainly as fermentation and sour or cheesy off odors and/or off tastes. Provided the shelf life of the product has been estimated correctly, spoilage changes do not occur before the sell-by day. However, in poultry products, our study set out to characterize of potent spoilage LAB and/or poor production and to identify these spoilage LAB to the species level. Severe quality faults (5, 7, 26, 27) have occurred in product recalls.

The consumption of marinated, ready-to-cook, raw meat products has been increasing in Europe. As easily consumed and low-fat food, they are favored by many consumers. In this study, we describe and characterize an unusual spoilage type of MA-packaged, tomato-marinaded, raw broiler meat strips. The product was manufactured at a modern large-scale processing plant, and normally, good quality was maintained at the time of spoilage. Based on these results, the dominant species of the spoilage population was considered novel, and we propose the name *Leuconostoc gasicomitatum* sp. nov. for it.

* Corresponding author. Mailing address: Department of Food and Environmental Hygiene, Faculty of Veterinary Medicine, University of Helsinki, P.O. Box 57, FIN-00014 Helsinki University of Helsinki, Finland. Phone: 358-9-19149705. Fax: 358-9-19149718. E-mail: kjorkroth@helsinki.fi.
Description of the product and pH, sensory, and microbiological analysis. The product was manufactured from raw, skinned broiler meat, which was cut into strips, marinated with the marinade, and packaged under MA as ca. 500-g consumer packages. The marinade contained plant oil, tomatoes, paprika, cayenne,

TABLE 1. *Leuconostoc sensu stricto* reference strains

Species	Strain
<i>L. carnosum</i>	LMG 11498 ^{Ta}
<i>L. citreum</i>	LMG 11417
<i>L. fallax</i>	CCUG 30061 ^{Tb}
<i>L. gelidum</i>	LMG 9850 ^T
<i>L. lactis</i>	CCUG 30064 ^T
	LMG 7940
<i>L. mesenteroides</i> ssp. <i>premoris</i>	LMG 13562
<i>L. mesenteroides</i> ssp. <i>dextranicum</i>	LMG 7954
	LMG 11318
<i>L. mesenteroides</i> ssp. <i>mesenteroides</i>	LMG 7939
<i>L. pseudomesenteroides</i>	LMG 11482 ^T
	LMG 11483
	LMG 11499

^a BCCM/LMG Belgian Coordinated Collections of Microorganisms.
^b Culture Collection of the University of Gothenburg.

salt, protein hydrosylates, starch and modified starch, natural aromatics, preservatives, and a buffering additive. The pH of the normal product varied from 4.0 to 4.5 with pH 4.5 set as the optimal target value. The expected shelf life was 60 days, with the day of manufacture regarded as day 0.

Six unopened packages showing clear bulging were analyzed on the shelf life. LAB were enumerated from serial 10-fold dilutions (Oxoid, Basingstoke, United Kingdom) and Rogosa lactobacillus agar (Orion Diagnostica, Espoo, Finland) as described by Korkeala et al. (3). The plates were incubated at 25 C in an anaerobic jar and then generating kit (Oxoid) for 5 days. The pH was measured directly on homogenized samples. Evaluation of odor, color, appearance, and spoiled product was performed by three trained judges, as described by Lindroth (21).

Bacterial strains and the use of strains in different phases of the study. A total of 120 spoiled packages recovered from the six spoiled packages (20 isolates from each package) were purified. During the course of the study, strains were assessed in the different phases of the study as described below. The 120 spoilage isolates were all subjected to basic phenotypic characterization and ribotyping. The ribopatterns were compared with the corresponding patterns in the LAB database of the Department of Food and Environmental Hygiene, University of Helsinki, Helsinki, Finland. These comprise patterns of the spoilage LAB in the genus *Lactobacillus*, *Leuconostoc*, *Enterococcus*, and *Weissella* (4, 6, 7, 25). Before Southern blotting, the restriction endonuclease analysis (REA) patterns of the main spoilage species were inspected visually. From these strains, those with different REA patterns, four strains representing both pattern types (two from each) were chosen for further taxonomic studies. These isolates were given the following strain numbers: LMG 18811, LMG 18812, LMG 18813, and LMG 18889. The strains presented in Table 1 were used during the more detailed taxonomic dealing with the main spoilage species, and the LAB ribotyping database contained the ribopatterns of these strains.

All of the strains were maintained in MRS broth (Difco, Detroit, Mich.) at 27 C and cultured using MRS broth or MRS agar (Oxoid).

Phenotypic characterization. All 120 isolates were Gram stained, and tested, streaked on Rogosa lactobacillus agar, and studied for the production of gas from glucose (44). Further phenotypic characterization of the main spoilage species was done with strains LMG 18811, LMG 18812, LMG 18813, and LMG 18889. Production of ammonia from arginine was determined by the method of Briggs (11). Dextran formation was studied on agar containing 5% sucrose (20). Fermentation of carbohydrates was determined by the API 50 CHL Lactobacillus identification system (Biomérieux, Marcy l'Etoile, France). The ability to produce different lactic acid isomers was tested enzymatic method (48) utilizing Boehringer Mannheim GmbH (Mannheim, Germany) and -lactate dehydrogenases. The four strains were also tested for growth in MRS broth at 4, 10, 15, and 37 C. No growth was observed or at least for 21 days.

Enzymatic activity. The proteolytic activity of *L. gasicomitatum* nov. isolates was tested on MRS agar supplemented with sterile skim milk (2% concentration). API ZYM (Biomérieux) was also used for the characterization of the enzymatic activities of the LMG 18811 and LMG 18812 strains.

Peptidoglycan analysis. Preparation of cell walls and determination of peptidoglycan structure of LMG 18811 was carried out by the methods of Schleifer and Kandler (41) with the modification of using thin-layer chromatography on cellulose sheets instead of paper chromatography. Briefly, freeze-dried cell walls were hydrolyzed in 100 C for 16 h (total hydrolysate) and 45 min (partial hydrolysate). Diamino acids were identified from total hydrolysate by one-dimensional chromatography in the system methanol-pyridine-water-10 N HCl (320:40:70:10 [vol/vol/vol/vol]). Peptides from total and partial hydrolysates were identified by two-dimensional chromatography in the systems published by Schleifer and Kandler (41), by their mobilities and staining characteristics with ninhydrin. The resulting "fingerprints" were compared with known peptidoglycan structures. Whole-cell protein analysis. The similarity of the main spoilage species (strains LMG 18811, LMG 18812, LMG 18813, and LMG 18889) to *Leuconostoc sensu stricto* species (Table 1) was studied by means of whole-cell protein analysis. Strains were grown for 5 days on MRS agar at 25 C in a microaerobic atmosphere (approximately 5% O₂, 10% CO₂, and 85% N₂). Preparation of cellular protein extracts and polyacrylamide gel electrophoresis were performed as described previously (36). Briefly, discontinuous gels were run overnight at constant temperature in a vertical slab apparatus. The separation gel was long and contained 12% total acrylamide (the monomer solution contained 2.67% cross-linking in 0.375 M Tris-HCl [pH 8.8] and sodium dodecyl sulfate). The stacking gel was 12 mm long and contained total acrylamide with 2.67% cross-linking in 0.375 M Tris-HCl [pH 8.8] and sodium dodecyl sulfate). The stacking gel was 12 mm long and contained total acrylamide (the monomer solution contained 30% total acrylamide with 2.67% cross-linking in 0.125 M Tris-HCl [pH 6.8] and 0.1% sodium dodecyl sulfate). Protein bands were stained with Coomassie blue R-250 in 50% (v/v) methanol and 10% (vol/vol) acetic acid. These conditions allowed the separation of proteins and peptides in the molecular weight range of 14,000 to 116,000.

Isolation of DNA, REA, and 16 and 23S rDNA RFLP (ribotyping). EcoRI, and HindIII restriction enzymes (New England Biolabs, Beverly, Mass.) were used for ribotyping. DNA was isolated by the guanidium thiocyanate method of Pitcher et al. (35) as modified by Bax and Korkeala (3) by the use of lysozyme and mutanolysin (Sigma, St. Louis, Mo.) treatment. Restriction endonuclease treatment of DNA was done as specified by the manufacturer (New England Biolabs), and REA was performed as described previously (3). Before Southern blotting, the REA patterns were inspected in order to obtain preliminary information about clonal variation. Genomic blots were made using a vacuum device (Vacugene; Pharmacia, Uppsala, Sweden), and the rDNA probe for ribotyping was labeled by reverse transcription (AMV-RT [Promega, Madison, Wis.] and Dig DNA labeling kit [Roche Molecular Biochemicals, Mannheim, Germany]) as previously described (Korkeala et al. (9)). Membranes were hybridized at 68 C as described by Bax and Korkeala (3).

Pattern analysis. The ClaI, EcoRI, and HindIII ribopatterns were compared with the corresponding patterns in the previously established LAB database. For ribotyping analysis, ribopatterns were scanned using a Hewlett-Packard Jet 4c/T scanner and analyzed using the BioNumerics 1.0 software package (Applied Maths, Kortrijk, Belgium). The similarities between pairs of patterns were assessed by Dice coefficient correlation, and unweighted pair-group method with arithmetic averages clustering was used for the construction of dendrograms.

Whole-cell protein profiles were scanned using a 2202 UltraScan laser densitometer (LKB, Bromma, Sweden). The densitometric analysis, normalization, and correlation of the protein profiles were performed with the GelComp software package (Applied Maths). Numerical analysis was performed using the BioNumerics 1.0 software package. The similarities between all pairs of patterns were expressed by the Pearson product moment correlation coefficient converted to a percent value.

Different types of banding patterns were integrated in a single database using the BioNumerics 1.0 software package. In these combined analyses, equal weight was given to each of the three banding patterns.

The 16S rRNA gene sequence analysis. The 16S rRNA gene was amplified with a universal primer pair (45): primer A (5'-GAGTATGATCTGGCTCAG3') and primer B (5'-GAAAGGAGGTGATCCAGCC3'). Two strains, LMG 18811 and LMG 18812, were studied. They represented the two groups detected in ribotyping. Chromosomal DNA was isolated as for ribotyping. Amplification was performed using 200 ng of chromosomal DNA as a template. The PCR mixture contained 500 U of Taq polymerase (Promega), 100 U of Taq polymerase buffer (BioLabs), 5% sucrose (20), and of nucleotide mixture (dATP, dCTP, dTTP, and dGTP; 2.5 mM each), 4 mM of MgCl₂ (25 mM), 1.25 M of primers A and B (120 μM), and 10 U of Taq polymerase adjusted to 1.0 U added to yield a total reaction volume of 100 μl.

Sequencing of the purified PCR product (Quantum Prep PCR Kleen spin columns, Bio-Rad Laboratories, Hercules, Calif.) was performed by Sanger dideoxynucleotide chain termination method using an ABI PRISM sequencer (Perkin-Elmer Corp., Norwalk, Conn.) according to the manufacturer's recommendations. Sequencing was performed as two long reactions, and complementary sequences were joined by the DNASIS program (Hitachi Software, Yokohama, Japan).

Phylogenetic analysis was performed by using the GeneCompar 2.0 software package (Applied Maths). The consensus sequence and the sequences of strains

TABLE 2. Recovery of LAB on MRS and Rogosa selective Lactobacillus agar and pH values analyzed from six spoiled packages showing clear bulging due to gas formation

Package no.	No. of LAB (CFU/g)		pH
	MRS	Rogosa selective Lactobacillus agar	
1	23 10 ¹⁰	13 10 ¹⁰	4.9
2	43 10 ¹⁰	43 10 ¹⁰	4.8
3	43 10 ¹⁰	13 10 ¹⁰	4.7
4	13 10 ¹⁰	23 10 ⁸	4.8
5	13 10 ¹⁰	33 10 ¹⁰	4.8
6	93 10 ¹⁰	33 10 ¹⁰	5.0

belonging to the same phylogenetic group (retrieved from the National Center for Biotechnology Information GenBank data library) were aligned. The accession numbers of the 16S rDNA sequences used are as follows: *Leuconostoc argentinum* LMG 18543^T, AF175403; *Leuconostoc carnosum* LMG 11498^T, X95997; *Leuconostoc citreum* LMG 9824^T, X53963 and S78399; *Leuconostoc fallax* LMG 13177^T, S63851; *Leuconostoc gelidum* LMG 9850^T, S63851; *Leuconostoc lactis* LMG 8894^T, M23031 and M23032; *Leuconostoc mesenteroides* subsp. cremoris LMG 6909^T, M23034; *Leuconostoc mesenteroides* subsp. mesenteroides LMG 6893^T, M23035; *Leuconostoc pseudomesenteroides* LMG 6852^T, X95979; and *Weissella paramesenteroides* LMG 6852^T, X95982.

DNA base composition and DNA-DNA hybridization was isolated from two spoilage isolates (LMG 18811 and LMG 18813) and the type strain NCFB 2775, and the *Leuconostoc mesenteroides* subsp. dextranictype strain DSM 20484. *L. gelidum* was selected because it had the highest similarity to the main spoilage species according to some phenotyping schemas, RFLP analysis, whole-cell protein analysis, and 16S rDNA sequence analysis. *L. gelidum* was chosen on the basis of API characterization results, and it was also used as a control species in the present study.

For large-scale DNA isolation, the modified (3) guanidium thiocyanate method of Pitcher et al. (35) was scaled up 10-fold. Cells from 200 ml of well-grown MRS broth culture were used for each isolation batch. In one batch was dissolved overnight in 1 ml of TE 10:1 (10 mM Tris, 1 mM EDTA, pH 8.0). RNase A (Sigma) was added to provide a concentration of 125 mg/ml, and the solution was incubated at 37 °C with gentle shaking for 1 h. Following the 1-h incubation, proteinase K (Sigma) was added to provide a concentration of 1 mg/ml, and incubation at 37 °C was continued for at least 6 h. DNA was isolated as described by Pitcher (35) and dissolved in SSC 13.5 SSC 1 is 0.15 M NaCl plus 0.015 M sodium citrate). When dissolved, the SSC concentration of a sample was adjusted to 20 SSC.

Purified DNA was dialyzed twice overnight at 4 °C using a 12,000–14,000 Da-pore-size membrane (Mediatech International Ltd., London, United Kingdom). The first dialysis was carried out against DEPC (10 mM), and the second was carried out against SSC. DNA was fragmented two times in a French pressure cell (SML Aminco; Colora Messtechnik GMBH, Lorch, Germany) at about 13,500 Pa. Before reassociation, it was dialyzed overnight at 4 °C against SSC.

The DNA base composition (moles percent) was estimated by the thermal-denaturation method (12), and the DNA homology values were determined from renaturation rates using a Gilford Response spectrophotometer (Giba Corning Diagnostics Corp., Gilford Systems, Oberlin, Ohio).

Nucleotide sequence accession numbers for approximately 1,500-bp sequences of the 16S ribosomal genes of strains LMG 18811 and LMG 18813 have been deposited in the GenBank data library with accession numbers AF231131 and AF231132, respectively.

RESULTS

Microbiological and sensorial qualities of the products according to Table 2 shows the results of microbial enumeration on MRS and Rogosa selective Lactobacillus agars and corresponding pH values obtained from the six packages. An increase in the pH of the product, very atypical for LAB spoilage, was detected. Instead of the normal pH values, ranging from 4.2 to 4.4, values from 4.7 to 5.0 were detected. All of the packages were deemed unfit for human consumption by all three judges. They were all described as clearly bulged, and the smell of the product was described as pungent and very unpleasant. The consistency and texture of the product was, however, normal, and no color changes were visible.

LAB population associated with the spoiled product shows the division of the 120 isolates into different and groups of species based on the LAB ribotyping database. An organism possessing typical lower molecular bands leuconostoc HindIII ribopatterns (Fig. 1C) was found to dominate (67 of 120) in the product. These isolates were gram-positive, catalase-negative oval cocci, produced glucose, and did not grow on Rogosa agar. They possessed identical ribopatterns, showing, however, two different the REA patterns. The distribution of the isolates between these two REA types was almost even.

The two other major species associated with the product were *Lactobacillus curvatus* (32 of 120) and *Lactobacillus sakei* (16 of 120). Isolates possessing ribotypes identical to *L. sakei* and *L. curvatus* patterns in the database (no new patterns were detected) were gram-positive rods or coccioid. The ones grown on Rogosa agar, were catalase negative, and did not produce gas from glucose. Three of the 120 isolates identified as *Leuconostoc sensu stricto* species, *Leuconostoc summatum* and *L. gelidum*. They all shared identical ribopatterns with the corresponding *Leuconostoc* type strain, were oval cocci, produced gas from glucose, and did not grow on Rogosa selective Lactobacillus agar. Twelve isolates could not be identified with the existing ribotyping database. They were gram-positive rods growing on Rogosa selective Lactobacillus agar, produced gas from glucose, and shared identical ribopatterns. They did not have any similarity to the ribopatterns of *Leuconostoc brevis*, *Leuconostoc buchneri*, *Leuconostoc colliformis*, *Leuconostoc fermentum*, *Leuconostoc fructivorans*, *Leuconostoc hilgardii*, or *L. gelidum* strains. This was also the case with respect to the patterns of *Carnobacterium divergens*, *Carnobacterium piscium*, *Carnobacterium mobile*, and *Carnobacterium valinartum* strains.

Phenotypic reactions of the main spoilage species, LMG 18811, LMG 18812, LMG 18813, and LMG 18889 strains showed typical reactions for the genus *Leuconostoc*. They did not produce ammonia from arginine, did not grow in the presence of 6.5 to 12% NaCl, and synthesized only lactic acid from glucose. They all grew at 4 and 15 °C but not at 25 °C. Growth was already slower at 30 °C, and during the study it was observed as an optimum temperature for growth on MRS. All four strains produced excessive slime from sucrose and fermented arabinose, ribose, glucose, fructose, mannose, a-methyl-glucoside, acetyl-glucosamine, esculetin, cellobiose, maltose, melibiose, sucrose, trehalose, gentiobiose, turanose, and 5-keto-gluconate. LMG 18811, LMG 18813, and LMG 18889 also fermented galactose and gluconate. Glycerol, erythritol, xylitol, xylose, adonitol, a-methyl-xyloside, sorbose, rhamnose, dulcitol, inositol, mannitol, sorbitol, a-methyl-mannoside, amygdalin, arbutin,

TABLE 3. Species division of 120 LAB isolates originating from packages (20/package) of MA-packaged raw marinated broiler meat products according to the EcoRI, and HindIII ribopattern database

Package no.	No. of isolates					
	<i>Leuconostoc</i> spp.	<i>L. carnosum</i>	<i>L. gelidum</i>	<i>L. sakei</i>	<i>L. curvatus</i>	Unidentified
1	4	1	2	4	5	4
2	9			4	5	2
3	10			3	5	2
4	15			4		1
5	12			1	6	1
6	7				11	2
Total	57	1	2	16	32	12

FIG. 1. Ribopatterns and numerical analysis of the patterns presented as dendrograms. Patterns and dendrograms (A) and (B), by using HindIII (C) restriction enzymes are shown. The left sides of the banding patterns show large molecular sizes (23 Kbp), and the right (50 Kbp) show small sizes (10 Kbp). The dendrograms show the similarity values between the patterns. The dendrogram (A) shows that the Argentine strain showed ribopatterns identical to those of the Chilean strain. Scales from 30 to 100 show percentile similarity values for the patterns.

TABLE 4. Main differences in sugar fermentation between *L. carnosum*, *L. gelidum*, and *L. gasicomitatum* nov. sp. n.

Sugar	Fermentation		
	<i>L. carnosum</i>	<i>L. gelidum</i>	<i>L. gasicomitatum</i>
Amygdalin	2	1	2
L-Arabinose	2	1	1
Arbutin	2	1	2
Raffinose	2	1	1
D-Xylose	2	1	1
Salicin	2/1 ^c	1	2

^a Reactions adapted from Shaw and Harding (43).
^b 2, no fermentation; 1, fermentation takes place.
^c Most strains do not ferment salicin.

salicin, lactose, inulin, melezitose, starch, glycogen, xyli-
 tose, tagatose, fucose, arabinol, and 2-keto-gluconate were
 not fermented. The API Chem System identified these iso-
 lates with an extremely good identification level (99.9%) as
 the key carbohydrate fermentation reactions among the
 phylogenetically associated *L. gelidum* and the
 strains representing the main spoilage group.

Peptidoglycan type of the main spoilage species. Cell walls
 of LMG 18811 contain, besides muramic acid and glucosamine,
 the amino acids lysine, glutamic acid, and alanine in a molar
 ratio of 1:1:4, respectively. The partial hydrolysate were
 compatible only with type A₃ Lys-Ala-Ala.

Enzymatic activities. None of the *L. gasicomitatum* nov. iso-
 lates changed the appearance of the skim milk on MRS agar.
 According to API ZYM analysis, both LMG 18811 and LMG 18812
 showed the presence of alkaline phosphatase, lipase, esterase,
 esterase-C, acid phosphatase, and naphthol-AS-BI-phospho-
 hydrolase activities.

Numerical analyses of the main spoilage species based on
 the dendrograms and banding patterns of the main spoilage
 species and the reference strains *C. casei*, *C. coli*, and
HindIII ribotypes, respectively. Figure 2 shows a dendro-
 gram obtained by combining the pattern information of all
 ribotypes into one numerical analysis. The result of the nu-
 merical analysis of the whole-cell protein patterns is shown
 in Figure 3, and the combined information from all of the ribo-
 patterns and whole-cell protein analysis is presented as a dendro-
 gram in Figure 4.

The three spoilage isolates and the reference strains *C. casei*,
C. coli, and *HindIII* ribotypes formed distinct clusters in the
 dendrograms based on the *HindIII* ribotypes (Fig. 1C) and the
 protein patterns (Fig. 3), indicating that these techniques
 generated species-specific patterns. In the dendrograms gener-
 ated from numerical comparison of *HindIII* and *EcoRI* ribotypes,
 only *Citrobacterium pseudomesenteroides* and the spoilage iso-
 lates formed distinct species-specific clusters (Fig. 1C).
 When equal weight is given to all three types of ribo-
 patterns, the analysis combining this information resulted in
 distinct species-specific clusters (Fig. 2). The *L. gelidum*
 type strain had the highest similarity to the spoilage iso-
 lates in the numerical analyses of combined ribo-
 patterns (Fig. 2) as well as in the whole-cell protein pat-
 terns (Fig. 3). In respect to the subdivision of the genus
Citrobacterium, the numerical analyses performed correlated
 with the interspecies division of the genus. The type strain
 of *C. casei* was found to possess the same ribopatterns as
 the LMG 7940 strain, and in the dendrogram based on the
 numerical analysis of whole-cell protein patterns, the
 LMG 7940 strain formed a tight cluster with the other strains
 (Fig. 3). Such an association was only seen among strains of a
 single species.
 Phylogenetic analyses based on 16S rDNA sequences shows
 the sequence homologies of strains LMG 18811 and LMG 18812
 compared with the *Leuconostoc sensu stricto* spe-

FIG. 2. Dendrogram obtained by combining the pattern information of *HindIII* and *HindIII* ribotypes into one numerical analysis. The scale from 40 to 100 shows percentile similarity values.

FIG. 3. Numerical analysis of whole-cell protein patterns presented as a dendrogram. The scale from 60 to 100 shows percentile si

cies. The two strains, LMG 18811 and 18812, showed DNA-DNA reassociation: LMG 18811 LMG 18812, 100%; sequence homology of 99.3%, and the highest homology with LMG 18811 3 L. gelidum LMG 18297T, 22%; LMG 188123 and 98.8%, respectively, was exhibited with the type L. gelidum LMG 18297^T, 34%; and LMG 188123 L. mesen- teroides dextranicum LMG 6908^T, 33%. The DNA G1 C

DNA base composition and DNA-DNA hybridization contents of strains LMG 18811 and LMG 18812 are 37 and 38 sults. The following DNA homology values were obtained with L. gelidum, respectively.

FIG. 4. Combined information from EcoRI, and HindIII ribopatterns and whole-cell protein patterns presented as a dendrogram. The strain showed ribopatterns identical to LMG 7940. The scale from 40 to 100 shows percentile similarity values.

TABLE 5. Homology values for a 1,491-nucleotide region of 16S rDNA

Species	Homology (%) to:											
	1	2	3	4	5	6	7	8	9	10	11	12
1. <i>Leuconostoc</i> sp. strain LMG 18811	100											
2. <i>Leuconostoc</i> sp. strain LMG 18812	99.3	100										
3. <i>L. argentinum</i> LMG 18534 ^T	97.1	97.1	100									
4. <i>L. carnosum</i> LMG 11498 ^T	98.3	98.5	97.5	100								
5. <i>L. citreum</i> LMG 9824 ^T	97.5	97.6	98.7	97.4	100							
6. <i>L. fallax</i> LMG 13177 ^T	93.6	93.7	92.4	94.3	94.5	100						
7. <i>L. gelidum</i> LMG 9850 ^T	98.8	99.0	97.1	98.3	97.4	93.5	100					
8. <i>L. lactium</i> LMG 8894 ^T	97.3	97.6	99.3	97.3	98.5	93.6	97.6	100				
9. <i>L. mesenteroides</i> ssp. <i>premorium</i> LMG 6909 ^T	97.7	98.0	97.7	97.8	97.7	94.6	97.9	98.2	100			
10. <i>L. mesenteroides</i> ssp. <i>mesenteroides</i> LMG 6893 ^T	97.7	98.0	97.5	97.8	97.7	94.6	97.9	98.2	100	100		
11. <i>L. pseudomesenteroides</i> LMG 1482 ^T	97.9	98.0	97.8	97.8	97.5	94.6	98.0	98.0	99.5	99.5	100	
12. <i>W. paramesenteroides</i> LMG 9852 ^T	91.2	91.4	90.3	91.0	91.2	92.4	90.9	91.3	91.8	91.8	91.7	

DISCUSSION

Gaseous deterioration caused by LAB has mainly been associated with highly acidic foods, such as fermented vegetables (16, 29) or acetic acid preserves (6, 24), but it may also affect meat products (10). Even though LAB have also been found as the dominant spoilage organisms in vacuum- or MA-packaged poultry products (34, 39, 40), the strains have never been identified to the species level. Gaseous deterioration as a LAB spoilage type in vacuum- or MA-packaged poultry products has not been reported previously. It was not surprising to find *L. sakei* and *L. curvatum* strains in the poultry product studied here. These species are very typical for all meat products (3, 4, 26, 28; Holzapfel and Gerber, Abstr. 32nd Eur. Meet. Meat Res. Workers) and might have been a major component of the spoilage population detected in the previous studies dealing with vacuum- or MA-packaged poultry products (17, 33, 39, 40). *L. carnosum*, *L. gelidum* and *L. lactium* are also quite common species occurring in vacuum- or MA-packaged cold-stored meat products (43, 49). Two species occurring in the spoiled poultry product were unusual LAB species for meat products. *U. comitatus* sp. nov. was the main spoilage species characterized in this study, and the identification of the gram-positive rod shaped organism will be carried out as a separate study. In addition to the novel species, this LAB spoilage also showed unique properties. Normally, in a case of clear acid spoilage, the pH of the product decreases due to lactic acid formation, but in this case an increase of pH was detected. This type of LAB spoilage was first reported by Meyer (50) for some canned fish marinades. He called it "protein swell" and distinguished it from "carbohydrate swell," where increased CO₂ and CO₂ formation result from heterofermentative utilization of glucose. In protein swell, proteins are decomposed by proteolytic enzyme action, and the subsequent decarboxylation of amino acids leads to enhanced CO₂ production. Therefore, the LAB having an effect on gas production in protein swell may also possess homofermentative glucose metabolism. An increase in acidity related to protein swell has been attributed to production of ammonia by bacterial deamination of amino acids.

Protein swell has also been reported to affect stuffed olives (19), but to our knowledge there are no reports of this type of spoilage affecting any type of meat product. The previous studies of protein swell have assigned the main component of the spoilage LAB with the decarboxylase activity to *U. comitatus* (19, 30) and considered protein hydrolysis to be due to endogenous fish enzymes. The initial hydrolysis of muscle proteins has also been attributed to endogenous enzymes, mainly cathepsins, and the bacterial activity was associated with the degradation of oligopeptides and

amino acids (32, 46). The proteolytic systems of various related LAB are poorly known, and the ability of *L. curvatum* to degrade myofibrillar proteins has only recently been studied (15, 37). These species have been shown to possess peptidase activity and also to express strong amino acid metabolism (15, 33, 37, 38), and even though they were not major components of the spoilage population, they may have played a major role in this case. *Leuconostocs* have not yet been detected as dominant species in protein swell, which makes their predominance in *U. comitatus* species produce gas during normal glucose fermentation, and in this case, the extreme bloom may have resulted from complicated interaction between various LAB species and the endogenous muscle-associated enzymes. Whether the *U. comitatus* component alone, or in association with the endogenous muscle proteinases, could cause the gaseous spoilage remains unknown. There are no data on the proteolytic systems of meat-associated *Leuconostocs*. The main component did not show proteolytic activity on the skim milk-supplemented MRS agar due to the substrate specificity of proteolytic systems. Complicated techniques should be used for the evaluation of the proteolytic effect on myofibrillar proteins. In the product, the LAB counts were exceptionally high (10⁹ g⁻¹). Two factors, the marinade and the small rise in pH, have played major roles in facilitating the growth of LAB. The marinade had a tomato base, which contains growth stimulating factors. The plant was simultaneously processed in poultry strips in other marinades, such as honey based, but the carbohydrate-marinated product showed gaseous deterioration. The carbohydrates and protein hydrolysates in the marinade may have provided nutrients facilitating pronounced gaseous spoilage. The spoilage problem was overcome by stabilizing the pH with another type of additive. Apparently this could have an effect on the growth of *Leuconostocs*, which are generally as acid tolerant as *Lactobacillus* species. *Leuconostoc sensu stricto* comprises *L. argentinum*, *L. carnosum*, *L. citreum*, *L. gelidum*, *L. lactium*, *L. mesenteroides* (three subspecies), *L. premoium*, *L. pseudomesenteroides* and *L. sakei* showing 97 to 99% 16S rDNA sequence homology. In addition, an atypical *Leuconostoc* strain assigned to 95% 16S rDNA homology with the other sensu stricto species, has been described. Our results show high sequence homology between *Leuconostoc sensu stricto* and *U. comitatus*, the main spoilage species, clearly assigning it to the genus. The highest 16S rDNA sequence homology (98.8%) was displayed with *L. gelidum* and the lowest (93.6 and 93.7%). According to these data

L. gasicomitatum sp. nov. is in the same evolutionary branch (42). One of these branches contains *L. gelidum* and *L. carnosum*. *L. gelidum* and *L. carnosum* contain another cluster and *L. lactis* and the *L. pseudomesenteroides*. In addition to the 16S rDNA sequence homology, whole-cell protein analysis and combined 16 and 23S rDNA RFLP showed that the species-specific clusters *L. gelidum* possess the highest similarity to *L. gasicomitatum* never reflect the phylogenetic branching of the 16S rDNA of *L. gelidum* sp. nov. According to the phenotyping schema of Villafraña et al. (47), this species should be regarded as *L. gasicomitatum* whereas protein patterns and the combined 16 and 23S rDNA RFLP API CHL analysis identified *L. gasicomitatum* and *L. gelidum* as different even though both techniques provide specific clustering. The consensus of both techniques revealed *L. gasicomitatum* and *L. pseudomesenteroides* as another type of dendrogram (Fig. 4). This shows clearly that the spoilage strains clearly represent the different values of different numerical analyses of species. Considering all of the results, status as a new species is comparable and that these techniques should be used for species-level identification and not for deduction. It is difficult to distinguish between *L. carnosum* and *L. gelidum* and as can be seen from the phenotypic reactions, identification of the new species follows the same lines. These three species have similar growth temperature characteristics and share the same doglycan type, and only some of the carbohydrate fermentations. E. Falsen, curator of the Culture Collection University of Gothenburg (personal communication) also observed similarity between *L. gasicomitatum* and *L. lactis* within *Leuconostoc* species (13), these reactions are not, however, absolute. The conserved nature of the gene *arg* in *L. gasicomitatum* and *L. lactis* type strains and also similar API profiles (API rapid ID 32 strep, API 50 CHL, and API ZYM). Identification based on sequence comparison of complete 16S rDNA of *L. gasicomitatum* strain is authentic. Therefore, DNA-DNA reassociation has been considered to be the only reliable method to distinguish (99.3%) that the two type strains show also supports this from *L. gelidum* (13).

The polyphasic approach used in this study showed clearly that *L. gasicomitatum* possess analysis for the characterization of strains. Using the highest similarity to *L. gelidum* The low homology merical analysis of total cellular proteins is a generally DNA-DNA reassociation experiments clearly depicted tool for speciation of bacteria, and we have used ribotyping for LAB identification (4, 5, 6, 7) considered to be a distinct species for with good results. Whole-cell protein analysis, and *HindIII* ribotyping, provided species-specific cluster patterns for *L. gasicomitatum* and *L. pseudomesenteroides* for the *Leuconostoc* reference strains and the new taxon (*L. gasicomitatum* sp. nov.). *L. gasicomitatum* and *L. pseudomesenteroides* (Fig. 1) are similar to *L. gasicomitatum* and *L. pseudomesenteroides* (Fig. 1) are similar to *L. gasicomitatum* and *L. pseudomesenteroides*. Gram-positive, nonmotile, and non-spore-forming molecular-weight fragments were obtained (Fig. 1) are similar to *L. gasicomitatum* and *L. pseudomesenteroides*. Colonies are small, projecting the numerical analysis to errors due to the mobility of these fragments. The *E. coli* is a facultative anaerobe; produces gas from glucose. More than 95% of other (Fig. 1A) and thus were not optimal for numerical analysis. Arginine is not hydrolyzed. The locations of the rDNA genes in many of the *Leuconostoc* species seem to be very conserved, providing the presence of 6.5 to 12% NaCl. The peptidoglycan type little variation between different strains. In our study, *L. gasicomitatum* and *L. pseudomesenteroides* differ with *L. carnosum* 29 different macrorestriction sites, fructose, mannose, methyl-glucoside, acetate, and patterns showed the same ribotype (7). Here also, *L. gasicomitatum* and *L. pseudomesenteroides* possess different REA patterns yielded the same ClA, EcoRI, and *HindIII* ribotypes. Due to the highly conserved were fermented. Some isolates ferment galactose and *HindIII* ribopatterns, ribotyping is a good tool for identification, but none of the three enzymes allowed good strain typing results. It can be concluded that *L. gasicomitatum* and *L. pseudomesenteroides* are closely related and analysis of protein patterns and *HindIII*-based ribopatterns can both be used for the identification of species. The xylitol, xylose, tagatose, fucose, arabinol, gluconate only limiting factor currently associated with these approaches is the lack of well-characterized reference strains in this case especially in respect to *L. gasicomitatum*.

The G1C content of the type strain is 37%, determined by the integral-denaturation method. Isolated from MA-pasteurized, tomato-marinated broiler meat strips showing whole-cell protein patterns results in clusters which correlate with phylogenetic branches based on 16S rDNA homology. When the phylogenetic tree of the type strain corresponds to that of the species *L. gasicomitatum* is placed under more precise scrutiny, three evolutionary branches are distinguished on the basis of 16S rDNA and phosphatase, and naphthol-AS-BI-phosphohydrolase.

tivities are also present. Ferments galactose and glucosaminoglycans, and mayonnaise and salad dressings. App 21: 67-71. 25. Lyons, U., J. Björkroth, and H. Korkeala. 1989. Characterisation of lactic acid bacteria from spoiled, vacuum-packaged, cold-smoked rainbow trout by ribotyping. *Int. J. Food Microbiol.* 5: 171-81.

ACKNOWLEDGMENTS

We are indebted to the Academy of Finland for financially supporting K. Johanna Björkroth's work and to the Fund for Scientific Research—Flanders (Belgium) for the positions of P. Vandamme as a postdoctoral research fellow and P. De Vos as a senior research associate.

S. Ekstrand and I. Lubecki are acknowledged for their skilled technical assistance. We also thank Häme for their kind help in the denomination of the novel species.

REFERENCES

1. Allen, J. R., and E. M. Foster. 1960. Spoilage of vacuum-packed sliced processed meats during refrigerated storage. *J. Food Res.* 25: 19-22.
2. Alm, F., I. Erichsen, and N. Molden. 1970. The effect of vacuum packaging on some sliced processed meat products as judged by organoleptic and bacteriological analysis. *Food Technol.* 10: 20-23.
3. Björkroth, J., and H. Korkeala. 1992a. Evaluation of *Lactobacillus sake* contamination in vacuum packaged sliced cooked meat products by ribotyping. *J. Food Prot.* 59: 398-401.
4. Björkroth, J., and H. Korkeala. 1992a. rRNA gene restriction patterns as a characterization tool for *Lactobacillus sake* strains producing ropy slime. *Int. J. Food Microbiol.* 10: 293-302.
5. Björkroth, J., and H. Korkeala. 1992a. Use of rRNA gene restriction patterns to evaluate the lactic acid bacterium contamination of vacuum-packaged sliced cooked whole-meat product in a meat processing plant. *App Environ. Microbiol.* 58: 448-453.
6. Björkroth, J., and H. Korkeala. 1992a. Characterization of *Lactobacillus fructivorans* spoilage in ketchup. *J. Food Prot.* 65: 508-509.
7. Björkroth, K. J., P. Vandamme, and H. J. Kiehl. 1991. Identification and characterization of *Leuconostoc carnosum* associated with production and spoilage of vacuum-packaged, sliced, cooked ham. *Appl. Environ. Microbiol.* 57: 3313-3319.
8. Blickstad, E., and G. Mörby. 1988. The microbial flora of smoked pork loin and frankfurter sausage stored in different gas atmospheres at 4°C. *Bacteriol. Rev.* 54: 45-56.
9. Blumberg, H. M., J. A. Kielbauch, and I. K. Wadsworth. Molecular epidemiology of *Yersinia enterocolitica* infections: use of chromosomal DNA restriction fragment length polymorphism of rRNA genes. *J. Food Microbiol.* 29: 2368-2374.
10. Borch, E., M.-L. Kant-Muermans, and Y. L. Ho. 1988. Bacterial spoilage of meat and cured meat products. *Int. J. Food Microbiol.* 1: 21-30.
11. Briggs, M. 1953. The classification of lactobacilli by means of physiological tests. *J. Appl. Bacteriol.* 16: 545-556.
12. De Ley, J., H. Cattoir, and A. Reyniers. 1960. The quantitative measurement of DNA hybridization from renaturation rates. *Eur. J. Biochem.* 1: 133-142.
13. Dellaglio, F., L. M. T. Dicks, and S. L. Brown. 1995. The genus *Leuconostoc* p. 235-271. In B. J. B. Wood and W. Holzappel (ed.), *The genera of lactic acid bacteria*. Blackie Academic and Professional, Glasgow, United Kingdom.
14. Dykes, G. A., T. E. Cloete, and A. van der Merwe. 1994. Identification of *Leuconostoc* species associated with the spoilage of vacuum-packaged Vienna sausages by DNA-DNA hybridization. *Food Microbiol.* 17: 271-274.
15. Fadda, S., Y. Sanz, G. Vignolo, M.-C. Arsitoy, G. Oliveri, and F. Toldrano. Hydrolysis of pork muscle sarcoplasmic proteins by *Lactobacillus curvatus* and *Lactobacillus sakei*. *Appl. Environ. Microbiol.* 65: 578-584.
16. Fleming, H. P. 1982. Fermented vegetables. *Econ. Microbiol.* 2: 227-238.
17. Gill, C. O., J. C. L. Harrison, and N. D. Pennie. 1981. The storage life of chicken carcasses packaged under carbon dioxide. *Int. J. Food Microbiol.* 1: 51-58.
18. Grimont, F., and P. A. D. Grimont. 1986. Ribosomal ribonucleic acid genes as restriction as potential taxonomic tools. *Ann. Inst. Pasteur/Microbiol.* 165: 165-175.
19. Harmon, S. M., D. A. Kautter, and C. McClure. 1990. Spoilage of anchovy-stuffed olives by heterofermentative lactobacilli. *J. Food Safety* 11: 205-210.
20. Harrigan, W. F., and M. E. McCann. 1976. Laboratory methods in food and dairy microbiology. Academic Press, New York, N.Y.
21. Korkeala, H., and S. Lindroth. 1987. Differences in microbial growth in the surface layer and center of vacuum-packaged cooked ring sausage. *Food Microbiol.* 4: 105-110.
22. Korkeala, H., T. Alanko, and S. Lindroth. 1989. Shelf-life of vacuum-packaged cooked ring sausages at different chill temperatures. *Food Microbiol.* 2: 237-247.
23. Korkeala, H. J., and K. Tikkanen. 1997. Microbiological spoilage and contamination of vacuum-packaged cooked sausages: a review. *J. Food Prot.* 60: 724-731.
24. Kurzman, C. P., R. Rogers, and C. W. Hesseltine. 1971. Microbiological contamination of vacuum-packaged processed meats. *Food Microbiol.* 1: 217-224.
25. Lyons, U., J. Björkroth, and H. Korkeala. 1989. Characterisation of lactic acid bacteria from spoiled, vacuum-packaged, cold-smoked rainbow trout by ribotyping. *Int. J. Food Microbiol.* 5: 171-81.
26. Ma'keala P., and H. Korkeala. 1987. Lactobacillus contamination of cooked ring sausages as a source of spoilage lactic acid bacteria. *Int. J. Food Microbiol.* 5: 95-96.
27. Ma'keala P., H. Korkeala, and J. Laitinen. 1990. Raw materials of cooked ring sausages as a source of spoilage lactic acid bacteria. *J. Food Microbiol.* 35: 167-172.
28. Ma'keala P., U. Schillinger, H. Korkeala, and W. H. Ross. 1992. Classification of ropy slime producing lactic acid bacteria based on DNA-homology, and identification of *Lactobacillus sakei*, *Leuconostoc amelibiosum* as dominant spoilage organisms in meat products. *Int. J. Food Microbiol.* 16: 167-172.
29. McFeeters, R. F., H. P. Fleming, and M. A. Daniels. 1956. Lactic acid degradation and brined cucumber bloating. *J. Food Res.* 30: 160-162.
30. Meyer, V. 1956. Die Bestimmung der Bombage-Arten bei Fischkonserven. *Fischwirtschaft.* 22: 224.
31. Mol, J. H. H., J. E. A. Hietbring, H. W. M. Mollen, and J. Van Tinteren. 1987. Observations on the microflora of vacuum packaged sliced cooked meat products. *J. Appl. Bacteriol.* 63: 371-397.
32. Molly, K., D. I. Demeyer, G. Johansson, M. Raemaekers, M. Ghisteln, and I. Geener. 1997. The importance of meat enzymes in ripening and flavour generation in dry fermented sausages: first results of a European project. *Food Chem.* 59: 539-545.
33. Montel, M. C., M. P. Seronine, R. Talon, and M. Hees. 1994. Purification and characterization of a dipeptidase from *Lactobacillus sakei*. *Appl. Environ. Microbiol.* 60: 837-839.
34. Ozbas, Z. Y., H. Vural, and S. A. Altin. 1997. Effects of modified atmosphere and vacuum-packaging on the growth of spoilage and inoculated pathogenic bacteria on fresh poultry. *Fleischwirtschaftl.* 71: 14-16.
35. Pacher, D. G., N. A. Saunders, and R. J. Owen. 1992. Rapid extraction of bacterial genomic DNA with guanidium thiocyanate. *Lett. Appl. Microbiol.* 15: 151-156.
36. Pot, B., P. Vandamme, and K. Kersters. 1984. Analysis of electrophoretic whole-organism protein fingerprints, p. 173-183. In D. F. Toldrano and A. G. Donnel (ed.), *Modern microbiological methods*. Chemical methods of microbiology. John Wiley & Sons Ltd., Chichester, United Kingdom.
37. Sanz, Y., S. Fadda, G. Vignolo, M.-C. Arsitoy, G. Oliveri, and F. Toldrano. Hydrolysis of muscle myofibrillar proteins by *Lactobacillus curvatus* and *Lactobacillus sakei*. *J. Food Microbiol.* 10: 115-125.
38. Sanz, Y., F. Mulholland, and F. Toldrano. 1988. Purification and characterization of a tripeptidase from *Lactobacillus sakei*. *Agric. Food Chem.* 36: 349-353.
39. Sawaya, W. N., A. S. Abu-Ruwaida, Z. H. Baroon, M. S. Khalafawi, and E. Murad. 1993. Shelf-life of vacuum-packaged eviscerated broiler carcasses under simulated market storage conditions. *J. Food Safety* 14: 221-224.
40. Sawaya, W. N., A. S. Abu-Ruwaida, A. J. Hussain, M. S. Khalafawi, and E. Murad. 1993. Shelf-life of eviscerated broiler carcasses as affected by vacuum-packaging and potassium sorbate. *Lebensm. Wiss. Technol.* 16: 525-528.
41. Schleifer, K. H., and O. Kahnt. 1972. Peptidoglycan types of bacterial cement cells and their taxonomic implications. *Bacteriol. Rev.* 36: 410-417.
42. Schleifer, K. H., and W. Ludwig. 1980. Phylogenetic relationships of lactic acid bacteria, p. 7-11. In B. J. B. Wood and W. Holzappel (ed.), *The genera of lactic acid bacteria*. Blackie Academic and Professional, Glasgow, United Kingdom.
43. Shattock, G., and C. D. Hardin. 1989. *Leuconostoc gelidum* nov. and *Leuconostoc carnosum* nov. from chill-stored meats. *Int. J. Syst. Bacteriol.* 39: 217-223.
44. Smittle, R. B., and M. C. Cirigliano. 1982. *Leuconostoc* in salad dressings, p. 976-983. In van der Zandt and D. F. Splittstoesser (ed.), *Compendium of methods for microbiological examination of foods*. American Public Health Association, Washington, D.C.
45. Stackebrandt, E., and W. Liesche. 1986. Nuclear acids and classification of bacteria. *J. Biol. Chem.* 261: 152-159.
46. Goodfellow, A. G., and A. G. O'Donnell (ed.), *Handbook of new bacterial systematics*. Academic Press, London, United Kingdom.
47. Verplaetse, L. 1994. Influence of raw meat properties and processing temperature on aroma quality of raw fermented meat products. *Imp.* 45-65. Proceedings of the 40th International Congress on Meat and Technol. ICOMST, The Hague, The Netherlands.
48. Willianni, F., G. Moschetti, G. Blaiotta, and P. Caporaso. 1997. Characterization of *Leuconostoc mesenteroides* by analysis of soluble whole-cell protein pattern, DNA fingerprinting and restriction of ribosomal DNA. *J. Food Microbiol.* 8: 2578-588.
49. Von Krush, U., and A. Lombardi. 1982. Schnelltest zum qualitativen Nachweis von Laktid (L) und D (D) Milchsäurebakterien. *Milchwissenschaften.* 37: 167-168.
50. Essanjari, J., and B. Raj. 1994. Prevalence and biological control of bacteria producing psychrotrophic leuconostocs associated with spoilage of packaged processed meats. *J. Food Prot.* 17: 209-217.
51. Yoshizumi, H. 1975. A malo-lactic bacterium and its growth factor, p. 102. In J. G. Carr, C. V. Curring, and C. C. Whiting (ed.), *Lactic acid bacteria in beverages and foods*. Academic Press, London, United Kingdom.