

 Open access • Journal Article • DOI:10.1016/J.JBANKFIN.2019.105687

Sparse portfolio selection via the sorted ℓ_1 -Norm — [Source link](#)

Philipp J. Kremer, Sangkyun Lee, Małgorzata Bogdan, Małgorzata Bogdan ...+1 more authors

Institutions: [EBS University of Business and Law](#), [Hanyang University](#), [Lund University](#), [University of Wrocław](#) ...+1 more institutions

Published on: 01 Jan 2020 - [Journal of Banking and Finance](#) (Elsevier)

Topics: [Portfolio optimization](#), [Portfolio](#), [Project portfolio management](#) and [Optimization problem](#)

Related papers:

- [A generalized approach to sparse and stable portfolio optimization problem](#)
- [Large-Scale Regularized Portfolio Selection Via Convex Optimization](#)
- [Robust minimum variance portfolio with L-infinity constraints](#)
- [Solving norm constrained portfolio optimization via coordinate-wise descent algorithms](#)
- [Sparse Markowitz Portfolio Selection by Penalty Methods](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/sparse-portfolio-selection-via-the-sorted-l1-norm-1wapzc0f7o>

Journal Pre-proof

Sparse Portfolio Selection via the sorted ℓ_1 - Norm

Philipp J. Kremer, Sangkyun Lee, Ma Igorzata Bogdan, Sandra Paterlini

PII: S0378-4266(19)30261-4
DOI: <https://doi.org/10.1016/j.jbankfin.2019.105687>
Reference: JBF 105687



To appear in: *Journal of Banking and Finance*

Received date: 26 September 2018

Accepted date: 1 November 2019

Please cite this article as: Philipp J. Kremer, Sangkyun Lee, Ma Igorzata Bogdan, Sandra Paterlini, Sparse Portfolio Selection via the sorted ℓ_1 - Norm, *Journal of Banking and Finance* (2019), doi: <https://doi.org/10.1016/j.jbankfin.2019.105687>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier B.V.

Sparse Portfolio Selection via the sorted ℓ_1 - Norm

Philipp J. Kremer^{a,b,*}, Sangkyun Lee^{c,*}, Małgorzata Bogdan^d, Sandra Paterlini^e

^a*Prime Capital AG, Frankfurt am Main, Germany*

^b*EBS Universität für Wirtschaft und Recht, Wiesbaden, Germany*

^c*Hanyang University ERICA, Ansan, Republic of Korea*

^d*University of Wrocław, Wrocław, Poland*

^e*University of Trento, Italy*

Abstract

We introduce a financial portfolio optimization framework that allows to automatically select the relevant assets and estimate their weights by relying on a sorted ℓ_1 -Norm penalization, henceforth SLOPE. To solve the optimization problem, we develop a new efficient algorithm, based on the Alternating Direction Method of Multipliers. SLOPE is able to group constituents with similar correlation properties, and with the same underlying risk factor exposures. Depending on the choice of the penalty sequence, our approach can span the entire set of optimal portfolios on the risk-diversification frontier, from minimum variance to the equally weighted. Our empirical analysis shows that SLOPE yields optimal portfolios with good out-of-sample risk and return performance properties, by reducing the overall turnover, through more stable asset weight estimates. Moreover, using the automatic grouping property of SLOPE, new portfolio strategies, such as sparse equally weighted portfolios, can be developed to exploit the data-driven detected similarities across

*Corresponding authors

Email addresses: philipp.kremer@ebs.edu (Philipp J. Kremer), sangkyun@hanyang.ac.kr (Sangkyun Lee)

¹The views expressed in this article are those of the authors, and do not necessarily reflect the views of Prime Capital AG.

assets.

Keywords: Portfolio Management, Markowitz Model, Sorted ℓ_1 -Norm
Regularization, Alternating Direction Method of Multipliers

1. Introduction

The development of successful asset allocation strategies requires the construction of portfolios that perform well out-of-sample, provide diversification benefits, and are cheap to maintain and monitor. The problem is then one of statistical model selection and estimation, i.e. the identification of the assets in which to invest and the determination of the optimal weight for each asset.² In 1952, Harry Markowitz laid the foundation for the modern portfolio theory by introducing the mean-variance optimization framework. Assuming that asset returns are normally distributed, such model requires only two input estimates: the vector of expected returns and the expected covariance matrix of the assets. Solving the quadratic optimization problem, by minimizing the portfolio expected risk, for a given level of expected return, the investor can then find the optimal portfolio allocation. Although Markowitz's model has been widely criticized, it is the backbone of the vast majority of portfolio optimization frameworks and is still largely used in practice, especially in fintech companies as part of their robo-advisory (see e.g. Kolm et al. (2014)). One of the major shortcomings of the mean-variance approach is the fact that opti-

²Another stream of literature investigates the utilization of norm penalties in portfolio selection from a behavioral perspective, in which the investor tries to model a simplified version of a complex investment processes. In such context, sparsity allows to simplify the model at hand by focusing the attention on the relevant variables and thereby taking into account the mental cost of processing data. Using the LASSO or the SLOPE penalty still results in tractable models, not NP-Hard ones, which allow a natural way to model investors preference for simpler representation of the world, in which many features are eliminated and sparsity can model dynamic attention to features of the environment. For the behavioral perspective, we refer the interested reader to Gabaix (2014) and references therein.

17 mized weights are highly sensitive to estimation errors and to the presence of mul-
 18 ticollinearity in the inputs. In particular, it is acknowledged that estimating the
 19 expected returns is more challenging, than just focusing on risk minimization and
 20 thereby looking for the portfolios with minimum risk, i.e. the so-called global min-
 21 imum variance portfolios (GMV) (Merton, 1980; Chopra and Ziemba, 1993; Jagan-
 22 nathan and Ma, 2003). But even in the GMV set-up, the sample covariance matrix
 23 might exhibit estimation error that can easily accumulate, especially when dealing
 24 with a large number of assets (Michaud, 1989; Ledoit and Wolff, 2003; DeMiguel and
 25 Nogales, 2009; Fan et al., 2012). Furthermore, multicollinearity and extreme obser-
 26 vations often leads to undesirable and unrealistic extreme long and short positions,
 27 which can hardly be implemented in practice, due to regulatory and short selling
 28 constraints (Shefrin and Statman, 2000; DeMiguel et al., 2009b; Boyle et al., 2012;
 29 Roncalli, 2013). An ideal portfolio then has: a) conservative asset weights, which are
 30 stable in time, to avoid high turnover and transaction costs, and b) still promotes
 31 the right amount of diversification, while being able to control the total amount of
 32 shorting.

33 A natural approach to solve this problem is to extend the Markowitz optimization
 34 framework, by using a penalty function on the weight vector, typically given by the
 35 norm, and whose intensity is controlled by a tuning parameter λ . Probably, the
 36 most recent successful approach, using convex penalty functions, is the Least Abso-
 37 lute Shrinkage and Selection Operator (LASSO) introduced by Tibshirani (1996).
 38 The LASSO framework typically relies on adding to the Markowitz formulation a
 39 penalty proportional to the ℓ_1 -Norm³ on the asset weight vector (Brodie et al., 2009;

³Let $\mathbf{w} = [w_1, w_2, \dots, w_k]'$ be the portfolio weight vector, then the ℓ_q -Norm is defined as: $\|\mathbf{w}\|_q = (\sum_{i=1}^k |w_i|^q)^{\frac{1}{q}}$, with $0 < q < \infty$. If $q = 1$, then $\ell_1 = \sum_{i=1}^k |w_i|$ (LASSO), while for $q = 2$ we have $\|\mathbf{w}\|_2 = (\sum_{i=1}^k w_i^2)^{1/2}$ (RIDGE). Note that ℓ_q with $0 < q < 1$ is not a norm but a quasi norm.

40 DeMiguel et al., 2009a; Carrasco and Noumon, 2012; Fan et al., 2012). DeMiguel
 41 et al. (2009a) provide a general framework that nests regularized portfolio strategies
 42 based on the ℓ_1 -Norm with the approaches introduced by Ledoit and Wolff (2003)
 43 and Jagannathan and Ma (2003). Furthermore, the authors advocate their superior
 44 performance in an out-of-sample setting. Brodie et al. (2009) and Fan et al. (2012)
 45 show that the LASSO (a) results in constraining the gross exposures, (b) can be
 46 used to implicitly account for transaction costs, and (c) sets an upper bound on the
 47 portfolio risk depending just on the maximum estimation error of the covariance ma-
 48 trix. Moreover, the shrinkage covariance estimation of Jagannathan and Ma (2003),
 49 obtained by adding a no-short sale constraint (the so-called GMV long-only (GMV-
 50 LO)), can be considered a special case of the LASSO.

51 Despite its appealing properties, the LASSO has reported shortcomings of (a) large
 52 biased coefficient values (Gasso et al., 2010; Fastrich et al., 2015), of (b) reduced
 53 recovery of sparse signals when applied to highly dependent data, like crisis periods
 54 (Giuzio and Paterlini, 2016), and of (c) randomly selecting among equally correlated
 55 coefficients (Bondell and Reich, 2008). Moreover, it is ineffective in the presence of
 56 no short selling (i.e. $w_i \geq 0$) and an imposed budget constraint (i.e., $\sum_{i=1}^k w_i = 1$),
 57 as the ℓ_1 -Norm is then just equal to 1.

58 To overcome these limitations, we extend the literature on convex regularization
 59 methods in various ways: First, we introduce the *Sorted ℓ_1 Penalized Estimator*
 60 (SLOPE), as a new penalty function within the mean-variance portfolio optimiza-
 61 tion framework. The SLOPE penalty takes the form of a sorted ℓ_1 - Norm, in
 62 which each asset weight is penalized individually using a vector of tuning parame-
 63 ters, $\boldsymbol{\lambda}_{SLOPE} = (\lambda_1, \lambda_2, \dots, \lambda_k)$, with $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k \geq 0$ and whereas $\boldsymbol{\lambda}_{SLOPE}$
 64 is decreasing, attributing the largest weight to the highest regularization parameter,
 65 such that SLOPE penalizes the weights according to their rank magnitude. This

66 leads to an octagonal shape of the penalty in a 2-dimensional setting (see Figure 1)
 67 that combines the two favorable properties of the ℓ_∞ -Norm and the ℓ_1 -Norm⁴, which
 68 is the grouping of variables (i.e. some asset weights are assigned the same coefficient
 69 value), and the singularity at the origin, respectively. Our work shows that, opposed
 70 to the LASSO, in portfolio optimization and together with an added budget con-
 71 straint (i.e. $\sum_{i=1}^k w_i = 1$), SLOPE continues to shrink the active weights, even when
 72 short sales are restricted (i.e. $w_i \geq 0, \forall i = 1, \dots, k$). Consequently, it spans the
 73 diversification frontier from the GMV-LO up to the equally weighted (EW) portfolio,
 74 as λ_{SLOPE} goes to infinity. Together with the feature of grouping equally correlated
 75 assets, the penalty provides an increased flexibility for the investor in creating indi-
 76 vidual trading strategies, as opposed to state-of-the-art shrinkage methods.
 77 Second, we introduce a new optimization algorithm to solve the mean-variance port-
 78 folio problem with the sorted ℓ_1 regularization and linear constraints on the asset
 79 weights. The algorithm uses the ideas of variable splitting and the Alternating Direc-
 80 tion Method of Multipliers (ADMM) framework (Powell, 1969; Hestenes, 1969; Boyd
 81 et al., 2011). Using a mathematically equivalent reformulation of the original prob-
 82 lem, the algorithm can use existing implementations of proximal operators (Parikh
 83 and Boyd, 2014), associated with the ℓ_1 , the sorted ℓ_1 , and even other regularizers.
 84 Furthermore, Appendix C shows that the ADMM provides a more efficient alter-
 85 native for solving the LASSO optimization problem, than the state-of-art Cyclic
 86 Coordinate Descend (CyCoDe) algorithm.
 87 Third, we are, to our knowledge, the first to investigate the properties of SLOPE un-
 88 der a realistic factor model, which assumes that all assets can be represented as linear
 89 combination of a small number of hidden risk factors, as e.g. in Fan et al. (2008).

⁴Given a weight vector \mathbf{w} with k elements, the $\ell_\infty = \|\mathbf{w}\|_\infty = \max(w_1, \dots, w_k)$.

90 In the set-up of classical multiple regression, in which the explanatory variables are
91 assumed independent, Bogdan et al. (2013, 2015) and Su and Candès (2016) provide
92 extensive evidence of SLOPE’s superior model selection and estimation properties.
93 Further evidence for these properties are provided by the results of Bellec et al.
94 (2016a) and Bellec et al. (2016b), which show that contrary to LASSO, SLOPE is
95 asymptotically optimal for the general class of design matrices satisfying the modi-
96 fied Restricted Eigenvalue condition.

97 Moreover, Bondell and Reich (2008) and Figueiredo and Nowak (2014) investigate
98 the properties of SLOPE and its predecessor OSCAR (Octagonal Shrinkage and Se-
99 lection Operator, Bondell and Reich (2008)) in the situation, when regressors are
100 strongly correlated. Bondell and Reich (2008) apply OSCAR to agricultural data,
101 showing that the method successfully forms predictive clusters, which can then be
102 analyzed according to their individual characteristics. Figueiredo and Nowak (2014)
103 illustrate the “clustering” properties of the *ordered weighted ℓ_1 - Norm* (OWL) in
104 the linear regression framework with strongly correlated predictors, providing fur-
105 ther simulation and theoretical results. However, none of these works addresses the
106 interesting situation, in which the correlation structure results from the dependency
107 of the explanatory variables on a few hidden factors and on financial real-world data.
108 Recently, Xing et al. (2014) applied the OSCAR to the mean-variance portfolio op-
109 timization, together with a linear combination of the ℓ_1 - and the ℓ_∞ -Norms. They
110 advocate the method for its ability to identify portfolios that attain higher Sharpe Ra-
111 tios and lower turnovers compared to those resulting from traditional approaches like
112 the GMV and the GMV-LO portfolios. However, they do not point out the clumping
113 property of the OSCAR. With SLOPE, we consider a generalized framework that
114 nests the GMV, the GMV-LO, the LASSO, the ℓ_∞ - Norm and the approach of Xing
115 et al. (2014).

116 In this paper, we analyze the properties of SLOPE, with both simulated and real
117 world data. The simulations show that SLOPE reduces the estimation errors in the
118 portfolio weights and groups assets depending on the same risk factors together. This
119 grouping behavior then allows the investor to select individual constituents from the
120 clusters, for example based on her preferences and asset-specific properties, enabling
121 her to develop new investment strategies such as SLOPE-EW, which we introduce
122 in Section 4.1.

123 For the real world data analysis, we use monthly returns of the 10- and 30-Industry
124 portfolios (Ind), as well as the 100 Fama French (FF) portfolios formed on Size and
125 Book-to-Market, covering the period from 1970 to 2017. Furthermore, we consider
126 daily returns of the S&P 500 (SP500) from 2004 to 2016. Our results show that the
127 risk of the SLOPE portfolio is comparable to or smaller than the risk of the LASSO
128 portfolio. Also, we observe that SLOPE outperforms the LASSO, yielding better
129 risk- and weight diversification measures. In fact, the sorted ℓ_1 - Norm is able to span
130 the entire risk-diversification frontier, starting from the GMV, via the GMV-LO up
131 to the EW. The investor can then select the portfolio with the risk-diversification
132 trade-off that best fits her preferences.

133 The above mentioned characteristics establish SLOPE as a new attractive portfolio
134 construction alternative, capable of controlling short sales and identifying groups of
135 assets. It thereby offers the possibility to implement individual views, which goes
136 beyond the standard statistical shrinkage or regularization approaches.

137 The paper is structured as follows: Section 2 introduces our methodology and dis-
138 cusses the properties of SLOPE. Section 3 analyses the behavior of SLOPE in simu-
139 lated environments, while Section 4 focuses on the empirical results. Section 5 con-
140 cludes.

141 **2. Sparse Portfolio Selection via the Sorted ℓ_1 -Norm**

142 Given k jointly normally distributed asset returns R_1, \dots, R_k , with expected value
 143 vector $\boldsymbol{\mu} = [\mu_1, \dots, \mu_k]'$ and covariance matrix $\boldsymbol{\Sigma}$, a generalization to the Markowitz
 144 (1952) portfolio selection problem can be stated as the following optimization:

$$\min_{\boldsymbol{w} \in \mathbb{R}^k} \frac{\phi}{2} \boldsymbol{w}' \boldsymbol{\Sigma} \boldsymbol{w} - \boldsymbol{\mu}' \boldsymbol{w}, \quad \text{subject to} \quad \sum_{i=1}^k w_i = 1 \quad (1)$$

145 where $\sigma_p^2 = \boldsymbol{w}' \boldsymbol{\Sigma} \boldsymbol{w}$ is the portfolio risk, $\boldsymbol{\mu}' \boldsymbol{w}$ is the portfolio return and $\phi > 0$ is the
 146 coefficient of relative risk aversion (Markowitz, 1952; Fan et al., 2012; Li, 2015).

147 Despite the advantage of being a quadratic optimization problem, the generalized
 148 Markowitz model is often criticized, as it leads to extreme and unstable optimal
 149 portfolio weights.

150 One approach to circumvent instability and extreme estimates is to modify the opti-
 151 mization problem (1), by adding a penalty function, $\rho_\lambda(\boldsymbol{w})$.⁵ In typical applications,
 152 the penalty is a non-decreasing function of \boldsymbol{w} and leads to shrinking an estimate of
 153 the weight vector towards zero. The shrinkage stabilizes the weight estimates and, in
 154 case when $\rho_\lambda(\boldsymbol{w})$ has a singularity at zero, promotes the sparsity by shrinking some
 155 coordinates of the estimated vector to 0. An additional parameter λ controls the
 156 impact of the penalty and thereby the amount of shrinkage applied to the weights
 157 vector and the level of sparsity. Additionally, the penalty function allows to take
 158 into account a prior knowledge of an investor, who can assign smaller penalty terms

⁵Another stream of literature focusses on directly shrinking the moments of the distribution, as opposed to adding a norm penalty on the weight vector to the model in (1). However, as have been shown by DeMiguel et al. (2009a), adding a norm constraint on the weight vector to the portfolio optimization is equal to shrinking the extreme estimates in the covariance matrix. For elaborations on directly improving the inputs to the Markowitz optimization, the interested reader is referred to Ledoit and Wolff (2003, 2004a), Jorion (1986) and references therein.

159 to selected important assets. The optimization problem can be stated as:

$$\min_{\mathbf{w} \in \mathbb{R}^k} \frac{\phi}{2} \mathbf{w}' \boldsymbol{\Sigma} \mathbf{w} - \boldsymbol{\mu}' \mathbf{w} + \rho_\lambda(\mathbf{w}) \quad \text{s.t.} \quad \sum_{i=1}^k w_i = 1 \quad (2)$$

160 The simplest approach is the LASSO, which considers as a penalty function the ℓ_1 -
 161 Norm of the asset weights vector ($\rho_\lambda(\mathbf{w}) = \lambda \times \sum_{i=1}^k |w_i|$, with λ being a scalar).
 162 The resulting optimization problem is still convex, while promoting model selection
 163 and estimation in a single step. From a financial perspective, LASSO is interpreted
 164 as a gross exposure constraint (i.e. a constraint on the total amount of shorting)
 165 or a way to account for transaction costs (Brodie et al., 2009). However, it is not
 166 effective in the presence of both a budget ($\sum_{i=1}^k w_i = 1$), and a no-short selling (i.e.,
 167 $w_i \geq 0$) constraint, as the ℓ_1 -norm is then simply equal to 1.

168 Following, we propose a more general approach that within a single optimization
 169 algorithm allows us to encompass the original LASSO, the OSCAR of Bondell and
 170 Reich (2008), and the combination of ℓ_1 and ℓ_∞ penalties, as proposed in Xing et al.
 171 (2014).

172 In fact, we penalize the weights vector by considering as $\rho_\lambda(\mathbf{w})$ the sorted ℓ_1 -Norm,
 173 defined as:

$$\rho_\lambda(\mathbf{w}) := \sum_{i=1}^k \lambda_i |w|_{(i)} = \lambda_1 |w|_{(1)} + \lambda_2 |w|_{(2)} + \dots + \lambda_k |w|_{(k)} \quad (3)$$

s.t. $\lambda_1 \geq \lambda_2 \geq \dots \lambda_k \geq 0$ and $|w|_{(1)} \geq |w|_{(2)} \geq \dots |w|_{(k)}$,

174 where $|w|_{(i)}$ denotes the i th largest element in absolute value of the vector \mathbf{w} . The
 175 sorted ℓ_1 -Norm was originally introduced in Bogdan et al. (2013, 2015) to con-
 176 struct the Sorted ℓ_1 Penalized Estimator for the selection of explanatory variables

177 in the multiple regression model. It was also developed independently by Zeng and
 178 Figueiredo (2014) as Ordered Weighted ℓ_1 Norm (OWL). To our knowledge, this is
 179 the first work in financial portfolio selection that applies SLOPE and discusses its
 180 grouping properties, while also introducing a new optimization algorithm.


181 *2.1. Geometric Interpretation*

182 Compared to most of the other regularization methods, SLOPE does not rely
 183 on a single tuning parameter λ , but rather on a non-increasing sequence $\boldsymbol{\lambda}_{SLOPE} =$
 184 $(\lambda_1, \lambda_2, \dots, \lambda_k)$, with $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k \geq 0$. This sequence is aligned to the sorted
 185 weight vector, such that the largest absolute weight is penalized with the largest
 186 tuning parameter. Consequently, the sequence of λ parameters gives a natural in-
 187 terpretation of importance to the asset weights, besides providing full flexibility in
 188 recapturing the profiles of the ℓ_1 - and ℓ_∞ - Norms, as well as of their linear combina-
 189 tions. Figure 1 shows a simple set-up with two assets and the respective shapes of
 190 spheres (i.e. the set points for which $\rho_\lambda(\mathbf{w}) = c$) that we obtain, depending on how
 191 the sequence $\boldsymbol{\lambda}_{SLOPE} = (\lambda_1, \lambda_2)$ is chosen. As shown in Panel (a), if $\lambda_1 = \lambda_2 > 0$ the
 192 SLOPE sphere coincides with the well studied diamond shape of the LASSO penalty.
 193 Through its singularity at the origin, the LASSO promotes sparse solutions that set
 194 one of the two assets' weights exactly equal to zero. On the other hand, choosing
 195 $\lambda_2 = 0$ and $\lambda_1 > 0$, yields the regularization term of the ℓ_∞ -Norm. The respective
 196 shape, as shown in Panel (b), takes the form of a square and promotes the grouping
 197 of variables, i.e. it encourages solutions under which both asset weights are assigned
 198 exactly the same value.

199 Given these two extreme cases, Panel (c) of Figure 1 shows the octagonal shape of
 200 SLOPE, obtained by using a decreasing sequence of lambda values, with $\lambda_1 > \lambda_2 > 0$.
 201 By choosing different tuning parameter sequences, the penalty allows to approximate

202 a variety of norms between the ℓ_1 and the ℓ_∞ , combining the properties of the Lasso
 203 and the ℓ_∞ penalties and due to its singularity, is either able to set some weights
 204 exactly equal to zero, and/or to assign the same value to some of the other weights.
 205 Furthermore, it approximates the shape of the already well studied RIDGE penalty,
 206 which corresponds to a circle in the 2-dimensional set-up, and is even able to reach
 207 one of RIDGE's special solutions, i.e. the equally weighted portfolio, which is ob-
 208 tained, when RIDGE's penalty parameter approaches infinity. Although RIDGE
 209 is still convex, the shape of the penalty does not promote sparsity among the co-
 210 efficients, leading to undesirable portfolios with a large number of active positions
 211 (Carrasco and Noumon, 2012; DeMiguel et al., 2009a). Thus, the choice of the
 212 lambda sequence for SLOPE provides the investor with the flexibility to choose any
 213 of these shapes of the unit sphere and of the corresponding mode of shrinking the
 214 dimension of the weight vector.

Figure 1: Geometric Representation of Penalty Functions



For two asset weights $\mathbf{w} = [w_1 \ w_2]'$, the figure shows the unit spheres for different SLOPE sequences: (a) the LASSO ℓ_1 sphere, when $\lambda_1 = \lambda_2 > 0$, (b) the ℓ_∞ sphere, when $\lambda_1 > \lambda_2 = 0$ and (c) the SLOPE sphere, when $\lambda_1 > \lambda_2 > 0$. The dashed lines in (c) represent the diamond shape of the LASSO and the RIDGE ℓ_2 -balls, respectively.

215

216 In portfolio optimization, a budget constraint that requires the weights of the port-


217 folio to sum to one, is imposed. Consequently, we discuss how the penalties behave
 218 in the presence of such an additional constraint. Figure 2 plots the SLOPE penalty,
 219 together with the LASSO and the RIDGE penalty for a universe of two assets and
 220 under the condition that $w_1 + w_2 = 1$. Furthermore, we consider the penalty func-
 221 tions in the presence of short sales (gray area) and no short sales (white area).

222 In Figure 2, we can see that the LASSO (shown in black) is only effective when
 223 short sales are permitted, while the presence of the budget constraint makes it in-
 224 effective in the long-only area. In contrast, the RIDGE attains its minimum for an
 225 equally weighted portfolio, and when short sales are restricted. Similarly, the SLOPE
 226 penalty (shown in red) also reaches its minimum at the equally weighted solution
 227 (i.e., $w_1 = w_2 = 0.5$). Still, to control for monitoring and transaction costs of fi-
 228 nancial assets, we prefer SLOPE over the RIDGE estimator, because it can promote
 229 sparsity by exploiting the singularities.

230

231 Figure 3 plots the contours of the objective function, together with those of the
 232 SLOPE spheres for the two asset case, and when we do not impose a budget con-
 233 straint (i.e. $\sum_{i=1}^k w_i = 1$), as well as considering orthogonal and correlated designs.
 234 As noted before, if only $\lambda_2 > 0$, SLOPE always has singular points when one of the
 235 asset weights is equal to zero, thereby promoting sparsity. When $\lambda_1 > \lambda_2 > 0$, that
 236 is, the sequence is monotonically decreasing, then SLOPE has additional singular
 237 points, which correspond to $|w_1| = |w_2|$. This is an appealing property in the pres-
 238 ence of correlated data. Specifically, as Panel (b) shows, strong correlation between
 239 assets lead to the same weights and thereby grouping. This is consistent with port-
 240 folio theory, as it is known that, if assets have all the same correlation coefficients,
 241 as well as identical means and variances, the EW is the unique optimal portfolio.
 242 SLOPE then allows us to automatically group assets with similar correlation.

Figure 2: Penalty Functions in a Two Asset Universe with Budget Constraint




The figure plots the SLOPE coefficient alongside the LASSO ($\ell_1 - Norm$) and the RIDGE penalty ($\ell_2 - Norm$), for a two asset case and under the condition that $w_1 + w_2 = 1$.

243

244 For our simulation analysis and empirical investigations, SLOPE requires us to define
 245 a specific form of the sequence of $\boldsymbol{\lambda}_{SLOPE} = (\lambda_1, \lambda_2, \dots, \lambda_k)$. For that, we use the
 246 decreasing sequence of quantiles of the standard normal distribution, as in Bogdan
 247 et al. (2013) and Bogdan et al. (2015), with $\lambda_i = \alpha \Phi^{-1}(1 - q_i)$, $\forall i = 1, \dots, k$, where
 248 Φ is the cumulative distribution function of the standard normal distribution and
 249 $q_i = i \times \theta / 2k$, and in which $\theta = 0.01$, regulates how fast the sequence of lambda
 250 parameters is decreasing. Bogdan et al. (2013) and Bogdan et al. (2015) have shown
 251 that in orthogonal design this sequence controls the False Discovery Rate in a multiple
 252 testing framework.⁶

⁶We investigated different sequences of lambda parameters, including changing values of θ , as well as a linear decreasing sequence and obtained qualitative similar results. Consequently, we

Figure 3: Sorted ℓ_1 -Norm Penalty without Budget Constraint

The figure plots in Panel (a) and (b), the Sorted ℓ_1 -Norm Penalty (SLOPE) in a 2-dimensional setting, considering orthogonal design and correlated design, respectively.

253 2.2. Optimization Algorithm

254 In this section, we describe our solution algorithm, which is based on equivalent
 255 reformulations of the Alternating Direction Method of Multipliers (ADMM) approach
 256 (see Appendix A for details).

choose the exponentially decreasing sequence, as proposed by Bogdan et al. (2013) and Bogdan et al. (2015). Still, future research on how to choose the sequence of lambda parameters is currently high on our agenda.

Algorithm 1 ADMM Algorithm

-
- 1: Input: Expected value vector $\boldsymbol{\mu} \in \mathbb{R}^k$, covariance matrix $\Sigma \in \mathbb{R}^{k \times k}$.
 - 2: Initialize $\boldsymbol{w}^0 \in \mathbb{R}^k$, $\boldsymbol{v}^0 \in \mathbb{R}^k$, $\boldsymbol{\alpha}^0 \in \mathbb{R}^k$, $\beta^0 \in \mathbb{R}$ and $j = 0$.
 - 3: Given a stopping threshold value $\tau > 0$.
 - 4: **while** $\mathcal{G}(\boldsymbol{w}^j, \boldsymbol{v}^j, \boldsymbol{\alpha}^j, \beta^j) > \tau$ **do**
 - 5: Update \boldsymbol{w}^j , \boldsymbol{v}^j , $\boldsymbol{\alpha}^j$, and β^j as follows,

$$\begin{cases} \boldsymbol{w}^{j+1} &= \arg \min_{\boldsymbol{w}} \mathcal{L}_{\eta}(\boldsymbol{w}, \boldsymbol{v}^j; \boldsymbol{\alpha}^j, \beta^j) = (\phi \Sigma + \eta(I + \boldsymbol{e}\boldsymbol{e}'))^{-1}(\boldsymbol{\mu} - \boldsymbol{\alpha}^j - \beta^j \boldsymbol{e} + \eta(\boldsymbol{v}^j + \boldsymbol{e})) \\ \boldsymbol{v}^{j+1} &= \arg \min_{\boldsymbol{v}} \mathcal{L}_{\eta}(\boldsymbol{w}^{j+1}, \boldsymbol{v}; \boldsymbol{\alpha}^j, \beta^j) = \text{prox}_{\lambda/\eta}(\boldsymbol{w}^{j+1} + (1/\eta)\boldsymbol{\alpha}^j) \\ \boldsymbol{\alpha}^{j+1} &= \boldsymbol{\alpha}^j + \eta(\boldsymbol{w}^{j+1} - \boldsymbol{v}^{j+1}) \\ \beta^{j+1} &= \beta^j + \eta(\boldsymbol{e}'\boldsymbol{w}^{j+1} - 1) \end{cases} \quad (4)$$

6: $j = j + 1$

7: **end while**

257 In Algorithm 1, the quantity $\mathcal{G}(\boldsymbol{w}^j, \boldsymbol{v}^j, \boldsymbol{\alpha}^j, \beta^j)$ represents the primal-dual gap which
 258 converges to the zero value when the iterates $\boldsymbol{w}^j, \boldsymbol{v}^j, \boldsymbol{\alpha}^j, \beta^j$ approaches the optimal
 259 quantities (see Appendix B for details). In the presence of the no-short selling con-
 260 straint, we consider a slightly different formulation to (2) with an extra constraint
 261 that $\boldsymbol{w} \geq 0$. In this case, we can use the Algorithm 1 almost as it is, except that the
 262 w update in (4) is modified as follows,

$$\boldsymbol{w}^{j+1} = \arg \min_{\boldsymbol{w} \geq 0} \mathcal{L}_{\eta}(\boldsymbol{w}, \boldsymbol{v}^j; \boldsymbol{\alpha}^j, \beta^j) = \max \{ (\phi \Sigma + \eta(I + \boldsymbol{e}\boldsymbol{e}'))^{-1}(\boldsymbol{\mu} - \boldsymbol{\alpha}^j - \beta^j \boldsymbol{e} + \eta(\boldsymbol{v}^j + \boldsymbol{e})), 0 \} \text{ ,} \quad (5)$$

263 where the minimizer is obtained by adding a simple clipping operation, since $L_{\eta}(\boldsymbol{w}, \boldsymbol{v}^j;$
 264 $\boldsymbol{\alpha}^j, \beta^j)$ is a convex function in \boldsymbol{w} .

265 Our algorithm can also be used to solve the LASSO optimization problem, which

266 is a specific instance of SLOPE. In Appendix C, we provide a direct comparison of
 267 our algorithm to the state-of-art Cyclic Coordinate Descent (CyCoDe) for LASSO,
 268 considering a simulated constant correlation model.

269

270 **Bounds on the Objective Function.** To solve the mean-variance problem, as
 271 stated in (2), the investor needs to provide an estimate of the true covariance matrix
 272 of asset returns Σ and of the true mean μ , which are in the most simplest form given
 273 by the sample covariance matrix $\hat{\Sigma}$ and the sample mean $\hat{\mu}$, respectively. However, $\hat{\Sigma}$
 274 and $\hat{\mu}$ might be prone to substantial estimation errors and highly sensitive to outliers.
 275 Let us define $M(\Sigma, \mu) = \frac{\phi}{2} \mathbf{w}' \Sigma \mathbf{w} - \mathbf{w}' \mu$, where \mathbf{w} is the vector of weights returned
 276 by SLOPE. Now, observe that the Sorted ℓ_1 -Norm satisfies $\rho_\lambda(\mathbf{w}) \geq \lambda_k \|\mathbf{w}\|_1$. Thus,
 277 as $\lambda_k > 0$, we have $\|\mathbf{w}\|_1 \leq c$, with $c = \frac{\rho_\lambda(\mathbf{w})}{\lambda_k}$, and simple calculations following the
 278 results of Fan et al. (2012) for LASSO, yield:

$$|M(\hat{\Sigma}, \hat{\mu}) - M(\Sigma, \mu)| \leq \frac{\phi}{2} \|\hat{\Sigma} - \Sigma\|_\infty \rho_\lambda^2(\mathbf{w}) / \lambda_k^2 + \|\hat{\mu} - \mu\|_\infty \rho_\lambda(\mathbf{w}) / \lambda_k \quad (6)$$

279 where $\|\hat{\Sigma} - \Sigma\|_\infty$ and $\|\hat{\mu} - \mu\|_\infty$ are the maximum component-wise estimation errors
 280 for the covariance matrix and the expected return.

281 This result implies that the difference between the objective functions for the esti-
 282 mated and true vector of parameters decreases, as we restrict the Sorted ℓ_1 -Norm of
 283 the weight vector. It is also important to observe that, due to the budget constraint,
 284 a higher weight on the penalty sets an upper bound on the total amount of short
 285 sales in the portfolio, as $\rho_\lambda(\mathbf{w}) \geq \lambda_k \|\mathbf{w}\|_1 = \lambda_k (\mathbf{w}^+ + \mathbf{w}^-)$, with $\mathbf{w}^+ - \mathbf{w}^- = 1$, where
 286 $\mathbf{w}^+ = \sum_{w_i \geq 0} w_i$ and $\mathbf{w}^- = \sum_{w_i < 0} w_i$ are the gross amount of long and short positions,
 287 respectively.

288 3. Simulation Analysis

289 This section investigates the effect of SLOPE on the model risk, the sparsity and
 290 the grouping properties, by considering simulated data. The purpose of the simula-
 291 tions is to investigate the properties of our new penalty, when the data generating
 292 mechanism is completely known, so that the results can be compared to the so-called
 293 *oracle* solution. Furthermore, and as is it widely acknowledged that the estimation
 294 errors in $\boldsymbol{\mu}$ are much larger than in $\boldsymbol{\Sigma}$, we focus on a risk minimization framework.
 295 Assuming $\boldsymbol{\Sigma}$ to be known, we can use the alternative formulation of SLOPE and
 296 define:

$$\mathbf{w}_{opt} = \arg \min_{\mathbf{w}: \sum_{i=1}^k w_i=1, \rho_{\lambda(\mathbf{w})} \leq c} \mathbf{w}' \boldsymbol{\Sigma} \mathbf{w} \quad \text{and} \quad \hat{\mathbf{w}}_{opt} = \arg \min_{\mathbf{w}: \sum_{i=1}^k w_i=1, \rho_{\lambda(\mathbf{w})} \leq c} \mathbf{w}' \hat{\boldsymbol{\Sigma}} \mathbf{w} \quad (7)$$

297 whereas \mathbf{w}_{opt} and $\hat{\mathbf{w}}_{opt}$ are the theoretical optimal and the empirical optimal weights
 298 vector, respectively. We then define the *empirical* portfolio risk as $\widehat{Risk}(\hat{\mathbf{w}}_{opt}) =$
 299 $\hat{\mathbf{w}}_{opt}' \hat{\boldsymbol{\Sigma}} \hat{\mathbf{w}}_{opt}$, the *actual* portfolio risk as $Risk(\hat{\mathbf{w}}_{opt}) = \hat{\mathbf{w}}_{opt}' \boldsymbol{\Sigma} \hat{\mathbf{w}}_{opt}$ and the *oracle*
 300 portfolio risk as $Risk(\mathbf{w}_{opt}) = \mathbf{w}_{opt}' \boldsymbol{\Sigma} \mathbf{w}_{opt}$, respectively. Following the proof of Theo-
 301 rem 1 of Fan et al. (2012), we can easily show that when $\lambda_k > 0$, the pair differences
 302 between the three measures are upper bounded by:

$$|Risk(\hat{\mathbf{w}}_{opt}) - Risk(\mathbf{w}_{opt})| \leq 2c^2 \|\hat{\boldsymbol{\Sigma}} - \boldsymbol{\Sigma}\|_{\infty}, \quad (8)$$

$$|Risk(\hat{\mathbf{w}}_{opt}) - \widehat{Risk}(\hat{\mathbf{w}}_{opt})| \leq c^2 \|\hat{\boldsymbol{\Sigma}} - \boldsymbol{\Sigma}\|_{\infty}, \quad (9)$$

$$|Risk(\mathbf{w}_{opt}) - \widehat{Risk}(\hat{\mathbf{w}}_{opt})| \leq c^2 \|\hat{\boldsymbol{\Sigma}} - \boldsymbol{\Sigma}\|_{\infty} \quad (10)$$

303 The three risk measures then allow us to extract different information: The empirical
 304 risk is the only one that is known, as it is estimated from our in-sample data. The
 305 actual risk is the one, to which the investor is truly exposed to, when using the

306 estimated optimal weights ($\hat{\boldsymbol{w}}_{opt}$). Finally, the oracle risk is the risk the investor
 307 could only obtain, if $\boldsymbol{\Sigma}$ is known. As the SLOPE penalty becomes more binding,
 308 when $\boldsymbol{\lambda}$ "increases", the three risk measures align. In the following section, we
 309 investigate how increasing the SLOPE penalty allows to reduce the estimation error
 310 and to avoid its accumulation in the portfolio risk.

311 Assume that the return of an asset is represented by a linear combination of r risk
 312 factors. Furthermore, let t be the number of observations, k be the number of assets,
 313 and $\boldsymbol{F}_{t \times r} = [\boldsymbol{f}_1 \ \boldsymbol{f}_2 \ \dots \ \boldsymbol{f}_r]$, where \boldsymbol{f}_i is the $t \times 1$ vector of returns of the i^{th} risk
 314 factor. Moreover, let $\boldsymbol{B}_{r \times k}$ be the loading matrix for the individual risk factors.
 315 Then, the $t \times k$ matrix of asset returns from the Hidden Factor Model (i.e. \boldsymbol{R}_{HF})
 316 can be represented as:

$$\boldsymbol{R}_{HF} = \boldsymbol{F} \times \boldsymbol{B} + \boldsymbol{\epsilon} \quad (11)$$

317 where $\boldsymbol{\epsilon}$ is a $t \times k$ matrix of error terms.

318 For our first illustration of the performance of SLOPE, we generate the data using
 319 the following simplified scenario:

- 320 • $t = 50$, $k = 12$, $r = 3$,
- 321 • the risk factors f_1, \dots, f_3 are independent from the multivariate standard
 322 normal $N(0, I_{r \times r})$ distribution, with $I_{r \times r}$ being an identity matrix,
- 323 • the vectors of error terms ϵ_i , $i = 1, \dots, k$, for each asset are independent
 324 from each other, as well as from each of the risk factors and come from the
 325 multivariate normal distribution $N(0, 0.05 \times I_{r \times r})$
- 326 • the loadings matrix $\boldsymbol{B}_{r \times k}$ is made of exactly four copies of each of the following

327 columns: $[0.77 \ 0.64 \ 0]'$, $[0.9 \ 0 \ -0.42]'$ and $[0 \ 0.31 \ 0.64]'$.

328 In this way, we generate three different groups that have the same exposure to the
329 same two risk factors and are thus strongly correlated.⁷

330 Finally, given (11), the covariance matrix of the assets Σ_{HF} is given by:

$$\Sigma_{HF} = \mathbf{B}'\mathbf{B} + 0.05 \times I_{k \times k}. \quad (12)$$


331 After generating our $t \times k$ matrix \mathbf{R}_{HF} of asset returns from (11), we can then estimate
332 Σ_{HF} , using the sample covariance estimate $\hat{\Sigma}_{HF}$.⁸ Figure 4 shows the correlation
333 matrix resulting from (12), illustrating that our simulation scenario explicitly models
334 a block correlation environment, with strong correlation among each of the assets
335 having the same underlying risk factor exposures, and low to negative correlations
336 between the assets with a different underlying factor structure. Following, we inves-
337 tigate the behavior of SLOPE and the LASSO with respect to portfolio risk, and
338 when we increase the value of the tuning parameter.

339 Unlike the LASSO, SLOPE requires us to define a decreasing sequence of $\lambda_{SLOPE} =$
340 $(\lambda_1, \lambda_2, \dots, \lambda_k)$. As pointed out in Section 2.1, we use the decreasing sequence of
341 quantiles of the standard normal distribution, as in Bogdan et al. (2013) and Bog-
342 dan et al. (2015), with $\lambda_i = \alpha \Phi^{-1}(1 - q_i)$, $\forall i = 1, \dots, k$, where Φ is the cumulative
343 distribution function of the standard normal distribution and $q_i = i \times \theta/2k$, and in
344 which $\theta = 0.01$, regulates how fast the sequence of lambda parameters is decreasing.

⁷For the robustness of our results, we tested SLOPE in various set-ups, with qualitatively similar results. Due to space limitations, we report only the most interesting one. The results of the remaining simulations are available from the authors upon request.

⁸We explicitly restrict us to use the sample covariance estimate, as opposed to an alternative shrinkage or factor based estimate, to investigate SLOPE's ability to account for estimation errors in the optimization.

Figure 4: Hidden Factors Correlation Matrix




The figure plots the correlation matrix, based on the modeled Hidden Factor Structure, considering a universe of 12 assets, of which 4 are always exposed to exactly two out of the three risk factors in the market.

345 In our simulations, we vary the scaling parameter α so that the first element of the
 346 sequence $\lambda_1 = \alpha\Phi^{-1}(1 - q_1)$ is equal to a grid of 100 log-spaced values between 10^{-5}
 347 and 10^2 . Note that in the case of the LASSO, we only choose one lambda param-
 348 eter, which then remains constant for all assets. Throughout the paper, we always
 349 choose $\lambda_{LASSO} = \lambda_1$. This choice favors sparser solutions for the LASSO, since for
 350 the remaining $k - 1$ assets its penalty is larger than that of SLOPE.

351 Figure 5 plots the resulting risk and weight profile for the minimum variance opti-
 352 mization, when we solve (2) separately with the LASSO and the SLOPE penalties for
 353 the grid of 100 lambda parameters, and considering Σ_{HF} and the sample covariance
 354 estimate $\hat{\Sigma}_{HF}$, respectively. In particular, Panels (a) and (b) show the risk profile of
 355 the LASSO and SLOPE, i.e. the actual, the oracle, and the empirical risk, together
 356 with the results of the GMV, the GMV-LO and the EW portfolios. For both, the
 357 oracle and the actual solution, Panels (c) and (d) display on top the number of active
 358 weights together with the number of groups, that is the number of distinct coeffi-

Figure 5: Hidden Factors Minimum-Variance Profile




The figure shows the Hidden Factor minimum-variance risk profile for the LASSO and the SLOPE, including in Panel (a) and (b) their actual, empirical and oracle risk profiles, together with that of the GMV, the GMV-LO and the EW solutions. Furthermore, Panel (c) and (d) display the number of active weights, together with the grouping profile (top) and the total amount of shorting (bottom). All values are computed based on a Hidden Factor Structure, with three risk factors and considering for the exponentially decreasing sequence of lambda parameters, a grid of 100 log spaced starting points for λ_1 from 10^{-5} (i.e. x-value = 1) to 10^2 (i.e. x-value = 100).

359 cients, while on the bottom, it shows the amount of shorting (i.e. w^-). The grey
 360 surface indicates the no-short-sale-area (i.e. $w_i \geq 0 \forall i = 1, \dots, k$). Figure 5 shows
 361 that for a tuning parameter equal to zero, which corresponds to the GMV solution,
 362 the empirical risk is about 1.3 times lower than the actual risk (Panels (a) and (b)),

363 with 12 active positions (Panel (c)) and slightly under 100% short sales (Panel (d)).
 364 This can be interpreted as evidence that in over-fitted models the estimation error in
 365 $\widehat{\Sigma}_{HF}$ strongly affects the estimation of the asset weights. As here neither the LASSO
 366 nor the SLOPE penalty are binding, estimation errors can enter unhindered into the
 367 optimization. Michaud (1989) describes this phenomenon as “error maximization”,
 368 in which the ill-conditioned covariance estimates are amplified through the optimiza-
 369 tion, leading to extreme long and short portfolio weights. Moving along the grid of λ
 370 parameters from the left to the right, Panels (c) and (d) show that the two penalties
 371 reduce the total amount of shorting in the oracle and the actual portfolio.
 372 As we move from the GMV towards the GMV-LO, the actual, oracle, and empirical
 373 risk of the LASSO and the SLOPE align. This effect was first observed and theo-
 374 retically motivated by Fan et al. (2012), showing that the portfolio risk evolves in
 375 a U-shape, in which risk first decreases before increasing again, due to the restric-
 376 tion of short sales. With the observations above, we extend the results of Fan et al.
 377 (2012), showing that the U-shaped behavior of the portfolio risk is not the only pos-
 378 sible one. Especially when the dependence among the assets is positive, the tighter
 379 constraint in terms of short sales shrinks the optimization search space of feasible
 380 solutions, making it impossible to exploit the optimal diversification benefits. This
 381 leads to a higher portfolio risk when reaching the GMV-LO. The investor also reaches
 382 the maximum sparsity, that is the maximum number of coefficients equal to zero, at
 383 this point. For the LASSO, increasing the tuning parameter beyond this point does
 384 not alter the allocation any further, as the regularization penalty is constant and
 385 equal to 1.
 386 This is different for SLOPE: in fact, Figure 6 shows the evolution of the portfolio
 387 weights for both the oracle and the actual solution, considering both the LASSO and
 388 the SLOPE penalty. As before, the grey surface indicates the no-short-sale-area.

From Figure 6, we can observe two important characteristics of SLOPE: First,

Figure 6: Hidden Factors Minimum-Variance Weight Profiles



The figure shows the weight profile of the oracle (top) and actual (bottom) solution of the LASSO and the SLOPE penalty, considering a minimum variance setup. All values are computed based on a Hidden Factor Structure, with three risk factors and considering for the exponentially decreasing sequence of lambda parameters, a grid of 100 log spaced starting points for λ_1 from 10^{-5} (i.e. x-value = 1) to 10^2 (i.e. x-value = 100). Equally colored weights characterize assets with the same underlying factor exposure.

389

390 while the LASSO shrinks the weights up until the no short sale area, all non-zero
 391 coefficients still receive a different weight, independent of their underlying factor ex-
 392 posures. SLOPE, on the other hand, is able to identify the three distinct types of
 393 securities, consistent with the true model, and groups them together, by assigning
 394 the same coefficient values to them. This provides information about the dependence
 395 structure among the assets, and gives the investor the flexibility to select from the
 396 groups the assets, which best fit her individual preferences. Not surprisingly, the
 397 oracle risk starts to form groups among the securities even before entering into the
 398 no short sale area, while the actual weights can only capture the underlying structure
 399 much later, and when we already impose a larger tuning parameter value. Second,
 400 and different to the LASSO, increasing the lambda parameters past the point of
 401 the GMV-LO, the octagonal shape of the penalty pushes the solution towards the

402 equally weighted portfolio. That is, the aforementioned grouping effect increases,
 403 and all weights - even those that were shrunk towards zero - are assigned the
 404 same coefficient value of $\frac{1}{k}$. Given that the equally weighted portfolio is only optimal
 405 when all assets have the same risk and return characteristics, in our example, this
 406 allocation results in higher portfolio risk when compared to the GMV-LO or GMV
 407 portfolios.

408 Summing up, SLOPEs properties allows investors to set up sophisticated asset alloca-
 409 tion strategies, exploiting its grouping property, like SLOPE-EW, which we introduce
 410 in Section 4.

411 4. Empirical Analysis

412 4.1. Set up and Data

413 This section studies the out-of-sample performance of the SLOPE procedure in
 414 a minimum variance framework (see i.e. Jagannathan and Ma (2003); Brodie et al.
 415 (2009); DeMiguel et al. (2009a); Giuzio and Paterlini (2016)) and compare it with
 416 state-of-the-art portfolio selection methods, such as the EW, the GMV, the GMV-
 417 LO, the equal risk contribution (ERC), the RIDGE and the LASSO portfolio. Fur-
 418 thermore, we examine two extensions to our standard SLOPE procedure: (1) SLOPE
 419 with an added long-only constraint (SLOPE-LO) and (2) a portfolio in which we uti-
 420 lize SLOPE-LO's selection and grouping ability (SLOPE-EW). For the latter, the
 421 portfolio is initialized in $t = 1$, keeping for each data set only the first G groups with
 422 the largest estimated parameter values active, while setting the remaining weights
 423 equal to zero. Afterwards, the portfolio is re-scaled, such that the weights sum again
 424 to one. At each subsequent t , we then rebalance the portfolio, if there is a statistically
 425 significant difference in the covariance matrices, at $\alpha = 0.1$, to the last re-balance

426 date.⁹

427 In the following analysis, we consider four data sets, including the monthly log-return
 428 observations for the 10- and 30 Industry Portfolios (Ind), the 100 Fama French (FF)
 429 portfolios, formed on Size and Book-to-Market, as well as the daily returns of the
 430 SP500. The monthly portfolio values are taken from Kenneth French’s Homepage¹⁰
 431 and span the period from January 1970 to January 2017 ($T = 565$ monthly obser-
 432 vations). The daily return data are obtained from Datastream, covering the period
 433 from 31.12.2004 to 31.01.2016 ($T = 2890$ daily observations). Table 1 reports the
 434 descriptive statistics. As shown by the skewness and the kurtosis values, all of them
 435 exhibit the typical return time series characteristics, including fat tails and slight
 436 asymmetry.

Table 1: Descriptive Statistics of the Dataset

Dataset	T	k	$\hat{\mu}$	$\hat{\sigma}$	\widehat{med}	\widehat{min}	\widehat{max}	\widehat{skew}	\widehat{kurt}	<i>period</i>	<i>freq.</i>
10Ind	565	10	0.099	0.043	0.012	-0.211	0.156	-0.476	5.077	01/1970 - 01/2017	Monthly
30Ind	565	30	0.010	0.048	0.012	-0.255	0.179	-0.507	5.749	01/1970 - 01/2017	Monthly
100FF	565	100	0.011	0.052	0.015	-0.262	0.241	-0.551	5.600	01/1970 - 01/2017	Monthly
SP500	2890	443	0.000	0.014	0.000	-0.107	0.109	-0.418	13.234	12/2004 - 01/2016	Daily

The table reports descriptive summary statistics for the 10 Industry Portfolios, the 30 Industry Portfolios, the 100 Fama French Portfolios and the S&P 500, respectively. Reported are for the daily (monthly) data: the number of observations (T), the number of constituents(k), the mean ($\hat{\mu}$), the standard deviation ($\hat{\sigma}$), the median (\widehat{med}), the minimum (\widehat{min}), the maximum (\widehat{max}), the skewness (\widehat{skew}), the kurtosis (\widehat{kurt}), the period that the data set covers (*period*) and the frequency (*freq.*).

437

438 To evaluate the portfolios in an out-of-sampling setting, we rely on a rolling window
 439 approach with a window size of $\tau = 120$ monthly observations for the 10Ind, the

⁹We use the MBox test (see e.g. Stevens (1992)) to test, if there is a significant difference in the covariance matrices. Furthermore, we perform robustness tests for $\alpha = 0.01, 0.05$, which are available from the authors upon request.

¹⁰<http://mba.tuck.dartmouth.edu/pages/faculty/ken.french/>

440 30Ind, and the 100FF, as well as $\tau = 500$ daily observations for the SP500.¹¹ All
 441 portfolios are re-balanced monthly, discarding the oldest and including the most
 442 recent observations, allowing for a total of $t = 445$ ($t = 115$) out-of-sample returns
 443 for the monthly (daily) data.

444 The rolling window approach for the daily data works as follows: the first τ return
 445 observations are used to estimate $\hat{\Sigma}_t$, according to the shrinkage approach by Ledoit
 446 and Wolff (2004b). Then, $\hat{\Sigma}$ is used as the input to compute the optimal weight
 447 vector $\hat{\boldsymbol{w}}_t$. The resulting portfolio is assumed to be held for the following 21 days.
 448 At $t + 1$, the k constituents' returns over this monthly period, \boldsymbol{R}_{t+1} , are used to
 449 compute the out-of-sample portfolio return as: $R_{p,t+1} = \hat{\boldsymbol{w}}_t \boldsymbol{R}_{t+1}$. In the next step,
 450 we roll the data window forward, dropping the last and adding the most recent
 451 21 observations to our training set. We then estimate a new weight vector, which
 452 determines our portfolio holdings and the out-of-sample return for the next month.
 453 This process is repeated until the end of the data set is reached. The same procedure
 454 is applied to the Industry and Fama French portfolios, though the window is rolled
 455 forward by one monthly observation instead of 21 daily observations.

456 Finally, and given that each data set consists of a different number of assets, we keep
 457 for our trading strategy, SLOPE-EW, and depending on the respective data set the
 458 following first G groups active: 10Ind - the first 4 groups; 30Ind - the first 2 groups;
 459 100Ind - the first 5 groups; SP500 - the first 5 groups.¹²

¹¹To test the robustness of our results, we account for different window sizes of $\tau = 250, 750$ and 1000 daily observations, and make the results available upon request. The obtained results are qualitative similar.

¹²Note: The stated number of active groups G for each data set are selected, such that we always obtain the portfolios with the lowest out-of-sample variance, given $\alpha = 0.1$ and when compared to portfolios that would include more or less groups, respectively. Results for different number of included groups G , as well as different significance levels, i.e. $\alpha = 0.05$ or 0.01 , are available from the author upon request.

460 For all portfolios, the optimal weights vector, $\hat{\boldsymbol{w}}_t$, depends on the choice of the op-
461 timal λ parameter value. To select the optimal tuning parameter, we consider a
462 grid of 100 log-spaced values of λ between $10^{-7.5}$ and 10^1 , from which we choose
463 $\lambda_{RIDGE} = \lambda_{LASSO} = \lambda_1 = \alpha\Phi^{-1}\left(1 - \frac{0.01}{2k}\right)$. The remaining elements $i = 2, \dots, k$ of
464 the λ sequence for SLOPE are as before, equal to $\lambda_i = \alpha\Phi^{-1}\left(1 - \frac{0.01i}{2k}\right)$.
465 Among the 100 lambda values, we select the optimal tuning parameter for the
466 RIDGE, the LASSO and the SLOPE, such that we obtain a portfolio with approx-
467 imately 10% of the GMV's short positions. Note that for SLOPE, as we increase
468 the tuning parameter, beyond the GMV-LO solution, we would move along the no-
469 short sale area towards the EW portfolio (see Figure 5). Therefore, we also compute
470 SLOPE-LO to explicitly exploit the grouping feature that predominates in the long-
471 only area, and select the lambda value, which provides us with a portfolio that has
472 the largest number of groups. To guarantee that all our portfolios can also be imple-
473 mented in practice, all weights that are smaller in absolute value than the threshold
474 of 0.05% are set to zero. Furthermore, we incorporate a transaction cost (TC) regime
475 of $TC = 10\text{bps}^{13}$, whereas the costs are proportional to the turnover and considered
476 to be the same for selling and buying securities.¹⁴
477 Given the optimal portfolio vector $\hat{\boldsymbol{w}}_t$ at time t , we compute the out-of-sample mean

¹³1 bps = 0.01% = 0.0001

¹⁴For the robustness of our results, we also consider regimes of no (TC = 0bps) and high trans-
action costs (TC = 50bps). They show that naturally, high turnover strategies like the GMV suffer
with regard to returns and the SR in the higher cost regimes. On the other hand, SLOPE portfolios
show a nearly steady performance for all data sets and when considering the different TC regimes.
All results are available upon request from the authors.

478 and the out-of-sample standard deviation, defined as:

$$\hat{\mu}_p = \frac{1}{t} \sum_{i=1}^t \hat{\mathbf{w}}_t \mathbf{R}_{t+1} \quad (13)$$

$$\hat{\sigma}_p = \sqrt{\frac{1}{t-1} \sum_{i=1}^t (\hat{\mathbf{w}}_t \mathbf{R}_{t+1} - \hat{\mu}_p)^2} \quad (14)$$

479 from which we construct the out-of-sample Sharpe Ratio (SR) as:

$$\widehat{SR} = \frac{\hat{\mu}_p}{\hat{\sigma}_p} \quad (15)$$

480 To evaluate whether the \widehat{SR} and $\hat{\sigma}_p^2$ of any portfolio is statistically different from our
481 SLOPE procedure, we use the tests developed by Ledoit and Wolf (2008) and Ledoit
482 and Wolf (2011), respectively.

483 As frequent re-balancing of a portfolio is costly, we complement our analysis by
484 computing the turnover of each portfolio, defined as:

$$\widehat{TO} = \frac{1}{t} \sum_{i=1}^t \|\hat{\mathbf{w}}_{t+1} - \hat{\mathbf{w}}_t^+\|_1, \quad (16)$$

485 whereas $\hat{\mathbf{w}}_t^+$ is the weight vector right before rebalancing at $t + 1$ and considering
486 the changes in the assets prices. Consequently, the TO for the EW can be non-zero,
487 as $\hat{\mathbf{w}}_t^+ \neq \hat{\mathbf{w}}_{t+1} = 1/k$ (DeMiguel et al., 2009a).

488 Furthermore, we include the following diversification measures: the Diversification
489 Ratio (DR), the weight (WDiv) and the risk diversification (RDiv) measures. The
490 DR is defined as the ratio of the weighted asset volatility to the overall portfolio

491 volatility:

$$\widehat{\text{DR}} = \frac{\sum_{i=1}^k \hat{w}_i \hat{\sigma}_i}{\hat{\sigma}_p}, \quad (17)$$

492 where $\hat{\sigma}_i$ is the i -th asset's estimated volatility, $\hat{\sigma}_p$ is the estimated portfolio volatility,
 493 for which the investor typically prefers a higher value (Choueifaty and Coignard,
 494 2008).

495 Finally, both the WDiv and RDiv measure the concentration of the portfolio in terms
 496 of weights and risk (Maillard et al., 2010; Roncalli, 2013). The WDiv ranges from $\frac{1}{k}$
 497 for a perfectly concentrated portfolio up to 1 for the equally weighted portfolio. It
 498 is computed according to:

$$\widehat{\text{WDiv}} = \frac{1}{k \times \sum_{i=1}^k \hat{w}_i^2} \quad (18)$$

499 On the other hand, we obtain the RDiv by substituting the weights for the risk
 500 contribution, defined as $\widehat{RC}_i = \hat{w}_i \times \partial_{w_i} \sigma(\hat{\mathbf{w}}_i)$, where $\partial_{w_i} \sigma(\hat{\mathbf{w}}_i)$, defines the marginal
 501 contribution to risk (MRC) of asset i , that is the first derivative of the portfolio
 502 variance with respect to portfolio weight w_i . The MRC measures the sensitivity of
 503 the portfolio variance, given a change in the i -th asset. The RDiv takes a value of 1 for
 504 the equally-weighted risk contributions (ERC) portfolio, which is least concentrated
 505 in terms of risk contributions and $\frac{1}{k}$ for a portfolio which is fully concentrated on one
 506 asset:

$$\widehat{\text{RDiv}} = \frac{1}{k \times \sum_{i=1}^k \widehat{RC}_i^2} \quad (19)$$

507 Summing up, we prefer values close to one for the WDiv and the RDiv (Cazalet
508 et al., 2014).

509 4.2. Empirical Results

510 Table 2 reports the annualized out-of-sample volatility, the annualized out-of-
511 sample SR, the number of active positions, and the turnover for the 10Ind, 30Ind, the
512 100FF, and the SP500, using a window size of $\tau = 120$ ($\tau = 500$) observations with
513 monthly re-balancing and $TC = 10$ bps. We indicate portfolios that are statistically
514 different from our SLOPE procedure at the 10%, 5% and 1% level, given the test
515 for the difference in the SR and the volatility, following Ledoit and Wolf (2008)
516 and Ledoit and Wolf (2011).

Table 2: Risk- and Return Measures

	Vol. (in %)				Sharpe Ratio				AP				Turnover			
	10Ind	30Ind	100FF	SP500	10Ind	30Ind	100FF	SP500	10Ind	30Ind	100FF	SP500	10Ind	30Ind	100FF	SP500
EW	14.491	16.257	17.509	20.238	0.776**	0.656***	0.677***	0.205	10.000	30.000	100.000	443.000	0.049	0.057	0.056	0.077
GMV	10.910	9.152	6.058	11.497	1.102*	1.283***	3.124***	0.057**	9.982	29.885	99.469	434.377	0.125	0.273	0.852	2.748
GMV-LO	11.473	11.214	13.134	10.825	1.012	0.998***	0.954***	0.389	5.371	8.562	9.220	27.553	0.064	0.074	0.101	0.238
ERC	13.578	15.029	16.906	17.948	0.840*	0.730***	0.714***	0.233	10.000	30.000	100.000	443.000	0.048	0.054	0.055	0.076
RIDGE	11.907	12.241	13.219	11.393	0.978	0.955	1.069***	0.490	9.989	29.824	98.171	408.474	0.061	0.078	0.109	0.212
LASSO	11.364	10.781	10.853	9.505	1.028	1.046	1.421***	0.572	6.755	12.301	18.371	130.211	0.079	0.104	0.184	0.434
SLOPE	11.352	10.822	10.977	9.643	1.024	1.047	1.382	0.534	7.027	13.231	22.598	145.552	0.078	0.101	0.172	0.409
SLOPE - LO	11.689	11.865	13.709	11.981	0.950**	0.946**	0.917***	0.344	7.299	18.465	34.616	129.632	0.119	0.217	0.409	0.590
SLOPE - EW	12.539	12.904	14.390	11.017	0.908**	0.929*	0.858***	0.645	6.330	6.128	17.022	76.386	0.052	0.055	0.057	0.284

The table reports the out-of-sample Risk and Return Measures for the 10-, 30-, and 100-Portfolios (SP500), considering a window size of $\tau = 120$ monthly ($\tau = 500$ daily) observations and re-balancing the portfolio every month over the period from 01/1970 to 01/2017 (from 12/2004 to 01/2016). Furthermore, we consider a transaction cost of 10bps, which is proportional to the turnover and is assumed to be the same for selling and buying securities. Reported are: The annualized out-of-sample volatility, the annualized out-of-sample Sharpe Ratio, the number of active positions (AP), and the average total turnover. Furthermore, we report the significance for the test of the difference in the volatility and the SRs with regard to SLOPE, at the 10%, 5% and 1% level with *, ** and ***, respectively.

517

518 Looking at the values for the out-of-sample volatility in Table 2, we observe that no
519 portfolio is statistically different from our new SLOPE procedure, across any of the
520 data sets. Still, SLOPE yields consistently lower variance than any of the EW, ERC,

521 RIDGE or GMV-LO portfolios. Furthermore, for the SP500, SLOPE and LASSO
522 perform best, reporting the smallest variance among all strategies. Especially for
523 the SP500, the number of observations in the window size is only marginally bigger
524 than the size of our investment universe, the estimated covariance matrix is degen-
525 erated and our estimates are very prone to estimation error. Therefore, and even
526 using the shrunk covariance matrix, SLOPE and LASSO are still able to reduce
527 extreme weight estimates. Simultaneously, we explicitly select for the LASSO and
528 the SLOPE, a portfolio with a moderate amount of short sales, making it possible
529 to still exploit diversification benefits. Hence, the resulting allocation has a smaller
530 variance, as compared to the GMV-LO.

531 Furthermore, the values for the out-of-sample SR, establish SLOPE among the best
532 performing portfolios, across all datasets, with some results being statistically sig-
533 nificant. For example, SLOPE is able to statistically significantly outperform the
534 EW, challenging its widely reported characteristic of a tough benchmark to beat
535 (DeMiguel et al., 2009b).

536 Beside reducing the overall portfolio variance, our goal is to construct sparse port-
537 folios with a low turnover. For that, reconsider that the EW always invests naively
538 in all constituents and thus has the highest possible number of active positions.
539 Similar values are obtained for the ERC, which aims at equalizing the risk contribu-
540 tion of each asset to the overall portfolio risk. The GMV, as being highly sensitive
541 to even small changes in the underlying data structure, typically resulting in ex-
542 treme positions (see i.e. Michaud (1989)), has the highest turnover values among
543 the non-regularization strategies. The RIDGE, on the other hand, results in more
544 stable asset allocations, despite not setting any asset weight exactly equal to zero.
545 Although both strategies should invest in all assets, the reported number of active
546 positions are slightly reduced, due to our imposed threshold of 5%.

547 Compared to the strategies above, our new SLOPE procedure is able to promote
548 sparse solutions and to reduce the overall portfolio turnover, consistently reporting
549 lower turnover values than the LASSO.

550 Of special interest is also the performance of SLOPE-EW. In general, our new SLOPE
551 procedure provides the investor with a large amount of flexibility, as with an increased
552 lambda value the penalty starts to form groups among assets, assigning to them the
553 same coefficient value. This is of special interest for investors, who want to move be-
554 yond the property of statistical shrinkage, and who want to include in their portfolio
555 construction process any form of financial indicator, like among others fundamen-
556 tal multiples (i.e. Price/Earnings, Dividends/Earnings), accounting values (i.e., Net
557 Income, Free Cash Flow) or other quantitative measures (i.e., Value-at-Risk or Ex-
558 pected Shortfall). With SLOPE-EW, we construct a simple strategy that selects,
559 out of the formed groups, those which carry assets that are the most important with
560 regard to minimizing the overall portfolio variance. Still, other strategies could be
561 easily developed.

562 Table 2 shows that SLOPE-EW performs best in reducing the variance for large
563 asset universes, i.e. for the SP500, even outperforming the initial SLOPE-LO port-
564 folio. This result provides two insights: First, using SLOPE-EW, we can eliminate
565 assets from the portfolio that rather increase the portfolio variance, as opposed to
566 reducing it, and second, by eliminating the groups according to the weight mag-
567 nitude, we might conclude that SLOPE assigns assets to the groups according to
568 their importance in reducing the overall variance. Finally, SLOPE-EW ranks among
569 the portfolios with the smallest number of active positions, and reports the lowest
570 turnover value, across all sparse portfolio methods.

571

572 Table 3 complements our risk and return analysis, reporting the DR, the WDiv, and

Table 3: Diversification Measures


	DR				WDiv				RDiv			
	10Ind	30Ind	100FF	SP500	10Ind	30Ind	100FF	SP500	10Ind	30Ind	100FF	SP500
EW	1.270	1.343	1.212	1.675	1.000	1.000	1.000	1.000	0.933	0.935	0.958	0.894
GMV	1.255	1.362	0.958	3.147	0.197	0.078	0.013	0.012	0.197	0.078	0.013	0.012
GMV-LO	1.289	1.414	1.299	1.944	0.320	0.150	0.062	0.032	0.320	0.150	0.062	0.032
ERC	1.300	1.382	1.225	1.728	0.935	0.914	0.963	0.880	1.000	1.000	1.000	1.000
RIDGE	1.330	1.457	1.256	1.920	0.540	0.430	0.262	0.248	0.577	0.440	0.188	0.120
LASSO	1.289	1.415	1.237	2.221	0.309	0.143	0.044	0.062	0.301	0.132	0.030	0.029
SLOPE	1.295	1.426	1.247	2.213	0.319	0.155	0.054	0.071	0.312	0.144	0.056	0.032
SLOPE - LO	1.315	1.457	1.295	1.936	0.417	0.287	0.209	0.206	0.437	0.319	0.221	0.219
SLOPE - EW	1.289	1.314	1.294	1.808	0.403	0.181	0.118	0.170	0.408	0.182	0.120	0.166

The table reports the diversification measures for the 10-, 30-, and 100- Portfolios (SP500 Portfolios), considering a window size of $\tau = 120$ monthly ($\tau = 500$ daily) observations and re-balancing the portfolio every month over the period from 01/1970 to 01/2017 (from 12/2004 to 01/2016). Reported are: The Diversification Ratio (DR), the Weight Diversification (WDiv) and the Risk Diversification (RDiv) measures.

573 the RDiv. As the EW invests equally in all assets, it achieves, by definition, the best
574 values for the WDiv, with similar values reported for the ERC. As the ERC aims to
575 equalize the contribution to portfolio risk from each asset, it also reports the highest
576 values for the RDiv. SLOPE-LO and SLOPE consistently outperform the LASSO
577 across all datasets for the WDiv and the RDiv. Except for the SP500, this is also
578 true for the DR, while a higher value for the LASSO only results due to the lower
579 variance, as reported in Table 2. It should be pointed out that SLOPE does not only
580 frequently outperform the LASSO, but also provides flexibility with regard to the
581 diversification measures. For that, Figure 7 plots the weight- and risk diversification
582 measure against the attainable portfolio volatility for the LASSO and the SLOPE,
583 together with the other portfolio strategies and considering the first window size of
584 $\tau = 120$ observations for the 10Ind.

585 For both frontiers, the full grid of lambda parameters for the LASSO enables the
586 investor to select only a combination between the GMV and the GMV-LO solution.
587 SLOPE, on the other hand, is able to span a much larger set of portfolios, beginning

Figure 7: Risk and Weight Diversification Frontier



The figure shows on the left the weight diversification and on the right the risk diversification frontier, both reporting on the x-axis the portfolio volatility and on the y-axis the risk and weight diversification measure, respectively. Considered are the first window size of $\tau = 120$ months for the 10Ind. Plotted are the resulting combinations for the GMV, the GMV-LO, the EW, the ERC, as well as the different combinations for the LASSO and the SLOPE procedure, considering a range of lambda values from $10^{-7.5}$ to 10^1 .

588 from the GMV, via the GMV-LO up to the EW. The investor can thus control the
 589 trade-off between diversification and volatility out of a much larger set of portfolios,
 590 to find the allocation that best suits her individual preferences.

591

592 5. Conclusion

593 This paper extends the literature on financial regularization by introducing SLOPE
 594 to the Markowitz portfolio optimization, discussing its properties and testing its per-
 595 formance with regard to risk and return on simulated and real world data.

596 SLOPE relies on a sorted ℓ_1 -Norm, whose intensity is controlled by a decreasing
 597 sequence of λ parameters and which penalizes the assets by their rank, provid-
 598 ing a natural interpretation of importance. To solve the penalized mean-variance
 599 optimization, we propose a novel algorithm based on the Alternating Direction

600 Method of Multipliers (ADMM). When applied to the LASSO, which is a specific
601 case of SLOPE, this algorithm provides the same accuracy as the state-of-the-art
602 CyCoDe, but is superior with regard to computing time, especially when the as-
603 set universe is large.

604 The simulated hidden risk factor analysis shows that SLOPE has the advantage of
605 still being active in the no short sales area and given an imposed budget constraint.
606 Furthermore, SLOPE can automatically identify assets with the same underlying risk
607 factor exposure and group them together, by assigning the same coefficient value to
608 them. This property is especially desirable for investor planning to incorporate their
609 individual views into the optimization, by selecting assets from these groups accord-
610 ing to a specific financial characteristic or individual preferences. We exploit such
611 property by building a simple investment strategy, SLOPE-EW.

612 Moreover, we investigate the performance of SLOPE for four major data sets to
613 other state-of-art portfolio methods in an out-of-sample setting, considering a rolling
614 window approach, and re-balancing the portfolio every month.

615 Our results show that SLOPE is able to achieve equal and even better out-of-sample
616 portfolio volatilities and SR, when compared to the LASSO. Although, only part of
617 the differences are statistically significant, SLOPE is able to construct sparse portfo-
618 lios with reduced turnover. This especially applies to situations with a large amount
619 of estimation error, for example when considering the SP500. Furthermore, our
620 SLOPE-EW portfolio results in very sparse portfolios with even lower turnover than
621 state-of-the-art methods and at the same time maintains a comparable performance.
622 Additionally, SLOPE reports improved values for the DR, the WDiv and the RDiv,
623 while the shape of the penalty extends the frontier of attainable portfolios, ranging
624 from the GMV via the GMV-LO, up to the EW portfolio. This enables the investor
625 to select among them the one that provides her with the desired volatility- and di-

626 versification trade-off.

627 The results establish SLOPE as a valid alternative to state-of-art methods by creating
628 sparse portfolios with a reduced turnover rate, improved risk- and weight diversifi-
629 cation, and a high degree of flexibility in the portfolio construction process.

630 A natural extension to our study is to investigate, how different sequences of lambda
631 parameters would impact the risk and portfolio allocation, and whether the investor
632 should choose them according to the underlying correlation regime of the stock mar-
633 ket or his own prior beliefs on the assets.

634 Appendix A. Derivation of the ADMM Algorithm

635 In order to facilitate the application of proximal operators involving ρ_λ , we first
636 reformulate (2) - (3) into the following form:

$$\min_{\mathbf{w} \in \mathbb{R}^k, \mathbf{v} \in \mathbb{R}^k} \frac{\phi}{2} \mathbf{w}' \boldsymbol{\Sigma} \mathbf{w} - \boldsymbol{\mu}' \mathbf{w} + \rho_\lambda(\mathbf{v}) \quad \text{s.t.} \quad \mathbf{w} = \mathbf{v}, \quad \sum_{i=1}^k w_i = 1, \quad (\text{A.1})$$

637 where $\rho_\lambda(\mathbf{w}) := \sum_{i=1}^k \lambda_i |w|_{(i)}$ is the sorted ℓ_1 -Norm corresponding to the sequence
638 $\boldsymbol{\lambda}_{SLOPE} = (\lambda_1, \dots, \lambda_k)'$ satisfying $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k \geq 0$. To solve (A.1), we design an
639 ADMM (for details, see e.g. Boyd et al. (2011)) algorithm, which is based on using
640 the augmented Lagrangian function of (A.1) and on partial updates for the primal
641 variables. In our case the associated augmented Lagrangian is given as:

$$\begin{aligned} \mathcal{L}_\eta(\mathbf{w}, \mathbf{v}; \boldsymbol{\alpha}, \beta) &= \frac{\phi}{2} \mathbf{w}' \boldsymbol{\Sigma} \mathbf{w} - \boldsymbol{\mu}' \mathbf{w} + \rho_\lambda(\mathbf{v}) + \boldsymbol{\alpha}'(\mathbf{w} - \mathbf{v}) + \beta(\mathbf{e}' \mathbf{w} - 1) \\ &\quad + \frac{\eta}{2} \{ \|\mathbf{w} - \mathbf{v}\|^2 + (\mathbf{e}' \mathbf{w} - 1)^2 \}, \end{aligned} \quad (\text{A.2})$$

642 where $\boldsymbol{\alpha} \in \mathbb{R}^k$, $\beta \in \mathbb{R}$, $\mathbf{e}_{k \times 1} = (1, \dots, 1)'$, $\mathbf{I}_{k \times k}$ is the identity matrix, and $\eta > 0$ is

643 a penalty parameter. Compared to the Lagrangian L_0 without the penalty term,
 644 the augmented Lagrangian \mathcal{L}_η with $\eta > 0$ brings the benefit that the dual objective
 645 $g_\eta(\boldsymbol{\alpha}, \beta) := \inf_{\mathbf{w}, \mathbf{v}} L_\eta(\mathbf{w}, \mathbf{v}; \boldsymbol{\alpha}, \beta)$ becomes differentiable without requiring further
 646 assumptions on the primal objective (e.g., strict convexity).
 647 The ADMM algorithm consists of the updates:

$$\begin{cases} w^{j+1} &= \arg \min_{\mathbf{w}} \mathcal{L}_\eta(\mathbf{w}, \mathbf{v}^j; \boldsymbol{\alpha}^j, \beta^j) = (\phi \boldsymbol{\Sigma} + \eta(I + \mathbf{e}\mathbf{e}'))^{-1}(\boldsymbol{\mu} - \boldsymbol{\alpha}^j - \beta^j \mathbf{e} + \eta(\mathbf{v}^j + \mathbf{e})) \\ v^{j+1} &= \arg \min_{\mathbf{v}} \mathcal{L}_\eta(\mathbf{w}^{j+1}, \mathbf{v}; \boldsymbol{\alpha}^j, \beta^j) = \text{prox}_{\lambda/\eta}(w^{j+1} + (1/\eta)\boldsymbol{\alpha}^j) \\ \alpha^{j+1} &= \alpha^j + \eta(w^{j+1} - v^{j+1}) \\ \beta^{j+1} &= \beta^j + \eta(\mathbf{e}'w^{j+1} - 1), \end{cases} \quad (\text{A.3})$$

648 where $\text{prox}_{\lambda/\eta}(\mathbf{z}) := \arg \min_{\mathbf{v}} \frac{1}{2}\|\mathbf{v} - \mathbf{z}\|_2^2 + \rho_{\lambda/\eta}(\mathbf{v})$ is the proximal operator of the
 649 Sorted ℓ_1 -Norm, corresponding to the sequence λ/η , provided e.g. in Bogdan et al.
 650 (2013, 2015). The updates regarding α and β are due to the gradient ascent applied to
 651 the dual objective $g_\eta(\boldsymbol{\alpha}, \beta) := \inf_{\mathbf{w}, \mathbf{v}} L_\eta(\mathbf{w}, \mathbf{v}; \boldsymbol{\alpha}, \beta)$, where $\nabla_{\boldsymbol{\alpha}} g_\eta(\boldsymbol{\alpha}, \beta) = w^{j+1} - v^{j+1}$
 652 and $\nabla_{\beta} g_\eta(\boldsymbol{\alpha}, \beta) = \mathbf{e}'w^{j+1} - 1$. The first iterates $\mathbf{w}^0, \mathbf{v}^0, \boldsymbol{\alpha}^0, \beta^0$ of the procedure (4)
 653 are typically initialized as the zero vectors.

654 Appendix B. Primal-Dual Gap

655 The stopping criterion for our algorithm is based on the Primal-Dual Gap, which we
 656 estimate as follows. First, taking the infimum over (\mathbf{w}, \mathbf{v}) of the Lagrangian, we get
 657 the dual objective,

$$g(\boldsymbol{\alpha}, \beta) = \inf_{\mathbf{w}} \frac{\phi}{2} \mathbf{w}' \boldsymbol{\Sigma} \mathbf{w} - (\boldsymbol{\mu} - \boldsymbol{\alpha} - \beta \mathbf{e})' \mathbf{w} - \beta - \rho_{\lambda}^*(\boldsymbol{\alpha}). \quad (\text{B.1})$$

658 From the optimality condition for the infimum over \mathbf{w} , we have

$$\mathbf{w}^* = \phi^{-1} \boldsymbol{\Sigma}^{-1} (\boldsymbol{\mu} - \boldsymbol{\alpha} - \beta \mathbf{e}). \quad (\text{B.2})$$

659 Also,

$$\rho_{\lambda}^*(\boldsymbol{\alpha}) = \sup_{\mathbf{v}} \{ \boldsymbol{\alpha}^T \mathbf{v} - \rho_{\lambda}(\mathbf{v}) \} = \begin{cases} 0 & \text{if } \boldsymbol{\alpha} \in C_{\lambda} \\ +\infty & \text{o.w.} \end{cases} \quad (\text{B.3})$$

660 where $C_{\lambda} := \{ \mathbf{v} : \mathbb{R}^k : \rho_{\lambda}^D(\mathbf{v}) \leq 1 \}$ is the unit sphere defined in the dual norm $\rho_{\lambda}^D(\cdot)$
661 of $\rho_{\lambda}(\cdot)$. Plugging-in these, we get the dual problem

$$\max_{\boldsymbol{\alpha}, \beta} -\frac{1}{2\phi} (\boldsymbol{\mu} - \boldsymbol{\alpha} - \beta \mathbf{e})' \boldsymbol{\Sigma}^{-1} (\boldsymbol{\mu} - \boldsymbol{\alpha} - \beta \mathbf{e}) - \beta \quad \text{s.t. } \boldsymbol{\alpha} \in C_{\lambda}. \quad (\text{B.4})$$

662 Then we can estimate the primal-dual gap as follows using (B.2),

$$\begin{aligned} \mathcal{G}(\mathbf{w}^*, \mathbf{v}^*, \boldsymbol{\alpha}^*, \beta^*) &= \frac{1}{2} \phi (\mathbf{w}^*)' \boldsymbol{\Sigma} \mathbf{w}^* - \boldsymbol{\mu}' \mathbf{w}^* + \rho_{\lambda}(\mathbf{w}^*) + \frac{1}{2\phi} (\boldsymbol{\mu} - \boldsymbol{\alpha}^* - \beta^* \mathbf{e})' \boldsymbol{\Sigma}^{-1} (\boldsymbol{\mu} - \boldsymbol{\alpha}^* - \beta^* \mathbf{e}) + \beta^* \\ &= -(\boldsymbol{\alpha}^* + \beta^* \mathbf{e})' \mathbf{w}^* + \beta^* + \rho_{\lambda}(\mathbf{w}^*) \end{aligned} \quad (\text{B.5})$$

663 given the dual feasibility of $\boldsymbol{\alpha}^*$, i.e., $\rho_{\lambda}^D(\boldsymbol{\alpha}^*) \leq 1$. Here, \mathbf{w}^* , \mathbf{v}^* , $\boldsymbol{\alpha}^*$, β^* can be
664 generated from the procedure (4), and, due to strong duality, the duality gap becomes
665 zero when these iterates are optimal to the problem (A.1). Therefore we can stop our
666 algorithms when the duality gap of the current iterates becomes sufficiently small.

667 Appendix C. ADMM vs. Cyclic Coordinate Descend

668 In this section, we use the ADMM algorithm to solve the minimum-variance opti-
669 mization with an ℓ_1 Norm (which is a specific instance of our new SLOPE penalty)

670 and compare its performance to the the Cyclic Coordinate Descend algorithm (Cy-
671 CoDe).

672 The CyCoDe algorithm is considered state-of-art and has found various applications
673 in solving norm constrained optimization problems (see i.e. Fastrich et al. (2014),
674 Yen (2015)). The algorithm works by optimizing the weights along one coordinate
675 direction, while holding all other weights constant. Although there is no general
676 rule on how the CyCoDe updates the weight vector, we follow the procedure of Yen
677 (2015) and update the weights cyclical, that is we first fix w_i , $i = 2, \dots, k$ and find a
678 new solution for w_1 that is closer to its optimal solution w^* . In a next step, we fix
679 w_i , $i = 1, 3, \dots, k$ and find a value for w_2 that is again closer to the optimal one w^* .
680 Given a starting criteria \mathbf{w}^0 for the weight vector, the Lagrange parameter, γ , for
681 the budget constraint and a trade-off parameter, θ , for μ and σ^2 , Algorithm 2 shows
682 the pseudo code for the CyCoDe.

Algorithm 2 Cyclic Coordinate Descend

```

1: Initialize  $\mathbf{w}^0$  and  $j = 0$ 
2: while convergence criteria is not met do
3:   for  $i = 1$  to  $k$  do
4:      $w_i = ST(\gamma - z_i, \lambda) \times (2 \times \sigma_i^2)^{-1}$ 
5:     where  $ST$  is the soft-thresholding function and  $z_i = 2 \sum_{j \neq i}^k w_j \sigma_{ij} - \theta \mu_i$ 
6:   end for
7:    $j = j + 1$ 
8: end while

```

683 To evaluate the performance of the two algorithms, we first draw a random sample

684 of size n for k assets from a multivariate normal $\mathbf{X} \sim MVN(0, \Sigma)$, where Σ :

$$\Sigma_{ij} = \begin{cases} 1, & i = j, \\ \rho, & i \neq j, \end{cases} \quad (\text{C.1})$$

685 and for which we choose $\rho = 0.2$ and 0.8 , respectively. Then, we solve the minimum
 686 variance problem given in (2) and subject to the ℓ_1 - Norm on the weight vector,
 687 using as an input for Σ the shrunken covariance matrix, introduced by Ledoit and
 688 Wolff (2004b).

689 We initialize both algorithms with a soft starting point \mathbf{w}^0 , that is (1) $\mathbf{w}_i^0 = \frac{1}{k} \forall i =$
 690 $1, \dots, k$, and (2) $\mathbf{w}_i^0 = \frac{a_i}{\sum_{i=1}^k a_i}$, with $a_i \sim U(0, 1) \forall i = 1, \dots, k$, and repeat the above
 691 procedure 100 times, using for both algorithms a tolerance stopping point of 10^{-7} .

692 All computations are performed in Matlab 2016a on a Lenovo T430, with Windows
 693 7, an Intel i7-3520M with 2.90 GHZ and 8 GB of RAM.

694 Table C.4 and C.5 display the minimum and the median of the objective function
 695 values, together with the median amount of shorting, the median time in seconds
 696 used for each algorithm to solve the 100 simulations and the median absolute weight
 697 difference¹⁵, considering as soft starting criteria an equally weighted and a random
 698 portfolio weight vector, respectively.¹⁶

699 The tables show that both algorithms reach the same global minimum and median
 700 objective function value and the same amount of shorting for the low correlation
 701 environment, regardless of the chosen lambda value and whether we consider the

¹⁵The difference in the weights is computed as: $\sum |\mathbf{w}^{ADMM} - \mathbf{w}^{CyCoDe}|$, where \mathbf{w}^{ADMM} and \mathbf{w}^{CyCoDe} are the optimal weights obtained with the ADMM and the CyCoDe algorithm, respectively.

¹⁶Due to space limitations, we have restricted ourselves to report the above mentioned measures. Further results, including the standard deviation of the objective function value and the median number of active positions are available upon request to the authors.

Table C.4: Simulation Results - Equal Weights

				$\lambda = 4.03 \times 10^{-6}$					$\lambda = 5.65 \times 10^{-4}$					$\lambda = 7.91 \times 10^{-2}$				
ρ	n	p	Algo	Min	Med	Short	Time	W.Diff.	Min	Med	Short	Time	W.Diff.	Min	Med	Short	Time	W.Diff.
0.2	500	100	CyCoDe	0.14	0.16	0.51	0.66	5×10^{-7}	0.14	0.16	0.49	0.62	5×10^{-7}	0.23	0.25	0.00	0.18	7×10^{-8}
			ADMM	0.14	0.16	0.51	0.01		0.14	0.16	0.49	0.01		0.23	0.25	0.00	0.01	
	500	250	CyCoDe	0.09	0.11	2.13	13.63	8×10^{-6}	0.09	0.11	2.02	12.87	6×10^{-6}	0.21	0.24	0.00	0.94	8×10^{-8}
			ADMM	0.09	0.11	2.13	0.09		0.09	0.11	2.02	0.09		0.21	0.24	0.00	0.03	
	1000	500	CyCoDe	0.09	0.10	3.46	117.69	3×10^{-5}	0.09	0.11	3.23	116.29	2×10^{-5}	0.22	0.24	0.00	5.58	1×10^{-7}
			ADMM	0.09	0.10	3.46	0.66		0.09	0.11	3.23	0.64		0.22	0.24	0.00	0.17	
0.8	500	100	CyCoDe	0.55	0.64	3.39	11.67	2×10^{-3}	0.55	0.65	3.30	11.23	2×10^{-3}	0.73	0.83	0.00	1.37	8×10^{-7}
			ADMM	0.55	0.64	3.39	0.06		0.55	0.65	3.30	0.05		0.73	0.83	0.00	0.03	
	500	250	CyCoDe	0.34	0.42	10.98	35.33	8×10^{-1}	0.35	0.43	10.46	34.75	8×10^{-1}	0.67	0.82	0.00	6.03	1×10^{-6}
			ADMM	0.34	0.42	10.94	0.58		0.35	0.43	10.47	0.56		0.67	0.82	0.00	0.11	
	1000	500	CyCoDe	0.36	0.42	16.49	109.37	2.1	0.38	0.44	15.44	107.64	1.8	0.75	0.83	0.00	37.20	2×10^{-6}
			ADMM	0.36	0.42	16.34	3.96		0.38	0.43	15.33	3.76		0.75	0.83	0.00	0.61	

The table reports, for the Cyclic Coordinate Descend (CyCoDe) and the Alternating Direction Method of Multipliers (ADMM), the simulation results to the penalized minimum variance problem given in (2), considering six data sets drawn from a multivariate normal distribution, with $\rho = 0.2$ and $\rho = 0.8$, respectively, and using the equally weighted portfolio as a soft starting point. Stated are across all 100 simulations: the minimum (Min) and the median (Med) value of the objective function, the median value of the total amount of shorting (Short) the median time in seconds needed to compute the solution (Time) and the average weight difference (W.Diff.).

702 equally weighted or the random weight vector as the soft starting point. This also
703 applies to the low dimensional data set, when the correlation is set to $\rho = 0.8$. When
704 $p = 500$ for $\rho = 0.8$, the ADMM reports a lower amount of shorting for the first two
705 lambda values. This holds regardless of how we choose the soft starting point. This
706 difference might also explain the discrepancy in the weight vectors, which is reported
707 to be the highest for these two data sets. Still, the difference in the resulting weight
708 vectors is modest and amounts to an average of 10^{-6} for both low correlation envi-
709 ronments, and to 10^{-4} , for the first two high correlation environments and regardless
710 on how we choose the soft starting point.

711 Most notably, the ADMM outperforms the CyCoDe, with regard to the median time
712 in seconds used to compute the solution for all six data sets. This difference is not
713 negligible: the ADMM uses on average 0.265 seconds in the low correlation environ-
714 ment across all lambdas and all starting criteria, while the CyCoDe is slower by a
715 factor of more than 100, using on average 28.88 seconds. This also applies to the high

716 correlation environment, with the ADMM finding the solution, by taking on average
 717 2.65 seconds and the CyCoDe using 38.98 second. Finally, and for both algorithms,
 718 selecting the random weight vector as a starting point results in longer computing
 719 times, as opposed to using the equally weighted solution.

Figure C.8 plots the computing times needed for the CyCoDe and the ADMM for

Table C.5: Simulation Results - Random Weights


			$\lambda = 4.03 \times 10^{-6}$					$\lambda = 5.65 \times 10^{-4}$					$\lambda = 7.91 \times 10^{-2}$					
ρ	n	p	Algo	Min	Med	Short	Time	W.Diff	Min	Med	Short	Time	W.Diff	Min	Med	Short	Time	W.Diff
0.2	500	100	CyCoDe	0.13	0.16	0.49	0.46	5×10^{-7}	0.13	0.16	0.47	0.44	4×10^{-6}	0.22	0.25	0.00	0.13	7×10^{-8}
			ADMM	0.13	0.16	0.49	0.01		0.13	0.16	0.47	0.01		0.23	0.25	0.00	0.01	
	500	250	CyCoDe	0.08	0.10	2.12	10.26	8×10^{-6}	0.08	0.10	2.02	10.02	6×10^{-6}	0.19	0.23	0.00	0.74	8×10^{-8}
			ADMM	0.08	0.10	2.11	0.07		0.08	0.10	2.02	0.07		0.19	0.23	0.00	0.02	
	1000	500	CyCoDe	0.08	0.10	3.50	111.66	3×10^{-5}	0.09	0.10	3.28	112.50	2×10^{-5}	0.22	0.24	0.00	5.31	1×10^{-7}
			ADMM	0.08	0.10	3.50	0.52		0.09	0.10	3.28	0.51		0.22	0.24	0.00	0.15	
0.8	500	100	CyCoDe	0.55	0.64	3.30	8.02	2×10^{-3}	0.56	0.64	3.21	7.86	2×10^{-3}	0.72	0.82	0.00	0.89	8×10^{-7}
			ADMM	0.55	0.63	3.30	0.03		0.55	0.64	3.21	0.03		0.72	0.82	0.00	0.02	
	500	250	CyCoDe	0.33	0.41	10.77	31.54	8×10^{-1}	0.35	0.42	10.34	32.05	8×10^{-1}	0.68	0.81	0.00	5.35	1×10^{-6}
			ADMM	0.33	0.41	10.75	0.55		0.35	0.42	10.33	0.53		0.68	0.81	0.00	0.10	
	1000	500	CyCoDe	0.36	0.40	16.42	111.10	2.193	0.37	0.42	15.38	111.70	1.99	0.76	0.82	0.00	38.7	1.81
			ADMM	0.36	0.40	16.37	3.89		0.37	0.42	15.36	3.69		0.76	0.82	0.00	0.60	

The table reports, for the Cyclic Coordinate Descent (CyCoDe) and the Alternating Direction Method of Multipliers (ADMM), the simulation results to the penalized minimum variance problem given in (2), considering six data sets drawn from a multivariate normal distribution, with $\rho = 0.2$ and $\rho = 0.8$, respectively, and using the equally weighted portfolio as a soft starting point. Stated are across all 100 simulations: the minimum (Min) and median (Med) value of the objective function, the median value of the total amount of shorting (Short) the median time in seconds needed to compute the solution (Time) and the average weight difference (W.Diff.).

720

721 both the EW and Random weight vector initialization, considering the two corre-
 722 lation regimes and varying the number of parameters that have to be estimated.
 723 Clearly the ADMM consistently shows a superior performance, by only using a frac-
 724 tion of the time of the CyCoDe. Furthermore, we can observe that both algorithms
 725 are also invariant to the selection of the soft starting point. Only the CyCoDe shows
 726 a slight difference for parameter values above $k = 450$, signaling that for the CyCoDe
 727 an EW portfolio results in finding the optimal solution faster.

Figure C.8: Computation Times for CyCoDe and ADMM



The figure shows the average computation times needed for the CyCoDe and ADMM algorithm, depending on the correlation regime, the number of parameters and the soft start criterion. All values are based on 100 simulations, considering a constant correlation set-up.

728 Appendix D. Portfolio Selection Models

Equally Weighted Portfolio. The equally weighted portfolio is considered as one of the toughest benchmarks to beat (see, i.e. DeMiguel et al. (2009b)), and naively distributes the wealth equally among all constituents, such that with k assets:

$$w_i = \frac{1}{k} \quad \forall i = \{1, \dots, k\}, \quad (\text{D.1})$$

729 where w_i is the weight of asset i . The EW ignores both the variances, the covari-
730 ances and the return of the assets, and is the optimal portfolio on the mean-variance
731 efficient frontier, when we assume that all three are the same.

732

Norm-Constrained Minimum Variance Portfolio. Reconsider the formulation of the mean-variance problem in (1). By disregarding the mean in the optimization,

we obtain the Global Minimum Variance Portfolio (GMV), given by:

$$\min_{\mathbf{w} \in \mathbb{R}^k} \sigma_p^2 = \mathbf{w}' \Sigma \mathbf{w} \quad s.t. \sum_{i=1}^k w_i = 1, \quad \forall i = \{1, \dots, k\}, \quad (\text{D.2})$$

However, this formulation is prone to estimation errors, and unstable portfolio weights. To circumvent these problems, we extend the framework in (D.2) by adding a penalty function $\rho_\lambda(\mathbf{w})$ on the weight vector. For LASSO, we add a ℓ_1 - Norm to the formulation in (D.2), such that:

$$\rho_\lambda(\mathbf{w}) = \lambda \times \sum_{i=1}^k |w_i| \quad (\text{D.3})$$

where λ is a regularization parameter that controls the intensity of the penalty. Besides LASSO, we also consider the RIDGE penalty, which adds an ℓ_2 -Norm on the weight vector to the formulation in (D.2), and that takes the form of:

$$\rho_\lambda(\mathbf{w}) = \lambda \times \sum_{i=1}^k w_i^2 \quad (\text{D.4})$$

733 As opposed to the LASSO, the RIDGE is not singular at the origin and thus does
 734 not promote sparse solutions. Still, imposing the ℓ_2 - Norm on the portfolio problem
 735 is equal to adding an identity matrix, weighted by the regularization parameter λ
 736 to the inverse of the variance-covariance matrix, i.e. $(\Sigma^{-1} + \lambda \mathbf{I})$, where \mathbf{I} is the
 737 $k \times k$ identity matrix. This leads to more numerical stability and makes the RIDGE
 738 penalty especially appealing in environments that suffer from multicollinearity (Zou
 739 and Hastie, 2005).

740

Equal Risk Contribution Portfolio. Finally, we consider the Equal Risk Con-

tribution (ERC) portfolio, which aims to equalize the marginal risk contributions of the assets to the overall portfolio risk. That is, given that portfolio variance can be decomposed as:

$$\sigma_p^2 = \sum_{i=1}^k \sum_{j=1}^k w_i w_j \sigma_{ij} = \sum_{i=1}^k w_i \sum_{j=1}^k w_j \sigma_{ij} \quad (\text{D.5})$$

the marginal contribution to the portfolio risk for asset i is given as:

$$c_i^{var} = w_i \sum_{j=1}^k w_j \sigma_{ij} = w_i (\boldsymbol{\Sigma} \mathbf{w})_i \quad \text{with} \quad \sum_{i=1}^k c_i^{var} = \sigma_p^2 \quad (\text{D.6})$$

where $(\boldsymbol{\Sigma} \mathbf{w})_i$ denotes the i^{th} row of the product of $\boldsymbol{\Sigma}$ and \mathbf{w} (Roncalli, 2013). As the marginal risk is dependent on the portfolio weight magnitude, the ERC portfolio has no analytically solution and must be obtained numerically, by solving:

$$\min_{\mathbf{w} \in \mathbb{R}^N} \sum_{i=1}^k \left(\frac{w_i (\boldsymbol{\Sigma} \mathbf{w})_i}{\sigma_p^2} - \frac{1}{k} \right)^2 \quad \text{s.t.} \quad \sum_{i=1}^k w_i = 1, \quad 0 \leq w_i \leq 1 \quad \forall i \in \{1, 2, \dots, k\} \quad (\text{D.7})$$

741 The ERC favors assets with lower volatility, lower correlation with other assets, or
 742 both, and is less sensitive to small changes in the covariance matrix as compared to
 743 the GMV portfolio (Kremer et al., 2018). Furthermore, (Maillard et al., 2010) show
 744 that the volatility of the ERC is between that of the EW and the GMV, and that it
 745 coincides with the latter, when both, correlations and SRs, are assumed to be equal
 746 (Maillard et al., 2010).

747 Acknowledgement

748 Małgorzata Bogdan acknowledges the grant of the Polish National Center of
 749 Science Nr 2016/23/B/ST1/00454, and together with Sandra Paterlini further ac-

750 knowledge ICT COST Action IC1408 from CRoNoS. Sangkyun Lee acknowledges
751 the support of the National Research Foundation of Korea (NRF) grant funded by
752 the Korea government (MSIP; Ministry of Science, ICT & Future Planning) (No.
753 2017R1C1B5018367).

754 References

755 Bellec, P., Lecué, G., Tysbakov, A., 2016a. Bounds on the prediction error of pe-
756 nalized least squares estimators with convex penalty. arXiv:1609.06675, to appear
757 in Modern Problems of Stochastic Analysis and Statistics, Festschrift in honor of
758 Valentin Konakov.

759 Bellec, P., Lecué, G., Tysbakov, A., 2016b. Slope meets lasso: improved oracle
760 bounds and optimality. arXiv:1605.08651, 1–29.

761 Bogdan, M., van den Berg, E., Sabatti, C., Su, W., Candes, E., 2015. Slope - adaptive
762 variable selection via convex optimization. Annals of Applied Statistics 9 (3), 1103–
763 1140.

764 Bogdan, M., van den Berg, E., Su, W., E.J., C., 2013. Statistical estimation and
765 testing via the ordered ℓ_1 norm. arXiv:1310.1969, 1–46.

766 Bondell, H., Reich, B., March 2008. Simultaneous regression shrinkage, variable se-
767 lection, and supervised clustering of predictors with oscar. Biometrics 64 (1), 115–
768 123.

769 Boyd, S., Parikh, N., Chu, E., Peleato, B., Eckstein, J., 2011. Distributed optimiza-
770 tion and statistical learning via the alternating direction method of multipliers.
771 Foundations and Trends in Machine Learning 3 (1), 1–122.

- 772 Boyle, P., Garlappi, L., Uppal, R., Wang, T., 2012. Keynes meets Markowitz: The
773 trade-off between familiarity and diversification. *Management Science* 58 (2), 253–
774 272.
- 775 Brodie, J., Daubechies, I., DeMol, C., Giannone, D., Loris, D., 2009. Sparse and sta-
776 ble markowitz portfolios. *Proceedings of the National Academy of Science* 106 (30),
777 12267–12272.
- 778 Carrasco, M., Noumon, N., 2012. Optimal portfolio selection using regularization.
779 Working Paper University of Montreal, 1–52.
- 780 Cazalet, Z., Grison, P., Roncalli, T., 2014. The smart beta indexing puzzle. *The*
781 *Journal of Index Investing* 5 (1), 97–119.
- 782 Chopra, V., Ziemba, W., 1993. The effect of errors in means, variances, and co-
783 variances on optimal portfolio choice. *Journal of Portfolio Management* 19 (2),
784 6–11.
- 785 Choueifaty, Y., Coignard, Y., 2008. Toward maximum diversification. *Journal of*
786 *Portfolio Management* 34 (4), 40–51.
- 787 DeMiguel, V., Garlappi, L., Nogales, F., Uppal, R., 2009a. A generalized approach
788 to portfolio optimization: Improving performance by constraining portfolio norm.
789 *Management Science* 55 (5), 798–812.
- 790 DeMiguel, V., Garlappi, L., Nogales, F., Uppal, R., 2009b. Optimal versus naive
791 diversification: How inefficient is the $1/n$ portfolio strategy? *Review of Financial*
792 *Studies* 22 (5), 1915–1953.
- 793 DeMiguel, V., Nogales, F., 2009. Portfolio selection with robust estimation. *Opera-*
794 *tions Research* 55 (5), 798–812.

- 795 Fan, J., Fan, Y., Lv, J., 2008. High dimension covariance matrix estimation using
796 a factor model. *Journal of Econometrics* 147 (1), 186–197.
- 797 Fan, J., Zhang, J., You, K., 2012. Vast portfolio selection with gross-exposure con-
798 straint. *Journal of the American Statistical Association* 107 (498), 592–606.
- 799 Fastrich, B., Paterlini, S., Winker, P., 2014. Cardinality versus q-norm constraints
800 for index tracking. *Quantitative Finance* 14 (11), 2019–2032.
- 801 Fastrich, B., Paterlini, S., Winker, P., 2015. Constructing optimal sparse portfolios
802 using regularization methods. *Computational Management Science* 12 (3), 417–
803 434.
- 804 Figueiredo, M., Nowak, R., 2014. Sparse estimation with strongly correlated variables
805 using ordered weighted ℓ_1 regularization. Working Paper, 1–15.
806 URL [arXiv:1409.4005](https://arxiv.org/abs/1409.4005)
- 807 Gabaix, X., 2014. A sparsity-based model of bounded rationality. *The Quarterly*
808 *Journal of Economics* 129, 1661–1710.
- 809 Gasso, G., Rakotomamonjy, A., Canu, S., 2010. Recovering sparse signals with a
810 certain family of non-convex penalties and dc programming. *IEEE Transactions*
811 *on Signal Processing* 57 (12), 4686–4698.
- 812 Giuzio, M., Paterlini, S., 2016. Un-diversifying during crises: Is it a good idea? FRB
813 of Cleveland Working Paper (16-28), 1–38.
- 814 Hestenes, M. R., 1969. Multiplier and gradient methods. *Journal of Optimization*
815 *Theory and Applications* 4 (5), 303–320.

- 816 Jagannathan, R., Ma, T., August 2003. Risk reduction in large portfolios: Why
817 imposing the wrong constraints helps. *The Journal of Finance* 58 (4), 1651–1683.
- 818 Jorion, P., 1986. Bayes-stein estimator for portfolio analysis. *The Journal of Financial*
819 *and Quantitative Analysis* 21 (3), 279–292.
- 820 Kolm, P., Tütüncü, R., Fabozzi, F. J., 2014. 60 years of portfolio optimization:
821 Practical challenges and current trends. *European Journal of Operational Research*
822 234 (2), 356–371.
- 823 Kremer, P., Talmaciu, A., Paterlini, S., 7 2018. Risk minimization in multi-factor
824 portfolios: What is the best strategy? *Annals of Operations Research* 266 (1-2),
825 255–291.
- 826 Ledoit, O., Wolf, M., 2008. Robust performance hypothesis testing with the sharpe
827 ratio. *Journal of Empirical Finance* 15 (5), 850–859.
- 828 Ledoit, O., Wolf, M., 2011. Robust performance hypothesis testing with the variance.
829 *Wilmott Technical Paper* 2011 (55), 86–89.
- 830 Ledoit, O., Wolff, M., 2003. Improved estimation of the covariance matrix of stock
831 returns with an application to portfolio selection. *Journal of Empirical Finance*
832 10 (5), 603–621.
- 833 Ledoit, O., Wolff, M., 2004a. Honey, i shrunk the covariance matrix. *Journal of*
834 *Portfolio Management* 30 (4), 110–119.
- 835 Ledoit, O., Wolff, M., 2004b. A well-conditioned estimator for large-dimensional
836 covariance matrices. *Journal of Multivariate Analysis* 88 (2), 365–411.

- 837 Li, J., 2015. Sparse and stable portfolio selection with parameter uncertainty. *Journal*
838 *of Business and Economic Statistics* 33 (3), 381–392.
- 839 Maillard, S., Roncalli, T., Teiletche, J., 2010. The properties of equally weighted risk
840 contribution portfolios. *The Journal of Portfolio Management* 36 (4), 60–70.
- 841 Markowitz, H., March 1952. Portfolio selection. *The Journal of Finance* 7 (1), 77–91.
- 842 Merton, R. C., 1980. On estimating the expected return on the market: An ex-
843 ploratory investigation. *Journal of Financial Economics* 8 (4), 323–361.
- 844 Michaud, R., 1989. The Markowitz optimization enigma: is 'optimized' optimal?
845 *Financial Analyst Journal* 45 (1), 31–42.
- 846 Parikh, N., Boyd, S., 2014. Proximal algorithms. *Foundations and Trends in Opti-*
847 *mization* 1 (3), 127–239.
- 848 Powell, M. J. D., 1969. A method for nonlinear constraints in minimization problems.
849 In: Fletcher, R. (Ed.), *Optimization*. Academic Press, New York, pp. 283–298.
- 850 Roncalli, T., 2013. *Introduction to Risk Parity and Budgeting*. Chapman &
851 Hall/CRC Financial Mathematics Series.
- 852 Shefrin, H., Statman, M., 2000. Behavioral portfolio theory. *Journal of Financial and*
853 *Quantitative Analysis* 35 (2), 127–151.
- 854 Stevens, J., 1992. *Applied multivariate Statistics for Social Sciences*, 2nd Edition.
855 New-Jersey:Lawrance Erlbaum Associates Publishers, pp. 260 - 269.
- 856 Su, W., Candès, E., 2016. Slope is adaptive to unknown sparsity and asymptotically
857 minimax. *Annals of Statistics* 44 (3), 1038–1068.

- 858 Tibshirani, R., 1996. Regression shrinkage and selection via the lasso. *Journal of the*
859 *Royal Statistical Society* 58 (1), 267–288, series B.
- 860 Xing, X., Hub, J., Yang, Y., 2014. Robust minimum variance portfolio with ℓ_∞
861 constraints. *Journal of Banking and Finance* 46, 107–117.
- 862 Yen, Y.-M., 2015. Sparse weighted norm minimum variance portfolio. *Review of*
863 *Finance* 20 (3), 1259–1287.
- 864 Zeng, X., Figueiredo, M., 2014. Decreasing weighted sorted l_1 regularization. *IEEE*
865 *Signal Processing Letters* 21 (10), 1240–1244.
- 866 Zou, H., Hastie, T., 2005. Regularization and variable selection via the elastic net.
867 *Journal of the Royal Statistical Society B* 67 (2), 301–320.