



# Spatiotemporal pattern of Covid-19 outbreak in Turkey

Neşe Aral · Hasan Bakır

Accepted: 18 April 2022 / Published online: 16 June 2022  
© The Author(s), under exclusive licence to Springer Nature B.V. 2022

**Abstract** The earliest case of Covid-19 was documented in Wuhan city of China and since then the virus has been spreading throughout the globe. The aim of this study is to evaluate the clusters of Covid-19 among the provinces in Turkey and to examine whether the clustering pattern has changed after the country's lockdown strategy. The spatial dependence of Covid-19 in 81 provinces of Turkey was examined by spatial analysis between February 8 and June 28, 2021. Global and Local Moran's I and  $G_i^*$  were employed to measure the global and local spatial autocorrelation degrees. The geographical distribution of Covid-19 in the provinces of Turkey showed a strong spatial autocorrelation while the spatial structure of the clusters varied by weeks. The findings of the study show that the complete lockdown carried out in Turkey has been quite effective in mitigating Covid-19. The importance of spatial relations in preventing the spread of the disease in Turkey has also been demonstrated in this context.

**Keywords** Covid-19 · Turkey · Spatial statistics · Spatial autocorrelation · Spatiotemporal pattern

## Introduction

The novel coronavirus disease first appeared in Wuhan, China. As a result of the examination of a group of patients, the virus was diagnosed on January 13, 2020. While the fight against the first pandemic of the century continues, public health and national economies continue to suffer serious damage (Arauzo-Carod, 2021). Different measures have been implemented by most countries to fight against the coronavirus. People have been locked down in cities, production has stopped, schools have been closed, and restrictions have been imposed on people's movements. While the pandemic has been threatening people's health and causing loss of lives, an economic downturn began to emerge on the other hand. In fact, as highlighted by the United Nations, the Covid-19 pandemic means much more than a medical crisis; it is an economic, a social and a human crisis (UN, 2020). During this pandemic, the main concern for each nation is to keep their populations healthy. However, the disease also had an impact on national economies. Hence, during this unusual and challenging time of Covid-19, every country has to deal with two major issues: Healthcare and economics (Sharma et al., 2021). Undetected spread of the virus shocked governments, scientists and international

---

N. Aral (✉)  
Department of Econometrics, Faculty of Economics  
and Administrative Sciences, Bursa Uludag University,  
Bursa, Turkey  
e-mail: nesearal@uludag.edu.tr

H. Bakır  
Department of International Trade, Vocational School  
of Social Sciences, Bursa Uludag University, Bursa,  
Turkey  
e-mail: hasanbakir@uludag.edu.tr

organizations. In addition to its impact on public health, it caused major economic distress. Therefore, these changes will cause a long-run international economic downturn. As we mention in this paper, health systems of most countries haven't been able to contain the evolution and the persistence of the virus (Kalla et al., 2021).

The whole world is fighting this pandemic, which shows the highest number of daily cases around the globe and losses of the last century, while the numbers affected by the pandemic differ by countries. For example, as a country with high population density, Turkey has experienced high percentages of incidence and fatality. Additionally, new cases and deaths related to Covid-19 increase Daily. The first news of death from Turkey came on March 17, 2020 (T.C.S.B, 2021). More than forty-nine thousand citizens of the Republic of Turkey died since the day Turkey met with the first Covid-19 case. In the face of the daily increase in registered cases in Turkey, the government and the public authorities took a wide range of measures that included the entire country at the end of April, and a complete lockdown was implemented in Turkey from April 29, 2021 until May 17, 2021. In this research, we aim to investigate the spatial distribution of the virus and the effect of complete lockdown measures on the spatial structure of the clusters.

## Review of the literature

After the break of coronavirus, studies have used statistical methods to determine the nexus between cases, the factors of transmission and the determination of clusters (Appiah-Otoo & Kursah, 2021). As the characteristics of Covid-19 are original and wild, it spread very quickly. Therefore, The World Health Organization has consistently used spatial data analysis to control infectious diseases (Esri, 2020; Nasiri et al., 2021). Within the scope of the research of measures that can be taken to prevent the pandemic, many researchers in different countries have revealed the effects of spatial factors on the fast spread of coronavirus (Adegboye et al., 2021; Casado-Aranda et al., 2021; Castro et al., 2021; Gupta et al., 2021; Li et al., 2021; Liu et al., 2021; Maiti et al., 2021; Mansour et al., 2021; Nasiri et al., 2021; Rubino et al., 2020; Sarkar et al., 2021; Shariati et al., 2020; Tang et al., 2020; Tao et al., 2020; Vaz, 2021; Xiong et al.,

2020). Ramírez-Aldana et al. (2020) determined that the number of Covid-19 cases in Iranian provinces is spatially related. Similarly Arauzo-Carod (2021) showed that in Catalonia (Spain), positive case numbers were associated with nearby settlements, through spatial autocorrelation. Eryando et al. (2020) stated that Covid-19 was not observed randomly in Indonesia, but that the neighboring provinces that were connected and five other cities with the highest number of cases were adjacent to each other. Ghosh and Cartone (2020) determined the spatial impact of the virus that emerged in different regions of Italy. Again, researchers similarly highlighted the importance of spatial relation in the spread and prevention of the epidemic in Oman (Al-Kindi et al., 2020), Kuwait (Alkhamis et al., 2020), South Korea (Kim & Castro, 2020), Iran (Pourghasemi et al., 2020), China (Kang et al., 2020), Lebanon (El Deeb, 2021), and Rio de Janeiro (Ferreira et al., 2020). In these studies, which take place in literature, the importance of spatial heterogeneity has been underlined and it has been determined that local characteristics affect the spread of coronavirus. The efficiency of health policies has been associated with coordination at both local and national levels. Geographical information about the pandemic helps to ensure coordination at national and local levels. It has been stated that maintaining health standards, preventing social tension and achieving fewer work interruptions can be accomplished this way (Ghosh & Cartone, 2020).

In the light of this information, the goal of this research is to survey the clusters of the pandemic among the provinces of Turkey and to examine whether the clustering structure has changed after the country's lockdown strategy. The course of the change in Covid-19 cases in Turkey during the period of study by provinces can also be examined in the timeline. Selected accordingly, the period of study offers the scope to examine the spatial distribution of Covid-19 in Turkey's provinces in a comparative way including the complete lockdown period and the previous period.

## Methodology

Data showing the number of coronavirus cases by province, corresponding to a weekly population of 100,000 began to be published as of February 15,

2021 in Turkey. The study uses the number of cases by province corresponding to a population of 100,000 for 20 weeks (8 February–28 June). The summary of the characteristics of the research variables are presented in Table 1. The number of coronavirus cases, which increased within weeks during the period of this study, reached its peak in the 11th week and then started to decline.

If the empirical study is developed on spatial data, or if the relations studied are “spatial” in nature, it becomes essential to use spatial analysis that can “appropriately” incorporate location and spatial arrangement into the study (Anselin, 1999). In this context, spatial autocorrelation was employed to examine whether Covid-19 has noteworthy global or local spatial autocorrelation features (Xiong et al., 2020). Global or local measures of spatial autocorrelation have been used to investigate the spatial patterns and define hot spots and cold spots of the epidemic in many researches (Alves et al., 2021; Castro et al., 2021; Das et al., 2021; Ghosh & Cartone, 2020; Islam et al., 2021; Kuznetsov & Sadovskaya, 2021; Maithani et al., 2020; Maroko et al., 2020). Spatial statistical

methods have also been employed in previous years to comprehend spatial patterns of epidemic risks (Bailey, 2001; Bailey et al., 2011; Cao et al., 2010; Cromley, 2003; Ge et al., 2016; Hu et al., 2013; Tsai et al., 2009; Wang et al., 2008; Wen et al., 2006).

Spatial autocorrelation, a concept that is of use in defining the field of spatial analysis, is at the center of studies using spatial statistics (Getis, 2008; Ord & Getis, 2001). Spatial autocorrelation measurements and tests are divided into local and global scales depending on the scale and scope of the analysis (Anselin, 1995).

The Moran’s I statistic was employed to determine the global spatial autocorrelation characteristics of Covid-19 (Anselin, 1988; Moran, 1948, 1950):

$$I = \frac{n}{\sum_{i=1}^n \sum_{j \neq i}^n w_{ij}} \frac{\sum_{i=1}^n \sum_{j=1}^n w_{ij} (x_i - \bar{x})(x_j - \bar{x})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

Similar to the correlation coefficient, Moran’s I score varies in the range of [−1 + 1]. (Anselin, 1996; Griffith, 2003).

**Table 1** Descriptive statistics of data

Variables	DDate	Moran’s I	Mean	Min	Max	SD
Week 1	8th to 14th February	0.488044	59.4881	7.82	228.02	45.44546
Week 2	15th to 21st February	0.414519	84.4057	3.21	228.40	50.11229
Week 3	20th to 26th February	0.434957	69.7683	2.29	301.76	61.72887
Week 4	27th February to 5th March	0.427239	85.1765	5.35	348.36	74.27887
Week 5	6th to 12th March	0.405174	99.7656	2.11	458.53	83.95166
Week 6	13th to 19th March	0.419647	121.4172	8.18	508.97	90.96018
Week 7	20th to 27th March	0.496801	171.0694	13.57	586.94	113.65164
Week 8	27th March to 2nd April	0.555422	243.2402	23.43	678.72	146.62483
Week 9	3rd to 9th April	0.623690	335.3253	32.17	882.13	184.64584
Week 10	10th to 16th April	0.638943	395.8872	47.60	962.98	196.72092
Week 11	17th to 23rd April	0.667480	397.3954	55.04	854.75	174.88728
Week 12	24th to 30th April	0.674243	260.6160	46.12	532.02	108.19413
Week 13	1st to 7th May	0.586110	188.7370	36.08	359.99	74.79200
Week 14	8th to 14th May	0.485420	106.2531	26.96	192.28	40.53047
Week 15	15th to 21st May	0.358145	82.3342	15.68	163.72	32.80929
Week 16	22nd to 28th May	0.296495	67.1184	12.47	162.31	29.07410
Week 17	29th May to 4th June	0.257387	57.8652	7.95	110.55	24.90257
Week 18	5th to 11th June	0.271462	52.5607	7.66	129.51	25.45958
Week 19	12th to 18th June	0.312857	49.9973	5.97	116.18	25.12250
Week 20	19th to 25th June	0.326617	47.3296	4.92	136.52	25.89888
Entire period	8th February to 25th June	0.648244	147.6085	23.51	292.38	63.64626

Source: Author

Local spatial autocorrelation was used to determine local spatial autocorrelation features of the coronavirus.

The Local Moran's I statistic is calculated as follows (Anselin, 1995):

$$I_i = (x_i - \bar{x}) \cdot \sum_{j \in J_i}^n w_{ij} (x_j - \bar{x})$$

Similar to Moran's I, the G test is another sign of local spatial autocorrelation. In this test proposed by Getis and Ord (1992) and then further detailed in Ord and Getis (1995), values near  $-1$  indicate clustering of similar values. Values near  $1$  indicate divergence (i.e., random models), which means that this test is inversely linked with Moran's I test. G is calculated as follows (Getis & Aldstadt, 2004; Getis & Ord, 2010):

$$G = \frac{\sum_{i=1}^n \sum_{j=1}^n w_{ij} x_i x_j}{\sum_{i=1}^n \sum_{j=1}^n x_i x_j}; \quad \forall j \neq i$$

With the G test, four values are obtained. First, the p-value is examined. If the obtained p-value is found to be statistically small and significant, it indicates that the observation values are clustered spatially, otherwise, the observation values are randomly distributed. If the p-value suggests a cluster, look at the z-score. If the z-score is positive, it indicates that high values of the observation values tend to cluster in the analyzed area. Otherwise, if the z-score is found to be negative, it indicates that low observation values tend to cluster (Getis & Ord, 2010).

The Getis-Ord  $G_i^*$  is another primarily used test to measure local spatial autocorrelation. The  $G_i^*$  statistic has been calculated to define hot spots and cold spots of coronavirus.  $G_i^*$  is calculated as follows:

$$G_i^* = \frac{\sum_{j=1}^n w_{ij} x_j - \bar{x} \sum_{j=1}^n w_{ij}}{S \sqrt{\frac{n \sum_{j=1}^n w_{ij}^2 - (\sum_{j=1}^n w_{ij})^2}{n-1}}}$$

Here;

$$S = \sqrt{\frac{\sum_{j=1}^n x_j^2}{n} - (\bar{x})^2}$$

$G_i^*$ , similar to the G test, yields two measures, the z-score and the p-value. The resulting z-score reports where locations with high or low observations show a tendency to cluster in space. Unlike Local Moran's and Local Geary statistics, the Getis-Ord method does not consider spatial outliers (Anselin, 1995, 2019; Getis & Ord, 1992, 1996; Ord & Getis, 1995).

## Results and discussion

In the study, Moran's I coefficients were calculated first in order to analyze the relationship between Covid-19 cases in any province and Covid-19 cases in neighboring provinces. Spatial autocorrelation indicates whether Covid-19 numbers are associated with close locations. Accordingly, it is observed that the spatial autocorrelation of Covid-19 cases is positive across the country. Moran's I values for Covid-19 were recorded in the range of 0.257387–0.674243, confirming the presence of spatial structure. While Moran's I statistics tended to increase in the period under consideration, it reached its highest value in the 12th week. As of the 13th week (during the complete lockdown period), it started to trend downwards, but the spatial correlation was still in clustering characteristic. At the 17th week, Moran's I values started to rise again (Table 1).

Figure 1 shows the results of cluster analysis of provinces in Turkey, which is created to examine whether the observation values show regionally significant spatial clustering or scattering. According to LISA, LL clusters are observed in eastern and south-eastern provinces in all periods. In these regions, the patterns of LL clusters tend to have a continuous distribution. While the main clusters of HH were frequently seen in the northern regions in the first 4 weeks, they started to appear in the west from the 5th week and intensified over time. While the clustering trend was similar in the 7th, 8th and 9th weeks, the northern clustering disappeared at the 10th week. As of the complete lockdown (13th week), the pattern of the distribution started to differ. The number of significant clusters observed in the 16th week

decreased considerably and started to rise again from the 17th week. While the clusters of cases seen by weeks tend to rise and rapidly spread over other regions, they tend to decrease during the complete lockdown period. Within this framework, it is possible to see that the effect of the measures taken during this period is positive.

The  $G_i^*$  statistic has been calculated to identify the clusters of Covid-19 (hot and cold spots). The  $G_i^*$  utilizes the data of each province infected by Covid-19 and analyzes the local situation, comparing it with the global situation in the neighboring provinces (Bhunia et al., 2021). Figure 2 shows the prevalence of Covid-19 coded by  $G_i^*$  statistics based on selected weekly data from March 8 to June 28, 2021. Accordingly, it is possible to detect the hot and cold spot regions of Covid-19. In this context, clusters of cold spots and hot spots were detected over the weeks in Turkey with a confidence level of 95%.

According to the results of the hot spot analysis, 8 cities turn up hot spots in the first week. While the distribution of hot spots in this region was similar for 4 weeks, Çanakkale province, located in the Marmara region, was included in the hot spots in the 5th week. It spread to Tekirdağ and Kırklareli in the 6th week, and was observed in Edirne, Yalova, Istanbul, Bursa and Kocaeli in the 7th week. Hot spots rapidly covered most of the cities, and these provinces have become important hot spots. From the 11th week, it has spread to neighboring provinces. In the 16th week, the number of hot spots decreased, and it started to rise again from the 17th week.


Again, in this context, the positive effect of complete lockdown is seen. Researchers in different countries have revealed the effects of lockdowns in reducing the spread of Covid-19 (Acter et al., 2020; Alcântara et al., 2020; Bourdin et al., 2021; Cheng et al., 2020; Kraemer et al., 2020; Saha et al., 2021). Therefore, the importance of spatial measures in preventing the spread of the pandemic emerges. Thanks to these measures, both the cost of lockdown will decrease, and the spread of Covid-19 can also be prevented more quickly. Thus, the public reaction against lockdowns will also ease, relieving public administrators as well.

## Conclusion and policy recommendations


In Turkey, the first case of coronavirus was diagnosed on March 11, 2020, and during the past year, the number of cases has exceeded five million, while the number of deaths has exceeded forty-nine thousand. On the other hand, different numbers of cases at the provincial level brought up the examination of the geographical distribution of Covid-19. In this context, the necessity of using spatial statistics has emerged. Thus, the efficiency of local decisions will increase and it will be possible to fight against the pandemic more strongly. In this direction, with the help of spatial statistics, this study aims to examine the spatial distribution of the virus and to investigate the effect of complete lockdown measures on the spatial model of the clusters. As a result of the analyses made in this framework, a positive spatial autocorrelation has been determined in Turkey. The relationship between provinces was measured and the spatial clustering of Covid-19 was determined. These findings revealed the significant effects of the infection spreading among provinces. This reveals the importance of geographical location in the fight against the pandemic. Identifying hot spot Covid-19 areas may be beneficial in supervising regional Covid-19 prevention programs (Bhunia et al., 2021). In the light of the results obtained, it is important for policy makers to take measures to prevent the spread of the pandemic for provinces with hot spots because these cities are the riskiest in terms of the virus spreading, while these provinces also have more spatial interactions with their neighbors. Therefore, measures should be highlighted and control should be strengthened in cities with hot spots and their neighbors (Zeren & Yilanci, 2020).

However, with the complete lockdown period, it was determined that the spatial clusters of Covid-19 decreased and the spatial relations between the provinces weakened. The spatial relations observed at the local level will enable decision makers to produce more successful solutions in reply to the pandemic. The spread of the pandemic can be prevented by taking measures for the provinces where the said relations are observed. Therefore, measures taken specifically in provinces with high risk will both prevent the spread of the virus and will have a lower economic cost compared to a nationwide lockdown.






**Fig. 1** Regional level spatial clustering of Covid-19 for 20 weeks and entire period in Turkey. Source: Author



**Fig. 1** (continued)

This research, which tries to reveal the spatial relationships of Covid-19, has some limitations. One of the limitations in question is that the data set does not include the number of cases and deaths. For this reason, the spatiotemporal spread pattern of Covid-19

could not be examined by using the number of deaths. Also, since the data began to be published as of February 2021, the spatiotemporal characteristics of the epidemic could not be examined for the whole process. In addition, since the data obtained are related



**Fig. 2** Clustering of Covid-19 for 20 weeks and entire period in Turkey. Source: Author






Fig. 2 (continued)

to the provinces, analyzes at lower levels could not be performed. In future studies, when the data is available, more detailed results can be obtained by using the number of deaths, analyzing at the district level or by separating rural and urban areas. It is thought that these analyzes will be beneficial in preventing the spread of the epidemic.

## References

- Acter, T., Uddin, N., Das, J., Akhter, A., Choudhury, T. R., & Kim, S. (2020). Evolution of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) as coronavirus disease 2019 (COVID-19) pandemic: A global health emergency. *Science of the Total Environment*, *730*, 138996. <https://doi.org/10.1016/j.scitotenv.2020.138996>
- Adegboye, O. A., Adekunle, A. I., Pak, A., Gayawan, E., Leung, D. H., Rojas, D. P., Elfaki, F., McBryde, E. S., & Eisen, D. P. (2021). Change in outbreak epicentre and its impact on the importation risks of COVID-19 progression: A modelling study. *Travel Medicine and Infectious Disease*, *40*(January), 101988. <https://doi.org/10.1016/j.tmaid.2021.101988>
- Al-Kindi, K. M., Alkharusi, A., Alshukaili, D., Al Nasiri, N., Al-Awadhi, T., Charabi, Y., & El Kenawy, A. M. (2020). Spatiotemporal assessment of Covid-19 spread over Oman using GIS techniques. *Earth Systems and Environment*, *4*(4), 797–811. <https://doi.org/10.1007/s41748-020-00194-2>
- Alcântara, E., Mantovani, J., Rotta, L., Park, E., Rodrigues, T., Carvalho, F. C., & Filho, C. R. S. (2020). Investigating spatiotemporal patterns of the covid-19 in São Paulo state, Brazil. *Geospatial Health*, *15*(2), 201–209. <https://doi.org/10.4081/gh.2020.925>
- Alkhamis, M. A., Al Youha, S., Khajah, M. M., Ben Haider, N., Alhardan, S., Nabeel, A., Al Mazeedi, S., & Al-Sabah, S. K. (2020). Spatiotemporal dynamics of the COVID-19 pandemic in the State of Kuwait. *International Journal of Infectious Diseases*, *98*, 153–160. <https://doi.org/10.1016/j.ijid.2020.06.078>
- Alves, J. D., Abade, A. S., Peres, W. P., Borges, J. E., Santos, S. M., & Scholze, A. R. (2021). Impact of COVID-19 on the indigenous population of Brazil: A geo-epidemiological study. *Epidemiology and Infection*, *149*(e185), 1–11. <https://doi.org/10.1017/S0950268821001849>
- Anselin, L. (1988). *Spatial econometrics: Methods and models*. Kluwer Academic Publishers.
- Anselin, L. (1995). Local indicators of spatial association—LISA. In *Geographical analysis* (vol. 27, Issue 2). <https://doi.org/10.1111/j.1538-4632.1995.tb00338.x>
- Anselin, L. (1996). The Moran scatter plot as an ESDA tool to assess local instability in spatial association. In H. J. Fischer & M. M. Scholten (Eds.), *Spatial analytical perspectives on GIS: GISDATA* (4th ed., pp. 111–125). New York: CRC Press.
- Anselin, L. (1999). The future of spatial analysis in the social sciences. *Annals of GIS*, *5*(2), 67–76. <https://doi.org/10.1080/10824009909480516>
- Anselin, L. (2019). A local indicator of multivariate spatial association: Extending Geary's *c*. *Geographical Analysis*, *51*(2), 133–150. <https://doi.org/10.1111/gean.12164>
- Appiah-Otoo, I., & Kursah, M. B. (2021). Modelling spatial variations of novel coronavirus disease (COVID-19): Evidence from a global perspective. *GeoJournal*. <https://doi.org/10.1007/s10708-021-10427-0>
- Arauzo-Carod, J. M. (2021). A first insight about spatial dimension of COVID-19: Analysis at municipality level. *Journal of Public Health (oxford, England)*, *43*(1), 98–106. <https://doi.org/10.1093/pubmed/fdaa140>
- Bailey, P. E., Keyes, E. B., Parker, C., Abdullah, M., Kebede, H., & Freedman, L. (2011). Using a GIS to model interventions to strengthen the emergency referral system for maternal and newborn health in Ethiopia. *International Journal of Gynecology and Obstetrics*, *115*(3), 300–309. <https://doi.org/10.1016/j.ijgo.2011.09.004>
- Bailey, T. C. (2001). Spatial statistical methods in health. *Cadernos De Saúde Pública*, *17*(5), 1083–1098. <https://doi.org/10.1590/S0102-311X2001000500011>
- Bhunia, G. S., Roy, S., & Shit, P. K. (2021). Spatio-temporal analysis of COVID-19 in India: a geostatistical approach. *Spatial Information Research*, *29*(5), 661–672. <https://doi.org/10.1007/s41324-020-00376-0>
- Bourdin, S., Jeanne, L., Nadou, F., & Noiret, G. (2021). Does lockdown work? A spatial analysis of the spread and concentration of Covid-19 in Italy. *Regional Studies*, *55*(7), 1182–1193. <https://doi.org/10.1080/00343404.2021.1887471>
- Cao, Z. D., Zeng, D. J., Zheng, X. L., Wang, Q. Y., Wang, F. Y., Wang, J. F., & Wang, X. L. (2010). Spatio-temporal evolution of Beijing 2003 SARS epidemic. *Science China Earth Sciences*, *53*(7), 1017–1028. <https://doi.org/10.1007/s11430-010-0043-x>
- Casado-Aranda, L.-A., Sánchez-Fernández, J., & Bastidas-Manzano, A.-B. (2021). Tourism research after the COVID-19 outbreak: Insights for more sustainable, local and smart cities. *Sustainable Cities and Society*, *73*, 103126. <https://doi.org/10.1016/j.scs.2021.103126>
- Castro, R. R., Santos, R. S. C., Sousa, G. J. B., Pinheiro, Y. T., Martins, I. M., Pereira, M. L. D., & Silva, R. A. R. (2021). Spatial dynamics of the COVID-19 pandemic in Brazil. *Epidemiology and Infection*, *149*(e60), 1–9. <https://doi.org/10.1017/S0950268821000479>
- Cheng, C., Zhang, T., Song, C., Shen, S., Jiang, Y., & Zhang, X. (2020). The coupled impact of emergency responses and population flows on the COVID-19 pandemic in China. *GeoHealth*, *4*(12), e2020GH000332. <https://doi.org/10.1029/2020GH000332>
- Cromley, E. K. (2003). GIS and disease. *Annual Review of Public Health*, *24*, 7–24. <https://doi.org/10.1146/annurev.publhealth.24.012902.141019>
- Das, A., Ghosh, S., Das, K., Basu, T., Dutta, I., & Das, M. (2021). Living environment matters: Unravelling the spatial clustering of COVID-19 hotspots in Kolkata megacity, India. *Sustainable Cities and Society*, *65*, 102577. <https://doi.org/10.1016/j.scs.2020.102577>

- El Deeb, O. (2021). Spatial autocorrelation and the dynamics of the mean center of Covid-19 infections in Lebanon. *Frontiers in Applied Mathematics and Statistics*, 6, 1–15. <https://doi.org/10.3389/fams.2020.620064>
- Eryando, T., Sipahutar, T., & Rahardiantoro, S. (2020). The risk distribution of Covid-19 in Indonesia: A spatial analysis. *Asia-Pacific Journal of Public Health*, 32(8), 450–452. <https://doi.org/10.1177/1010539520962940>
- Esri. (2020). *Mapping epidemics: From SARS, Zika, and Ebola to the pandemic of COVID-19*. <https://www.esri.com/about/newsroom/blog/maps-that-mitigate-epidemics/>
- Ferreira, D., Ferreira, P., Oliveira, P., Ribeiro, J., Goncalves, E., & Papa, A. (2020). Temporal and spatial characteristics of the spread of COVID-19 in Rio de Janeiro state and city. *MedRxiv*. <https://doi.org/10.1101/2020.05.13.20101113>
- Ge, L., Zhao, Y., Zhou, K., Mu, X., Yu, H., Wang, Y., Wang, N., Fan, H., Guo, L., & Huo, X. (2016). Spatio-temporal pattern and influencing factors of Hemorrhagic Fever with Renal Syndrome (HFRS) in Hubei province (China) between 2005 and 2014. *PLoS ONE*, 11(12), e0167836. <https://doi.org/10.1371/journal.pone.0167836>
- Getis, A., & Ord, J. K. (1996). Local spatial statistics: an overview. In P. Longley & M. Batty (Eds.), *Spatial analysis modelling in a GIS environment* (pp. 269–285). Wiley.
- Getis, A. (2008). A history of the concept of spatial autocorrelation: A geographer's perspective. *Geographical Analysis*, 40(3), 297–309. <https://doi.org/10.1111/j.1538-4632.2008.00727.x>
- Getis, A., & Aldstadt, J. (2004). Constructing the spatial weights matrix using a local statistic. *Geographical Analysis*, 36(2), 90–104. <https://doi.org/10.1111/j.1538-4632.2004.tb01127.x>
- Getis, A., & Ord, J. K. (1992). The analysis of spatial association by use of distance statistics. *Geographical Analysis*, 24(3), 189–206. <https://doi.org/10.1111/j.1538-4632.1992.tb00261.x>
- Getis, A., & Ord, J. K. (2010). The analysis of spatial association by use of distance statistics. In L. Anselin & S. J. Rey (Eds.), *Springer*. Springer.
- Ghosh, P., & Cartone, A. (2020). A Spatio-temporal analysis of COVID-19 outbreak in Italy. *Regional Science Policy and Practice*, 12(6), 1047–1062. <https://doi.org/10.1111/rsp3.12376>
- Griffith, D. (2003). *Spatial autocorrelation and spatial filtering: Gaining understanding through theory and scientific visualization*. Springer Science & Business Media.
- Gupta, D., Biswas, D., & Kabiraj, P. (2021). COVID-19 outbreak and Urban dynamics: Regional variations in India. *GeoJournal*. <https://doi.org/10.1007/s10708-021-10394-6>
- Hu, B., Gong, J., Zhou, J., Sun, J., Yang, L., Xia, Y., & Ibrahim, A. N. (2013). Spatial-temporal characteristics of epidemic spread in-out flow—Using SARS epidemic in Beijing as a case study. *Science China Earth Sciences*, 56(8), 1380–1397. <https://doi.org/10.1007/s11430-012-4479-z>
- Islam, M. S., Tusher, T. R., Roy, S., & Rahman, M. (2021). Impacts of nationwide lockdown due to COVID-19 outbreak on air quality in Bangladesh: A spatiotemporal analysis. *Air Quality, Atmosphere and Health*, 14(3), 351–363. <https://doi.org/10.1007/s11869-020-00940-5>
- Kalla, M. I., Lahmar, B., Geullouh, S., & Kalla, M. (2021). Health geo-governance to assess the vulnerability of Batna, Algeria to COVID-19: The role of GIS in the fight against a pandemic. *GeoJournal*. <https://doi.org/10.1007/s10708-021-10449-8>
- Kang, D., Choi, H., Kim, J. H., & Choi, J. (2020). Spatial epidemic dynamics of the COVID-19 outbreak in China. *International Journal of Infectious Diseases*, 94, 96–102. <https://doi.org/10.1016/j.ijid.2020.03.076>
- Kim, S., & Castro, M. C. (2020). Spatiotemporal pattern of COVID-19 and government response in South Korea (as of May 31, 2020). *International Journal of Infectious Diseases*, 98, 328–333. <https://doi.org/10.1016/j.ijid.2020.07.004>
- Kraemer, M. U. G., Yang, C.-H., Gutierrez, B., Wu, C.-H., Klein, B., Pigott, D. M., du Plessis, L., Faria, N. R., Li, R., Hanage, W. P., Brownstein, J. S., Layan, M., Vespignani, A., Tian, H., Dye, C., Pybus, O. G., & Scarpino, S. V. (2020). The effect of human mobility and control measures on the COVID-19 epidemic in China. *Science*, 368(6490), 493–497. <https://doi.org/10.1126/science.abb4218>
- Kuznetsov, A., & Sadovskaya, V. (2021). Spatial variation and hotspot detection of COVID-19 cases in Kazakhstan, 2020. *Spatial and Spatio-Temporal Epidemiology*, 39, 100430. <https://doi.org/10.1016/j.sste.2021.100430>
- Li, B., Peng, Y., He, H., Wang, M., & Feng, T. (2021). Built environment and early infection of COVID-19 in urban districts: A case study of Huangzhou. *Sustainable Cities and Society*, 66, 102685. <https://doi.org/10.1016/j.scs.2020.102685>
- Liu, Y., Pei, T., Song, C., Chen, J., Chen, X., Huang, Q., Wang, X., Shu, H., Wang, X., Guo, S., & Zhou, C. (2021). How did human dwelling and working intensity change over different stages of COVID-19 in Beijing? *Sustainable Cities and Society*, 74, 103206. <https://doi.org/10.1016/j.scs.2021.103206>
- Maithani, S., Nautiyal, G., & Sharma, A. (2020). Investigating the effect of lockdown during COVID-19 on land surface temperature: study of Dehradun City, India. *Journal of the Indian Society of Remote Sensing*, 48(9), 1297–1311. <https://doi.org/10.1007/s12524-020-01157-w>
- Maiti, A., Zhang, Q., Sannigrahi, S., Pramanik, S., Chakraborti, S., Cerda, A., & Pilla, F. (2021). Exploring spatiotemporal effects of the driving factors on COVID-19 incidences in the contiguous United States. *Sustainable Cities and Society*, 68, 102784. <https://doi.org/10.1016/j.scs.2021.102784>
- Mansour, S., Al Kindi, A., Al-Said, A., Al-Said, A., & Atkinson, P. (2021). Sociodemographic determinants of COVID-19 incidence rates in Oman: Geospatial modelling using multiscale geographically weighted regression (MGWR). *Sustainable Cities and Society*, 65, 102627. <https://doi.org/10.1016/j.scs.2020.102627>
- Maroko, A. R., Nash, D., & Pavilonis, B. T. (2020). COVID-19 and inequity: A comparative spatial analysis of New York City and Chicago hot spots. *Journal of Urban Health*, 97(4), 461–470. <https://doi.org/10.1007/s11524-020-00468-0>
- Moran, P. A. P. (1948). The interpretation of statistical maps. *Journal of the Royal Statistical Society: Series B (methodological)*, 10(2), 243–251. <https://doi.org/10.1111/j.2517-6161.1948.tb00012.x>

- Moran, P. A. P. (1950). Notes on continuous stochastic phenomena. *Biometrika*, 37(1/2), 17–23. <https://doi.org/10.2307/2332142>
- Nasiri, R., Akbarpour, S., Zali, A., Khodakarami, N., Boochani, M., Noory, A., & Soori, H. (2021). Spatio-temporal analysis of COVID-19 incidence rate using GIS: a case study—Tehran metropolitan, Iran. *GeoJournal*. <https://doi.org/10.1007/s10708-021-10438-x>
- Ord, J. K., & Getis, A. (1995). Local Spatial Autocorrelation Statistics: Distributional Issues and an Application. *Geographical Analysis*, 27(4), 286–306. <https://doi.org/10.1111/j.1538-4632.1995.tb00912.x>
- Ord, J. K., & Getis, A. (2001). Testing for local spatial autocorrelation in the presence of global autocorrelation. *Journal of Regional Science*, 41(3), 411–432. <https://doi.org/10.1111/0022-4146.00224>
- Pourghasemi, H. R., Pouyan, S., Heidari, B., Farajzadeh, Z., Fallah Shamsi, S. R., Babaei, S., Khosravi, R., Etemadi, M., Ghanbarian, G., Farhadi, A., Safaeian, R., Heidari, Z., Tarazkar, M. H., Tiefenbacher, J. P., Azmi, A., & Sadeghian, F. (2020). Spatial modeling, risk mapping, change detection, and outbreak trend analysis of coronavirus (COVID-19) in Iran (days between February 19 and June 14, 2020). *International Journal of Infectious Diseases*, 98, 90–108. <https://doi.org/10.1016/j.ijid.2020.06.058>
- Ramírez-Aldana, R., Gomez-Verjan, J. C., Bello-Chavolla, O. Y., & Peterson, T. (2020). Spatial analysis of COVID-19 spread in Iran: Insights into geographical and structural transmission determinants at a province level. *PLoS Neglected Tropical Diseases*, 14(11), e0008875. <https://doi.org/10.1371/journal.pntd.0008875>
- Rubino, I., Coscia, C., & Curto, R. (2020). Identifying spatial relationships between built heritage resources and short-term rentals before the Covid-19 pandemic: Exploratory perspectives on sustainability issues. *Sustainability*, 12(11), 4533. <https://doi.org/10.3390/su12114533>
- Saha, J., Mondal, S., & Chouhan, P. (2021). Spatial-temporal variations in community mobility during lockdown, unlock, and the second wave of COVID-19 in India: A data-based analysis using Google's community mobility reports. *Spatial and Spatio-Temporal Epidemiology*, 39, 100442. <https://doi.org/10.1016/j.sste.2021.100442>
- Sarkar, S. K., Ekram, K. M. M., & Das, P. C. (2021). Spatial modeling of COVID-19 transmission in Bangladesh. *Spatial Information Research*. <https://doi.org/10.1007/s41324-021-00387-5>
- Shariati, M., Mesgari, T., Kasraee, M., & Jahangiri-rad, M. (2020). Spatiotemporal analysis and hotspots detection of COVID-19 using geographic information system (March and April, 2020). *Journal of Environmental Health Science and Engineering*, 18(2), 1499–1507. <https://doi.org/10.1007/s40201-020-00565-x>
- Sharma, N., Yadav, S., Mangla, M., Mohanty, A., Satpathy, S., Mohanty, S. N., & Choudhury, T. (2021). Geospatial multivariate analysis of COVID-19: A global perspective. *GeoJournal*. <https://doi.org/10.1007/s10708-021-10520-4>
- T.C.S.B. (2021). *T.C. Sağlık Bakanlığı*.
- Tang, W., Liao, H., Marley, G., Wang, Z., Cheng, W., Wu, D., & Yu, R. (2020). The changing patterns of coronavirus disease 2019 (COVID-19) in China: A tempogeographic analysis of the severe acute respiratory syndrome coronavirus 2 epidemic. *Clinical Infectious Diseases*, 71(15), 818–824. <https://doi.org/10.1093/cid/ciaa423>
- Tao, R., Downs, J., Beckie, T. M., Chen, Y., & McNelley, W. (2020). Examining spatial accessibility to COVID-19 testing sites in Florida. *Annals of GIS*, 26(4), 319–327. <https://doi.org/10.1080/19475683.2020.1833365>
- Tsai, P.-J., Lin, M.-L., Chu, C.-M., & Perng, C.-H. (2009). Spatial autocorrelation analysis of health care hotspots in Taiwan in 2006. *BMC Public Health*, 9(1), 464. <https://doi.org/10.1186/1471-2458-9-464>
- UN. (2020). *United Nations*. <https://www.un.org/en/>
- Vaz, E. (2021). Covid-19 in Toronto: A spatial exploratory analysis. *Sustainability (switzerland)*, 13(2), 498. <https://doi.org/10.3390/su13020498>
- Wang, J. F., Christakos, G., Han, W. G., & Meng, B. (2008). Data-driven exploration of “spatial pattern-time process-driving forces” associations of SARS epidemic in Beijing, China. *Journal of Public Health*, 30(3), 234–244. <https://doi.org/10.1093/pubmed/fdn023>
- Wen, T.-H., Lin, N. H., Lin, C.-H., King, C.-C., & Su, M.-D. (2006). Spatial mapping of temporal risk characteristics to improve environmental health risk identification: A case study of a dengue epidemic in Taiwan. *Science of the Total Environment*, 367(2–3), 631–640. <https://doi.org/10.1016/j.scitotenv.2006.02.009>
- Xiong, Y., Wang, Y., Chen, F., & Zhu, M. (2020). Spatial statistics and influencing factors of the COVID-19 epidemic at both prefecture and county levels in Hubei Province, China. *International Journal of Environmental Research and Public Health*, 17(11), 3903. <https://doi.org/10.3390/ijerph17113903>
- Zeren, F., & Yilanci, V. (2020). Analysing spatial patterns of the Covid-19 outbreak in Turkey. *Bingöl Üniversitesi İktisadi Ve İdari Bilimler Fakültesi Dergisi*, 4(2), 27–40.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.