

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Species interactions and climate change: How the disruption of species co-occurrence will impact on an avian forest guild

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1727539> since 2020-02-13T14:33:36Z

Published version:

DOI:10.1111/gcb.14953

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

1
2
3
4
5
6
7
8
9
10
11
12
13

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

[[Global Change Biology](#), 2019, DOI: 10.1111/gcb.13964]

The definitive version is available at:

La versione definitiva è disponibile alla URL:

<https://onlinelibrary.wiley.com/doi/full/10.1111/gcb.14953>

14 **Species interactions and climate change: how the disruption of species co-occurrence will**
15 **impact on an avian forest guild**

16

17 **Running Head:** Climate change and interactions in birds

18

19 Mattia Brambilla^{a,b}, Davide Scridel^{a,c}, Gaia Bazzi^b, Luca Ilahiane^b, Aaron Iemma^a, Paolo Pedrini^a,
20 Enrico Bassi^d, Radames Bionda^e, Luigi Marchesi^a, Fulvio Genero^f, Norbert Teufelbauer^g, Remo
21 Probst^g, Al Vrezec^h, Primož Kmeclⁱ, Tomaž Miheličⁱ, Giuseppe Bogliani^c, Hans Schmid^j, Giacomo
22 Assandri^{a,c}, Renato Pontarini^k, Veronika Braunisch^l, Raphaël Arlettaz^l, Dan Chamberlain^m

23

24

25 Mattia Brambilla and Davide Scridel should be considered joint first author

26

27 ^a Museo delle Scienze, Sezione Zoologia dei Vertebrati, Trento, Italy

28 ^b Fondazione Lombardia per l'Ambiente, Settore biodiversità e aree protette, Seveso (MB), Italy

29 ^c Dipartimento di Scienze della Terra e dell'Ambiente, Università degli Studi di Pavia, Pavia, Italy

30 ^d Parco Nazionale dello Stelvio, Bormio (SO), Italy

31 ^e Parco Naturale Alpe Veglia e Devero – Alta Valle Antrona, Varzo, Italy

32 ^f Riserva Naturale Regionale del Lago di Cornino, Cornino (UD), Italy

33 ^g BirdLife Austria, Wien, Austria

34 ^h National Research Institute, Ljubljana, Slovenia

35 ⁱ DOPPS BirdLife Slovenia, Ljubljana, Slovenia

36 ^j Schweizerische Vogelwarte, Sempach, Switzerland

37 ^k Progetto Lince Italia, Tarvisio (UD), Italy

38 ^l University of Bern, Institute of Ecology and Evolution, Division of Conservation Biology, Bern,

39 Switzerland

40 ^m Dipartimento di Scienze della Vita e Biologia dei Sistemi, Università di Torino, Turin, Italy

41

42 **Contact information**

43 corresponding author: Mattia Brambilla, phone: +39 02 80616123, email:

44 brambilla.mattia@gmail.com

45

46

47 **Abstract:**

48 Interspecific interactions are crucial in determining species occurrence and community assembly.

49 Understanding these interactions is thus essential for correctly predicting species' responses to

50 climate change. We focussed on an avian forest guild of four hole-nesting species with differing

51 sensitivities to climate, that show a range of well-understood reciprocal interactions, including

52 facilitation, competition and predation. We modelled the potential distributions of black

53 woodpecker and boreal, tawny and Ural owl, and tested whether the spatial patterns of the more

54 widespread species (excluding Ural owl) were shaped by interspecific interactions. We then

55 modelled the potential future distributions of all four species, evaluating how the predicted changes

56 will alter the overlap between the species' ranges, and hence the spatial outcomes of interactions.

57 Forest cover/type and climate were important determinants of habitat suitability for all species.

58 Field data analysed with N-mixture models revealed effects of interspecific interactions on current

59 species abundance, especially in boreal owl (positive effects of black woodpecker, negative effects

60 of tawny owl). Climate change will impact the assemblage both at species and guild-levels, as the

61 potential area of range-overlap, relevant for species interactions, will change in both proportion and

62 extent in the future. Boreal owl, the most climate-sensitive species in the guild, will retreat, and the

63 range-overlap with its main predator, tawny owl, will increase in the remaining suitable area:

64 climate change will thus impact on boreal owl both directly and indirectly. Climate change will

65 cause the geographical alteration or disruption of species interaction networks, with different

66 consequences for the species belonging to the guild and a likely spatial increase of competition
67 and/or intraguild predation. Our work shows significant interactions and important potential
68 changes in the overlap of areas suitable for the interacting species, which reinforce the importance
69 of including relevant biotic interactions in predictive climate change models for increasing forecast
70 accuracy.

71

72 **Keywords:** biotic interactions, citizen science, global warming, SDM, Strigidae, woodpeckers

73 **Introduction**

74 The importance of environmental factors such as climate, topography and land-cover in dictating
75 species distributions is well recognized in the literature (e.g. Hawkins et al., 2003; Elith &
76 Leathwick, 2009) and it is the basis of correlative species distribution models (SDMs). These
77 models, also known as environmental niche models (ENMs), have represented one of the most
78 frequent applications in ecology, biogeography and conservation over recent decades (see e.g.
79 Engler et al., 2017). In addition to the above-mentioned environmental factors, interspecific
80 interactions can also be crucial in determining species occurrence over different spatial scales (Wisz
81 et al., 2013), as well as in structuring biological communities (Bertness & Callaway, 1994). This
82 underlies the growing interest in macroecological models that include or evaluate biotic interactions
83 (Dormann et al., 2018). In fact, biotic interactions have been hypothesized from macroecological
84 patterns (Pollock et al., 2014), as well as used to improve distribution predictions for interacting
85 species (Araújo & Luoto, 2007; Heikkinen, Luoto, Virkkala, Pearson, & Körber, 2007).

86 One of the greatest recent challenges for ecologists is to predict the likely consequences of
87 climate change on species, communities and ecosystems (Valiente-Banuet et al., 2015): proper
88 forecasting is essential for species conservation and the maintenance of functioning ecosystems.
89 (Groves et al., 2012). A crucial point that severely complicates the assessment of the potential
90 impacts of climate change on wildlife is represented by its indirect effects via changes in biotic
91 interactions (Tylianakis, Didham, Bascompte, & Wardle, 2008), to the point that considering biotic
92 interactions is essential to correctly predict species' responses to climate change (Van der Putten,
93 Macel, & Visser, 2010). Climate change may disrupt trophic webs by altering the distribution of
94 species acting as key resources, competitors and predators, or by shifting phenologies of interacting
95 organisms, ultimately causing important changes in the nature of relationships between species
96 (Blois, Zarnetske, Fitzpatrick, & Finnegan, 2013; Kubelka et al., 2018; Van der Putten et al., 2010).

97 Facilitation (a positive interaction whereby one species promotes the occurrence of another) and
98 intraguild predation are two particular biotic interactions that have been found to be very important

99 for predicting the occurrence of several species (Heikkinen et al., 2007; Holt & Huxel, 2007), but
100 have received little attention in terms of how they could be altered by the influence of climate
101 change (but see (Bateman, Vanderwal, Williams, & Johnson, 2012). In fact, variation in climatic
102 conditions may impact on facilitation relationships as well as on intraguild predation (e.g. Rogers et
103 al., 2018), with potentially cascading effects over the entire system (Barton & Schmitz, 2009).

104 In this study, we focus on an avian forest guild of four hole-nesting species with different types
105 of reciprocal interactions, ranging from facilitation to competition and predation. The distribution of
106 the model species we considered is partially limited by climate, and in particular by temperature.
107 On this basis, our study system offers an unprecedented opportunity to evaluate the potential effect
108 of climate change on species distributions and, in particular, on different types of interaction among
109 species in the studied guild. On the basis of well-established interspecific relationships, we
110 formulated an *a priori* interaction scenario, that accounts for the effects of co-occurrence in this
111 guild. Then, using a large sample size and relevant environmental predictors, measured at a
112 biologically meaningful spatial scale, we modelled potential species distributions. Successively, we
113 evaluated whether the co-occurrence patterns that could be hypothesized on the basis of both
114 environmental suitability (according to species-specific SDMs), and whether potential biotic
115 interactions of the three more widespread species within the Alpine region were consistent with real
116 occurrence data (under current climatic conditions) gathered through dedicated fieldwork. Finally,
117 we modelled the potential future distribution of the study species according to the forecast future
118 climate, to provide an estimate of the potential impact of climate change on (co)occurrence and
119 hence on likely interspecific interactions.

120

121

122 **Material and methods**

123 **Study system**

124 The forest guild we investigated included four avian hole-nesting species, black woodpecker
125 *Dryocopus martius*, boreal owl *Aegolius funereus*, tawny owl *Strix aluco* and Ural owl *Strix*
126 *uralensis*. The study area encompassed the Alpine region, i.e. the Alps and the surrounding areas
127 across seven European countries (Fig. 1). This iconic mountain system harbours all the model
128 species, although they display rather different breeding distribution patterns. The Ural owl is limited
129 to the eastern portion of the study area, whereas the other three species occur over most of the Alps,
130 showing different associations with elevation belts and with forest types.

131

132 [Figure 1 approximately here]

133

134 Black woodpecker, the only diurnal taxon among the modelled species, is tied to mature forests,
135 with large stems and availability of dead wood and ant-rich habitats (Brambilla & Saporetti, 2014;
136 Karimi, Moradi, Rezaei, Brambilla, & Ghadimi, 2018; Pirovano & Zecca, 2014) over a wide
137 elevation gradient, from sea level to c.2000 m asl. This species has shown a dramatic increase in the
138 Alps, and colonization of previously unoccupied lowland forests (Nardelli et al., 2015). This
139 expansion is most likely due to an increase in forest quality and extent (Nardelli et al., 2015; see
140 also Mikusiński, 1995), yet no study has investigated the potential impact of climate change on this
141 species, with the exception of a paper suggesting future retraction in central Europe (Vos et al.,
142 2008).

143 The Alps are a climate *refugium* for boreal owl, representing a relict portion of the former range
144 that the species occupied in a colder past (Brambilla et al., 2015). According to this, climate change
145 will likely impact on boreal owl by reducing the suitability of most of its current range as a
146 consequence of increasing temperatures (Brambilla et al., 2017; Scridel et al., 2017) and/or by
147 altering its preferred breeding habitat type (coniferous or mixed forests) (Brambilla et al., 2015;

148 Hartl-Meier et al., 2014).

149 Tawny owl is a generalist species with a wide niche and distribution, occurring over most of
150 Europe in forest, farmland and also urban habitats, and occupying a broad climatic gradient (Francis
151 & Saurola, 2004; Vrezec & Tome, 2004a; Marchesi et al., 2006). In the Alps, the species is currently
152 expanding its distribution towards higher elevation, most likely due to milder climates (pers. obs.)
153 similar to the northwards expansion observed at higher latitudes, a response to warmer winters and
154 reduced snow cover (Francis & Saurola, 2004).

155 Ural owl, the largest of the owls studied here, is widely distributed in northern Eurasia (Konig,
156 Weick, & Becking, 1999). Towards the south it occurs mostly in mountain areas, inhabiting
157 intermediate elevations in the eastern Alps, particularly in mixed forests with mature trees and
158 clearings (Benussi & Genero, 2008; Rassati, 2006; Vrezec & Mihelič, 2013; Vrezec & Tome,
159 2004a). However, this pattern might be due to the lack of mature forest stands in the lowlands
160 because of intensive logging, since the Ural owl is relatively abundant in preserved mature forest
161 stand fragments in lowlands (Vrezec & Mihelič, 2013). Recent observations indicate that this
162 species is expanding in montane as well as in lowland forest areas in different parts of its southern
163 range in Europe (Bashta, 2009; Vrezec, 2019). So far, only a single study (Huntley, Green,
164 Collingham, & Willis, 2007) has evaluated the potential effect of climate change on its distribution
165 in central-southern Europe.

166 These four species represent an ideal set of interspecific interactions (Fig. 2) for testing the
167 potential disrupting effect of climate change. Black woodpecker facilitates the occurrence of tawny
168 and, especially, boreal owl, providing the great majority of nest cavities for the latter and potential
169 nesting sites for the former (Brambilla et al., 2013; Gustin, Brambilla, & Celada, 2010). Tawny owl
170 is one of the main predators of boreal owl (Konig et al., 1999; Mikkola, 1976). Ural owl can predate
171 both tawny and boreal owls (Mikkola, 1983); competitive exclusion of tawny owls from areas
172 occupied by Ural owls has been reported from northern (Korpimaki, 1986), southern (Vrezec &
173 Tome, 2004a) and eastern Europe (Kajtoch, Źmihorski, & Wieczorek, 2015). Even if Ural owl have

174 been reported to negatively affect breeding density and reproductive success of boreal owl in
175 northern Europe (Hakkarainen & Korpimäki, 1996), the competitive exclusion exerted by Ural owl on
176 tawny owl in the eastern Alps benefits the smaller boreal owl. Ural owl exerts a much lower
177 predation pressure on boreal owl than on tawny owl, and the sites free of tawny owl created by Ural
178 owl occurrence are regularly occupied by boreal owl (Vrezec & Tome, 2004b).

179

180 [Figure 2 approximately here]

181

182

183 **Data collection**

184 Two different datasets were used for this study. For distribution modelling, we gathered already
185 existing, georeferenced occurrence data, fulfilling the following requirements: spatial accuracy
186 equal or higher than 2 km, period 2000-2017, records within the breeding season of the target
187 species (March-June), or data with an associated atlas code indicating breeding or territorial
188 behaviour. Data were collected both via research projects and citizen science initiatives: i) during
189 previous surveys carried out within the framework of different projects (e.g. (Brambilla et al., 2015,
190 2017; Mihelič et al., 2019; Vrezec & Mihelič, 2013; Vrezec & Tome, 2004a, 2004b), ii) via online
191 platforms (www.ornitho.ch, www.ornitho.at, www.ornitho.it), after official requests specifying the
192 aims of the study. Data were from the study area and from neighbouring sites (i.e., areas
193 surrounding the study region, within the countries investigated; see Suppl. Mat.). A few occurrence
194 points of black woodpecker, located at high elevations (>2000m asl) outside the breeding habitat of
195 the species, were discarded as non representative of the environmental contexts used by the species
196 for reproduction. The final dataset used for modelling comprised 41911 records and included the
197 following sample sizes (number of 2km x 2km cells occupied by each species): 9323 for black
198 woodpecker, 1207 for boreal owl, 5791 for tawny owl, and 436 for Ural owl.

199 The second dataset was used for testing the current effects of interspecific interactions on the

200 presence of three of the species, boreal owl, tawny owl and black woodpecker, given the
201 simultaneous effects of environmental suitability and species co-occurrence. These data were
202 collected by means of dedicated surveys, carried out in northern Italy (in Lombardy region and
203 Trento province), during March-June 2017. Surveys consisted of point counts carried out in the
204 morning (for black woodpecker) and on the same day at dusk/night (for owls). Points were located
205 along several different valleys (see Fig. 1), at an average nearest neighbour distance of ~1100 m
206 (with a minimum of ~450 m in the case of different sides of the same relief), set according to local
207 morphology to avoid double counting of the same individuals as well as to avoid large, unsurveyed
208 tracts of valley. However, there were some general differences in the spacing of points because of
209 the variable geomorphology of the study sites. In Lombardy, the 122 survey sites had an average
210 distance between neighbouring points of c. 1.3 km, and most points were separated by at least 1 km,
211 except when placed on different sides of the same mountain massifs. Within the Trento province
212 study sites, neighbouring points were sometimes located at closer distances (average distance ~850
213 m, minimum ~450), because of the complex valley morphology in the survey sites. At each point,
214 after 10 min of listening to spontaneous vocalizations, if the target species was/were not recorded,
215 we broadcast territorial calls (taken from Roché & Chevereau (2000) of males (owls; playback
216 order: boreal, tawny) and drumming (woodpecker) for one minute (stopping immediately after any
217 contacts), and listened again for four minutes. 218 points were surveyed for black woodpecker (58
218 once, 99 twice, 51 three times). Of these, 192 points were also surveyed for owls (57 once, 91
219 twice, 44 three times). The estimated position of all individuals of target species was recorded on
220 aerial photographs or other detailed maps to avoid double counting from the same or neighbouring
221 points. Ural owl does not occur within the test area, but there is a strong evidence base from
222 intensive fieldwork in the eastern Alps of its interactions with the other species of the guild (Vrezec,
223 2019; Vrezec & Tome, 2004b, 2004a).

224

225 **Modelling current and future distributions**

226 To model the current and likely future distribution of the target species, we used environmental
227 niche models, which combine the occurrence data of a species with a set of environmental
228 predictors (including e.g. climatic, land-use/land-cover and topographical variables) to estimate the
229 suitability of a given area for the study species. We considered a grid composed of 2 km x 2 km
230 cells, covering the entire Alpine region. For each cell, we estimated the proportional cover of the
231 most representative land-use/land-cover types (from CORINE CLC; European Environment
232 Agency, 2016), tree density according to (Moreno, Neumann, & Hasenauer, 2017), and the average
233 value for global solar radiation in May (derived from a 30-m DEM and calculated in GRASS 7.04;
234 Neteler et al., 2012) and climatic variables (from CHELSA database; Karger et al., 2017). We then
235 removed from the environmental predictors those occurring only rarely within the study area
236 (identified by means of visual plotting of each predictor), and the most intercorrelated ones based
237 on CORVIF ($G_{VIF} < 16$) (Zuur, Ieno, Walker, Saveliev, & Smith, 2009); the resulting set of
238 variables is summarised in Table S1.

239 We adopted a maximum entropy approach by developing MaxEnt models (Jane Elith et al., 2011;
240 Phillips, Anderson, & Schapire, 2006) in R (R Development Core Team, 2016), using the package
241 ENMeval (Muscarella et al., 2014). We discarded all duplicates, i.e. records occurring within a 2km
242 x 2km cell already having a given species' record. We built models considering the effectively
243 sampled area, by restricting background points ($N = 10\ 000$) to cells with at least one record of any
244 of the target species. In this way, the background corresponded to the visited areas and
245 corresponding environmental characteristics. We built models limiting the type of species-habitat
246 relationships to linear and quadratic, to avoid overfitting; simpler models have to be preferred when
247 it is necessary to expand model outcomes over different areas or temporal scenarios (Brambilla,
248 Pedrini, Rolando, & Chamberlain, 2016). However, for black woodpecker, we also included hinge
249 relationships as the simpler model was not precise enough in terms of correspondence between
250 predicted distribution and current knowledge about real occurrence within the study region. For
251 each species, occurrence data were partitioned into two groups, according to a checkerboard scheme

252 (“checkerboard1” in ENMeval) with each of the units aggregating four original (2km x 2km) cells.
253 This allowed testing model validity over independent datasets, assessing model robustness and
254 enhancing generalizability. Model validity was checked by evaluating variations in discriminatory
255 power (AUC – Area Under the Curve of the receiver operating characteristic (ROC) plot) over the
256 two different data partitions (bins), and by checking omission rates on test data, which had been
257 reclassified using two threshold-dependent metrics, i.e. 10% training omission rates, and training
258 omission for minimum training presence (i.e. lowest suitability at occurrence sites used for training
259 the model); omission rates larger than the expected values suggest overfitting (Muscarella et al.,
260 2014).

261 We tested different values of the regularization multiplier: each model was trained with eight
262 different values (from 0.5 to 4), and then the one with the lowest AIC was selected. Successively, all
263 variables unlikely to be important for species’ distribution (i.e. with both permutation importance
264 and percentage contribution < 1) were discarded, and the model was run again with different values
265 of the regularization multiplier, until we obtained a most supported model with no variables
266 showing both permutation importance and percentage contribution < 1. The raw model outcome
267 was reclassified by means of a logistic transformation to allow an easier interpretation (Elith et al.,
268 2011). The final logistic output of the model was then reclassified into three-class maps of
269 suitability: unsuitable, partly suitable, and suitable. This reclassification was made on the basis of
270 some widely adopted thresholds, generally used for binary reclassification of MaxEnt models, i.e.
271 the 10th percentile and the *maximum training sensitivity plus specificity threshold* (Engler, Rödder,
272 Stiels, & Förchler, 2014; Liu, Berry, Dawson, & Person, 2005; Liu, White, & Newell, 2013).
273 Values between 0 and the lowest of such thresholds were considered as unsuitable, values between
274 the thresholds as partly suitable, and values above the highest one, as definitely suitable. To
275 calculate potential range overlap between species and changes in range extent between current and
276 future conditions for each species, we considered all sites with suitability higher than the lower
277 threshold as potentially occupied by a species (thus both partly suitable and definitely suitable were

278 treated as potentially occupied). Finally, for all species, we removed all those locations at an
279 elevation higher than 2200m asl from suitable and partly suitable areas. In fact, even if some cells
280 around that elevation could be predicted as suitable for the target species, the occurrence of the
281 mature forests required by them is very unlikely at such an elevation in the Alps, and will be rather
282 unlikely to reach it in the near future.

283 Distribution models were then projected over future scenarios of climate change, derived from
284 the HADGEM model, under the worst scenario (representative concentration pathway RCP 8.5,
285 IPCC, 2013), with future climate conditions (for 2050) taken from the Worldclim database
286 (www.worldclim.org; Fick & Hijmans, 2017). We selected a single, pessimistic, scenario because i)
287 we were interested in exploring the potential effect of climate change on a guild of interacting
288 species rather than in obtaining several alternative predictions, e.g. for planning or conservation
289 purposes, and ii) scenarios with larger changes are becoming unfortunately increasingly probable
290 (Peters et al., 2013).

291

292 **Testing the effects of interspecific interactions**

293 To check whether interspecific interactions have the potential to affect the model species, we carried
294 out a field test considering current patterns of co-occurrence of three of our species in a sample of
295 sites in the Italian Alps.

296 We tested for the effect of tawny owl abundance on the number of boreal owls at survey points,
297 and of the effect of black woodpecker occurrence on the local abundance of both tawny and boreal
298 owl. N-mixture models, developed using the package ‘unmarked’ (Fiske & Chandler, 2011) in R (R
299 Development Core Team, 2016), were employed for evaluating the effect of species co-occurrence
300 and environmental suitability on the latent abundance of the target species, while taking into
301 account the potential variation in detectability and hence imperfect detection. Under this approach,
302 repeated counts in a set of sites are used to estimate simultaneously the detectability and the
303 abundance of individuals at survey sites (Ficetola et al., 2018; Royle, 2004). We evaluated the

304 factors that can affect the species' local density by modelling the latent abundance of each species.

305 We assumed population closure because we focused on a single breeding season.

306 As factors potentially affecting the observation process (and hence detection), we tested time of
307 day, survey date, disturbance as a three-level categorical factor (absent; weak – some far or faint
308 noises; strong – close noise or human activities potentially affecting species detection by the
309 observer or even species behaviour), and wind, a three-level categorical factor (calm - Beaufort
310 scale 0-1; weak - Beaufort scale 2-3; moderate - Beaufort scale 4-5). For boreal owl, the number of
311 calling tawny owls was also considered as a variable potentially affecting detection. Instead of
312 entering several abiotic and habitat factors potentially determining species' abundance into the
313 model, we used the environmental suitability produced by the respective MaxEnt models for each
314 species (taking the value of the model cell including the surveyed point), and the maximum
315 abundance of tawny owl recorded at a site for boreal owl. We also tested for a positive effect of
316 occurrence of black woodpecker at a survey point as a proxy for nest-site availability for both owl
317 species. All continuous variables were standardized before the analyses for a better comparison of
318 their relative effects (Cade, 2015; Schielzeth, 2010). For each species, we then developed models
319 based on all possible variable combinations, and ranked them based on the AICc (Akaike's
320 Information Criterion for small sample size), using the package 'MuMIn' (Bartoń, 2016), and
321 checked whether the co-occurrence factors were selected in the most supported models. We
322 repeated the analysis excluding the points surveyed in the Trento province, which often were closer
323 to each other, considering only the points surveyed in Lombardy (see under "Data collection") to
324 check for consistency in the model results when only well-spaced points were included.

325

326

327 **Results**

328

329 **Current distribution and overlap**

330 For all modelled species, the discriminatory power over the two data partitions was nearly identical
331 (all differences < 0.013), omission rates at test sites according to the 10% training threshold showed
332 values invariably close (0.09-0.12) to the expected one (0.10) on both bins (data partitions), and
333 similarly omission values on both bins were always close to zero, as expected (< 0.005); most
334 importantly, the predicted distribution matched well the known breeding range. Summary statistics
335 for models are shown in Table S2. For Ural owl, suitable sites were predicted to occur not only
336 within the current range of the species in the eastern Alps, but also in the central and western part of
337 the Alpine region, although much more sparsely (Fig. 3).

338

339 [Figure 3 approximately here]

340

341 All species were associated with higher forest cover, although for boreal owl, this positive
342 association was relevant only for coniferous forest, and for tawny owl, only deciduous forest (Table
343 1). Furthermore, tawny owl was negatively associated with coniferous forest. Both Ural owl and
344 tawny owl also showed evidence of greater habitat suitability at intermediate cover of variables that
345 can be considered as proxies for open or semi-open habitats within forests (grassland, sparsely
346 vegetated areas, complex cultivation patterns, discontinuous urban fabric). In terms of climatic
347 variables, there was a clearer distinction in the response of the four species. Black woodpecker
348 exhibited a wide thermal niche, with average annual temperature from a few degrees below zero up
349 to 15°C, whilst boreal owl inhabited a cooler part of the temperature gradient, preferring values
350 between -2° and 5°C. Tawny owl was the only species showing a positive linear effect of average
351 temperature on habitat suitability. Habitat suitability for Ural owl was positively affected by annual
352 temperature range and annual precipitation, but negatively by precipitation seasonality (Table 1).

353

354

[Table 1 approximately here]

355

356 The overlap in current modelled distributions was highest for boreal owl and black woodpecker
357 (99% of the former species' range overlapping with the latter), followed by tawny and boreal owls
358 (25%), tawny and Ural owls (15%), boreal and Ural owls (12%), boreal-tawny-Ural owls (6%).

359

360 **Effect of interactions**

361 The N-mixture models revealed important effects of observation covariates on owl detection (see
362 Tables S3 and S4) and, most importantly, a positive effect of environmental suitability (as
363 calculated by MaxEnt models) on species abundance for both owl species considered in the field
364 surveys, with a particularly strong effect for tawny owl (for boreal owl, the variable ranking was
365 slightly different according to the dataset used; Tables S3 and S4). A positive effect of black
366 woodpecker occurrence was found for boreal owl, but not for tawny owl. A negative effect of the
367 abundance of tawny owl on that of boreal owl was also suggested by the models; for boreal owl, the
368 effect of variables describing interspecific interactions was particularly important (Table S3).
369 Notably, all boreal owls occurring in sites where black woodpecker was not detected during the
370 2017 survey, occupied sites predicted to be suitable for the woodpecker by the MaxEnt model, thus
371 potentially offering nest-sites excavated by the woodpecker in previous seasons. The test of
372 interaction effects therefore strongly supported the importance of interspecific interactions for
373 boreal owl, but not for tawny owl.

374

375 **Future distributions**

376 Predicted future environmental suitability for the target species is displayed in Fig. S2 and changes
377 are summarized in Table 2. The distribution of boreal owl will be substantially affected by climate
378 change, with a range reduction and especially a contraction towards higher elevation. Ural owl is

379 predicted to gain suitable areas in Austria and in the central and western Alps, where it is currently
380 absent. Tawny owl will likely show a range expansion towards higher elevations, especially in the
381 central Alps. Black woodpecker distribution will likely undergo only minor changes, even under the
382 rather 'extreme' climate change scenario considered.

383

384 [Table 2 approximately here]

385

386 The potential overlap between the interacting species will change in the future (Fig. 4; see Table
387 S5 for absolute variation). Almost the entire range of boreal owl is predicted to be suitable for black
388 woodpecker both in current and future conditions (Fig. 5); the decrease in extent of potential
389 overlap, due to owl contraction (Fig. 4 and Table S5) will not impact on the potential interactions.
390 The predicted overlap between boreal owl and its main intraguild predator, tawny owl, will show a
391 marked (proportional) increase (from 25% to 54% of boreal owl potential range) and, importantly,
392 will increase in the central Alps (one of the strongholds for boreal owl under a changing climate),
393 because milder climates will enable tawny owl to expand its potential distribution in Alpine valleys.
394 The likely overlap between boreal and Ural owls will decrease in absolute terms, but will keep
395 nearly stable in relative share over the boreal owl range. The areas where Ural owl occurrence could
396 benefit boreal owl (i.e. those with potential occurrence of tawny owl) will decrease (from 6% to 2%
397 of boreal owl range). Finally, the overlap between tawny owl and its competitor/intra-guild
398 predator, the Ural owl, will probably increase over most of the Alps.

399

400

401 [Figure 4 approximately here]

402 [Figure 5 approximately here]

403

404

405 **Discussion**

406 Modification of species interactions in terms of spatial or functional patterns is a potentially crucial,
407 but usually overlooked, consequence of climate change on biological communities. To our
408 knowledge, this is one of the few studies that has investigated patterns of co-occurrence for an
409 interacting guild of birds in response to climate change. We have demonstrated the importance of
410 these interactions on the current distribution of the focal species based on field surveys (boreal owl,
411 tawny, black woodpecker) and empirical evidence (Ural owl). Our combined analyses indicated that
412 both environmental variables (habitat and climate) and interactions with other species were
413 important predictors of species occurrence. The models predicted the current species distribution
414 well and therefore were used to evaluate the potential disruption of the interaction network (via
415 changes in spatial co-occurrence) in this guild in response to future climatic alterations. We have
416 shown that changes to the interaction network are likely to have highly variable effects depending
417 on the particular species, but for boreal owl at least, a spatial increase in areas with negative
418 interactions (without compensatory increases in areas with positive interactions) is very likely to
419 have net negative effects in the future.

420

421 **Large-scale environmental predictors of species occurrence**

422 Environmental correlates of habitat suitability for all model species were coherent with the
423 biological and ecological requirements found in the literature (e.g. Lundberg, 1980; Vrezec &
424 Bertoneclj, 2018; Vrezec & Tome, 2004b). Considering the link with climate, black woodpecker
425 and tawny owl showed the broadest thermal niche, consistent with their wider distribution over the
426 Palearctic. Habitat suitability for tawny owl in the Alps is linearly and positively affected by
427 temperature and thus a positive outcome of the temperature increase could be expected for that
428 species. The recent increase of tawny owl at higher elevations observed in several sites in the Alps
429 (all authors, pers. obs.) confirms this pattern. While temperature changes in the Alpine region are
430 unlikely to severely impact black woodpecker and Ural owl, a strong effect could be expected for

431 boreal owl. For the woodpecker, a possible minor shift towards upper elevations could be expected
432 under extreme scenarios, which is consistent with the broader distribution of the species, which is
433 much more abundant in mountain areas in southern Europe. Boreal owl has already been reported as
434 a climate-sensitive species in the Alps (Brambilla et al., 2015), its distribution at the European scale
435 appears strictly related to temperature (Brambilla et al., 2017), and it is among the cold-adapted
436 species undergoing population decline and range contraction in Europe (Korpimaki & Hakkarainen,
437 2012) and Italy (Scridel et al., 2017).

438

439 **Interspecific interactions, climate change and its consequences**

440 The relative abundance of boreal owl at sampling sites revealed the potential importance of co-
441 occurrence patterns on the species' distribution. Black woodpecker occurrence and abundance of
442 tawny owl were indeed even more important than environmental suitability *per se* for boreal owl in
443 the Central Italian Alps (see under 'Modelling issues' for further discussion), and likely also in the
444 wider Alpine region (Vrezec & Tome, 2004b). This means that, within this largely suitable belt,
445 interspecific dynamics play an important role in driving the occurrence of boreal owl, the species
446 most sensitive to interactions of those investigated. For tawny owl, the presence of black
447 woodpecker is less relevant as the former species has a greater flexibility in selecting suitable
448 cavities for breeding, which include woodpecker holes, but also a variety of old nests, rotten tree
449 trunks, other holes, ledges in rock cliffs and even buildings (Mikkola, 1983). For tawny owl, the
450 environmental suitability derived from MaxEnt models was a better predictor of abundance.

451

452 **Changes in the predicted 'room for interactions'**

453 According to our analysis, climate change will strongly impact on the investigated species both at
454 species level and in the form of community changes in interacting species resulting from
455 distribution shifts, as the area where species interactions are likely to occur is predicted to vary in
456 extent in the future. In most cases, the potential overlap between species range will decline, but it

457 will likely increase for tawny owl and Ural owl, potentially increasing the frequency of competitive
458 interactions between the species. The spatial relevance of the facilitator role of black woodpecker in
459 favour of boreal owl will probably remain unchanged.

460 Boreal owl was the most climate-sensitive species, and will retreat further into the mountains.
461 For this reason alone, the species will lose 65% of its habitat. Therefore, it is not surprising that,
462 while the absolute overlap with the tawny owl will decrease (-26%), the overlap will increase in the
463 area remaining suitable for boreal owl (from 25% to 54%). Since the potential overlap with the
464 black woodpecker and the Ural owl will remain roughly the same, climate change will have a
465 negative effect on the boreal owl both directly (via contraction of suitable areas) and indirectly, via
466 a likely increase in the overlap with tawny owl.

467 Interestingly, our modelling outcomes suggested a potential westwards expansion of Ural owl.
468 This species was until recently confined to the very eastern side of the Alps and eastern Europe
469 (Vrezec, 2009), but was more westerly distributed in historical times (Goffette, Denis, Pöllath, &
470 van Neer, 2016) and in recent decades it has colonized new areas, expanding its range towards the
471 central portion of the Alpine chain (Benussi & Genero, 2008, 2017; Nardelli et al., 2015; Rassati,
472 2006, 2017). Therefore, the modelled increase in suitability in the central and western Alps is fully
473 coherent with the current pattern of range expansion. Successful reintroduction projects recently
474 carried out in Lower Austria (Zink & Walter, 2018) further confirm environmental suitability of the
475 central-eastern Alps for the species, where the provisioning of nest-boxes, which compensates for
476 the widespread lack of nesting sites due to forest harvesting, could further favour the species'
477 westward expansion. The ongoing expansion of Ural owl, coupled with that of tawny owl, implies
478 an increasing potential overlap and thus likely increasing interactions between these two competing
479 species (Figs. 4 and 5).

480

481 **Modelling issues**

482 The distribution models we obtained (at a spatial scale highly representative of the territory

483 size/home range of the species) appeared rather robust for all species, with a high level of
484 consistency in discriminatory power over the two partitions of the dataset (Table S2). The resulting
485 predicted distributions were in line with the current range of target species in the Alps. Similarly,
486 the species-habitat relationships underlying the models were coherent with the knowledge of
487 species' ecology.

488 Other species interact with the target ones. These basically include prey, and especially voles
489 (Brommer, Pietiäinen, & Kolunen, 2002; Korpimäki & Hakkarainen, 2012; Vrezec, Saurola,
490 Avotins, Kocijančič, & Sulkava, 2018), as well as other predators, like goshawk *Accipiter gentilis*
491 and eagle owl *Bubo bubo* (Byholm, Burgas, Virtanen, & Valkama, 2012; Hakkarainen &
492 Korpimäki, 1996; Lõhmus, 2003; Mikkola, 1976, 1983; Sergio, Marchesi, Pedrini, & Penteriani,
493 2007). Such additional factors might further modulate the effect of competition and coexistence at a
494 finer scale (Ciach, 2008; Ciach & Czyżowicz, 2014).

495 Finally, local forest characteristics, potentially sensitive to human management and climate
496 change (Braunisch et al., 2014), can be important, especially for black woodpecker (Karimi et al.,
497 2018; Pirovano & Zecca, 2014). However, at broader scales they are unlikely to be relevant (see e.g.
498 Tjernberg, Johnsson, & Nilsson, 1993; Brambilla & Saporetto, 2014), as the ongoing range
499 expansion in a large part of the study area suggests (Gustin, Brambilla, & Celada, 2019). In
500 addition, we cannot completely exclude the possibility that the effect of black woodpecker and
501 tawny owl (positive and negative, respectively) on boreal owl occurrence, highlighted by N-mixture
502 models, could be due to an influence of some unmeasured habitat variable which has an effect on
503 both interacting species of a given species pair. However, we are confident that these effects are
504 likely mirroring true interaction effects. Black woodpeckers provide almost all boreal owl nest-sites
505 in the study area (Pedrini, Caldonazzi, & Zanghellini, 2005), and hence a potential unmeasured
506 habitat variable could also be represented by a better-than-average availability of cavities. On the
507 other side, tawny owls have been shown to significantly impact on boreal owl occurrence (Vrezec &
508 Tome, 2004b), and we are aware of several cases of local replacement in recent years (our pers. obs.

509 from Italy, Austria and Slovenia). Nevertheless, long-term studies that include more detailed habitat
510 descriptions would better elucidate whether the positive and negative relationships are consistent
511 with regard to temporal variation in the same habitat, and therefore some caution is needed in
512 interpreting the observed patterns.

513

514 **Conclusions**

515 Climate change will result in the disruption or alteration of species interaction networks (Blois et
516 al., 2013; Tylianakis et al., 2008). Several studies have investigated the potential impacts of
517 variations in climate on animal-plant networks (insects and host species, plants and their pollinators;
518 e.g. Schweiger et al., 2008; Gorostiague et al., 2018). However, fewer studies have dealt with
519 changes in interspecific relationships among vertebrates, with the main exception represented by
520 carnivorous mammals (e.g. Zielinski et al., 2017; Pandey & Papeş, 2018; Scully et al., 2018). Here,
521 we have shown how climate change will result in changes in the distribution overlap in a guild of
522 interacting species, with different consequences for the species belonging to this guild. The process
523 will likely result in an increase in areas where the target species will experience competition and/or
524 intraguild predation rates, due to a higher proportional range overlap between subordinate (boreal
525 and tawny owl) and dominant (tawny and Ural owl, respectively) species. In turn, these changes
526 will probably enhance the importance of interaction effects for those species at the regional scale.
527 The facilitation provided by black woodpecker to boreal owl (nest provision) would instead remain
528 substantially unchanged, but the latter species will likely be the most negatively affected by climate
529 change.

530 Our work modelled the potential effects of climate change on the distribution of an interacting
531 owl guild and of its main nest facilitator, the black woodpecker, and showed potential important
532 changes in the overlap of suitable areas for those interacting species. The analysis of current
533 patterns of abundance at the local scale confirmed the likely importance of interspecific
534 interactions. Taken together, our findings suggest that future predictions of species distribution

535 under climate change should include relevant biotic interactions to achieve higher forecast
536 accuracy; in particular, testing the relevance of interspecific interactions will facilitate the
537 interpretation of distribution models and the more reliable estimation of predicted range changes.

538

539

540

541 **Acknowledgements**

542 Swiss data were archived in www.ornitho.ch and provided by the Swiss Ornithological Institute.

543 Austrian data were archived in www.ornitho.at and provided by BirdLife Austria. Slovenian data
544 were in the most part archived in www.ptice.si/noags and provided by DOPPS BirdLife Slovenia.

545 Italian data were partly provided by www.ornitho.it. Data from Piedmont were partly provided by
546 GPSO; data from Trentino were largely provided by the MUSE/PAT webGIS developed under the
547 project LIFE+ TEN. We are very grateful to Fabrizio Florit, ASTORE FVG (board council and

548 members), Matteo de Luca, Matteo Skodler, Alessandro Bertoli, Enrico Benussi, Matteo Toller,

549 Adamo Moreno, and Regione Friuli-Venezia Giulia for providing data from this region. We are

550 grateful to Matteo Anderle, Alessandro Franzoi, Andrea Parisi, Piergiovanni Partel, Diego Rubolini,

551 Roberto Lardelli, and to all observers contributing data to the online platforms www.ornitho.at,

552 www.ornitho.ch, www.ornitho.it, www.ptice.si/noags. We are also very grateful to an anonymous

553 reviewer for helpful comments on a first draft of the manuscript. MB, GBa and LI were partly

554 supported by the project LIFE+ GESTIRE2020. Fieldwork in Trentino was partly supported under

555 provincial RDP fundings. Parco Naturale Paneveggio-Pale di San Martino partly supported this

556 research as part of DS's PhD. AV was supported by research core funding No. P1-0255 by the

557 Slovenian Research Agency.

558

559 **Author contributions**

560 MB conceived the idea, which was then developed by all authors. MB, DS, PP, GBa, LI, EB, RB,

561 LM, FG, NT, RP, AV, PK, TM, RP, GA, HS collected data on the field and/or through database
562 interrogation. DS processed bird data; DS and AI prepared environmental layers. MB ran the
563 analyses and all authors contributed to model checking/development and check final model
564 outcomes. MB with DS' help wrote a first draft of the ms. All authors critically contributed to the
565 final version.

566

577 **Table 1.** Environmental factors used to model species distributions that were selected in at least one
578 model (see Table S1 for full list), the relative importance of each factor (percentage
579 contribution/permutation importance) and short description of the effect (within brackets; relative to
580 the model including all the selected predictors) according to final models for each species.
581 Numerical codes for land cover variables represent CORINE categories. Symbols used for effects:
582 +: positive, -: negative, +/-: quadratic (hump-shaped), -/++: quadratic (U-shaped), +/- -: quadratic
583 (hump-shaped)/negative, 0: nearly null (very weak positive effect).

574

Variable	Description	Boreal owl	Tawny owl	Ural owl	Black woodpecker
bio_1	Annual Mean Temperature	76.60/81.27 (+/-)	31.93/54.68 (+)		16.91/20.23 (+/-)
bio_12	Annual Precipitation		1.47/2.67 (+)	17.61/2.80 (+)	14.17/36.82 (-)
bio_15	Precipitation Seasonality (Coefficient of Variation)	2.96/4.21 (-)	5.93/0.00 (-)	12.47/22.94 (-)	
bio_19	Precipitation of Coldest Quarter	2.64/0.37 (-/+)		4.05/0.00 (0)	
bio_7	Temperature Annual Range (BIO5-BIO6)	2.23/5.66 (-)	24.01/35.13 (-)	15.99/14.94 (+)	
solarMay	Global solar radiation for May		1.66/4.49 (+)	0.86/0.78 (+)	8.24/13.22 (+/-)
X15	2.2.1 Vineyards				0.64/0.66 (-)
X18	2.3.1 Pastures		1.02/0.43 (+)	0.61/0.22 (-)	0.20/0.69 (-)
X1.1	1.1.2 Continuous urban fabric			0.38/0.94 (+)	0.90/0.25 (-)
X2.1	1.1.1 Discontinuous urban fabric	0.29/1.05 (-)	0.85/0.77 (+)	1.03/0.94 (+)	1.60/0.0 (-)
X20	2.4.2 Complex cultivation patterns		0.25/1.21 (+)		0.65/3.33 (-)
X21	2.4.3 Land principally occupied by agriculture, with significant areas of natural vegetation			0.62/0.42 (-)	
X25	3.1.3 Mixed forest			4.75/0.00 (+)	
X26	3.2.1 Natural grasslands		9.44/0.00 (-)		
X31	3.3.2 Bare rocks		2.79/0.44 (-)		
X32	3.3.3 Sparsely vegetated areas	0.75/0.48 (-)	6.03/0.00 (-)	1.55/1.50 (-)	
X35	4.1.1 Inland marshes				1.61/2.97 (+)
X4	1.2.2 Road and rail networks and associated land				0.49/2.31 (-)
X40	5.1.1 Water courses				1.03/3.43 (+)
x2632_TCD_TCD_20m	Tree cover density	0.72/1.63 (+)		25.89/0.22 (-)	18.07/11.02 (+/-)
X2.2	x2632_TCD_FTY_20m Coniferous forest	13.83/5.33 (+)	14.62/0.19 (-)	5.16/17.55 (+)	33.63/2.00 (+)

575

576 **Table 2.** Predicted extent of suitable habitats in current and future conditions (RCP 8.5 scenario for
577 2050) for the target species.

578

Species	current extent of suitable area (km²)	future extent of suitable area (km²)	change (%)
boreal owl	85644	29988	-65
tawny owl	95200	120928	+27
Ural owl	20012	67200	+236
black woodpecker	160452	151564	-6

579

580

581

582

583 **Figure captions**

584

585 **Figure 1.** Study area. The darker the colour, the higher the elevation. The inset shows the location
586 of the point counts (yellow dots) used to test the interspecific effects on current distribution patterns
587 for boreal and tawny owl.

588

589 **Figure 2.** Schematic representation of the interspecific interactions characterizing the study guild of
590 forest birds. Larger arrows indicate stronger effects. The negative effect of Ural owl on tawny owl,
591 in combination with the negative effect of tawny owl on boreal owl, appears as facilitation for
592 boreal owl when viewed in isolation.

593

594 **Figure 3.** Modelled current distribution of boreal owl (upper left), tawny owl (upper right), Ural
595 owl (lower left) and black woodpecker (lower right). The darker the colour, the higher the
596 environmental suitability.

597

598 **Figure 4.** Predicted range overlap under current (yellow) and future climatic conditions (blue; in
599 green, areas with predicted overlap under both scenarios), between boreal owl and black
600 woodpecker (upper left), boreal owl and tawny owl (upper right), tawny owl and Ural owl (lower
601 left), boreal owl and Ural owl (lower right).

602

603 **Figure 5.** Graphical representation of predicted extent of overlap between species and relative
604 variation due to climate change. For each scenario (current and future), boreal owl range is depicted
605 in beige, and tawny owl range in brown. The proportional overlap with other species is shown in
606 each pie chart, in black (black woodpecker for boreal owl), brown (tawny owl for boreal owl), grey
607 (Ural owl for both species). For boreal owl, the overlap with both tawny and Ural owl is shown in
608 grey-brown. Pie chart size is proportional to the relative value of current (100%) and future species

609 range (35% for boreal owl, 127% for tawny owl).

610

- 612 Araújo, M. B., & Luoto, M. (2007). The importance of biotic interactions for modelling
613 species distributions under climate change. *Global Ecology and Biogeography*, 16(6),
614 743–753. <https://doi.org/10.1111/j.1466-8238.2007.00359.x>
- 615 Barton, B. T., & Schmitz, O. J. (2009). Experimental warming transforms multiple predator
616 effects in a grassland food web. *Ecology Letters*, 12(12), 1317–1325.
617 <https://doi.org/10.1111/j.1461-0248.2009.01386.x>
- 618 Bartoń, K. (2016). *MuMIn: Multi-Model Inference. R package version 1.15.6*. Retrieved from
619 <https://cran.r-project.org/web/packages/MuMIn/index.html>
- 620 Bashta, A.-T. (2009). Ural Owl *Strix uralensis* Population Dynamics and Range Expansion in
621 Western Ukraine. *Ardea*, 97(4), 483–487. <https://doi.org/10.5253/078.097.0412>
- 622 Bateman, B. L., Vanderwal, J., Williams, S. E., & Johnson, C. N. (2012). Biotic interactions
623 influence the projected distribution of a specialist mammal under climate change.
624 *Diversity and Distributions*, 18(9), 861–872. [https://doi.org/10.1111/j.1472-](https://doi.org/10.1111/j.1472-4642.2012.00922.x)
625 [4642.2012.00922.x](https://doi.org/10.1111/j.1472-4642.2012.00922.x)
- 626 Benussi, E., & Genero, F. (2008). Allocco degli Urali *Strix uralensis*: nidificazione in
627 cassetta-nido e aggiornamento della distribuzione in Italia. *Quaderni Di Birdwatching*,
628 10, https://www.ebnitalia.it/Qb/QB020/allocco_ur.htm. Retrieved from
629 https://www.ebnitalia.it/Qb/QB020/allocco_ur.htm
- 630 Benussi, E., & Genero, F. (2017). Nuovi dati sulla distribuzione e nidificazione di allocco
631 degli Urali *Strix uralensis* nel Tarvisiano (Alpi orientali). In S. G. Fasano & D. Rubolini
632 (Eds.), *Riassunti del XIX Convegno Italiano di Ornitologia. Torino, 27 settembre - 1*
633 *ottobre 2017* (pp. 28–29). Torino: Tichodroma. Monografie del Gruppo Piemontese Studi
634 Ornitologici.
- 635 Bertness, M. D., & Callaway, R. (1994). Positive interactions in communities. *Trends in*
636 *Ecology and Evolution*, 9(5), 191–193. [https://doi.org/10.1016/0169-5347\(94\)90088-4](https://doi.org/10.1016/0169-5347(94)90088-4)
- 637 Blois, J. L., Zarnetske, P. L., Fitzpatrick, M. C., & Finnegan, S. (2013). Climate change and
638 the past, present, and future of biotic interactions. *Science*, Vol. 341, pp. 499–504.
639 <https://doi.org/10.1126/science.1237184>
- 640 Brambilla, M., Bassi, E., Bergero, V., Casale, F., Chemollo, M., Falco, R., ... Vitulano, S.
641 (2013). Modelling distribution and potential overlap between Boreal Owl *Aegolius*
642 *funereus* and Black Woodpecker *Dryocopus martius*: implications for management and
643 monitoring plans. *Bird Conservation International*, 23(04), 502–511.
644 <https://doi.org/10.1017/S0959270913000117>
- 645 Brambilla, M., Bergero, V., Bassi, E., & Falco, R. (2015). Current and future effectiveness of
646 Natura 2000 network in the central Alps for the conservation of mountain forest owl
647 species in a warming climate. *European Journal of Wildlife Research*, 61(1), 35–44.
648 <https://doi.org/10.1007/s10344-014-0864-6>
- 649 Brambilla, M., Caprio, E., Assandri, G., Scridel, D., Bassi, E., Bionda, R., ... Chamberlain, D.

- 650 (2017). A spatially explicit definition of conservation priorities according to population
651 resistance and resilience, species importance and level of threat in a changing climate.
652 *Diversity and Distributions*, 23(7), 727–738. <https://doi.org/10.1111/ddi.12572>
- 653 Brambilla, M., Pedrini, P., Rolando, A., & Chamberlain, D. E. (2016). Climate change will
654 increase the potential conflict between skiing and high-elevation bird species in the Alps.
655 *Journal of Biogeography*, 43(11), 2299–2309. <https://doi.org/10.1111/jbi.12796>
- 656 Brambilla, M., & Saporetti, F. (2014). Modelling distribution of habitats required for different
657 uses by the same species: Implications for conservation at the regional scale. *Biological*
658 *Conservation*, 174, 39–46. <https://doi.org/10.1016/j.biocon.2014.03.018>
- 659 Braunisch, V., Coppes, J., Arlettaz, R., Suchant, R., Zellweger, F., & Bollmann, K. (2014).
660 Temperate Mountain Forest Biodiversity under Climate Change: Compensating Negative
661 Effects by Increasing Structural Complexity. *PLoS ONE*, 9(5), e97718.
662 <https://doi.org/10.1371/journal.pone.0097718>
- 663 Brommer, J. E., Pietiäinen, H., & Kolunen, H. (2002). Reproduction and Survival in a
664 Variable Environment: Ural Owls (*Strix Uralensis*) and the Three-Year Vole Cycle. *The*
665 *Auk: Ornithological Advances*, 119(2), 544–550. [https://doi.org/10.1642/0004-](https://doi.org/10.1642/0004-8038(2002)119[0544:raslav]2.0.co;2)
666 [8038\(2002\)119\[0544:raslav\]2.0.co;2](https://doi.org/10.1642/0004-8038(2002)119[0544:raslav]2.0.co;2)
- 667 Byholm, P., Burgas, D., Virtanen, T., & Valkama, J. (2012). Competitive exclusion within the
668 predator community influences the distribution of a threatened prey species. In *Ecology*
669 (Vol. 93). Retrieved from [https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.1890/12-](https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.1890/12-0285.1)
670 [0285.1](https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.1890/12-0285.1)
- 671 Cade, B. S. (2015). Model averaging and muddled multimodel inferences. *Ecology*, 96(9),
672 2370–2382. <https://doi.org/10.1890/14-1639.1>
- 673 Ciach, M. (2008). Abundance and distribution patterns of owls in Pieniny National Park,
674 Southern Poland. *Acta Zoologica Cracoviensia*, 48(1), 21–33.
675 <https://doi.org/10.3409/173491505783995680>
- 676 Ciach, M., & Czyżowicz, S. (2014). Abundance and distribution of owls Strigiformes in the
677 Pieniny Mountains National Park (southern Poland)-the pattern of changes in the
678 protected area after 10 years. In *Ornis Polonica* (Vol. 55). Retrieved from
679 http://www.ornis-polonica.pl/_pdf/OP_2014_2_083-095.pdf
- 680 Dormann, C. F., Bobrowski, M., Dehling, D. M., Harris, D. J., Hartig, F., Lischke, H., ...
681 Kraan, C. (2018). Biotic interactions in species distribution modelling: 10 questions to
682 guide interpretation and avoid false conclusions. *Global Ecology and Biogeography*.
683 <https://doi.org/10.1111/geb.12759>
- 684 Elith, J., & Leathwick, J. (2009). Species distribution models: ecological explanation and
685 prediction across space and time. *Annual Review of Ecology, Evolution*, . Retrieved from
686 <http://www.annualreviews.org/eprint/HWR4cusJrXYCSPZ9sUDj/full>
- 687 Elith, Jane, Phillips, S. J., Hastie, T., Dudík, M., Chee, Y. E., & Yates, C. J. (2011). A

- 688 statistical explanation of MaxEnt for ecologists. *Diversity and Distributions*, 17(1), 43–
689 57. <https://doi.org/10.1111/j.1472-4642.2010.00725.x>
- 690 Engler, J. O., Rödder, D., Stiels, D., & Förchler, M. I. (2014). Suitable, reachable but not
691 colonised: seasonal niche duality in an endemic mountainous songbird. *Journal of*
692 *Ornithology*, 155(3), 657–669. <https://doi.org/10.1007/s10336-014-1049-5>
- 693 Engler, J. O., Stiels, D., Schidelko, K., Strubbe, D., Quillfeldt, P., & Brambilla, M. (2017).
694 Avian SDMs: current state, challenges, and opportunities. *Journal of Avian Biology*, Vol.
695 48, pp. 1483–1504. <https://doi.org/10.1111/jav.01248>
- 696 European Environment Agency. (2016). Corine Land Cover 2012. Retrieved from
697 <https://www.eea.europa.eu/data-and-maps/data/external/corine-land-cover-2012> website:
698 <https://www.eea.europa.eu/data-and-maps/data/external/corine-land-cover-2012>
- 699 Ficetola, G. F., Barzaghi, B., Melotto, A., Muraro, M., Lunghi, E., Canedoli, C., ... Manenti,
700 R. (2018). N-mixture models reliably estimate the abundance of small vertebrates.
701 *Scientific Reports*, 8(1). <https://doi.org/10.1038/s41598-018-28432-8>
- 702 Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate
703 surfaces for global land areas. *International Journal of Climatology*.
704 <https://doi.org/10.1002/joc.5086>
- 705 Fiske, I. J., & Chandler, R. B. (2011). unmarked : An R Package for Fitting Hierarchical
706 Models of Wildlife Occurrence and Abundance. *Journal of Statistical Software*, 43(10),
707 1–23. <https://doi.org/10.18637/jss.v043.i10>
- 708 Francis, C. M., & Saurola, P. (2004). Estimating components of variance in demographic
709 parameters of Tawny Owls, *Strix aluco*. *Animal Biodiversity and Conservation*, 27(1),
710 489–502. <https://doi.org/57354>
- 711 Goffette, Q., Denis, M., Pöllath, N., & van Neer, W. (2016). Change in historical range of the
712 ural owl in Europe. *Belgian Journal of Zoology*, 146(1), 33–43.
- 713 Gorostiague, P., Sajama, J., & Ortega-Baes, P. (2018). Will climate change cause spatial
714 mismatch between plants and their pollinators? A test using Andean cactus species.
715 *Biological Conservation*, 226, 247–255. <https://doi.org/10.1016/j.biocon.2018.07.003>
- 716 Groves, C. R., Game, E. T., Anderson, M. G., Cross, M., Enquist, C., Ferdaña, Z., ... Shafer,
717 S. L. (2012). Incorporating climate change into systematic conservation planning.
718 *Biodiversity and Conservation*, 21(7), 1651–1671. <https://doi.org/10.1007/s10531-012-0269-3>
- 720 Gustin, M., Brambilla, M., & Celada, C. (2010). *Valutazione dello Stato di Conservazione*
721 *dell'avifauna italiana. Volume I. Non Passeriformes*. Ministero dell'Ambiente e della
722 Tutela del Territorio e del Mare, Lega Italiana Protezione Uccelli (LIPU).
- 723 Gustin, M., Brambilla, M., & Celada, C. (2019). *Conoscerli, proteggerli. Guida allo stato di*
724 *Conservazione degli uccelli in Italia*. Parma: LIPU/BirdLife Italia.

- 725 Hakkarainen, H., & Korpimäki, E. (1996). Competitive and Predatory Interactions among
726 Raptors: An Observational and Experimental Study. *Ecology*, 77(4), 1134–1142.
727 <https://doi.org/10.2307/2265582>
- 728 Hartl-Meier, C., Zang, C., Dittmar, C., Esper, J., Göttelein, A., & Rothe, A. (2014).
729 Vulnerability of Norway spruce to climate change in mountain forests of the European
730 Alps. *Climate Research*, 60(2), 119–132. <https://doi.org/10.3354/cr01226>
- 731 Hawkins, B. A., Field, R., Cornell, H. V., Currie, D. J., Guégan, J.-F. F., Kaufman, D. M., ...
732 Turner, J. R. G. (2003). Energy, water, and broad-scale geographic patterns of species
733 richness. *Ecology*, 84(12), 3105–3117. <https://doi.org/10.1890/03-8006>
- 734 Heikkinen, R. K., Luoto, M., Virkkala, R., Pearson, R. G., & Körber, J. (2007). Biotic
735 interactions improve prediction of boreal bird distributions at macro-scales. *Global*
736 *Ecology and Biogeography*, 16(6), 754–763. [https://doi.org/10.1111/J.1466-](https://doi.org/10.1111/J.1466-8238.2007.00345.X)
737 [8238.2007.00345.X](https://doi.org/10.1111/J.1466-8238.2007.00345.X)
- 738 Holt, R. D., & Huxel, G. R. (2007). Alternative prey and the dynamics of intraguild predation:
739 Theoretical perspectives. *Ecology*, 88(11), 2706–2712. <https://doi.org/10.1890/06-1525.1>
- 740 Huntley, B., Green, R. E., Collingham, Y. C., & Willis, S. G. (2007). A climatic atlas of
741 European breeding birds. In *A climatic atlas of European breeding birds*.
- 742 IPCC. (2013). Working Group I Contribution to the IPCC Fifth Assessment Report, Climate
743 Change 2013: The Physical Science Basis. In *Ipcc* (Vol. AR5).
744 <https://doi.org/10.1017/CBO9781107415324.Summary>
- 745 Kajtoch, Ł., Żmihorski, M., & Wieczorek, P. (2015). Habitat displacement effect between two
746 competing owl species in fragmented forests. *Population Ecology*, 57(3), 517–527.
747 <https://doi.org/10.1007/s10144-015-0497-y>
- 748 Karger, D. N., Conrad, O., Böhrer, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., ... Kessler,
749 M. (2017). Climatologies at high resolution for the earth's land surface areas. *Scientific*
750 *Data*, 4, 170122. <https://doi.org/10.1038/sdata.2017.122>
- 751 Karimi, S., Moradi, H. V., Rezaei, H. R., Brambilla, M., & Ghadimi, M. (2018). Fine-scale
752 habitat use by black woodpecker *Dryocopus martius*: A year-round study in the
753 Hyrcanian forest, Iran. *North-Western Journal of Zoology*, 14(1), 76–84.
- 754 König, C., Weick, F., & Becking, J. H. (1999). *Owls, a guide to the owls of the world*. Sussex:
755 Pica Press.
- 756 Korpimäki, E. (1986). Niche Relationships and Life-History Tactics of Three Sympatric *Strix*
757 Owl Species in Finland. *Ornis Scandinavica*, 17(2), 126–132.
758 <https://doi.org/10.2307/3676861>
- 759 Korpimäki, E., & Hakkarainen, H. (2012). The Boreal Owl. In *The Boreal Owl*.
760 <https://doi.org/10.1017/cbo9780511844164>
- 761 Kubelka, V., Šálek, M., Tomkovich, P., Végvári, Z., Freckleton, R. P., & Székely, T. (2018).

- 762 Global pattern of nest predation is disrupted by climate change in shorebirds. *Science*
763 (*New York, N.Y.*), 362(6415), 680–683. <https://doi.org/10.1126/science.aat8695>
- 764 Liu, C., Berry, P. M., Dawson, T. P., & Person, R. G. (2005). Selecting Thresholds of
765 Occurrence in the Predictions of Species Distribution. *Ecography*, 28(3), 385–393.
766 Retrieved from <http://onlinelibrary.wiley.com/doi/10.1111/j.0906-7590.2005.03957.x/full>
- 767 Liu, C., White, M., & Newell, G. (2013). Selecting thresholds for the prediction of species
768 occurrence with presence-only data. *Journal of Biogeography*, 40(4).
769 <https://doi.org/10.1111/jbi.12058>
- 770 Lõhmus, A. (2003). Do Ural owls (*Strix uralensis*) suffer from the lack of nest sites in
771 managed forests? *Biological Conservation*, 110(1), 1–9. [https://doi.org/10.1016/S0006-3207\(02\)00167-2](https://doi.org/10.1016/S0006-3207(02)00167-2)
- 772
- 773 Lundberg, A. (1980). Why Are the Ural Owl *Strix uralensis* and the Tawny Owl *S. aluco*
774 Parapatric in Scandinavia? *Ornis Scandinavica*, 11(2), 116.
775 <https://doi.org/10.2307/3675917>
- 776 Marchesi, L., Sergio, F., & Pedrini, P. (2006). Implications of temporal changes in forest
777 dynamics on density, nest-site selection, diet and productivity of Tawny Owls *Strix aluco*
778 in the Alps. *Bird Study*, 53(3), 310–318. <https://doi.org/10.1080/00063650609461447>
- 779 Mihelič, T., Kmecl, P., Denac, K., Koce, U., Vrezec, A., & Denac, D. (2019). *Atlas ptic*
780 *Slovenije. Popis gnezdilk 2002-2017* (T. Mihelič, P. Kmecl, K. Denac, U. Koce, A.
781 Vrezec, & D. Denac, Eds.). Ljubljana: DOPPS.
- 782 Mikkola, H. (1976). Owls killing and killed by other owls and raptors in Europe. *British*
783 *Birds*, 69(April), 144–154.
- 784 Mikkola, H. (1983). Owls of Europe. *A.D. & T. Poyser, Calton*, 397.
785 <https://doi.org/10.1111/j.1467-8691.2007.00421.x>
- 786 Mikusiński, G. (1995). Population trends in black woodpecker in relation to changes and
787 characteristics of European forests. *Ecography*, 18(4), 363–369.
788 <https://doi.org/10.1111/j.1600-0587.1995.tb00139.x>
- 789 Moreno, A., Neumann, M., & Hasenauer, H. (2017). Forest structures across Europe.
790 *Geoscience Data Journal*, 4(1), 17–28. <https://doi.org/10.1002/gdj3.45>
- 791 Muscarella, R., Galante, P. J., Soley-Guardia, M., Boria, R. A., Kass, J. M., Uriarte, M., &
792 Anderson, R. P. (2014). ENMeval: An R package for conducting spatially independent
793 evaluations and estimating optimal model complexity for <scp>Maxent</scp> ecological
794 niche models. *Methods in Ecology and Evolution*, 5(11), 1198–1205.
795 <https://doi.org/10.1111/2041-210X.12261>
- 796 Nardelli, R., Andreotti, A., Bianchi, E., Brambilla, M., Brecciaroli, B., Celada, C., ... Serra, L.
797 (2015). *Rapporto sull'applicazione della Direttiva 147/2009/CE in Italia: dimensione,*
798 *distribuzione e trend delle popolazioni di uccelli (2008-2013)*. Rome: ISPRA, Serie
799 Rapporti, 219/2015.

- 800 Neteler, M., Bowman, M. H., Landa, M., & Metz, M. (2012). GRASS GIS: A multi-purpose
801 open source GIS. *Environmental Modelling & Software*, *31*, 124–130.
802 <https://doi.org/10.1016/j.envsoft.2011.11.014>
- 803 Pandey, R., & Papeş, M. (2018). Changes in future potential distributions of apex predator
804 and mesopredator mammals in North America. *Regional Environmental Change*, *18*(4),
805 1223–1233. <https://doi.org/10.1007/s10113-017-1265-7>
- 806 Pedrini, P., Caldonazzi, M., & Zanghellini, S. (2005). Atlante degli Uccelli nidificanti e
807 svernanti in provincia di Trento. *Studi Trentini Di Scienze Naturali, Acta Biologica*, *80*,
808 suppl. 2.
- 809 Peters, G. P., Andrew, R. M., Boden, T., Canadell, J. G., Ciais, P., Le Quéré, C., ... Wilson, C.
810 (2013). The challenge to keep global warming below 2C. *Nature Climate Change*, *3*(1),
811 4–6. <https://doi.org/10.1038/nclimate1783>
- 812 Phillips, S. J., Anderson, R. P., & Schapire, R. E. (2006). Maximum entropy modeling of
813 species geographic distributions. *Ecological Modelling*, *190*(3), 231–259.
814 <https://doi.org/10.1016/j.ecolmodel.2005.03.026>
- 815 Pirovano, A. R., & Zecca, G. (2014). Black woodpecker dryocopus martius habitat selection
816 in the italian alps: Implications for conservation in Natura 2000 network. *Bird
817 Conservation International*, Vol. 24, pp. 299–315.
818 <https://doi.org/10.1017/S0959270913000439>
- 819 Pollock, L. J., Tingley, R., Morris, W. K., Golding, N., O'Hara, R. B., Parris, K. M., ...
820 Mccarthy, M. A. (2014). Understanding co-occurrence by modelling species
821 simultaneously with a Joint Species Distribution Model (JSDM). *Methods in Ecology
822 and Evolution*, *5*(5). <https://doi.org/10.1111/2041-210X.12180>
- 823 R Development Core Team. (2016). *A Language and Environment for Statistical Computing*.
824 Retrieved from <https://www.r-project.org>
- 825 Rassati, G. (2006). Primi dati su presenza, svernamento e nidificazione dell'Allocco degli
826 Urali *Strix uralensis* nel Friuli settentrionale (Alpi orientali). *Picus*, *32*, 1–3.
- 827 Rassati, G. (2017). Allocco degli Urali *Strix uralensis*: nuovi dati e quadro distributivo
828 italiano. In S. G. Fasano & D. Rubolini (Eds.), *Riassunti del XIX Convegno Italiano di
829 Ornitologia. Torino, 27 settembre - 1 ottobre 2017* (p. 38). Torino: Tichodroma.
830 Monografie del Gruppo Piemontese Studi Ornitologici.
- 831 Roché, J. C., & Chevereau, J. (2000). *Une guide sonore des oiseaux d'Europe et du Maghreb*.
832 Sittelle.
- 833 Rogers, T. L., Gouhier, T. C., & Kimbro, D. L. (2018). Temperature dependency of intraguild
834 predation between native and invasive crabs. *Ecology*, *99*(4), 885–895.
835 <https://doi.org/10.1002/ecy.2157>
- 836 Royle, J. A. (2004). N-Mixture Models for Estimating Population Size from Spatially
837 Replicated Counts. *Biometrics*, *60*(1), 108–115. <https://doi.org/10.1111/j.0006->

- 838 341X.2004.00142.x
- 839 Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients.
840 *Methods in Ecology and Evolution*, 1(2), 103–113. <https://doi.org/10.1111/j.2041->
841 210X.2010.00012.x
- 842 Schweiger, O., Settele, J., Kudrna, O., Klotz, S., & Kühn, I. (2008). Climate change can cause
843 spatial mismatch of trophically interacting species. *Ecology*, 89(12), 3472–3479.
844 <https://doi.org/10.1890/07-1748.1>
- 845 Scridel, D., Bogliani, G., Pedrini, P., Iemma, A., Von Hardenberg, A., & Brambilla, M. (2017).
846 Thermal niche predicts recent changes in range size for bird species. *Climate Research*,
847 73(3), 207–216. <https://doi.org/10.3354/cr01477>
- 848 Scully, A. E., Fisher, S., Miller, D. A. W., & Thornton, D. H. (2018). Influence of biotic
849 interactions on the distribution of Canada lynx (*Lynx canadensis*) at the southern edge of
850 their range. *Journal of Mammalogy*, 99(4), 760–772.
851 <https://doi.org/10.1093/jmammal/gyy053>
- 852 Sergio, F., Marchesi, L., Pedrini, P., & Penteriani, V. (2007). Coexistence of a generalist owl
853 with its intraguild predator: distance-sensitive or habitat-mediated avoidance? *Animal*
854 *Behaviour*, 74(6), 1607–1616. <https://doi.org/10.1016/j.anbehav.2006.10.022>
- 855 Tjernberg, M., Johnsson, K., & Nilsson, S. G. (1993). Density variation and breeding success
856 of the black woodpecker *Dryocopus martius* in relation to forest fragmentation. *Ornis*
857 *Fennica*, 70(3), 155–162.
- 858 Tylianakis, J. M., Didham, R. K., Bascompte, J., & Wardle, D. A. (2008). Global change and
859 species interactions in terrestrial ecosystems. *Ecology Letters*, 11(12), 1351–1363.
860 <https://doi.org/10.1111/j.1461-0248.2008.01250.x>
- 861 Valiente-Banuet, A., Aizen, M. A., Alcántara, J. M., Arroyo, J., Cocucci, A., Galetti, M., ...
862 Zamora, R. (2015). Beyond species loss: The extinction of ecological interactions in a
863 changing world. *Functional Ecology*, 29(3), 299–307. <https://doi.org/10.1111/1365->
864 2435.12356
- 865 Van der Putten, W. H., Macel, M., & Visser, M. E. (2010). Predicting species distribution and
866 abundance responses to climate change: why it is essential to include biotic interactions
867 across trophic levels. *Philosophical Transactions of the Royal Society of London. Series*
868 *B, Biological Sciences*, 365(1549), 2025–2034. <https://doi.org/10.1098/rstb.2010.0037>
- 869 Vos, C. C., Berry, P., Opdam, P., Baveco, H., Nijhof, B., O’Hanley, J., ... Kuipers, H. (2008).
870 Adapting landscapes to climate change: Examples of climate-proof ecosystem networks
871 and priority adaptation zones. *Journal of Applied Ecology*, 45(6), 1722–1731.
872 <https://doi.org/10.1111/j.1365-2664.2008.01569.x>
- 873 Vrezec, A. (2009). Melanism and plumage variation in macroura Ural Owl. *Dutch Birding*,
874 31, 159–170.
- 875 Vrezec, A. (2019). Kozača *Strix uralensis*. In T. Mihelič, P. Kmecl, K. Denac, U. Koce, A.

- 876 Vrezec, & D. Denac (Eds.), *Atlas ptic Slovenije. Popis gnezdilk 2002-2017* (pp. 214–
877 215). Ljubljana: DOPPS.
- 878 Vrezec, A., & Bertoncej, I. (2018). Territory monitoring of Tawny Owls *Strix aluco* using
879 playback calls is a reliable population monitoring method. *Bird Study*, 1–11.
880 <https://doi.org/10.1080/00063657.2018.1522527>
- 881 Vrezec, A., & Mihelič, T. (2013). The Ural Owl, *Strix uralensis macroura*, in Slovenia: an
882 overview of current knowledge on species ecology. *Rivista Italiana Di Ornitologia*, Vol.
883 82, pp. 30–37. <https://doi.org/10.4081/rio.2012.107>
- 884 Vrezec, A., Saurola, P., Avotins, A., Kocijančič, S., & Sulkava, S. (2018). A comparative study
885 of Ural Owl *Strix uralensis* breeding season diet within its European breeding range,
886 derived from nest box monitoring schemes. *Bird Study*, 65(sup1), S85–S95.
887 <https://doi.org/10.1080/00063657.2018.1553026>
- 888 Vrezec, A., & Tome, D. (2004a). Altitudinal segregation between Ural Owl *Strix uralensis* and
889 Tawny Owl *S. aluco*: Evidence for competitive exclusion in raptorial birds. *Bird Study*,
890 Vol. 51, pp. 264–269. <https://doi.org/10.1080/00063650409461362>
- 891 Vrezec, A., & Tome, D. (2004b). Habitat selection and patterns of distribution in a hierarchic
892 forest owl guild. *Ornis Fennica*, 81(3), 109–118.
- 893 Wisz, M. S., Pottier, J., Kissling, W. D., Pellissier, L., Lenoir, J., Damgaard, C. F., ...
894 Svenning, J. C. (2013). The role of biotic interactions in shaping distributions and
895 realised assemblages of species: Implications for species distribution modelling.
896 *Biological Reviews*, 88(1). <https://doi.org/10.1111/j.1469-185X.2012.00235.x>
- 897 Zielinski, W. J., Tucker, J. M., & Rennie, K. M. (2017). Niche overlap of competing
898 carnivores across climatic gradients and the conservation implications of climate change
899 at geographic range margins. *Biological Conservation*, 209, 533–545.
900 <https://doi.org/10.1016/J.BIOCON.2017.03.016>
- 901 Zink, R., & Walter, T. (2018). *Endbericht Habichtskauz Wiederansiedelung - 2015-2018*.
902 Wien.
- 903 Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed effects*
904 *models and extensions in ecology with R*. <https://doi.org/10.1007/978-0-387-87458-6>
905