

SPECTRAL INCLUSION FOR SUBNORMAL n -TUPLES

MIHAI PUTINAR

ABSTRACT. Let S be a subnormal operator on a Hilbert space and let N be its minimal extension. Then a celebrated theorem due to P. Halmos asserts that $\text{Sp}(N) \subset \text{Sp}(S)$, denoting by Sp the spectrum. This note contains a multidimensional version, with respect to Taylor's joint spectrum, of this spectral inclusion theorem.

Recently R. Curto [1] has extended Halmos' spectral inclusion theorem for subnormal operators to the case of n -tuples of doubly commuting subnormal operators. In this note we improve Curto's result by removing the double commutativity assumption.

Let $S = (S_1, \dots, S_n)$ be a subnormal n -tuple of commuting operators on a Hilbert space \mathcal{H} (i.e. there exists a commuting n -tuple of normal operators which extends S). Then there exists a unique, up to isometric isomorphism, minimal extension of S . Let $\text{Sp}(S, \mathcal{H})$ denote Taylor's joint spectrum of S on \mathcal{H} .

THEOREM. *Let S be a commuting subnormal n -tuple on \mathcal{H} and let N be its minimal normal extension to a Hilbert space \mathcal{K} . Then*

$$\text{Sp}(N, \mathcal{K}) \subset \text{Sp}(S, \mathcal{H}).$$

PROOF. It is enough to prove that $0 \notin \text{Sp}(S, \mathcal{H})$ implies $0 \notin \text{Sp}(N, \mathcal{K})$, or equivalently, by a Gelfand transformation argument, that $0 \notin \text{Sp}(|N|, \mathcal{K})$, where $|N|^2 = \sum_{i=1}^n N_i N_i^*$.

Suppose $0 \notin \text{Sp}(S, \mathcal{H})$. Then the operator $S: \mathcal{H}^n \rightarrow \mathcal{H}$ is onto, and after a homothety, one can suppose that

$$(1) \quad \begin{aligned} &(\forall) h \in \mathcal{H}, (\exists) h_1, \dots, h_n \in \mathcal{H} \text{ such that } \sum_{i=1}^n S_i h_i = h, \\ &\text{and } \sum_{i=1}^n \|h_i\|^2 \leq \|h\|^2. \end{aligned}$$

Take the spectral measure E of N and let \mathcal{L} be the space $E(\{z \mid |z| \leq 1/2n\})\mathcal{K}$, which reduces the operators N_i . If we prove that $\mathcal{L} \perp \mathcal{H}$, then, by the minimality of the extension N , \mathcal{L} must be 0, hence $|N|$ will be invertible.

Received by the editors December 17, 1982.

1980 *Mathematics Subject Classification.* Primary 47B20; Secondary 47A10, 47D99.

Key words and phrases. Subnormal operator, commuting n -tuple, Taylor's joint spectrum, minimal normal extension.

©1984 American Mathematical Society
0002-9939/84 \$1.00 + \$.25 per page

Let $l \in \mathcal{L}$ and $h \in \mathcal{H}$. By using (1) a finite number of times, one obtains

$$\begin{aligned}
 |\langle l, h \rangle| &= \left| \left\langle l, \sum_{I \leq i_1, \dots, i_p \leq n} S_{i_1} \cdots S_{i_p} h_{i_1 \dots i_p} \right\rangle \right| \\
 &= \left| \left\langle l, \sum N_{i_1} \cdots N_{i_p} h_{i_1 \dots i_p} \right\rangle \right| \leq \sum \|N_{i_1}^* \cdots N_{i_p}^* l\| \cdot \|h_{i_1 \dots i_p}\| \\
 &\leq \sum \| |N|^p l \| \cdot \|h_{i_1 \dots i_p}\| \leq \|l\| / (2n)^p \sum \|h_{i_1 \dots i_p}\| \\
 &\leq (\|l\| / (2n)^p) \sqrt{n^p} \left(\sum \|h_{i_1 \dots i_p}\|^2 \right)^{1/2} \leq \|l\| \cdot \|h\| (1/2\sqrt{n})^p.
 \end{aligned}$$

By passing to the limit when $p \rightarrow \infty$, $\langle l, h \rangle = 0$, and the proof is complete.

REFERENCES

1. R. E. Curto, *Spectral inclusion for doubly commuting subnormal n -tuples*, Proc. Amer. Math. Soc. **83** (1981), 730–734.
2. P. Halmos, *A Hilbert space problem book*, Van Nostrand, Princeton, N. J., 1967.

DEPARTMENT OF MATHEMATICS, NATIONAL INSTITUTE FOR SCIENTIFIC AND TECHNICAL CREATION (INCREST), Bd. PĂCIII 220, 79622 BUCHAREST, ROMANIA