

 Open access • Proceedings Article • DOI:10.1117/12.2238694

Spin Hall effect in 5d Au: W transition metal alloys(Conference Presentation)

— [Source link](#) 

P. Laczkowski, [Juan-Carlos Rojas-Sánchez](#), [W. Savero-Torres](#), [Nicolas Reyren](#) ...+10 more authors

Institutions: [Centre national de la recherche scientifique](#), [French Alternative Energies and Atomic Energy Commission](#)

Published on: 04 Nov 2016 - [Proceedings of SPIE](#) (International Society for Optics and Photonics)

Topics: [Spin Hall effect](#), [Spin polarization](#), [Quantum spin Hall effect](#), [Spin pumping](#) and [Spintronics](#)

Related papers:

- [Spin pumping and inverse spin Hall effect in platinum and other 5d metals: the essential role of spin-memory loss and spin-current discontinuities at interfaces](#)
- [Reciprocal spin Hall effects in conductors with strong spin-orbit coupling: a review](#)
- [Disentanglement of bulk and interfacial spin Hall effect in ferromagnet/normal metal interface](#)
- [Moderate Positive Spin Hall Angle in Uranium](#)
- [Room-temperature reversible spin Hall effect.](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/spin-hall-effect-in-5d-au-w-transition-metal-alloys-32siwpght>



HAL
open science

Spin Hall effect in AuW alloys

Piotr Laczkowski, Juan Rojas-Sánchez, Yu Fu, Williams Savero-Torres,
Nicolas Reyren, Cyrile Deranlot, Jean-Marie George, Henri Jaffrès, Lucien
Notin, Cyrile Beigné, et al.

► **To cite this version:**

Piotr Laczkowski, Juan Rojas-Sánchez, Yu Fu, Williams Savero-Torres, Nicolas Reyren, et al.. Spin Hall effect in AuW alloys. SPIE 9931, Spintronics IX, Aug 2016, San Diego, United States. pp. 993108, 10.1117/12.2238694 . cea-01511834

HAL Id: cea-01511834

<https://hal-cea.archives-ouvertes.fr/cea-01511834>

Submitted on 21 Apr 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Spin Hall effect in AuW alloys

Piotr Laczkowski^{1,2} *, Juan Carlos Rojas-Sánchez^{1,2}, Yu Fu², Williams Savero-Torres², Nicolas Reyren¹, Cyrile Deranlot¹, Jean-Marie George¹, Henri Jaffrès¹, Lucien Notin², Cyrile Beigné², Alain Marty², Patrick Warin², Jean-Philippe Attané², Laurent Vila² and Albert Fert¹


¹UMR/CNRS-Thales and Université Paris-Sud, 91767, Palaiseau, France

²Institut Nanosciences et Cryogénie, CEA and Université Grenoble Alpes, 38054 Grenoble, France

* piotr.laczkowski@gmail.com

The spin Hall effect (SHE) [1] allows for a reciprocal conversion between charge and spin currents using the spin orbit coupling which can be at the core of several promising spintronics devices. The spin orbit interaction is used to produce a transverse flow of spin or charge in response to a longitudinal excitation, these are the direct or inverse SHE. The spin Hall angle (SHA), the ratio of longitudinal and transverse electronic conductivities, is the characterizing parameter of this conversion. So far, large SHA have been reported in transition metals like Pt, Pd, W, Beta-Ta and in a few alloys with large spin orbit coupling impurities: CuIr, CuBi or CuPb [2].

In this presentation we will report on our study of the SHA in AuW alloys [3] which exhibits a non-monotonic relation with the W concentration. In the regime of diluted alloys the behaviour suggests a dominant side-jump contribution to the spin Hall resistivity, the SHA increasing with the W concentration. We will present experiments demonstrating the requirement of new spin-absorption model in lateral spin valves adapted to the case of strong spin absorption for a correct evaluation of the SHA. Altogether with complementary Ferromagnetic Resonance Spin-Pumping studies, it then leads to SHA as large as +15%. At higher W content the SHA sign is reversed, becoming negative as for pure W.



[1] J.E. Hirsch, *PRL* **83**, 1834 (1999).

[2] Y. Niimi *et al.*, *PRL* **106**, 126601 (2011), *PRL* **109**, 156602 (2012), *PRB* **89**, 054401 (2014).

[3] P. Laczkowski *et al.*, *APL* **104**, 142403 (2014). P L. et al arxiv