

Spontaneous Isotropy Breaking: A Mechanism for CMB Multipole Alignments


Christopher Gordon

University of Chicago (KICP)

Reference: Gordon, C., Hu, W., Huterer, D. and Crawford, T., [astro-ph/0509xx]

Possible Sources of directionality:

- One spatial dimension smaller than horizon (Tegmark, de Oliveira-Costa and Hamilton (2003)).
- One spatial dimension expanding at a different rate to the others (Jaffe, Banday, Eriksen, Gorski and Hansen (2005)).



Say quintessence had a gradient across the horizon in the z direction:

$$Q = Az + B$$

This gets mapped on to a sub-horizon modulation by the potential:

$$\begin{aligned} V &= V_0(1 + f \cos[Q/M_0]) \\ &= V_0(1 + f \cos[k_0 z + \delta]) \end{aligned}$$

Assume the field is light

$$\begin{aligned} \frac{\partial^2 V}{\partial Q^2} \frac{1}{H^2} &< \frac{\partial^2 V}{\partial Q^2} \frac{M_{\text{pl}}^2}{V} \\ &\approx f \left(\frac{M_{\text{pl}}}{M_0} \right)^2 \end{aligned}$$

Thus,

$$\frac{M_0}{M_{\text{pl}}} \gg f^{1/2}.$$

Then the field will be frozen. This gives rise to a sub-horizon density perturbation that has Fourier components

$$\frac{\delta\rho_Q}{\rho_Q}(\mathbf{k}) = \frac{f}{2}e^{i\delta}(2\pi)^3\delta(\mathbf{k} - k_0\hat{\mathbf{x}}_3) + \frac{f}{2}e^{-i\delta}(2\pi)^3\delta(\mathbf{k} + k_0\hat{\mathbf{x}}_3).$$

The curvature perturbation on co-moving hyper-surfaces is given by:

$$\begin{aligned}\zeta &= \zeta_i - \int_0^a \frac{da'}{a'} \frac{\delta p_T}{\rho_T + p_T} \\ &\approx \int_0^a \frac{da'}{a'} \frac{\delta\rho_Q}{\rho_m}.\end{aligned}$$

The matter density red-shifts as

$$\rho_m = \frac{\rho_Q \Omega_m}{a^3 \Omega_Q}.$$

So that

$$\zeta = \frac{a^3 \delta \rho_Q \Omega_Q}{3 \rho_Q \Omega_m}.$$

The Newtonian gravitational potential can be expressed in terms of ζ as

$$\Psi(\mathbf{k}, a) = \zeta - \frac{H}{a} \int_0^a \frac{da'}{H} \left(\zeta - \frac{\delta p_T}{\rho_T + p_T} \right) \approx \zeta - \frac{H}{a} \int_0^a \frac{da'}{H} \left(\zeta + \frac{\delta \rho_Q}{\rho_m} \right).$$

Then,

$$\Psi(\mathbf{k}, a) \approx \psi(2\pi)^3 \delta(\mathbf{k} - k_0 \hat{\mathbf{x}}_3) + \psi^*(2\pi)^3 \delta(\mathbf{k} + k_0 \hat{\mathbf{x}}_3)$$

where

$$\psi = -\frac{1}{3} \frac{\Omega_Q}{\Omega_m} \frac{f}{2} e^{i\delta} \left(a^3 - 4 \frac{H(a)}{a} \int \frac{da'}{H(a')} a'^3 \right).$$

The integrated Sachs Wolfe effect due to the dark energy perturbations is given by

$$\frac{\Delta T(\hat{\mathbf{n}})}{T} = fw(\hat{\mathbf{n}}) = \int 2\Psi'(\mathbf{x} = D\hat{\mathbf{n}}, \eta) d \ln a,$$

The multipole moments are given by

$$fw_\ell \equiv \int d\hat{\mathbf{n}} Y_{\ell m}^*(\hat{\mathbf{n}}) \frac{\Delta T}{T}(\hat{\mathbf{n}}).$$

With the Rayleigh expansion of a plane wave

$$\exp(i\mathbf{k} \cdot \mathbf{x}) = \sum_{\ell m} 4\pi i^\ell j_\ell(kD) Y_{\ell m}^*(\hat{\mathbf{k}}) Y_{\ell m}(\hat{\mathbf{n}})$$

they can be written as

$$fw_\ell = \int d \ln a \int \frac{d^3 k}{(2\pi)^3} 4\pi i^\ell j_\ell(kD) Y_{\ell m}^*(\hat{\mathbf{k}}) 2\Psi'(\mathbf{k}, \ln a)$$

Using the solution for the Newtonian gravitation potential gives

$$w_\ell = -\frac{\Omega_Q}{\Omega_m} s_\ell \sqrt{4\pi(2\ell+1)} \int d\ln a j_\ell(k_0 D) I(a) a^3,$$

$$s_\ell \equiv \cos \delta (-1)^{\ell/2} \delta_\ell^e + \sin \delta (-1)^{(\ell+1)/2} \delta_\ell^o,$$

where $\delta_\ell^e = 1$ if ℓ is even and 0 if ℓ is odd, and vice versa for δ_ℓ^o . Here we have defined

$$I(a) \equiv -\frac{8}{3a^3} \frac{d}{d\ln a} \left[\frac{H(a)}{a} \right] \int \frac{da'}{H(a')} a'^3 - \frac{2}{3}.$$

The spatial modulation projects onto an angular modulation with a weight given by the spherical Bessel function j_ℓ . For a superhorizon fluctuation

$$k_0 D = \frac{k_0}{H_0} H_0 D \sim \frac{k_0}{H_0} \ll 1$$

and so $j_\ell \propto (k_0/H_0)^\ell$.

What exactly needs to be fixed?

To test the quadrupole-octopole alignment, we take the normalized angular momentum (the t statistic of de Oliveira-Costa, Tegmark, Zaldarriaga, and A. Hamilton as generalized by Copi, Huterer, Schwarz, and Starkman.

$$(\Delta\hat{L})_\ell^2 \equiv \frac{\sum_{m=-\ell}^{\ell} m^2 |a_{\ell m}|^2}{\ell^2 \sum_m |a_{\ell m}|^2}.$$

The statistic

$$(\Delta\hat{L})^2 \equiv (\Delta\hat{L})_2^2 + (\Delta\hat{L})_3^2.$$

maximized over direction of the preferred axis captures both the alignment of the quadrupole and octopole and the planar nature of the octopole.

Then

$$\Pr [(\Delta \hat{L})^2 > (\Delta \hat{L})_{\text{WMAP}}^2 | \Lambda\text{CDM}] \approx 0.2\%$$

ℓ	m	$C_\ell^{\text{fid}} (\mu\text{K}^2)$	N_ℓ/C_ℓ	dipole	ΔL_{max}^2
2	0	1233	0.005	0.005	0.005
	1		0.005	0.013	0.003
	2		0.005	0.396	0.406
3	0	577	0.007	0.179	0.315
	1		0.007	0.079	0.003
	2		0.007	0.426	0.029
	3		0.007	2.155	2.560
4	0	322	0.009	3.929	1.641
	1		0.009	0.156	0.939
	2		0.009	0.052	0.508
	3		0.009	0.271	0.401
	4		0.009	0.406	0.180
5	0	202	0.011	0.035	0.014
	1		0.011	0.376	0.811
	2		0.011	0.035	1.254
	3		0.011	6.827	2.777
	4		0.011	0.077	2.769
	5		0.011	0.364	0.078

Multiplicative Modulation

$$T(\hat{\mathbf{n}}) \equiv A(\hat{\mathbf{n}}) + f[1 + w(\hat{\mathbf{n}})]B(\hat{\mathbf{n}}).$$

Multipole decomposition:

$$\begin{aligned} T(\hat{\mathbf{n}}) &= \sum_{lm} t_{lm} Y_{lm}(\hat{\mathbf{n}}), \\ A(\hat{\mathbf{n}}) &= \sum_{lm} a_{lm} Y_{lm}(\hat{\mathbf{n}}), \\ B(\hat{\mathbf{n}}) &= \sum_{lm} b_{lm} Y_{lm}(\hat{\mathbf{n}}), \\ w(\hat{\mathbf{n}}) &\equiv \sum_l w_l Y_{l0}(\hat{\mathbf{n}}). \end{aligned}$$

The assumption of statistical isotropy for the underlying fields A and B requires that their covariance matrices satisfy

$$\begin{aligned}\langle a_{lm}^* a_{l'm'} \rangle &= \delta_{ll'} \delta_{mm'} C_l^{aa}, \\ \langle a_{lm}^* b_{l'm'} \rangle &= \delta_{ll'} \delta_{mm'} C_l^{ab}, \\ \langle b_{lm}^* b_{l'm'} \rangle &= \delta_{ll'} \delta_{mm'} C_l^{bb}.\end{aligned}$$

However statistical isotropy is not preserved in the observed temperature field $T(\hat{\mathbf{n}})$. We then get the convolution

$$t_{lm} = a_{lm} + f b_{lm} + f \sum_{l_1 l_2} R_{lm}^{l_1 l_2} b_{l_2 m}$$

with a coupling matrix written in terms of Wigner 3j symbols

$$R_{lm}^{\ell_1 \ell_2} \equiv (-1)^m \sqrt{\frac{(2\ell+1)(2\ell_1+1)(2\ell_2+1)}{4\pi}} \\ \times \begin{pmatrix} \ell_1 & \ell_2 & \ell \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \ell_1 & \ell_2 & \ell \\ 0 & m & -m \end{pmatrix} w_{\ell_1}.$$

The covariance matrix between the multipole moments then becomes


$$\langle t_{\ell m}^* t_{\ell' m} \rangle = \delta_{\ell \ell'} [C_{\ell}^{aa} + 2fC_{\ell}^{ab} + f^2C_{\ell}^{bb}] \\ + f \sum_{\ell_1} [R_{\ell' m}^{\ell_1 \ell} (C_{\ell}^{ab} + fC_{\ell}^{bb}) + (\ell \leftrightarrow \ell')] \\ + f^2 \sum_{\ell_1 \ell'_1 \ell_2} R_{\ell m}^{\ell_1 \ell_2} R_{\ell'_1 m}^{\ell'_1 \ell_2} C_{\ell_2}^{bb}.$$


Assume,

$$w_l = w_2 \delta_{l2}.$$

and

$$C_l^{aa} = \begin{cases} C_l^{\text{fid}}, & \text{if } l > 3 \\ 0, & \text{if } l = 2 \text{ or } l = 3 \end{cases}$$
$$C_l^{ab} = 0$$
$$C_l^{bb} = \begin{cases} 0, & \text{if } l > 3 \\ C_l^{\text{fid}}, & \text{if } l = 2 \text{ or } l = 3 \end{cases}$$





$$\begin{aligned}
& p\left((\Delta\tilde{L})^2 > (\Delta\tilde{L})^2 | a_{\ell m}\right) \\
&= \int_{w_2, f} p\left((\Delta\tilde{L})^2 > (\Delta\tilde{L})^2, w_2, f | a_{\ell m}\right) dw_2 df \\
&= \int_{w_2, f} p\left((\Delta\tilde{L})^2 > (\Delta\tilde{L})^2 | w_2\right) p(w_2, f | a_{\ell m}) dw_2 df
\end{aligned}$$

Substituting in the previously evaluated quantities we get $p\left((\Delta\tilde{L})^2 > (\Delta\tilde{L})^2 | a_{\ell m}\right) = 0.07$, which is 28 times larger than the value for the fiducial model.

Conclusions

- WMAP data show anomalous alignments between $\ell = 2$ and $\ell = 3$.
- Superhorizon perturbations can lead to a subhorizon modulation.
- Possible in dark energy model but gives wrong modulation.
- Multiplicative modulation is needed.