



# Stability of higher-order nonlinear impulsive differential equations

Shuhong Tang<sup>a</sup>, Akbar Zada<sup>b</sup>, Shah Faisal<sup>b</sup>, M. M. A. El-Sheikh<sup>c</sup>, Tongxing Li<sup>d,e,\*</sup>

<sup>a</sup>*School of Information and Control Engineering, Weifang University, Weifang, Shandong 261061, P. R. China.*

<sup>b</sup>*Department of Mathematics, University of Peshawar, Peshawar 25000, Pakistan.*

<sup>c</sup>*Department of Mathematics, Faculty of Science, Menoufia University, Shebin El-Koom 32511, Egypt.*

<sup>d</sup>*LinDa Institute of Shandong Provincial Key Laboratory of Network Based Intelligent Computing, Linyi University, Linyi, Shandong 276005, P. R. China.*

<sup>e</sup>*School of Informatics, Linyi University, Linyi, Shandong 276005, P. R. China.*

Communicated by M. Bohner

---

## Abstract

For a higher-order nonlinear impulsive ordinary differential equation, we present the concepts of Hyers–Ulam stability, generalized Hyers–Ulam stability, Hyers–Ulam–Rassias stability, and generalized Hyers–Ulam–Rassias stability. Furthermore, we prove the generalized Hyers–Ulam–Rassias stability by using integral inequality of Grönwall type for piecewise continuous functions. These results extend related contributions to the corresponding first-order impulsive ordinary differential equation. Hyers–Ulam stability, generalized Hyers–Ulam stability, and Hyers–Ulam–Rassias stability can be discussed by the same methods. ©2016 All rights reserved.

*Keywords:* Hyers–Ulam stability, generalized Hyers–Ulam stability, Hyers–Ulam–Rassias stability, generalized Hyers–Ulam–Rassias stability, nonlinear impulsive differential equation, higher-order, Grönwall inequality.

*2010 MSC:* 34A37, 34D20.

---

## 1. Introduction

The stability theory is an important branch of the qualitative analysis of differential equations. In particular, for the stability of functional equations, Ulam [25] raised a question: “When can an approximate

---

\*Corresponding author

*Email addresses:* [wfxytang@163.com](mailto:wfxytang@163.com) (Shuhong Tang), [zadababo@yahoo.com](mailto:zadababo@yahoo.com) (Akbar Zada), [shahfaisal8763@gmail.com](mailto:shahfaisal8763@gmail.com) (Shah Faisal), [msheikh\\_1999@yahoo.com](mailto:msheikh_1999@yahoo.com) (M. M. A. El-Sheikh), [litongx2007@163.com](mailto:litongx2007@163.com) (Tongxing Li)

homomorphism from a group  $G_1$  to a metric group  $G_2$  be approximated by an exact homomorphism?"

To solve this question, assuming that  $G_1$  and  $G_2$  are Banach spaces and using a direct method, Hyers [7] brilliantly gave a partial answer. This result was then extended and improved by Aoki [2] and Rassias [22] who weakened the condition for the bound of the norm of Cauchy difference. For further details and discussion, see the monograph by Jung [11].

As far as we know, works by Obloza [17, 18] were among the first contributions dealing with the Hyers–Ulam stability of differential equations. Since then, Hyers–Ulam stability and Hyers–Ulam–Rassias stability of various classes of differential equations and differential operators have been explored by using a wide spectrum of approaches; see, e.g., [1, 4–6, 8–10, 12, 13, 16, 20, 21, 24, 34] and the references cited therein.

In recent years, Hyers–Ulam stability and Hyers–Ulam–Rassias stability of impulsive differential equations have always attracted interest of researchers; see, for instance, [3, 14, 19, 26, 27, 33]. One of the main reasons for this lies in the fact that, as pointed out by Lupulescu and Zada [15], Rogovchenko [23], Wang and Liu [28], Wang and Wu [29], and Wang et al. [30–32], impulsive differential equations arise in a number of applied problems in natural sciences and engineering. Note that the results reported in [3, 14, 19, 26, 27, 33] are concerned with several classes of first-order impulsive differential equations. Thereinto, Wang et al. [26] introduced four Ulam’s type stability (Hyers–Ulam stability, generalized Hyers–Ulam stability, Hyers–Ulam–Rassias stability, and generalized Hyers–Ulam–Rassias stability) concepts for a first-order impulsive ordinary differential equation. So far, to the best of our knowledge, Ulam’s type stability results of higher-order impulsive ordinary differential equations have not been studied yet.

It should be noted that research in this paper was strongly motivated by the recent contributions of Wang et al. [26]. Our principal goal is to analyze the Ulam’s type stability of the higher-order impulsive differential equation

$$\begin{cases} y^{(n)}(t) = F(t, y(t), y'(t), y''(t), \dots, y^{(n-1)}(t)), & t \in I' = I \setminus \{t_1, t_2, \dots, t_m\}, \\ \Delta y^{(i)}(t_k) = y^{(i)}(t_k^+) - y^{(i)}(t_k^-) = \Upsilon_k(y^{(i)}(t_k^-)), & i = 0, 1, \dots, n - 1 \text{ and } k = 1, 2, \dots, m, \\ y(t_0) = y_0, y'(t_0) = y_1, y''(t_0) = y_2, \dots, y^{(n-1)}(t_0) = y_{n-1}, \end{cases} \quad (1.1)$$

where  $n \geq 1$  is a natural number,  $I = [t_0, t_F]$ ,  $t_k$  satisfy  $0 \leq t_0 < t_1 < t_2 < \dots < t_m < t_{m+1} = t_F < +\infty$ ,  $F : B \rightarrow R$  is a continuous function on a closed ball  $B$  in  $I \times R^n$ ,  $\Upsilon_k : R \rightarrow R$  is a continuous function for each  $k$ ,  $y^{(i)}(t_k^+) = \lim_{\tau \rightarrow 0^+} y^{(i)}(t_k + \tau)$  and  $y^{(i)}(t_k^-) = \lim_{\tau \rightarrow 0^+} y^{(i)}(t_k - \tau)$  represent the right-sided and left-sided limits of  $y^{(i)}(t)$  at  $t_k$ , respectively.

## 2. Preliminaries

In this section, we present some definitions of Ulam’s type stability and auxiliary lemmas to prove our main results. Throughout this paper, we use the following spaces:

- $C(I, R)$  is the Banach space of all continuous functions from  $I$  to  $R$  with norm  $\|x\|_C = \sup\{|x(t)| : t \in I\}$ ;
- $PC(I, R)$  denotes the Banach space of all functions  $x : I \rightarrow R$  with norm  $\|x\|_{PC} = \sup\{|x(t)| : t \in I\}$  such that  $x \in C((t_k, t_{k+1}], R)$ ,  $k = 0, 1, \dots, m$  and there exist  $x(t_k^+)$  and  $x(t_k^-)$  satisfying  $x(t_k^-) = x(t_k)$ ,  $k = 1, 2, \dots, m$ ;
- $PC^n(I, R) = \{x : I \rightarrow R \mid x^{(i)} \in PC(I, R), i = 0, 1, \dots, n\}$  is the Banach space with norm  $\|x\|_{PC^n} = \max\{\|x^{(i)}\|_{PC} : i = 0, 1, \dots, n\}$ .

Let  $R^+ = [0, +\infty)$ ,  $\{y\} = (y, y', y'', \dots, y^{(n-1)})$ ,  $\epsilon > 0$ ,  $\mu \geq 0$ , and  $\theta \in PC(I, R^+)$  be nondecreasing. We focus on the following inequalities:

$$\begin{cases} |y^{(n)}(t) - F(t, \{y\})| \leq \epsilon, & t \in I', \\ |\Delta y^{(i)}(t_k) - \Upsilon_k(y^{(i)}(t_k^-))| \leq \epsilon, & i = 0, 1, \dots, n - 1 \text{ and } k = 1, 2, \dots, m, \end{cases} \quad (2.1)$$

$$\begin{cases} |y^{(n)}(t) - F(t, \{y\})| \leq \theta(t), & t \in I', \\ |\Delta y^{(i)}(t_k) - \Upsilon_k(y^{(i)}(t_k^-))| \leq \mu, & i = 0, 1, \dots, n - 1 \text{ and } k = 1, 2, \dots, m, \end{cases} \quad (2.2)$$

and

$$\begin{cases} |y^{(n)}(t) - F(t, \{y\})| \leq \epsilon\theta(t), & t \in I', \\ |\Delta y^{(i)}(t_k) - \Upsilon_k(y^{(i)}(t_k^-))| \leq \epsilon\mu, & i = 0, 1, \dots, n - 1 \text{ and } k = 1, 2, \dots, m. \end{cases} \quad (2.3)$$

In what follows, we introduce the concepts of Ulam’s type stability of (1.1).

**Definition 2.1.** Equation (1.1) is said to be Hyers–Ulam stable on  $I$  if there exists a real number  $K_{F,m} > 0$  such that, for every  $\epsilon > 0$  and for every solution  $y \in PC^n(I, R)$  of (2.1), there exists a solution  $x_0 \in PC^n(I, R)$  of (1.1) with

$$|y(t) - x_0(t)| < K_{F,m}\epsilon, \quad \text{for } t \in I.$$

**Definition 2.2.** Equation (1.1) is called generalized Hyers–Ulam stable on  $I$  if there is a function  $G_{F,m} \in C(R^+, R^+)$  with  $G_{F,m}(0) = 0$  such that, for every  $\epsilon > 0$  and for every solution  $y \in PC^n(I, R)$  of (2.1), there exists a solution  $x_0 \in PC^n(I, R)$  of (1.1) with

$$|y(t) - x_0(t)| < G_{F,m}(\epsilon), \quad \text{for } t \in I.$$

**Definition 2.3.** Equation (1.1) is termed Hyers–Ulam–Rassias stable on  $I$  with respect to  $(\theta, \mu)$  if there exists an  $M_{F,m,\theta} > 0$  such that, for every  $\epsilon > 0$  and for every solution  $y \in PC^n(I, R)$  of (2.3), there exists a solution  $x_0 \in PC^n(I, R)$  of (1.1) with

$$|y(t) - x_0(t)| < M_{F,m,\theta}\epsilon(\theta(t) + \mu), \quad \text{for } t \in I.$$

**Definition 2.4.** Equation (1.1) is said to be generalized Hyers–Ulam–Rassias stable on  $I$  with respect to  $(\theta, \mu)$  if there exists an  $L_{F,m,\theta} > 0$  such that, for every solution  $y \in PC^n(I, R)$  of (2.2), there exists a solution  $x_0 \in PC^n(I, R)$  of (1.1) with

$$|y(t) - x_0(t)| < L_{F,m,\theta}(\theta(t) + \mu), \quad \text{for } t \in I.$$

*Remark 2.5.* Definition 2.1  $\Rightarrow$  Definition 2.2; Definition 2.3  $\Rightarrow$  Definition 2.4; for  $\theta(t) = \mu = 1$ , Definition 2.3  $\Rightarrow$  Definition 2.1.

The following inequality is the well-known integral inequality of Grönwall type for piecewise continuous functions.

**Lemma 2.6.** *If*

$$x(t) \leq a(t) + \int_{t_0}^t b(s)x(s)ds + \sum_{t_0 < t_k < t} \xi_k x(t_k^-)$$

for  $t \geq t_0 \geq 0$ , where  $x, a, b \in PC([t_0, \infty), R^+)$ ,  $a$  is nondecreasing,  $b(t) > 0$ , and  $\xi_k > 0$ , then

$$x(t) \leq a(t) \prod_{t_0 < t_k < t} (1 + \xi_k) \exp\left(\int_{t_0}^t b(s)ds\right)$$

for  $t \geq t_0$ .

*Remark 2.7.* It follows directly from inequality (2.1) that a function  $y \in PC^n(I, R)$  satisfies (2.1) if and only if there is a function  $f \in PC(I, R)$  and a sequence  $f_k^i$  (which depend on  $y$ ) such that  $|f(t)| \leq \epsilon$  for  $t \in I$ ,  $|f_k^i| \leq \epsilon$  for  $i = 0, 1, \dots, n - 1$  and  $k = 1, 2, \dots, m$ , and

$$\begin{cases} y^{(n)}(t) = F(t, \{y\}) + f(t), & t \in I', \\ \Delta y^{(i)}(t_k) = \Upsilon_k(y^{(i)}(t_k^-)) + f_k^i, & i = 0, 1, \dots, n - 1 \text{ and } k = 1, 2, \dots, m. \end{cases}$$

*Remark 2.8.* If  $y \in PC^n(I, R)$  satisfies (2.1), then

$$\left| y^{(n-i)}(t) - \sum_{j=0}^{i-1} \frac{(t-t_0)^j y_{n-i+j}}{j!} - \sum_{j=1}^k \Upsilon_j(y^{(n-i)}(t_j^-)) - \int_{t_0}^t \frac{(t-s)^{i-1}}{(i-1)!} F(s, \{y\}) ds \right| \leq \left( \frac{(t-t_0)^i}{i!} + m \right) \epsilon,$$

where  $t \in I$  and  $i = 1, 2, \dots, n$ .

*Proof.* It follows from Remark 2.7 that, for  $t \in (t_k, t_{k+1}]$ ,

$$\begin{aligned} y^{(n-i)}(t) &= \sum_{j=0}^{i-1} \frac{(t-t_0)^j y_{n-i+j}}{j!} + \sum_{j=1}^k \Upsilon_j(y^{(n-i)}(t_j^-)) + \sum_{j=1}^k f_j^{n-i} + \int_{t_0}^t \frac{(t-s)^{i-1}}{(i-1)!} F(s, \{y\}) ds \\ &\quad + \int_{t_0}^{v_i=t} \dots \int_{t_0}^{v_2} \int_{t_0}^{v_1} f(s) ds. \end{aligned}$$

Therefore, we conclude that

$$\begin{aligned} &\left| y^{(n-i)}(t) - \sum_{j=0}^{i-1} \frac{(t-t_0)^j y_{n-i+j}}{j!} - \sum_{j=1}^k \Upsilon_j(y^{(n-i)}(t_j^-)) - \int_{t_0}^t \frac{(t-s)^{i-1}}{(i-1)!} F(s, \{y\}) ds \right| \\ &\leq \int_{t_0}^{v_i=t} \dots \int_{t_0}^{v_2} \int_{t_0}^{v_1} |f(s)| ds + \sum_{j=1}^k |f_j^{n-i}|, \end{aligned}$$

which implies that

$$\left| y^{(n-i)}(t) - \sum_{j=0}^{i-1} \frac{(t-t_0)^j y_{n-i+j}}{j!} - \sum_{j=1}^k \Upsilon_j(y^{(n-i)}(t_j^-)) - \int_{t_0}^t \frac{(t-s)^{i-1}}{(i-1)!} F(s, \{y\}) ds \right| \leq \left( \frac{(t-t_0)^i}{i!} + m \right) \epsilon.$$

The proof is complete. □

*Remark 2.9.* One can obtain similar remarks for solutions of inequalities (2.2) and (2.3). The details are left to the reader.

### 3. Main results

Define a closed ball  $B = I \times \prod_{i=0}^{n-1} [-M_i, M_i]$ , where  $M_i = \|y^{(i)}\|_{PC}$ . In this section, we prove the Ulam’s type stability of (1.1) with the condition

$$(t-t_0)^{n-1} |F(t, \{y\}) - F(t, \{z\})| \leq h(t) |y(t) - z(t)|, \tag{3.1}$$

where  $h : I \rightarrow R^+$  is an integrable function and the Lipschitz condition

$$|F(t, \{y\}) - F(t, \{z\})| \leq S_0 \sum_{i=0}^{n-1} |y^{(i)}(t) - z^{(i)}(t)|, \quad S_0 > 0 \text{ is a constant}, \tag{3.2}$$

respectively.

**Theorem 3.1.** *If*

(H<sub>1</sub>)  $F$  satisfies condition (3.1);

(H<sub>2</sub>)  $\Upsilon_k : R \rightarrow R$  and there exist constants  $M_k > 0$  such that  $|\Upsilon_k(x_1) - \Upsilon_k(x_2)| \leq M_k |x_1 - x_2|$  for  $k = 1, 2, \dots, m$  and  $x_1, x_2 \in R$ ;

(H<sub>3</sub>) there exists a nondecreasing function  $\theta \in PC(I, R^+)$  such that, for  $t \in I$  and for some  $\rho_\theta > 0$ ,

$$\int_{t_0}^t \theta(s) ds \leq \rho_\theta \theta(t),$$

then (1.1) has generalized Hyers–Ulam–Rassias stability on  $I$  with respect to  $(\theta, \mu)$ . If, in addition,

$$\frac{1}{(n-1)!} \int_{t_0}^{t_F} h(s) ds + \sum_{j=1}^m M_j < 1,$$

then (1.1) has a unique solution in  $PC^n(I, R) \cap PC(I, R)$ .

*Proof.* Let  $y \in PC^n(I, R)$  be a solution to (2.2). The exact solution  $x \in PC^n(I, R)$  of the initial value problem

$$\begin{cases} x^{(n)}(t) = F(t, \{x\}), & t \in I', \\ \Delta x^{(i)}(t_k) = \Upsilon_k(x^{(i)}(t_k^-)), & i = 0, 1, \dots, n-1 \text{ and } k = 1, 2, \dots, m, \\ x(t_0) = y_0, x'(t_0) = y_1, x''(t_0) = y_2, \dots, x^{(n-1)}(t_0) = y_{n-1}, \end{cases}$$

is given by

$$x(t) = \begin{cases} \sum_{j=0}^{n-1} \frac{(t-t_0)^j y_j}{j!} + \frac{1}{(n-1)!} \int_{t_0}^t (t-s)^{n-1} F(s, \{x\}) ds, & t \in [t_0, t_1], \\ \sum_{j=0}^{n-1} \frac{(t-t_0)^j y_j}{j!} + \Upsilon_1(x(t_1^-)) + \frac{1}{(n-1)!} \int_{t_0}^t (t-s)^{n-1} F(s, \{x\}) ds, & t \in (t_1, t_2], \\ \sum_{j=0}^{n-1} \frac{(t-t_0)^j y_j}{j!} + \sum_{j=1}^2 \Upsilon_j(x(t_j^-)) + \frac{1}{(n-1)!} \int_{t_0}^t (t-s)^{n-1} F(s, \{x\}) ds, & t \in (t_2, t_3], \\ \cdot \\ \cdot \\ \cdot \\ \sum_{j=0}^{n-1} \frac{(t-t_0)^j y_j}{j!} + \sum_{j=1}^m \Upsilon_j(x(t_j^-)) + \frac{1}{(n-1)!} \int_{t_0}^t (t-s)^{n-1} F(s, \{x\}) ds, & t \in (t_m, t_F]. \end{cases}$$

Similar as in Remark 2.8, an application of inequality (2.2) implies that, for  $t \in I$ ,

$$\left| y(t) - \sum_{j=0}^{n-1} \frac{(t-t_0)^j y_j}{j!} - \sum_{j=1}^k \Upsilon_j(y(t_j^-)) - \int_{t_0}^t \frac{(t-s)^{n-1}}{(n-1)!} F(s, \{y\}) ds \right| \leq (m + \rho_\theta^n)(\theta(t) + \mu).$$

Hence, for  $t \in (t_k, t_{k+1}]$ ,

$$\begin{aligned} |y(t) - x(t)| &\leq \left| y(t) - \sum_{j=0}^{n-1} \frac{(t-t_0)^j y_j}{j!} - \sum_{j=1}^k \Upsilon_j(y(t_j^-)) - \frac{1}{(n-1)!} \int_{t_0}^t (t-s)^{n-1} F(s, \{y\}) ds \right| \\ &\quad + \frac{1}{(n-1)!} \int_{t_0}^t (t-s)^{n-1} |F(s, \{y\}) - F(s, \{x\})| ds + \sum_{j=1}^k |\Upsilon_j(y(t_j^-)) - \Upsilon_j(x(t_j^-))| \\ &\leq (m + \rho_\theta^n)(\theta(t) + \mu) + \frac{1}{(n-1)!} \int_{t_0}^t h(s) |y(s) - x(s)| ds + \sum_{j=1}^k M_j |y(t_j^-) - x(t_j^-)|. \end{aligned}$$

By virtue of Lemma 2.6, we conclude that, for  $t \in I$ ,

$$|y(t) - x(t)| \leq (m + \rho_\theta^n)(\theta(t) + \mu) \prod_{t_0 < t_k < t} (1 + M_k) \exp\left(\frac{1}{(n-1)!} \int_{t_0}^t h(s) ds\right) \leq L_{F,m,\theta}(\theta(t) + \mu),$$

where

$$L_{F,m,\theta} = (m + \rho_\theta^n) \prod_{k=1}^m (1 + M_k) \exp\left(\frac{1}{(n-1)!} \int_{t_0}^{t_F} h(s) ds\right).$$

Therefore, (1.1) is generalized Hyers–Ulam–Rassias stable on  $I$  with respect to  $(\theta, \mu)$ .

**Uniqueness of solution.** For  $g \in PC(I, R)$ , define an operator  $\Lambda : PC(I, R) \rightarrow PC(I, R)$  by

$$(\Lambda g)(t) = \begin{cases} \sum_{j=0}^{n-1} \frac{(t-t_0)^j y_j}{j!} + \frac{1}{(n-1)!} \int_{t_0}^t (t-s)^{n-1} F(s, \{g\}) ds, & t \in [t_0, t_1], \\ \sum_{j=0}^{n-1} \frac{(t-t_0)^j y_j}{j!} + \Upsilon_1(g(t_1^-)) + \frac{1}{(n-1)!} \int_{t_0}^t (t-s)^{n-1} F(s, \{g\}) ds, & t \in (t_1, t_2], \\ \sum_{j=0}^{n-1} \frac{(t-t_0)^j y_j}{j!} + \sum_{j=1}^2 \Upsilon_j(g(t_j^-)) + \frac{1}{(n-1)!} \int_{t_0}^t (t-s)^{n-1} F(s, \{g\}) ds, & t \in (t_2, t_3], \\ \cdot \\ \cdot \\ \cdot \\ \sum_{j=0}^{n-1} \frac{(t-t_0)^j y_j}{j!} + \sum_{j=1}^m \Upsilon_j(g(t_j^-)) + \frac{1}{(n-1)!} \int_{t_0}^t (t-s)^{n-1} F(s, \{g\}) ds, & t \in (t_m, t_F]. \end{cases}$$

Clearly,  $\Lambda$  is well-defined. We show that  $\Lambda$  is a Picard operator on  $PC(I, R)$ . For this, let  $g_1, g_2 \in PC(I, R)$  and consider

$$\begin{aligned} |(\Lambda g_1)(t) - (\Lambda g_2)(t)| &= \left| \sum_{j=1}^k (\Upsilon_j(g_1(t_j^-)) - \Upsilon_j(g_2(t_j^-))) \right. \\ &\quad \left. + \frac{1}{(n-1)!} \int_{t_0}^t (t-s)^{n-1} (F(s, \{g_1\}) - F(s, \{g_2\})) ds \right| \\ &\leq \frac{1}{(n-1)!} \int_{t_0}^t (t-s)^{n-1} |F(s, \{g_1\}) - F(s, \{g_2\})| ds \\ &\quad + \sum_{j=1}^k |\Upsilon_j(g_1(t_j^-)) - \Upsilon_j(g_2(t_j^-))| \\ &\leq \frac{1}{(n-1)!} \int_{t_0}^t h(s) |g_1(s) - g_2(s)| ds + \sum_{j=1}^k M_j |g_1(t_j^-) - g_2(t_j^-)| \\ &\leq \left( \frac{1}{(n-1)!} \int_{t_0}^{t_F} h(s) ds + \sum_{j=1}^m M_j \right) \|g_1 - g_2\|_{PC}. \end{aligned}$$

Then,  $\Lambda$  is contractive with respect to  $\|\cdot\|_{PC}$ . By virtue of Banach contraction principle,  $\Lambda$  is a Picard operator. The unique fixed point of this operator is the unique solution of (1.1) in  $PC^n(I, R) \cap PC(I, R)$ . This completes the proof.  $\square$

*Remark 3.2.* Theorem 3.1 includes [26, Theorem 4.1] in the case where  $n = 1$  and  $h(t) = h_0 > 0$ .

**Theorem 3.3.** Assume that  $(H_2)$  and  $(H_3)$  are satisfied. If  $F$  satisfies condition (3.2), then (1.1) has generalized Hyers–Ulam–Rassias stability on  $I$  with respect to  $(\theta, \mu)$ .

*Proof.* Let  $y \in PC^n(I, R)$  be a solution to (2.2). For  $i = 1, 2, \dots, n$ , define a function by

$$x^{(n-i)}(t) = \begin{cases} \sum_{j=0}^{i-1} \frac{(t-t_0)^j y_{n-i+j}}{j!} + \frac{1}{(i-1)!} \int_{t_0}^t (t-s)^{i-1} F(s, \{x\}) ds, & t \in [t_0, t_1], \\ \sum_{j=0}^{i-1} \frac{(t-t_0)^j y_{n-i+j}}{j!} + \Upsilon_1(x(t_1^-)) + \frac{1}{(i-1)!} \int_{t_0}^t (t-s)^{i-1} F(s, \{x\}) ds, & t \in (t_1, t_2], \\ \sum_{j=0}^{i-1} \frac{(t-t_0)^j y_{n-i+j}}{j!} + \sum_{j=1}^2 \Upsilon_j(x(t_j^-)) + \frac{1}{(i-1)!} \int_{t_0}^t (t-s)^{i-1} F(s, \{x\}) ds, & t \in (t_2, t_3], \\ \cdot \\ \cdot \\ \cdot \\ \sum_{j=0}^{i-1} \frac{(t-t_0)^j y_{n-i+j}}{j!} + \sum_{j=1}^m \Upsilon_j(x(t_j^-)) + \frac{1}{(i-1)!} \int_{t_0}^t (t-s)^{i-1} F(s, \{x\}) ds, & t \in (t_m, t_F]. \end{cases}$$

Then, for  $t \in (t_k, t_{k+1}]$ ,

$$\begin{aligned} |y^{(n-i)}(t) - x^{(n-i)}(t)| &\leq \left| y^{(n-i)}(t) - \sum_{j=0}^{i-1} \frac{(t-t_0)^j y_{n-i+j}}{j!} - \sum_{j=1}^k \Upsilon_j(y^{(n-i)}(t_j^-)) \right. \\ &\quad \left. - \frac{1}{(i-1)!} \int_{t_0}^t (t-s)^{i-1} F(s, \{y\}) ds \right| \\ &\quad + \frac{1}{(i-1)!} \int_{t_0}^t (t-s)^{i-1} |F(s, \{y\}) - F(s, \{x\})| ds \\ &\quad + \sum_{j=1}^k |\Upsilon_j(y^{(n-i)}(t_j^-)) - \Upsilon_j(x^{(n-i)}(t_j^-))| \\ &\leq (m + \rho_\theta^n)(\theta(t) + \mu) + \frac{nS_0}{(i-1)!} \int_{t_0}^t (t-s)^{i-1} |y^{(n-i)}(s) - x^{(n-i)}(s)| ds \\ &\quad + \sum_{j=1}^k M_j |y^{(n-i)}(t_j^-) - x^{(n-i)}(t_j^-)|. \end{aligned}$$

Using Lemma 2.6, we deduce that, for  $i = 1, 2, \dots, n$  and  $t \in I$ ,

$$|y^{(n-i)}(t) - x^{(n-i)}(t)| \leq (m + \rho_\theta^n)(\theta(t) + \mu) \prod_{t_0 < t_k < t} (1 + M_k) \exp\left(\frac{S_0(t-t_0)^n}{(n-1)!}\right) \leq L_{F,m,\theta}(\theta(t) + \mu),$$

where

$$L_{F,m,\theta} = (m + \rho_\theta^n) \prod_{k=1}^m (1 + M_k) \exp\left(\frac{S_0(t_F - t_0)^n}{(n-1)!}\right).$$

Hence, (1.1) is generalized Hyers–Ulam–Rassias stable on  $I$  with respect to  $(\theta, \mu)$ . The proof is complete.  $\square$

*Remark 3.4.* Theorem 3.3 contains [26, Theorem 4.1] in the case when  $n = 1$ .

*Remark 3.5.* One can proceed as the same way to prove the Hyers–Ulam stability, generalized Hyers–Ulam stability, and Hyers–Ulam–Rassias stability of (1.1). The details are left to the reader.

## Acknowledgment

This research is supported by NNSF of P. R. China (Grant Nos. 61503171, 61403061, and 11447005), CPSF (Grant No. 2015M582091), NSF of Shandong Province (Grant No. ZR2012FL06), DSRF of Linyi University (Grant No. LYDX2015BS001), and the AMEP of Linyi University, P. R. China.

## References

- [1] C. Alsina, R. Ger, *On some inequalities and stability results related to the exponential function*, J. Inequal. Appl., **2** (1998), 373–380. 1
- [2] T. Aoki, *On the stability of the linear transformation in Banach spaces*, J. Math. Soc. Japan, **2** (1950), 64–66. 1
- [3] M. Gowrisankar, P. Mohankumar, A. Vinodkumar, *Stability results of random impulsive semilinear differential equations*, Acta Math. Sci. Ser. B Engl. Ed., **34** (2014), 1055–1071. 1
- [4] J. Huang, Q. H. Alqifiary, Y. Li, *Superstability of differential equations with boundary conditions*, Electron. J. Differential Equations, **2014** (2014), 8 pages. 1
- [5] J. Huang, S.-M. Jung, Y. Li, *On Hyers–Ulam stability of nonlinear differential equations*, Bull. Korean Math. Soc., **52** (2015), 685–697.
- [6] J. Huang, Y. Li, *Hyers–Ulam stability of linear functional differential equations*, J. Math. Anal. Appl., **426** (2015), 1192–1200. 1
- [7] D. H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. USA, **27** (1941), 222–224. 1
- [8] S.-M. Jung, *Hyers–Ulam stability of linear differential equations of first order*, Appl. Math. Lett., **17** (2004), 1135–1140. 1
- [9] S.-M. Jung, *Hyers–Ulam stability of linear differential equations of first order, III*, J. Math. Anal. Appl., **311** (2005), 139–146.
- [10] S.-M. Jung, *Hyers–Ulam stability of linear differential equations of first order, II*, Appl. Math. Lett., **19** (2006), 854–858. 1
- [11] S.-M. Jung, *Hyers–Ulam–Rassias Stability of Functional Equations in Nonlinear Analysis*, Springer Optim. Appl., New York, (2011). 1
- [12] Y. Li, Y. Shen, *Hyers–Ulam stability of nonhomogeneous linear differential equations of second order*, Int. J. Math. Math. Sci., **2009** (2009), 7 pages. 1
- [13] Y. Li, Y. Shen, *Hyers–Ulam stability of linear differential equations of second order*, Appl. Math. Lett., **23** (2010), 306–309. 1
- [14] Y. Liao, J. R. Wang, *A note on stability of impulsive differential equations*, Bound. Value Probl., **2014** (2014), 8 pages. 1
- [15] V. Lupulescu, A. Zada, *Linear impulsive dynamic systems on time scales*, Electron. J. Qual. Theory Differ. Equ., **2010** (2010), 30 pages. 1
- [16] T. Miura, S. Miyajima, S.-E. Takahasi, *A characterization of Hyers–Ulam stability of first order linear differential operators*, J. Math. Anal. Appl., **286** (2003), 136–146. 1
- [17] M. Obloza, *Hyers stability of the linear differential equation*, Rocznik Nauk.-Dydakt. Prace Mat., **13** (1993), 259–270. 1
- [18] M. Obloza, *Connections between Hyers and Lyapunov stability of the ordinary differential equations*, Rocznik Nauk.-Dydakt. Prace Mat., **14** (1997), 141–146. 1
- [19] C. Parthasarathy, *Existence and Hyers–Ulam stability of nonlinear impulsive differential equations with nonlocal conditions*, Electron. J. Math. Anal. Appl., **4** (2016), 106–115. 1
- [20] D. Popa, I. Raşa, *On the Hyers–Ulam stability of the linear differential equation*, J. Math. Anal. Appl., **381** (2011), 530–537. 1
- [21] D. Popa, I. Raşa, *Hyers–Ulam stability of the linear differential operator with nonconstant coefficients*, Appl. Math. Comput., **219** (2012), 1562–1568. 1
- [22] Th. M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc., **72** (1978), 297–300. 1
- [23] Yu. V. Rogovchenko, *Impulsive evolution systems: main results and new trends*, Dynam. Contin. Discrete Impuls. Systems, **3** (1997), 57–88. 1
- [24] S.-E. Takahasi, T. Miura, S. Miyajima, *On the Hyers–Ulam stability of the Banach space-valued differential equation  $y' = \lambda y$* , Bull. Korean Math. Soc., **39** (2002), 309–315. 1
- [25] S. M. Ulam, *A Collection of Mathematical Problems*, Interscience Publishers, New York-London, (1960). 1
- [26] J. R. Wang, M. Fečkan, Y. Zhou, *Ulam’s type stability of impulsive ordinary differential equations*, J. Math. Anal. Appl., **395** (2012), 258–264. 1, 3.2, 3.4
- [27] J. R. Wang, M. Fečkan, Y. Zhou, *On the stability of first order impulsive evolution equations*, Opuscula Math., **34** (2014), 639–657. 1
- [28] P. Wang, X. Liu,  *$\phi_0$ -Stability of hybrid impulsive dynamic systems on time scales*, J. Math. Anal. Appl., **334** (2007), 1220–1231. 1



- [29] P. Wang, M. Wu, *Practical  $\phi_0$ -stability of impulsive dynamic systems on time scales*, Appl. Math. Lett., **20** (2007), 651–658. 1
- [30] L. Wang, B. Yang, A. Abraham, *Distilling middle-age cement hydration kinetics from observed data using phased hybrid evolution*, Soft Comput., **20** (2016), 3637–3656. 1
- [31] L. Wang, B. Yang, Y. Chen, X. Zhang, J. Orchard, *Improving neural-network classifiers using nearest neighbor partitioning*, IEEE Trans. Neural Netw. Learn. Syst., (In press).
- [32] L. Wang, B. Yang, J. Orchard, *Particle swarm optimization using dynamic tournament topology*, Appl. Soft Comput., **48** (2016), 584–596. 1
- [33] A. Zada, S. Faisal, Y. Li, *On the Hyers–Ulam stability of first-order impulsive delay differential equations*, J. Funct. Spaces, **2016** (2016), 6 pages. 1
- [34] A. Zada, O. Shah, R. Shah, *Hyers–Ulam stability of non-autonomous systems in terms of boundedness of Cauchy problems*, Appl. Math. Comput., **271** (2015), 512–518. 1