

Staying cold through dinner: cold-climate bats rewarm with conspecifics but not sunset during hibernation

Zenon J. Czenze · Andrew D. Park ·
Craig K. R. Willis

Received: 10 October 2012 / Revised: 7 March 2013 / Accepted: 10 March 2013
© Springer-Verlag Berlin Heidelberg 2013

Abstract For temperate endotherms (i.e., mammals and birds) energy costs are highest during winter but food availability is lowest and many mammals depend on hibernation as a result. Hibernation is made up of energy-saving torpor bouts [periods of controlled reduction in body temperature (T_b)], which are interrupted by brief periodic arousals to normothermic T_b . What triggers these arousals in free-ranging hibernators is not well understood. Some temperate bats with intermittent access to flying insects during winter synchronize arousals with sunset, which suggests that, in some species, feeding opportunities influence arousal timing. We tested whether hibernating bats from a cold climate without access to food during winter also maintain a circadian rhythm for arousals or whether cues from conspecifics in the same cluster are more important. We used temperature telemetry to monitor skin temperature (T_{sk}) of free-ranging little brown bats (*Myotis lucifugus*) hibernating in central Manitoba, Canada, where temperatures from 22 October to 22 March were too cold for flying insects. We found no evidence bats synchronized arousals with photoperiod but they did arouse synchronously with other bats in the same cluster. Thus, in the northern part of their range where flying insects are almost never available during winter, little brown bats exhibit no circadian pattern to arousals. Warming synchronously with others could reduce the energetic costs of arousal for individuals or could reflect disturbance of torpid bats by cluster-mates.

Keywords Periodic arousal · Circadian rhythms · Social thermoregulation · Chiroptera · White-nose syndrome

Introduction

Many temperate animals have evolved physiological and behavioural mechanisms to survive the challenging winter months. Hibernating mammals, for example, spend the majority of their time during winter in torpor, a state of controlled reduction in body temperature (T_b) and metabolic rate (Geiser 2004). Long-term torpor provides enormous energetic benefits (Geiser 2004), and may provide additional benefits for some species by reducing exposure to predators and extending lifespan (Wilkinson and South 2002; Stawski and Geiser 2009). However, it also appears to result in physiological and ecological costs which limit the duration of torpor bouts. For example, the inability to drink while torpid results in dehydration stress (Thomas and Cloutier 1992; Thomas and Geiser 1997) and immune responses of torpid hibernators may be suppressed increasing susceptibility to pathogens (Luis and Hudson 2006; Moore et al. 2011; Prendergast et al. 2002). Presumably as a result of these costs of torpor, all hibernators exhibit periodic arousals throughout hibernation, during which they return to a normothermic T_b (e.g., Grahn et al. 1994; Jonasson and Willis 2012; Ruby et al. 2002).

Although periodic arousals are brief they account for up to 90 % of winter energy expenditure due to the high metabolic cost of thermoregulation at low ambient temperatures (T_a) (Geiser 2004; Thomas et al. 1990; Wang 1978). Hibernators like temperate zone bats, which cannot store a food cache, must survive exclusively on stored fat which places a limit to how many arousals they can perform (Thomas et al. 1990). As a result, they must balance

Communicated by G. Heldmaier.

Z. J. Czenze · A. D. Park · C. K. R. Willis (✉)
Department of Biology and Centre for Forest-Interdisciplinary
Research, University of Winnipeg, Winnipeg, MB, Canada
e-mail: c.willis@uwinnipeg.ca

the energetic, physiological and ecological costs and benefits of torpor and arousal. Presumably, the frequency of arousals must be timed carefully to offset the risk of exhausting fat stores prematurely (Humphries et al. 2003; Jonasson and Willis 2011, 2012; Racey and Swift 1981).

The mechanism that initiates periodic arousals is still unknown but one of several hypotheses is that circadian clocks may play a role (e.g., Malan 2010). The T_b of hibernating golden-mantled ground squirrels (*Callospermophilus lateralis*) fluctuates with circadian rhythms during deep torpor and individuals of this species are more likely to enter torpor at night and arouse during the day when exposed to a light/dark cycle (Grahn et al. 1994; Ruby et al. 2002). However, Ruby et al. (2002) found that the timing of entry into and arousal from torpor occurred at random times for animals housed under 24 h lights-on photoperiod. Free-ranging, cave-dwelling greater horseshoe bats (*Rhinolophus ferrumequinum*) and Natterer's bats (*Myotis nattereri*) from the United Kingdom (UK) receive little or no natural photoperiod cues in caves during winter. In spite of the lack of photoperiod, however, these species appear to maintain a circadian pattern to arousals (Hope and Jones 2012; Park et al. 2000). Hibernating bats of these species “warm up for dinner”, synchronizing arousals with sunset, likely to take advantage of flying insects during occasional mild winter nights (Hope and Jones 2012; Park et al. 2000).

Greater horseshoe and Natterer's bats hibernate in relatively mild climates where temperatures during winter occasionally exceed 10 °C. This is a threshold above which aerial insect abundance can increase dramatically (Jones et al. 1995), providing opportunities to forage. Even though foraging opportunities will be less common, bats from colder environments might also synchronize arousals with sunset. To our knowledge, the maintenance costs of neural mechanisms underlying a circadian clock are unknown. However, it seems likely these costs would be small relative to the benefits of even unlikely potential foraging opportunities, especially in mid- to late-winter when fat reserves are dwindling. Thus, bats might still synchronize arousals with sunset, as selection against arousal at dusk might be relatively weak.

Another trigger of arousal, specific to communal hibernators like bats, could be cues from conspecifics. Such cues could be beneficial if groups of individuals can reduce energetic costs of arousal via social thermoregulation (Ruf and Arnold 2000). Alpine marmots (*Marmota marmota*), for example, hibernate in groups of related individuals and synchronize arousals with burrow mates to save energy. These benefits presumably increase over-winter survival, particularly for young-of-the-year (YOY), and may enhance inclusive fitness via kin selection (Ruf and Arnold 2000). On the other hand, if arousals of different

individuals are not well synchronized, disturbance by normothermic cluster-mates could be detrimental to energy balance for some animals if it induces arousals more frequently than required to offset costs of torpor. For example, in bats, males may attempt to mate with torpid females during hibernation and these mating attempts may disturb females.

In contrast to most mammalian hibernators which spend the winter in well-insulated burrows, bats hibernate in caves and mines often in large, open-air chambers. These environments may reduce the potential for bats to use metabolic heat to influence their immediate microclimate and could explain the energetically conservative hibernation patterns of little brown bats (*Myotis lucifugus*), characterized by long torpor bouts, brief arousals and the expression of shallow heterothermy in the midst of arousals (Jonasson and Willis 2012). Thus, benefits of social thermoregulation for hibernating bats may be especially important. An individual-based population model, incorporating presumed thermoregulatory benefits of clustering for little brown bats predicted that 96 % of clustering bats should survive a typical winter compared to 73 % for solitary bats (Boyles and Brack 2009). This suggests that bats from the same cluster should exploit the energetic benefits of social thermoregulation and arouse synchronously with other individuals in the cluster.

We used temperature radio-telemetry to test two hypotheses about periodic arousals in hibernating bats. First, we tested whether arousal and photoperiod are synchronized in bats from a cold climate. We predicted that arousals would remain synchronized with sunset based on the hypothesis that maintaining neural mechanisms underlying a circadian clock would be inexpensive relative to potential benefits of obtaining even an unlikely meal during winter. Second, we tested whether individual bats would take advantage of social thermoregulation to reduce arousal costs. We predicted that individuals in the same cluster would arouse from torpor at the same time and that a majority of arousals would overlap in time. Alternatively, if bats tend to arouse from torpor on a similar timetable (e.g., on roughly the same days) but not at precisely the same times, this would suggest that disturbance by conspecifics could be influencing torpor-arousal cycles. Such a pattern might be observed if, for example, bats gradually disturb each other as they rewarm, resulting in a cascade of arousal within a cluster.

Methods

This study was conducted in Firecamp cave, near the town of Grand Rapids, in central Manitoba Canada (53°29'25"N 99°20'50"W). This cave is accessed by a sinkhole entrance

approximately 50 cm in diameter and consists of a bell-shaped chamber about 7 m deep. Between 30 and 60 bats over-winter in Firecamp cave from mid-September until mid-May. Bats in this cave all hibernate in one tight cluster within a crack 1 m long by 20 cm deep, on the ceiling of the main chamber.

All procedures were approved by the University of Winnipeg Animal Care Committee and by Manitoba Conservation Wildlife Scientific Permit WB0612. We entered the cave and handled bats during hibernation but to minimize disturbance we only entered once and limited the duration of our visit to less than 90 min. Twenty-four torpid bats were captured by hand on 20 Feb 2012, sexed, weighed to the nearest 0.01 g (Durascale-100, My Weigh, ON, Canada) and their forearm length was measured to the nearest 0.05 mm. We identified YOY based on unfused 3rd digit metacarpal-phalangeal joints, and tapered rather than “knobby” joints following Davis and Hitchcock (1965). In many parts of their range YOY little brown bats cannot be distinguished from adults by the time hibernation begins but in our northern study area we can readily identify at least some juveniles during swarming and even into hibernation (Jonasson and Willis 2011, 2012).

For small bats, skin temperature (T_{sk}) provides a good approximation of T_b during torpor (Audet and Thomas 1996; Barclay et al. 1996; Willis and Brigham 2003). We recorded T_{sk} by attaching temperature-sensitive radio-transmitters (0.64–0.80 g, BD-2NT; Holohil Systems Ltd, Carp, Ontario) using a non-toxic, latex-based adhesive (Osto-Bond, Montreal Ostomy, Vaudreuil, QC, Canada) after clipping a small (<1 cm²) patch of fur between the shoulders. Transmitters represented between 6.9 and 8.6 % of body mass, greater than the 5 % guideline suggested by Aldridge and Brigham (1988) but similar to the size which Jonasson and Willis (2012) demonstrated had no negative impact on body condition in hibernating little brown bats. We calibrated transmitters prior to use in a water bath at 5 °C increments from just above 0–45 °C against a thermometer traceable to the National Institutes of Standards and Technology. A data logging receiver (Lotek SRX600; Lotek Engineering Inc., Newmarket, Ontario, Canada) powered by a solar panel was deployed outside the cave and connected to an omnidirectional whip antenna positioned in the cave, within 60 cm of the cluster of bats. The receiver recorded T_{sk} of each individual at 6-min intervals. Each bat was recorded for 15 s and a full cycle of recordings took 360 s. We recorded T_a inside the cave using iButtons (DS1922L iButtons, Maxim Integrated Products, Dallas, TX, USA) coated in paraffin wax to dampen ultrasonic noise (Willis et al. 2009) and positioned at the base of the crack within 60 cm of where the bats roosted while hibernating.

We defined the phases of arousal following Jonasson and Willis (2012). Due to the large difference between normothermic T_{sk} and T_a in the cave, arousals were always readily obvious in the T_{sk} trace. Torpor bouts were defined as periods of reduced stable T_{sk} between the obvious warming and cooling phases associated with each arousal. Skin temperature measured using Holohil transmitters occasionally appears to increase slightly and gradually after initial entry into a torpor bout (e.g., Jonasson and Willis 2012). This appears to reflect an artifact of long periods at low temperature, which we could readily identify in our T_{sk} traces and which did not affect our analysis.

Warming was defined by an abrupt increase in torpid T_{sk} that resulted in an elevation of T_{sk} until it stabilized at a normothermic steady-state. Cold T_a in the cave appeared to increase ambient cooling of external transmitters, leading to slightly reduced T_{sk} values during arousal (Willis and Brigham 2003). The normothermic period was defined as the time between the end of this rapid warming period and the initiation of cooling, which was in turn defined as an abrupt decline in T_{sk} preceding steady-state torpor. We excluded data from the first 24 h after releasing the bats to avoid the influence of this disturbance on our results.

We recorded the date and time that arousals occurred from the T_{sk} trace and converted time of arousal from minutes before or after sunset into radians. We then used Rayleigh’s test for circular distributions to determine if arousal times were significantly different from a random distribution. All analyses were conducted in R version 2.10.1 (R Development Core Team 2009) and values are reported as the mean \pm SD. We assessed significance at the $P < 0.05$ level.

Results


We recorded 459 bat-days of T_{sk} data from 15 individuals over 73 days following transmitter attachment and release. Average body mass at capture was 9.4 ± 1.3 g and BCI was 0.26 ± 0.04 g/mm (Table 1). All bats exhibited long bouts of torpor averaging 15.3 ± 8.9 days with the longest bout lasting 35 days, interspersed with brief periodic arousals averaging 3.3 ± 1.1 h (Fig. 1). We recorded a total of 42 discrete torpor bouts and 29 arousals before the final transmitter battery failed on May 5, 2012.

We found no evidence of a relationship between arousal times and sunset. The circular frequency distribution of differences between arousal time and sunset time, expressed in radians, was no different from a random distribution ($Z = 0.65$, $P = 0.54$, mean vector = 0.15, $N = 15$ bats, $n = 29$ arousals, Fig. 2a, b). This analysis was technically pseudoreplicated as we recorded multiple arousals from some, but not all individuals. However, we still found no

Table 1 Sex, age class, morphometric measurements number of arousals recorded for hibernating *M. lucifugus* in Firecamp cave in Central Manitoba

ID	Demographic	Mass (g)	Forearm (mm)	BCI (g/mm)	No. of arousals	Duration of recording (days)
1	Female adult	10.3	39.84	0.26	2	33
2	Female adult	11.0	37.93	0.29	1	25
3	Female adult	12.4	38.37	0.32	2	31
4	Female adult	11.1	38.26	0.29	1	22
5	Female YOY	11.5	38.51	0.30	6	52
6	Female YOY	10.0	36.57	0.27	1	26
7	Female YOY	8.7	36.23	0.24	1	23
8	Female YOY	11.8	37.30	0.32	2	26
9	Male adult	9.3	38.46	0.24	4	73
10	Male adult	9.5	37.36	0.25	2	31
11	Male adult	8.8	38.70	0.23	1	24
12	Male adult	8.3	38.01	0.22	1	22
13	Male YOY	8.9	37.70	0.24	2	25
14	Male YOY	8.2	37.69	0.22	1	17
15	Male YOY	8.2	38.98	0.21	2	29

Note that we did not include each bat's first arousal associated with capture in our analysis
YOY young-of-the-year

Fig. 1 Skin temperature (upper trace) of three representative hibernating *M. lucifugus*. **a** Female adult, **b** Male young of the year, **c** female adult; lower trace shows the average cave temperature as measured by data loggers within 60 cm of the roosting bats

relationship between sunset and arousal time, when we limited the analysis to each individual's first arousal after capture ($Z = 1.2$, $P = 0.32$, mean vector = 0.28) or calculated the average arousal time for each individual ($Z = 0.41$, $P = 0.67$, mean vector = 0.16).

Although bats did not synchronize arousals with sunset, they did appear to synchronize arousals with each other. Twenty-one % (6/29) of all the arousals we observed occurred on only a single day (Day 17 of the study, Fig. 3) and 38 % (11/29) occurred on only 2 days. Moreover, only 38 % (11/29) of arousals occurred in isolation, on days when no other bats aroused. The distributions of days on

which each individual's first arousal occurred differed from a random distribution ($Z = 3.32$, $P = 0.036$, mean vector = 0.47, $N = 15$, $n = 15$) indicating that bats were more likely than predicted by chance to arouse on the same day.

In keeping with our prediction that individuals in the same cluster should arouse from torpor at the same time, many (but not all) bats aroused on the same days although not all bats aroused at precisely the same times. For example, the six arousals we detected on Day 17 each lasted 2.9 ± 0.9 h and were spread out over a 14.9-h period so that some bats had already re-entered torpor before




Fig. 2 **a** Histogram showing frequency of arousals from torpor versus time relative to sunset for 15 *M. lucifugus*. **b** Clock face representing the circular distribution of arousal times by individual *M. lucifugus* during hibernation. Average sunset time was 18:42 h \pm 0.4 ($Z = 0.65$, $P = 0.54$, mean vector = 0.15, $N = 15$ bats, $n = 29$ arousals)

other bats rewarmed. Of all the arousals that we observed occurring on the same days as at least one other arousal, 78 % (14/18) were completely or partially overlapping with the arousals of at least one other bat (Fig. 3). In other words, bats were normothermic at the same times as other individuals in their cluster.

Discussion

In contrast to previous studies of bats from warmer environments where winter foraging is a possibility (Park et al. 2000; Hope and Jones 2012) we found no evidence of a circadian rhythm associated with arousal. Similar to a study using bat detectors in a hibernaculum (Thomas 1993), we


found that bats did not synchronize arousals with sunset. The evidence showing that bats do arouse with sunset was obtained from the UK where T_a often exceeds 10 °C during winter and flying insects are likely to be available, at least occasionally during hibernation (Park et al. 2000). Thus, the maintenance of a circadian rhythm, and the ability to synchronize arousal with a prospective foraging opportunity could be highly beneficial, allowing bats to supplement their fat reserves and take advantage of these warm nights. In central Manitoba, T_a never exceeds 10 °C for 5 months of the year, and typically never exceeds freezing for at least 2–3 months (National Climate Data and Information Archive 2012). Therefore, flying insects are consistently unavailable for a much longer period during winter. Despite this limitation on foraging opportunities, we predicted that bats in our study would exhibit arousals linked with sunset because, presumably the cost of maintaining a circadian clock is relatively low and, especially as spring approaches, the potential for foraging opportunities does occasionally arise. For example, during late-winter (i.e., March) 2012, there were multiple nights during which temperatures were warm enough to support flying insects and foraging by bats (Czenze and Willis, unpublished data). However, our results suggest that, at least during mid- to late-winter in our study area, the costs of maintaining a daily rhythm appear to outweigh any possible benefits of winter foraging.

Although we found no daily pattern to arousals, we did find evidence that bats arouse synchronously with each other. We predicted that benefits of social thermoregulation would favour synchronous arousals, especially in this site as all bats in Firecamp cave hibernate exclusively in one cluster within a crack in the ceiling. Thus, in contrast to many little brown bat hibernacula, individuals in this cave have the potential to warm their local environment and presumably could also gain significant energetic benefits by arousing synchronously (Boyles et al. 2008; Boyles and Brack 2009).

Our results were consistent with the social thermoregulation hypothesis given that 78 % of all the arousals we observed overlapped with the arousals of at least one other individual and often more than one. Therefore, although bats did not appear to maximize benefits of social thermoregulation by all arousing at the same times, they almost certainly gained some energetic benefit. Mating attempts by males which occur during hibernation (Thomas et al. 1979) may also have influenced this synchrony. Interestingly, for the three sets of arousals we observed, each appeared to be initiated by a male bat, followed by multiple females. These arousals were not, however, completely synchronized. Given the tight energy budgets of hibernating bats, an interesting question for future studies is, why has selection not favoured mechanisms that allow all bats

Fig. 3 Skin temperature (upper trace) of six *M. lucifugus* showing the timing of arousals on Day 17 of the study.

a Female young of the year, **b** female young of the year, **c** male young of the year, **d** male adult, **e** female adult, **f** male adult. The lower flat trace shows the average cave temperature as measured by temperature data loggers within 60 cm of the roosting bats. Dashed lines indicate sunrise and sunset



in a cluster to precisely synchronize arousals to minimize rewarming costs?

Approximately, synchronous arousals by bats in the same microclimate within a hibernaculum might be explained by multiple individuals accumulating physiological imbalance at about the same rate. They might therefore arouse at about the same time to restore homeostasis. For example, assuming evaporative water loss (EWL) influences arousal timing, bats in the same cluster in our study may have experienced similar rates of EWL during torpor (Thomas and Cloutier 1992) and, therefore, may have almost simultaneously reached a point at which arousal was necessary to re-hydrate. All radiotagged bats aroused during handling on 20 February 2012 when we attached transmitters, so it is also possible that this disturbance-induced arousal resets the mechanism underlying arousal (i.e., allowed all individuals to drink or excrete metabolic wastes) increasing the chance that their subsequent arousals would be synchronized. However, given large between-individual variation in most physiological processes, including those likely to influence arousals (e.g., metabolic rate, EWL) it seems unlikely that this is the exclusive explanation for the similar arousal times of multiple individuals in our study.

Another possible explanation for the timing of arousals we observed is that warm bats disturb torpid conspecifics. Even non-tactile disturbance by human visitors to hibernacula can stimulate arousals in bats (Thomas et al. 1990) and, it is possible that movement by other bats within a hibernaculum or within a cluster could have similar effects.

This seems unlikely given strong selection pressure for bats to remain torpid for long enough during winter to conserve energy. Such a mechanism might be important if different individuals exhibit consistent individual differences in sensitivity to disturbance. For example, mated adult females may be especially resistant to disturbance because selection appears to favour mechanisms allowing females to save fat for spring reproduction (Jonasson and Willis 2011). At the other extreme, inexperienced juvenile bats may be more prone to disturbance during torpor and, to avoid exhausting fat reserves too quickly, may need to learn how to differentiate disturbances which warrant arousing (e.g., a potential predator) from those that do not (e.g., conspecifics).

If disturbance by bats can cause arousals, this could have consequences for bats hibernating in sites affected by white-nose syndrome (WNS). WNS is caused by cutaneous infection with the fungus *Geomyces destructans* (Blehert et al. 2009; Lorch et al. 2011) and infection causes bats to arouse from torpor too frequently during hibernation and prematurely exhaust energy reserves (Boyles and Willis 2010; Reeder et al. 2012; Warnecke et al. 2012). If active bats can induce arousals in other torpid individuals, the increased arousal frequency caused by infection with *G. destructans* could result in a positive feedback of arousals, accelerating the use of fat reserves even further. More work is needed to understand the physiological signals and environmental cues that trigger arousal in hibernators, especially in bats which are highly prone to disturbance and, in North America, now facing the threat of WNS.

This is the first study to examine the timing of arousal in free-ranging little brown bats, particularly in the extreme northern part of their range and provides, to our knowledge, the first evidence that bats in northern climates synchronize their arousals with cluster-mates rather than photoperiod. Future research on northern bats should aim to examine arousal patterns of bats in caves with multiple clusters, as well as the arousal patterns of related vs. unrelated bats from the same hibernacula, to improve understanding of mechanisms underlying this critical aspect of hibernation.

Acknowledgments We thank Dylan Baloun, Ally Menzies, Felix Martinez-Nuñez, Chonny Sayapheth and Adriana Suarez for help with fieldwork and Manitoba Conservation for lodging in the field. This research was supported by Grants to CKRW from the Natural Sciences and Engineering Research Council (Canada), the Canada Foundation for Innovation, the Manitoba Research and Innovation Fund, The Manitoba Hydro Forest Enhancement Program and The University of Winnipeg Chancellor's Research Chair in Wildlife Health.

References

- Aldridge H, Brigham RM (1988) Load carrying and maneuverability in an insectivorous bat: a test of the 5% "rule" of radio-telemetry. *J Mammal* 69:379–382
- Audet D, Thomas DW (1996) Evaluation of the accuracy of body temperature measurement using external radiotransmitters. *Can J Zool* 74:1778–1781
- Barclay RMR, Kalcounis MC, Crampton LH, Stefan C, Vonhof MJ, Wilkinson L, Brigham RM (1996) Can external radiotransmitters be used to assess body temperature and torpor in bats? *J Mammal* 77:1102–1106
- Blehert DS, Hicks AG, Behr M, Meteyer CU, Berlowski-Zier BM, Buckles EL, Coleman JTH, Darling SR, Gargas A, Niver R, Okoniewski JC, Rudd RJ, Stone WB (2009) Bat white-nose syndrome: an emerging fungal pathogen? *Science* 323:227
- Boyles JG, Brack JV (2009) Modeling survival rates of hibernating mammals with individual-based models of energy expenditure. *J Mammal* 90:9–16
- Boyles JG, Willis CKR (2010) Could localized warm areas in cold caves reduce mortality of hibernating bats affected by white-nose syndrome? *Front Ecol Environ* 8:92–98
- Boyles JG, Storm JJ, Brack JV (2008) Thermal benefits of clustering during hibernation—a field test of competing hypotheses on *Myotis sodalis*. *Funct Ecol* 22:632–6369
- Davis WH, Hitchcock HB (1965) Biology and migration of the bat, *Myotis lucifugus*, in New England. *J Mammal* 46:296–313
- Geiser F (2004) Metabolic rate and body temperature reduction during hibernation and daily torpor. *Annu Rev Physiol* 66:239–274
- Grahn DA, Miller JD, Houng VS, Heller HC (1994) Persistence of circadian rhythmicity in hibernating ground squirrels. *Am J Physiol Regul Integr Comp Physiol* 266:R1251–R1258
- Hope PR, Jones G (2012) Warming up for dinner: torpor and arousal in hibernating Natterer's bats (*Myotis nattereri*) studied by radio telemetry. *J Comp Physiol B* 182:569–578
- Humphries MM, Thomas DW, Kramer DL (2003) The role of energy availability in mammalian hibernation: a cost–benefit approach. *Physiol Biochem Zool* 76:165–179
- Jonasson KA, Willis CKR (2011) Changes in body condition of hibernating bats support the thrifty female hypothesis and predict consequences for populations with white-nose syndrome. *PLoS One* 6:e21061
- Jonasson KA, Willis CKR (2012) Hibernation energetics of little brown bats. *J Exp Biol* 215:2141–2149
- Jones G, Duvergé PL, Ransome RD (1995) Conservation biology of an endangered species: field studies of greater horseshoe bats. *Symp Zool Soc Lond* 67:309–324
- Lorch JM, Meteyer CU, Behr MJ, Boyles JG, Cryan PM et al (2011) Experimental infection of bats with *Geomyces destructans* causes white-nose syndrome. *Nature* 480:376–378
- Luis AD, Hudson PJ (2006) Hibernation patterns in mammals: a role for bacterial growth? *Funct Ecol* 20:471–477
- Malan A (2010) Is the torpor-arousal cycle of hibernation controlled by a nontemperature-compensated circadian clock? *J Biol Rhythms* 25:166–175
- Moore MS, Reichard JD, Murtha TD, Zahedi B, Fallier RM, Kunz TH (2011) Specific alterations in complement protein activity of little brown myotis (*Myotis lucifugus*) hibernating in white-nose syndrome affected sites. *PLoS One* 6:e27430
- National Climate Data and Information Archive. Environment Canada. www.climate.weatheroffice.gc.ca. Accessed 9 October 2012
- Park KJ, Jones G, Ransome RD (2000) Torpor, arousal and activity of hibernating greater horseshoe bats (*Rhinolophus ferrumequinum*). *Funct Ecol* 14:580–588
- Prendergast BJ, Freeman DA, Zucker I, Nelson RJ (2002) Periodic arousal from hibernation is necessary for initiation of immune responses in ground squirrels. *Am J Physiol* 282:1054–1062
- Racey PA, Swift SM (1981) Variations in gestation length in a colony of pipistrelle bats (*Pipistrellus pipistrellus*) from year to year. *J Reprod Fertil* 6:123–129
- R Development Core Team (2009) R: A language and environment for statistical computing R Foundation for Statistical Computing, Vienna, Austria ISBN 3- 900051-07-0, URL: <http://www.R-project.org>
- Reeder DM, Frank CL, Turner GG, Meteyer CU, Kurta A et al (2012) Frequent arousal from hibernation linked to severity of infection and mortality in bats with whitenose syndrome. *PLoS One* 7:e38920
- Ruby NF, Dark J, Burns DE, Heller HC, Zucker I (2002) The suprachiasmatic nucleus is essential for circadian body temperature rhythms in hibernating ground squirrels. *J Neurosci* 22:357–364
- Ruf T, Arnold W (2000) Mechanisms of social thermoregulation in hibernating alpine marmots (*Marmota marmota*). In: Heldmaier G, Klingenspor M (eds) Life in the cold: eleventh international hibernation symposium. Springer, Berlin, pp 81–94
- Stawski C, Geiser F (2009) Fat and fed: frequent use of summer torpor in a subtropical bat. *Naturwissenschaften* 97:29–35
- Thomas DW (1993) Lack of evidence for a biological alarm clock in bats (*Myotis* spp.) hibernating under natural conditions. *Can J Zool* 71:1–3
- Thomas DW, Cloutier D (1992) Evaporative water loss by hibernating little brown bats, *Myotis lucifugus*. *Physiol Zool* 65:443–456
- Thomas DW, Geiser F (1997) Periodic arousals in hibernating mammals is evaporative water loss involved? *Funct Ecol* 11:585–591
- Thomas DW, Fenton BM, Barclay RMR (1979) Social behavior of the little brown bat, *Myotis lucifugus*: I. mating behavior. *Behav Ecol Sociobiol* 6:129–136
- Thomas DW, Dorais M, Bergeron JM (1990) Winter energy budgets and cost of arousals for hibernating little brown bats, *Myotis lucifugus*. *J Mammal* 71:475–479

- Wang LCH (1978) Energetic and field aspects of mammalian torpor: the Richardson's ground squirrels. In: Wang LCH, Hudson JW (eds) *Strategies in cold*. Academic Press, New York, pp 109–145
- Warnecke L, Turner JM, Bollinger TK, Lorch JM, Misra V, Cryan PM, Wibbelt G, Blehert DS, Willis CKR (2012) Inoculation of a North American bat with European *Geomyces destructans* supports the novel pathogen hypothesis for the origin of white-nose syndrome. *Proc Natl Acad Sci USA* 109:6999–7003
- Wilkinson GS, South JM (2002) Life history, ecology and longevity in bats. *Aging Cell* 1:124–131
- Willis CKR, Brigham RM (2003) Defining torpor in free-ranging bats: experimental evaluation of external temperature-sensitive radiotransmitters and the concept of active temperature. *J Comp Physiol B* 173:379–389
- Willis CKR, Jameson J, Faure PA, Boyles JG, Brack JV, Cervone TH (2009) Thermocron iButton and iBBat temperature dataloggers emit ultrasound. *J Comp Physiol B* 179:867–874