

Stellar Population Models with Variable Element Abundance Ratios

Daniel Thomas, Claudia Maraston, Ralf Bender

MPE, Giessenbachstraße, 85748 Garching, Germany

Universitäts-Sternwarte, Scheinerstraße 1, 81679 München, Germany

Abstract. We present a comprehensive set of new generation stellar population models of Lick absorption line indices, which for the first time include element abundance ratios different from solar. We computed the 21 Lick indices CN₁, CN₂, Ca4227, G4300, Fe4383, Ca4455, Fe4531, C₂4668, H β , Fe5015, Mg₁, Mg₂, Mg_b, Fe5270, Fe5335, Fe5406, Fe5709, Fe5782, Na D, TiO₁, and TiO₂, in the wavelength range $4000 \lesssim \lambda \lesssim 6500 \text{ \AA}$. Models are provided with: $[\alpha/Fe] = 0.0, 0.3, 0.5$, $[\alpha/Ca] = -0.1, 0.0, 0.2, 0.5$, and $[\alpha/N] = -0.5, 0.0$; ages from 1 to 15 Gyr; total metallicities from 1/200 to 3.5 solar ($-2.25 \leq [Z/H] \leq 0.67$).

The models are based on the evolutionary synthesis technique described in Maraston (1998). The α/Fe enhanced mixtures are obtained by increasing the abundances of α -group elements and by decreasing the abundances of the Fe-peak elements, such that total metallicity is conserved. The impact from these element abundance variations on the absorption line indices is taken from Tripicco & Bell (1995), using an extension of the method introduced by Trager et al. (2000). Most importantly, we take into account that the empirical stellar libraries used to compute model indices follow the chemical enrichment history of the Milky Way, and are therefore biased towards super-solar α/Fe ratios at sub-solar metallicities. We corrected for this bias, so that the models presented here have well-defined α/Fe ratios at all metallicities.

We take particular care at calibrating the models with galactic globular clusters, for which ages, metallicities, and element abundance ratios are known from independent sources. Our α/Fe enhanced models with $[\alpha/Fe] = 0.3$ (and 12 Gyr age) perfectly reproduce the positions of the globular cluster data in the Mg₁-(Fe) diagram up to solar metallicities (see also Maraston et al. 2003). The total metallicities for the sample clusters that we derive from these indices are in excellent agreement with the Zinn & West (1984) metallicity scale. We point out that the latter most likely reflects total metallicity rather than iron abundance, because it is obtained essentially by averaging the abundances derived from the Mg triplet near 5175 Å and the Fe blend at 5270 Å. This aspect needs to be emphasized, as with the α/Fe enhanced models we are now in the position to distinguish total metallicity $[Z/H]$ and iron abundance $[Fe/H]$.

By means of our calibrated α/Fe enhanced models, we confirm that the index $[MgFe]$, suggested by González (1993) to balance α/Fe ratio effects, is almost independent of α/Fe . As it modestly decreases with increasing α/Fe , however, we define the slightly modified index $[MgFe]' \equiv \sqrt{Mg_b (0.72 \cdot Fe5270 + 0.28 \cdot Fe5335)}$ which is completely independent of α/Fe , and hence an even better tracer of total metallicity. We further show that the linear correlation between Mg₂ and metallicity at old ages derived

empirically by Brodie & Huchra (1990) is valid up to $\sim 1/3$ solar metallicity, but underpredicts Mg_2 indices at metallicities above that threshold.

It turns out to be hard to find indices that correlate with α/Fe as well as the intensively studied indices Mg_1 , Mg_2 , and Mg_b . Promising alternatives are the blue indices CN_1 and CN_2 that also increase with increasing α/Fe ratio, mainly because of an anti-correlation with Fe abundance. With the caveat that CN_1 and CN_2 are additionally sensitive to C and N abundances, they can be regarded to be complementary to the indices Mg_1 , Mg_2 , and Mg_b . Alternatives to the iron indices Fe5270 and Fe5335, the strengths of which decrease with increasing α/Fe ratio, are easier to find. The best cases are the indices Fe4383, Fe4531, Fe5015, and Fe5709.

The indices CN_1 , CN_2 , and Ca4227 of globular clusters are very interesting, particular cases. We find that the relatively strong CN features observed in globular clusters require models in which nitrogen is enhanced by a factor three relative to the α -elements, hence $[\alpha/N] = -0.5$. This is in agreement with early suggestions by D'Antona (2003), that stars in globular clusters are nitrogen enriched by a previous generation of stars. The good calibration of other indices like Mg_1 , Mg_b or $\langle Fe \rangle$ is not affected by a variation of the $[\alpha/N]$ ratio, as these indices are not sensitive to nitrogen abundance. We note that an enhancement of carbon abundance, instead, would lead to serious inconsistencies with Mg_1 . Interestingly, also Ca4227 is sensitive to nitrogen abundance, and the globular cluster data of this index are also best reproduced by the model with increased nitrogen abundance.

To conclude, the stellar population models presented here make it possible, for the first time, to study in detail individual element abundance ratios of unresolved stellar populations (Thomas et al. 2003b). In particular, total metallicity is now a well-defined quantity.

The models are published in Thomas et al. (2003a). Tables are available electronically via anonymous ftp at [ftp.mpe.mpg.de](ftp://ftp.mpe.mpg.de/people/dthomas/SSPs) in the directory `people/dthomas/SSPs`. They are also available via WWW by going to <ftp://ftp.mpe.mpg.de/people/dthomas/SSPs>.

References

- Brodie J., Huchra J., 1990, *ApJ*, 362, 503
- Da Costa, G. S., D'Antona, F. and Gratton, R. G. 2003, This volume, JD 4
- González, J. J. 1993, PhD thesis, University of Santa Cruz
- Maraston C. 1998, *MNRAS*, 300, 872
- Maraston C., Greggio L., Renzini A. et al. 2002, *A&A*, 400, 823
- Trager S. C., Faber S. M., Worthey G., González J. J. 2000, *AJ*, 119, 164
- Tripicco M. J., Bell R. A. 1995, *AJ*, 110, 3035
- Thomas, D., Maraston, C., Bender, R. 2003a, *MNRAS*, 339, 897
- Thomas, D., Maraston, C., Bender, R. 2003b, *MNRAS*, 343, 279
- Zinn R., West M. J. 1984, *ApJS*, 55, 45