

Stem Cell-Based Therapies for Ischemic Stroke: Preclinical Results and the Potential of Imaging-Assisted Evaluation of Donor Cell Fate and Mechanisms of Brain Regeneration

Peer-reviewed author version

GERVOIS, Pascal; WOLFS, Esther; RATAJCZAK, Jessica; DILLEN, Yorg; VANGANSEWINKEL, Tim; HILKENS, Petra; BRONCKAERS, Annelies; LAMBRICHTS, Ivo & STRUYS, Tom (2016) Stem Cell-Based Therapies for Ischemic Stroke: Preclinical Results and the Potential of Imaging-Assisted Evaluation of Donor Cell Fate and Mechanisms of Brain Regeneration. In: MEDICINAL RESEARCH REVIEWS, 36(6), p. 1080-1126.

DOI: 10.1002/med.21400

Handle: <http://hdl.handle.net/1942/22909>

APTARA	MED	med21400	Dispatch: July 9, 2016	CE: AFL
	Journal	MSP No.	No. of pages: 47	PE: XXXXX

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

Stem Cell-Based Therapies for Ischemic Stroke: Preclinical Results and the Potential of Imaging-Assisted Evaluation of Donor Cell Fate and Mechanisms of Brain Regeneration

Pascal Gervois, Esther Wolfs, Jessica Ratajczak, Yörg Dillen, Tim Vanganswinkel, Petra Hilkens, Annelies Bronckaers, Ivo Lambrechts, and Tom Struys

Morphology Research Group, Biomedical Research Institute, Hasselt University, Campus Diepenbeek, Bioville, Diepenbeek, Belgium

Published online in Wiley Online Library (wileyonlinelibrary.com).
DOI 10.1002/med.21400

Abstract: Stroke is the second most common cause of death and is a major cause of permanent disability. Given the current demographic trend of an ageing population and associated increased risk, the prevalence of and socioeconomic burden caused by stroke will continue to rise. Current therapies are unable to sufficiently ameliorate the disease outcome and are not applicable to all patients. Therefore, strategies such as cell-based therapies with mesenchymal stem cell (MSC) or induced pluripotent stem cell (iPSC) pave the way for new treatment options for stroke. These cells showed great preclinical promise despite the fact that the precise mechanism of action and the optimal administration route are unknown. To gain dynamic insights into the underlying repair processes after stem cell engraftment, noninvasive imaging modalities were developed to provide detailed spatial and functional information on the donor cell fate and host microenvironment. This review will focus on MSCs and iPSCs as types of widely used stem cell sources in current (bio)medical research and compare their efficacy and potential to ameliorate the disease outcome in animal stroke models. In addition, novel noninvasive imaging strategies allowing temporospatial in vivo tracking of transplanted cells and coinciding evaluation of neuronal repair following stroke will be discussed. © 2016 Wiley Periodicals, Inc. Med. Res. Rev., 00, No. 0, 1–47, 2016

Key words: ischemic stroke; mesenchymal stem cells; induced pluripotent stem cells; mechanisms of stem cell therapy; noninvasive imaging

Correspondence to: Tom Struys, Morphology Research Group, Biomedical Research Institute, Hasselt University, Campus Diepenbeek, Agoralaan, Bioville, Diepenbeek, Belgium.
E-mail: tom.struys@uhasselt.be.

1 2 • GERVOIS ET AL.

2 **1. INTRODUCTION**

3
4 Worldwide, stroke is the second single most common cause of death, accounting for 10–15%
5 of deaths each year.^{1,2} Moreover, stroke is an important cause of adult disability as 90% of
6 patients that survive from a stroke are left with a residual deficit.^{3,4} It might therefore be clear
7 that stroke-related public and insurance costs constitute a major burden on healthcare systems
8 worldwide.^{1,2} Combining the expectation that the amount of people over the age of 65 will
9 double by 2030, and that the risk of suffering a stroke doubles for each decade over the age
10 of 55, will even lead to a further increase in patient numbers with permanent disabilities and
11 socioeconomic burden.^{2,4–7}

12 Despite this increased incidence, current available therapies are unable to sufficiently ame-
13 liorate the disease outcome or are even not applicable for subgroups of patients due to many
14 contraindications as will be discussed below. Therefore, new therapeutic strategies are needed
15 for treating and preventing stroke that can be applied to patients with distinct risk profiles and
16 in a broader time frame, as time plays a crucial role in the treatment of acute ischemic stroke.
17 In addition to clinical advances in stroke management, cell-based therapies have emerged as
18 a potential candidate to promote functional recovery in patients suffering from stroke.⁸ De-
19 spite the promising results achieved with cell-based therapies in stroke, the host response, the
20 precise mechanisms of action of these therapies, and the fate of the donor cells remain largely
21 unknown.⁹ Therefore, noninvasive imaging modalities have been developed that are able to
22 provide detailed temporospatial and functional information on the donor cell fate, the host
23 microenvironment, and endogenous repair mechanisms,¹⁰ which will be discussed later.

24
25 **A. Pathophysiology of Stroke**

26
27 The pathophysiology of stroke can be defined as a neurologic dysfunction of vascular origin
28 with the sudden or rapid occurrence of symptoms and signs corresponding to the involvement
29 of focal areas in the brain.¹¹ Two different types of stroke can occur: ischemic stroke (80–85%)
30 and hemorrhagic stroke (15–20%). Ischemic stroke is most frequently caused by thromboem-
31 bolisms while hemorrhagic stroke most often results from vessel wall pathology associated with
32 hypertension and microaneurysms.¹² This review will only focus on ischemic stroke as the main
33 pathology.

34 In ischemic stroke, the blood supply to certain brain areas is compromised due to vascular
35 occlusion thereby causing several changes at the (sub)cellular level and ultimately tissue damage.
36 These cellular and molecular processes start with energy depletion followed by glutamate
37 release leading to glutamate-induced excitotoxicity, ion channel dysfunction, and free radical
38 production. These processes in turn disrupt the cellular membrane, damage mitochondria and
39 DNA, generate an immune response, and trigger necrotic and apoptotic cell death (Fig. 1).¹³
40 In the ischemic core, these cellular changes are irreversible.¹⁴ However, the tissue surrounding
41 the core, also termed as the ischemic penumbra, is functionally impaired but still viable.¹⁵
42 This area “at risk” is therefore considered as the main target for therapeutic interventions that
43 are believed to exert a protective effect in intervening with the cellular processes discussed
44 above.^{16,17} Using noninvasive imaging methods, the ischemic penumbra has been divided in
45 additional border zones characterized by different grades of hypoperfusion and varying risk of
46 progressing toward lost infarcted tissue if a proper treatment is not initiated (Fig. 1).^{18,19}

47 When considering therapies that are aimed to salvage the ischemic penumbra by restoring
48 perfusion, it is also important to take into account that restoring the blood flow in ischemic
49 tissue by thrombolytic treatment can lead to secondary damage by reperfusion injury.^{13,20}
50 This reperfusion injury is mediated by leukocyte infiltration through local disruption of the
51 blood–brain barrier (BBB) and accompanying matrix metalloprotease (MMP) production in




Figure 1. Areas at risk and pathophysiology of ischemic stroke. (1) Blood flow to focal areas of the brain is diminished by vascular occlusion by, for example, an embolism. The affected ischemic tissue can be divided into the ischemic core (C) where tissue damage is irreversible, the salvageable ischemic penumbra (P), and a zone of benign oligemia (B) where blood supply can be obtained by leptomeningeal collaterals. Additional border zones with different grades of hypoperfusion and varying risk of progressing toward unsalvageable tissue if a treatment is not initiated were identified with perfusion-weighted MRI. These areas are the core-penumbra border zone (BZ1) and the penumbra-benign oligemia zone (BZ2). (2) The cellular changes ultimately leading to cell death initiate with ATP depletion due to ischemia, followed by depolarization of the affected neurons that triggers (3) glutamate release. (4) Glutamate-induced excitotoxicity is mediated by an elevated sodium and calcium influx that causes cell swelling, a depolarization wave that will lead to damage in neighboring cells, activation of a cascade of enzymatic reactions ultimately leading to membrane and mitochondrial damage and ROS production, which will additionally damage mitochondria and DNA ultimately leading to cell death. (5) Necrotic/apoptotic neurons secrete inflammatory mediators that activate resting microglia and enhance neutrophil and macrophage infiltration. The effects of activated microglia vary and include migration toward and phagocytosis of damaged neurons and depending on the M1/M2 activation state of activated microglia, proinflammatory and/or anti-inflammatory mediators are released. Image was created using Servier Medical Art.

addition to stimulation of reactive oxygen species (ROS) production, thereby damaging the reperfused environment.^{13,14,20–22} In turn, the reperfused ischemic stroke lesion can transform into a petechial hemorrhage that does not influence the prognosis or it can transform into an intracerebral hematoma, which is associated with a poor outcome.^{22–24}

Due to the complexity of the molecular processes that are involved in the onset of stroke, but also in ischemic reperfusion injury, multiple strategies are considered for treating stroke. These strategies include both acute and long-term approaches. Acute therapies aim to salvage the ischemic penumbra and limit reperfusion injury, while long-term therapeutic strategies aim to reconstitute the lost tissue from the ischemic core, as will be discussed later.

B. Limitations and Potential Improvements of Available Therapies for Ischemic Stroke

Current therapies or approaches that have been proven to be effective in reducing the mortality rate and improving the functional outcome of acute ischemic stroke include the establishment of a specialized stroke care unit (SCU),²⁵ thrombolysis with tissue plasminogen activator (tPA),^{26,27} aspirin administration,²⁸ and decompressive surgery following ischemic stroke.²⁹ The most remarkable advance in stroke management that reduced the mortality and disability

1 4 • GERVOIS ET AL.

2 rate has been the establishment of a SCU, which is a separate physical space in general medical
3 wards with specialized and specifically dedicated trained staff.²⁵ Despite these advances
4 in stroke management, they can only be applied in a short therapeutic window. The FDA-
5 approved standard treatment for stroke, thrombolysis with tPA, is only applicable to less than
6 10% of stroke patients and ideally needs to be initiated within 3 hr after the onset of ischemia.
7 This low applicability rate of tPA is mainly associated with the higher risk of intracerebral
8 hemorrhage when tPA is administered longer than 3 hr after the onset of stroke symptoms.³⁰
9 Moreover, patients often do not recognize stroke-associated symptoms and only show up at the
10 hospital with advanced stroke symptoms³¹ or the symptoms are not immediately recognized
11 by the hospital staff that can be counteracted by a SCU.²⁵ Therefore, the presenting stroke
12 patient has often exceeded the safe 3-hr time window to be considered for tPA treatment. This
13 time window can be extended to 4.5 hr if low-risk patients are selected together with providing
14 extensive care due to the higher risk of secondary damage and increased mortality due to reper-
15 fusion injury.^{32,33} The major criteria to consider these low-risk patients included the absence
16 of cerebral hemorrhage or major infarction and they presented with acute ischemic stroke and
17 the symptoms started 3–4.5 hr before initiation of drug administration (for all inclusion and
18 exclusion criteria, see Table I in Ref. [34]).

19 Other approaches such as decompressive surgery and aspirin administration need to be
20 started within 48 hr after the onset of stroke. Moreover, the benefits of aspirin administration
21 are small while decompressive surgery is only applicable for patients whose stroke-associated
22 infarct region is caused by middle cerebral artery-related pathology, combined with malignant
23 space-occupying brain edema.³⁵

24 Various interventions are currently under investigation, which include extending the
25 time window for thrombolysis with desmoteplase or alteplase,^{36,37} ultrasound-enhanced
26 thrombolysis,³⁸ the creation of new thrombectomy devices,³⁹ stents, stent retrievers, and pro-
27 tective drugs.¹⁷ Asadi et al. provide an up to date, in depth overview of the different studies
28 using endovascular treatments, which can be used as an important supplementary therapy to
29 current intravenous thrombolysis.⁴⁰

30 However, when extending the therapeutic window for thrombolytic interventions, the is-
31 chemic brain still needs to be protected from reperfusion injury. Therefore, reducing reperfusion
32 injury is also a route that is currently being investigated as an additional procedure to be imple-
33 mented after thrombolytic therapy.²⁰ These approaches aim at reducing the local production of
34 ROS and BBB-damaging MMPs, or mediating the local immune response that would otherwise
35 lead to secondary damage.^{41–44} One of these approaches is therapeutic hypothermia⁴⁵ that has
36 been shown to decrease ischemic and reperfusion injury by influencing local excitotoxicity,
37 neuroinflammation, and ROS production.⁴⁶

38 While these therapies and novel interventions aim at mitigating the disease outcome, they
39 can only be applied in the first few hours or days after the onset of ischemic stroke.^{32,47} Patients
40 surviving stroke and not treated properly within this narrow time window are therefore often
41 left with permanent disabilities, associated with the focal areas of the brain that are affected.³²
42 In these patients, a therapy that can be applied weeks to months after stroke onset can be
43 beneficial. These therapies aim at restoring the lost neural tissue or stimulating brain plasticity
44 to improve the functional outcome but also muscle strengthening and physical conditioning
45 has been shown to improve the quality of life of patients with permanent disabilities.⁴⁸ Stem
46 cell based therapies have been shown to be a promising approach in achieving such results.⁴⁹

47
48
49 **C. Stem Cell Sources and Mechanisms of Action for Cell-Based Therapies**

50 When considering stem cells sources for a cell-based therapy in ischemic stroke, ex vivo ex-
51 panded and manipulated neural stem cells (NSCs) or neural precursor cells (NPCs) would be

Table 1. Overview of Preclinical Stroke Studies Using MSCs

Stem cell type	(Pre)differentiation and/or treatment	Species	Occlusion time	Time of transplantation	Cell dose and location of transplantation	Fate of transplanted cells	Outcome	Reference
h-DPSCs	No	Sprague-Dawley rats	2 hr	24 hr post-surgery	3×10^5 intrastriatal; 3×10^5 intracortical	2.3% migrates toward the stroke lesion. Differentiation toward astrocytes in preference to neurons	Improvement in forelimb sensorimotor function at 4 weeks posttreatment, mediated by paracrine effects	59
h-UMSCs	hUMSCs and hUMSCs cultured in neuronal conditioned medium	Sprague-Dawley rats	90 min	24 hr post-surgery	2.5×10^5 intracortical in two sites	36 Days survival, not quantified	Significant improvements in motor function, greater metabolic activity of cortical neurons, and better revascularization in the infarct cortex due to paracrine effects	71
h-UMSCs	No	Wistar rats	2 hr	24 hr post-surgery	2×10^5 intracortical	5 Weeks survival, <3% expressing neural markers	hUMSC accelerate neurologic recovery after stroke by promoting angiogenesis	130
h-BMSCs	No	SHR rats	Permanent	1 week post-surgery	7.5×10^4 in three different cortical sites	6 Weeks after grafting, donor cells expressed astrocyte, oligodendroglial, and neuronal markers. Functional integration was unlikely	Improved functional outcome, mediated by paracrine factors that are produced by the surviving donor cells	70
r-BMSC and h-BMSCs	Notch-induced BMSCs	Sprague-Dawley rats	1 hr	4-6 weeks post-surgery	6×10^4 intrastriatal in three sites	r-BMSCs show higher survival (15% vs. 7%) and differentiation than h-BMSCs	Improvement in locomotor and neurological function. Reduced loss of striatal perinifart cells.	137
h-ASCs	No	C57BL/6J mice	1.5 hr	1 week post-surgery	5×10^6 in the stroke lesion	Large percentage hASCs express MAP2. Low percentage of GFAP expression	Cognitive recovery and decrease in infarct size. Immunomodulation by decreasing the presence of Iba-1+ microglia and GFAP+ astrocytes	142
m-ASCs	No	C57BL/6J mice	Permanent	24 hr post-surgery	1.8×10^4 above the corpus callosum	Migration after 1 week, toward vessels, 5% survival after 4 weeks	Ischemia induces ASC-survival, migration toward the lesion and microvessels, differentiation into smooth muscle cells	138
r-BMSC	Hypoxic pretreatment (HP)	C57BL/6J mice	Permanent	24 hr post-surgery	1.0×10^6 intranasal	1.5 hr after administration, donor cells were observed in the ischemic cortex. No long-term follow up was performed	HP of BMSCs induced a higher expression of migration associated and significantly reduced infarct size and improved sensorimotor function compared to non-HP BMSCs	140
r-BMSCs	No	Wistar rats	90 min	1, 6, 24, or 48 hr post-surgery	1×10^6 into the carotid artery	q-dot nanocrystal marked BMSCs could be detected 7 days poststroke	Injecting BMSCs 24 hr after stroke had the most significant effect on graft survival/integration, infarct size reduction, and improvement of neurological function. SDF-1 and bFGF were upregulated	127
r-BMSCs	No	Wistar rats	2 hr	24 hr post-surgery	2×10^6 into the carotid artery	n/a	BMSCs facilitate axonal sprouting and remyelination in the cortical ischemic boundary zone and corpus callosum	150

Continued

Table I. Continued

Stem cell type	(Pre)differentiation and/or treatment	Species	Occlusion time	Time of transplantation reperfusion	Cell dose and location of transplantation	Fate of transplanted cells	Outcome	Reference
r-BMSCs	No	Wistar rats	2 hr	30 min after reperfusion	1 × 10 ⁶ into internal carotid artery	Magnetically labeled BMSCs could be detected with MRI	Magnetically labeled IA-delivered BMSCs could be detected with MRI and high cerebral engraftment rates are associated with impeded cerebral blood flow after injection	143
h-BMSCs	No	Wistar rats	90 min	24 hr post-surgery	1.1 × 10 ⁶ or 0.5 × 10 ⁶ into the external carotid artery	Transient localization of engrafted cells in the host brain	Localization of BMSCs in the brain but relocated to other organs 24 hr later. Increased radioactivity counts in the ipsilateral stroke hemisphere	144
h-BMSCs	No	Sprague-Dawley rats	75 min	24 hr, 4 days, and 7 days post-surgery	1 × 10 ⁶ into the carotid artery	Low survival, no expression of neuronal markers. Migration toward the lesion and secretion of BDNF	Time dependent functional recovery and cell distribution around the lesion. Mechanisms of action: neuroprotection, angiogenesis and enhancing reactive astrocytes, downregulation of MMP9	128
Autologous r-ASCs	No	Sprague-Dawley rats	90 min	3 days post-surgery	2 × 10 ⁶ into the carotid artery	1.5% of surviving cells expressed NeuN; 1% survival of transplanted cells	Improvement of neurological deficits, migration of donor cells to lesion; attenuation of astroglial activity, inhibition of apoptosis, and promotion of cellular proliferation	131
r-BMSCs	No	Aged Wistar rats	2 hr	24 hr post-surgery	2 × 10 ⁶ into the carotid artery	Donor cells survive up to 1 year and preferentially differentiate toward astrocytes	The beneficial effects of cell transplantation persisted for at least 1 year. Donor cells survived, differentiated toward astrocytes or neurons or colocalized with microglia and endothelial cells. Reduction of axonal loss and glial scar thickness	147
h-ASCs and r-ASCs	No	Sprague-Dawley rats	Permanent	30 min post-surgery	2 × 10 ⁶ h-ASCs or r-ASCs	No migration/implantation of donor cells was observed	Improved functional outcome; reduction in neuronal cell death. No reduction in lesion size. VEGF and synaptophysin was upregulated and GFAP was downregulated in the treated groups. No difference was observed between the h-ASC- and r-ASC-treated groups	151
h-BMSCs	Immortalized cells	Wistar rats	1 hr	24 hr post-surgery	3 × 10 ⁶ into the jugular vein	After 7 days no donor cells were detected	IV- transplanted human MSCs induced functional improvement, reduced infarct volume, and neuroprotection by providing IGF-1 and inducing neurotrophin expression in host brain	135

Continued

Table I. Continued

Stem cell type and h-BMSC-EVs	(Pre)differentiation and/or treatment	Species	Occlusion time	Time of transplantation	Cell dose and location of transplantation	Fate of transplanted cells	Outcome	Reference
h-BMSCs and h-BMSC-EVs	No	C57BL/6J mice	30 min	24 hr post-surgery Repeated after 3 and 5 days for h-BMSC-EVs	1 × 10 ⁶ BMSCs or EVs from 2 × 10 ⁶ BMSC in the femoral vein	n/a	Mice receiving EVs showed improved neurological function and long-term survival associated with improved angiogenesis and neurogenesis, which resembled BMSC responses.	129
r-BMSCs	No	Wistar rats	2 hr	30 min after reperfusion	1 × 10 ⁶ into the femoral vein	Magnetically labeled IV-delivered BMSCs could not be detected	Magnetically labeled IV-delivered BMSCs could not be detected	143
h-BMSCs	BMSCs, PlGF gene transfected MSCs	Sprague-Dawley rats	Permanent	6 hr post-surgery	1 × 10 ⁷ intravenously	LacZ-expressing PlGF-hBMSCs were found primarily in the penumbra and express NeuN (±10%) and GFAP (<17.23%)	hBMSCs and PlGF-transfected BMSCs improved angiogenesis, reduced the lesion size, and elicited functional improvement; the effect was more pronounced in PlGF-transduced BMSCs	136
r-BMSCs	BMSCs, CXCR4 gene transfected BMSCs, and siRNA-CXCR4 transfected BMSCs	Sprague-Dawley rats	2 hr	24 hr post-surgery	2 × 10 ⁶ into the femoral vein	Increase in CXCR4-BMSCs surrounding the infarct compared to nontransfected and siRNA-CXCR4 transfected BMSCs	CXCR4-transfected BMSCs increased the perinfarct capillary bed, reduced the infarct size, and improved the functional outcome compared to nontransfected and siRNA-CXCR4-transfected BMSCs	141
r-ASCs	No	Sprague-Dawley rats	3 hr	0, 12, and 24 hr after stroke onset	2 × 10 ⁶ intravenously	Migration toward the lesion, questionable differentiation toward endothelial cells	Reduction of infarct region. Improvement in sensorimotor function, upregulation of CXCR4 and SDF-1. Decreased apoptosis in infarct region	132
r-BMSCs and r-ASCs	No	Sprague-Dawley rats	60 min	30 min after reperfusion	2 × 10 ⁶ into the femoral vein	Migration of transplanted cells toward the lesion was not observed	BMSC and ASC administration improves functional recovery independent of reducing the infarct volume and cell migration. Treated groups show higher cell proliferation, oligodendrogenesis, synaptogenesis, and angiogenesis markers	60
m-BMSCs and m-ASCs	No	C57BL/6J mice	90 min	Immediately after reperfusion	1 × 10 ⁵ ASCs or BMSCs into the tail vein	n/a	ASC administration attenuated ischemic damage. Incomplete ASC incorporation in the brain. HGF and angiotensin-1 expression was significantly increased in ASC-treated mice compared with the BMSC group	139
h-BMSCs	No	Sprague-Dawley rats	90 min	7 days post-surgery	3.4 ± 1.2 × 10 ⁶ into the saphenous vein	Donor cells accumulate in the ischemic hemisphere, but also in the spleen and lungs	IV-injected 99mTc-HMPAO-labeled MSCs home to the ischemic lesion, but also accumulate in the lungs and the spleen	145

Continued

Table 1. Continued

Stem cell type	(Pre)differentiation and/or treatment	Species	Occlusion time	Time of transplantation	Cell dose and location of transplantation	Fate of transplanted cells	Outcome	Reference
h-BMSCs	No	Sprague-Dawley rats	Hours, no details	60 days post-surgery	4 × 10 ⁶ in the jugular vein	Donor cells preferentially migrate to the spleen, up to 12 days postinjection	Significant reduction in striatal and perinfarct area. Reduced loss of hippocampal neurons, significant reduction in MHC-II activated inflammatory cells in gray and white matter. TNF- α expression in the spleen was decreased	134
r-BMSCs	No	Sprague-Dawley rats	90 min	1, 4, or 7 days post-surgery	3 × 10 ⁶ intravenously	Cells transplanted 1 day after stroke migrated toward the cortex, cells transplanted after 4 days or 7 days migrated to the striatum	Functional recovery (mNSS score) was highest when cells were transplanted 1 day after surgery. This was correlated with a time-dependent expression of SDF-1 and MCP-1 between ischemic regions	133
r-BMSCs	No	Aged Wistar rats	2 hr	1 month after surgery	3 × 10 ⁶ intravenously	Preferential differentiation toward astrocytes (13%) over neurons (6%). Survival of donor cells was not quantified	Significant sensorimotor and general neurological recovery after cell compared with control animals. BMSC treatment reduced scar thickness, and increased the number of proliferating cells and oligodendrocyte precursors. SDF-1 is upregulated in the ischemic boundary zone after stroke. BMSCs express CXCR4	146
r-BMSCs from SHR-SP rats	No	Aged SHR-SP rats	Permanent	30 days before stroke onset	5 × 10 ⁵ into the tail vein	No direct transplantation, injected donor cells prior to MCAO	SHR-SP BMSCs transplantation increased microvasculature density in the perinfarct zone, reduced ischemic brain damage, and improved neurologic function. Rejuvenation of bone marrow from aged rats with young cells enhanced the ischemic response at the level of endothelial/vascular activation	152
h-UTCs	No	Aged Wistar rats	Permanent	24 hr post-surgery	1 × 10 ⁷ cells/kg into the tail vein	Very few donor cells present at lesion site, no reactivity for MAP2 or GFAP	IV administration of hUTC improved neurological functional recovery without reducing infarct size, increased progenitor cell proliferation and vessel density in the ischemic boundary zone, and enhanced synaptogenesis	148
r-BMSCs, h-BMSC	No	Aged Sprague-Dawley rats	3 hr	6 hr post-surgery	1 × 10 ⁶ cells/kg into the tail vein	1% migrates toward the lesion	Daily treatment with G-CSF improved neurological function. G-CSF + BMSC transplantation stimulated angiogenesis in the infarct core but did not further improve neurological function or infarct volume size	149

IGF-1, insulin-like growth factor 1; HGF, hepatocyte growth factor; SHR, spontaneous hypertensive rats; SHR-SP, stroke-prone SHR; ^{99m}Tc-HMPAO, ^{99m}Tc-hexamethylpropylene amine oxine; hUTCs, Human umbilical tissue derived cells; EV, extracellular vesicles; MCP-1, monocyte chemoattractant protein-1; TNF- α , tumor necrosis factor α ; prefix h, human; m, mouse; r, rat.


1
2 the ideal candidates to stimulate repair in the central nervous system (CNS) due to their neu-
3 rogenic differentiation potential and predisposition.^{50–53} Promising results have already been
4 achieved with human NSCs in animal models of neurological disorders, including stroke.^{8,54}
5 However, there is a need for alternative stem cell sources with regenerative potential due to
6 ethical considerations with regard to the isolation of NSCs from embryonic or fetal tissue
7 together with isolation and culturing complications of adult NSCs.^{55,56} These alternative stem
8 cells need to be able to reconstitute the lost tissue or stimulate endogenous repair. The most
9 promising alternatives for NSC that are of nonembryonic or nonfetal origin are mesenchymal
10 stem cells (MSCs), induced pluripotent stem cells (iPSCs), and bone marrow mononuclear cells
11 (BMMNCs). These stem cell sources have shown to possess regenerative effects on the brain
12 and allogeneic transplantation potential^{57–63} Moreover, these stem cell types can be obtained by
13 means of minimal invasive procedures thereby reducing donor site morbidity during isolation.
14 Although it remains a topic of debate whether MSCs possess NPC properties, several studies
15 have reported the ability of subtypes of MSCs to acquire neuronal features following exposure
16 to the proper environmental stimuli.^{64–66} In addition to the discussion of which stem cell source
17 is most suitable for stroke research, different animal models such as the middle cerebral artery
18 occlusion (MCAO) model or photothrombotic stroke model are available to induce stroke in
19 an experimental setting, each with their own strengths and weaknesses.^{67–69}

20 Multiple mechanisms have been proposed for stem cell mediated therapies, including brain
21 protection, cell replacement, immunomodulation, and promoting both brain plasticity and
22 angiogenesis in damaged brain regions (Fig. 2).⁴⁹ Interestingly, these mechanisms are mainly
23 thought to be mediated by the effect of the stem cell secretome on endogenous stem cells
24 and on the host microenvironment instead of directly replacing the lost cells,^{59,70,71} although
25 encouraging results have also been achieved with cell replacement studies.^{57,58} Therefore, the
26 transplanted cells can be seen as a vehicle for sustained growth factor delivery at the stroke
27 lesion, which can also respond dynamically to changes in the local microenvironment as will
28 be discussed next and into more detail in the following sections.

29 Stem cell mediated neuroprotective effects have been observed in in vitro and in vivo
30 models of neurological disorders.^{72–74} These neuroprotective effects are mainly attributed to
31 the soluble factors secreted by the stem cells. In addition to the development of protective
32 therapies, interventions aiming at the directed recruitment and differentiation of NSCs to the
33 site of injury are considered. It is known that NSCs are present in the subventricular zone
34 and dentate gyrus of the hippocampus in the adult brain.^{53,75,76} Moreover, following ischemic
35 stroke, endogenous NSCs differentiate into neurons and migrate toward the site of stroke injury
36 and contribute to brain repair.^{77,78} A determining factor in the directed migration of neurons
37 is stromal cell derived factor $\alpha 1$ (SDF-1) and its receptor CXCR4.^{78,79} This SDF-1/CXCR-4
38 axis has been shown to act as an inflammatory mediator after acute cerebral ischemia^{80,81} but
39 has also been shown to play an important role in CNS development,⁸² has a modulating effect
40 on different subsets of neurons,⁸² and has strong effects on cell migration, axon guidance, and
41 angiogenesis in the postacute phase of stroke.^{79,82}

42 Unfortunately, the endogenous repair by NSCs is insufficient to completely replace the
43 lost tissue. Therefore, in addition to exerting a protective effect on the brain, novel cell based
44 therapies are focusing on improving the recruitment of and repair by endogenous NSCs and
45 supporting cells.⁸ Direct cell replacement by neurons derived from stem cells themselves is also
46 a route that is being considered, although it is uncertain whether the transplanted stem cells
47 are able to survive and adequately integrate into the host brain.^{57,59} It has been suggested that
48 damaged areas in the brain can only be successfully reconstituted by the equivalent homotopic
49 neurons, which stresses that adequate pretransplantation targeted differentiation of stem cells
50 grafts toward specific types of neurons is required for direct cell replacement by the stem cells
51 themselves.^{83–85} For example, it has been shown that grafting cortical donor tissue into the

10 • GERVOIS ET AL.



COLOUR ONLINE,
B&W IN PRINT

Figure 2. Mechanisms of action of cell-based therapies in ischemic stroke. The poststroke microenvironment can be modulated by exogenously delivered stem cells by multiple mechanisms to trigger tissue repair. Stem cells can contribute to poststroke recovery by stimulating the migration of endogenous NSCs toward the stroke lesion, where proliferation and differentiation toward replacement neurons can be triggered. In addition, transplanted stem cells are thought to be able to replace the lost neurons themselves in addition to stimulating host NSCs. Moreover, the formation and attraction of new blood vessels toward the ischemic lesion and the stimulation of synaptogenesis and synaptoplasticity contributes to host repair. In addition to directly stimulating the formation of new brain tissue, the degradation of existing cells in, for example, the ischemic penumbra is inhibited by neuroprotective mechanisms such as ROS scavenging by the transplanted cells. Immunomodulatory effects are also observed and include the inhibition of neutrophil activation and migration, effector T-cell and B-cell inhibition, reducing the activation and attraction of peripheral dendritic cells, and stimulating the M2 microglial phenotype. These effects are predominantly caused by the soluble factors released by the stem cells, but also cell–cell interactions appear to play a role. Image was created using Servier Medical Art.

damaged motor cortex reestablished cortical and even subcortical circuitry,⁸⁴ a feature that was not observed when using heterotopical tissue such as occipital cortex.⁸⁶ More recently, a study by Michelsen et al. demonstrated that in vitro differentiated mouse embryonal stem cell (ESC) derived visual cortical neurons were able to reestablish connections with the damaged visual cortex with reciprocal axonal projections and synaptic integration.⁸³ Interestingly, grafting these cells in the damaged motor cortex or ESC-derived motor neurons in the damaged visual cortex did not lead to graft integration.⁸³ In addition, targeted differentiated iPSCs to pyramidal cortical neurons have been shown to integrate in the host circuitry after transplantation into the neonatal mouse brain⁸⁷ and have been used in preclinical stroke research,⁵⁷ as will be discussed later. Similarly, it has been suggested that potential donor cells for Parkinson's disease should be of the correct nigral dopaminergic neuron phenotype to improve functional engraftment with the appropriate targets.⁸⁵ Another theory in which cell-based therapies are believed to improve the functional outcome in stroke is by directly inducing brain plasticity after the ischemic insult. Although studies report the functionality of transplanted cells in the endogenous neuronal circuitry, these effects are thought to be mainly mediated by promoting the formation of new synapses between existing neuronal cells and not by functionally integrating into the host neuronal network.^{57, 58, 87, 88}

A key concept in the regeneration of lost tissue is establishing adequate blood supply to the regenerating tissue. Without proper vascularization that provides oxygen and nutrients, the newly formed neuronal tissue will be unable to survive. Previously, it has been shown that stem cells can form vascular structures in vitro and secrete proangiogenic factors that can positively influence the growth of blood vessels in vitro and in vivo.⁸⁹⁻⁹¹ Therefore, stimulating angiogenesis is another mechanism by which cell-based therapies can influence stroke outcome. Remarkably, revascularization appears to be the main mechanism by which BMMNCs are able to ameliorate the disease outcome.⁶¹⁻⁶³

In addition to protecting damaged neurons, restoring the lost neuronal circuitry and blood supply, stem cells have been shown to be able to mediate the immune response.^{92,93} The mechanisms of these immunomodulating properties include influencing the activation state of monocytes, natural killer cells, B cells, T cells, and neutrophils. Stem cells were also shown to mediate immunoglobulin release from plasma cells and upregulate the amount of regulatory T cells.⁹²⁻⁹⁴ However, it is important to take into account that in ischemic stroke, one of the most common causes of stroke-related morbidity is severe systemic immunosuppression, making patients susceptible to infections.⁴¹ Therefore, additional systemic immunosuppression by cell-based therapies could worsen stroke outcome. Fortunately, no adverse effects on systemic cytokine levels were observed following stem cell transplantation in a rat model of stroke.⁹⁴

Despite the promising results with BMMNCs in ischemic stroke from preclinical studies,^{61-63,95,96} in vitro evidence of the effect of BMMNCs on the above-mentioned mechanisms is scarce.^{97,98} Therefore, this review will focus on MSCs and iPSCs as readily available sources of stem cells and compare their efficacy and potential to ameliorate the disease outcome in animal models of ischemic stroke. In addition, novel imaging strategies allowing in vivo tracking of transplanted cells and noninvasive evaluation of brain repair following stroke will be discussed.

2. MESENCHYMAL STEM CELLS AS A THERAPY IN STROKE

MSCs, initially discovered in the bone marrow stromal cells (BMSCs) by Friedenstein et al. in the late 1960s,⁹⁹ were later found to be able to differentiate toward cells producing mesenchymal tissues including bone-forming osteoblasts, cartilage-producing chondroblasts, and adipocytes.¹⁰⁰ In addition to bone marrow, MSCs have been isolated from a varying range of other tissues including but not limited to adipose tissue (ASCs), Wharton's Jelly in the umbilical cord (UMSCs), umbilical cord blood, and dental tissues.¹⁰¹⁻¹⁰⁵ Additional research into the differentiation capacity of MSCs suggested that these cells were able to differentiate toward hepatocytes,¹⁰⁶ cardiomyocytes,¹⁰⁷ and neuron-like cells.¹⁰⁸ The presence of MSCs in various easily accessible and available donor tissues such as the dental pulp and adipose tissue makes MSCs a promising cell type for stem cell based therapies. However, the main problem in the extensive research with MSCs is the difficulty to compare study outcomes between different research groups. Research groups often have their own methods of isolating, expanding, and characterizing the cells, leading to diverging criteria to define MSCs.^{101-103,109,110} MSCs are a heterogeneous population that generally express the surface markers, CD29, CD44, CD90, CD117, and CD146, while they do not express CD34 and CD45 although subpopulations of CD45- and CD34-expressing MSCs were identified.¹¹¹

A. In vitro Evidence for the Regenerative and (Neuro)protective Potential of MSCS on the Brain

Despite interlaboratory differences in defining and culturing MSCs, researchers agreed on the multilineage differentiation potential^{100,101,112} of these stem cells and subsequently investigated

1 **12 • GERVOIS ET AL.**

2 the ability of these cells to transdifferentiate into neuronal or neural-like cells in order to obtain
3 a cell source to replace the lost tissue after ischemic stroke. Early studies that investigated the
4 neurogenic differentiation potential of MSCs were performed using BMSCs, but also hASCs
5 and dental tissue and umbilical cord derived stem cells were successfully differentiated to cells
6 with a neuronal phenotype, expressing markers such as neuronal nuclei (NeuN), microtubule-
7 associated protein 2 (MAP2), neural cell adhesion molecule, and synapsin I.^{66,108,113,114} Al-
8 though a consensus was not found between the differentiation protocols, epidermal growth
9 factor and basic fibroblast growth factor (bFGF) are thought to play an important role in in-
10 ducing MSCs toward a neuronal cell lineage.^{66,113} Subsequent maturation of the induced cells
11 was based on increasing intracellular cyclic adenosine monophosphate and protein kinase C sig-
12 naling or by specific growth factor administration.^{66,108,113–115} However, few studies performed
13 electrophysiological measurements on the differentiated cells but were able to report both
14 voltage-gated sodium and potassium currents that could be reversibly blocked by tetrodotoxin
15 and tetraethylammonium, respectively.^{66,108,113,115} Of these studies, only the studies by Wislet-
16 Gendebien et al.¹⁰⁸ and Gervois et al.⁶⁶ could demonstrate the ability of the differentiated cells
17 to generate a single action potential in neuronally differentiated BMSCs and human dental
18 pulp stem cells, respectively, demonstrating only incomplete neuronal differentiation.

19 In addition to the neuronal differentiation capacity of MSCs, researchers also investigated
20 the neuroprotective and regenerative potential of the MSC secretome. Hypoxia- and glutamate-
21 induced excitotoxicity assays were used as in vitro models for ischemic stroke. It was shown
22 that the MSC protect SH-SY5Y neuroblastoma cells against hypoxia- and glutamate-induced
23 excitotoxicity both in coculture assays and assays using conditioned medium of MSCs, suggest-
24 ing paracrine effects.^{72,74,116,117} Although the influence of the MSC secretome on NSC survival
25 and/or differentiation is not evaluated in vitro, the MSC secretome has been shown to stim-
26 ulate neurite outgrowth in dorsal root ganglia (DRG)^{118,119} and axotomized retinal ganglion
27 cells (RGCs),⁷³ and to enhance survival of these axotomized RGCs⁷³ and primary cortical¹²⁰
28 and dopaminergic neurons.¹²¹ Neurotrophins/growth factors that are secreted by MSCs include
29 glial-derived neurotrophic factor, neurotrophin-3 (NT-3), nerve growth factor (NGF), and brain-
30 derived neurotrophic factor (BDNF).^{66,73,118–122} These factors are believed to play an important
31 role in neurite outgrowth of DRGs^{118,119,122} and axotomized RGCs.⁷³ Moreover, these factors
32 are also suggested to protect RGCs from neurodegeneration after axotomization,⁷³ cortical
33 neurons from nitric oxide exposure and withdrawal of trophic support,¹²⁰ and dopaminergic
34 neurons from 6-hydroxy-dopamine.¹²¹

35 As mentioned previously, another key concept in promoting brain regeneration after is-
36 chemic stroke is stimulating revascularization of the regenerating tissue. Therefore, the influence
37 of the MSCs was not only investigated in (neuro)protective or neurite outgrowth assays but
38 also the ability of the MSCs to stimulate angiogenesis was evaluated. These studies showed
39 that MSCs are able to stimulate tube formation and endothelial cell migration, enhance wound
40 healing, and improve blood vessel formation in the chorioallantoic membrane assay.^{89,123,124}
41 These proangiogenic properties of MSCs were attributed to the soluble factors that are secreted
42 by the cells (Table I in Refs. [89] and [90]) Furthermore, it was shown that MSCs protect en-
43 dothelial cells against hypoxia-induced cell death.¹²⁵ Several studies also suggest that MSCs not
44 only promote angiogenesis by paracrine effects but that these cells are also able to differentiate
45 into endothelial cells (Table II in Ref. [89]).

46 Although neurogenic differentiated MSCs express neuronal markers, differentiation of
47 these cells toward mature neurons appears limited, as only immature electrophysiological pro-
48 files can be generated from these cells. Nonetheless, MSCs show great promise in vitro as
49 shown by the neuroprotective and proangiogenic effects of the MSC secretome. Therefore,
50 several studies have transplanted MSCs into animal models of ischemic stroke and evaluated
51 the outcome, as will be discussed below.

Table II. Preclinical Studies with Neurogenic Preadifferentiated iPSCs

Stem cell type	(Pre)differentiation and/or treatment	Species	Occlusion time	Time of transplantation	Cell dose and location of transplantation	Fate of transplanted cells	Outcome	Reference
h-iPSCs	iPSC-derived It-NES cells	Nude Rats	30 min	48 hr post-surgery	2 × 10 ⁵ intrastriatal or 1.5 × 10 ⁵ intracortical in two sites	>50% of grafted cells survive up to 4 months posttransplantation. Intrastriatal injected cells differentiate toward neurons in preference to astrocytes (72.9% vs. 1.9%) 4 months posttransplantation). Of the intracortical injected cells, 78.5% of cells in the graft core were NeuN positive while 13.4% were DCX-reactive in the periphery	Cells grafted into the striatum survive up to 4 months, differentiate to functional neurons, and receive synaptic input from host cells. Intracortical transplanted cells survive for 4 months and differentiate into functional neurons that receive synaptic input from host cells in T-cell-deficient rats without forming tumors	58
h-iPSCs	iPSC-derived It-NES cells	C57BL/6 Mice	30 min	1 Week post-surgery	1 × 10 ⁵ intrastriatal	Approximately 10% survival after 10 weeks; 78.5% HuD-positive cells found at graft core, 13.4% DCX-positive cells in the graft periphery	Improvement of motor function, independent of long-term graft survival mediated by enhancing endothelial plasticity. Remaining grafted cells differentiate toward neurons and integrate into the host brain	58
h-iPSCs	iPSC-derived It-NES cells fated into cortical neurons and nondifferentiated iPSC-derived It-NES cells	Sprague-Dawley and nude rats	30 min	48 hr post-surgery	1.5 × 10 ⁵ intracortical in two sites	Transplanted cells survive 2 months posttransplantation. Fated cells are fewer and proliferate less than nonfated cells but express a higher percentage of the mature neuronal marker NeuN (13.7% vs. 7.1%) and the cortical marker TBR1 (2.5% vs. 1.4%)	iPSCs-derived cortical neurons survive, differentiate to functional neurons, and improve neurological outcome after intracortical implantation without tumor formation in a rat stroke model	57

Continued

Table II. Continued

Stem cell type	(Pro)differentiation and/or treatment	Species	Occlusion time	Time of transplantation	Cell dose and location of transplantation	Fate of transplanted cells	Outcome	Reference
h-iPSCs	iPSC-derived lt-NES cells	Aged Sprague-Dawley rats	30 min	48 hr post-surgery	1.5×10^5 intracortical in two sites	49.2% of the grafted cells survive 8 weeks posttransplantation, 30% of transplanted cells express DCX in the periphery, 91.3% express HuD, and 19.6% express GABA outside the graft core	Cell-grafted showed increased sensorimotor function compared to vehicle-treated rats. Transplanted lt-NES cells expressed markers of neuroblasts, mature and GABAergic neurons. Microglia activation was diminished in lt-NES grafted rats. Neuronal loss was diminished after transplantation	170
h-iPSCs	iPSC-derived NPCs	Sprague-Dawley rats	90 min	1 Week post-surgery	1×10^5 intrastriatal	Survival not quantified, grafted cells express Sox2, nestin, Pax6, and extend MAP-2 expressing processes into the perilesional parenchyma	Grafted iPSC-NPCs initially exert trophic effects on host brain structures, followed by iPSC-NPCs integration into the host brain	169
h-iPSCs	iPSC-derived NSCs	Sprague-Dawley rats	2 hr	Immediately after reperfusion	1×10^6 intrastriatal	Transplanted cells survived and migrated into the damaged host tissue and express nestin (51.4%) and beta III tubulin (44.3%)	Engrafted cells survive, migrate, and differentiate toward neuronal cells. Transplanted cells improved behavioral and sensorimotor function	164
h-iPSCs	iPSC-derived NSCs	C57BL/6J mice	1 hr	24 hr post-surgery	1×10^5 intrahippocampal	Engrafted cells migrated toward the site of injury. Survival % not quantified	Improved motor and sensorimotor function in graft-receiving animals. Mechanisms of action include a decrease in proinflammatory markers, adhesion molecules, and microglial activation BBB damage was attenuated	172
h-iPSCs	iPSC-derived NPCs	Wistar rats	30 min	1 Week post-surgery	2.5×10^5 intracerebral	Double amount of cells in the graft of which 41% were beta III tubulin/MAP2 positive; 5% of the grafted cells expressed GFAP	No migration of cells toward the lesion. No significant difference in behavioral recovery. Tumor formation was not present	171
h-iPSCs	Unclear	C57BL/6N and NSG-mice	Photobotic stroke	1 Week post-surgery	1×10^5 iPSC in the stroke cavity with or without hyaluronic acid hydrogel	38% versus 30% survival of cells with or without hydrogel 1 week after transplantation in NSG-mice. Grafted cells form DCX-positive neuroblasts	No assessment of functional recovery. Tumor formation was not evaluated	173

lt-NES, long-term expandable neuroepithelial-like stem cells; NSG-mice, NOD scid gamma immunodeficient mice; prefix h, human; m = mouse.

B. MSCS as a Therapy for Stroke *In vivo*

Due to the encouraging *in vitro* results of MSCs in protecting damaged neurons and stimulating revascularization in addition to secreting multiple soluble factors, the potential of different subtypes of MSCs to ameliorate stroke outcome after transplantation was evaluated *in vivo* (Table I). These studies evaluated the functional outcome after transplantation with a variety of behavioral tests. These tests include global neurological assessments to evaluate the disease severity such as the Bederson test and the modified neurological severity score (mNSS). In addition, specific motor, sensorimotor, and cognitive tests were performed. For detailed information on behavioral and disease severity tests in animal models of stroke, see Schaar et al.¹²⁶ Taken together, the studies that report an improvement in general neurological, sensorimotor, motor, or cognitive function are described in Table I.

The proposed underlying mechanisms responsible for the improvement in stroke outcome suggested by the studies listed in Table I are diverse. One of the possibilities was that the transplanted cells migrated toward the stroke lesion and differentiated locally toward neurons, establishing new connections with the host environment,¹²⁷ although the majority of the *in vivo* studies using MSCs as a therapy for stroke support paracrine mediated brain regeneration.^{59, 128, 129} Regardless of the administration route, the fate of the transplanted cells was tracked using markers such as DiI,^{60, 130–133} DiR,¹³⁴ q-dot¹²⁷ or bromodeoxyuridine (BrdU)¹³⁵ incorporation prior to transplantation; LacZ¹³⁶ or GFP transduction;^{59, 70, 137–142} iron particle^{60, 143} or radionuclide labeling;^{144, 145} *in situ* hybridization with the Y chromosome when male donors were used in a female host^{146, 147}; or antibodies directed against human mitochondria or human nuclei when human MSCs were grafted in a rodent stroke model.^{71, 128, 148, 149} While the listed studies could observe improvement of stroke outcome after transplantation, the amount of engrafted cells that was present in the stroke lesion was limited. Pioneered by Zhao et al., intracranial transplantation of MSCs showed that MSCs migrated toward the brain infarct region and were able to survive in the host brain and promote functional recovery.⁷⁰ Additional studies showed that although they were present only in low numbers,^{59, 71, 130, 137, 138} the transplanted cells locally differentiated toward neural cells with a predisposition toward astroglial cells in preference to neurons.^{59, 130, 137} Despite the local delivery of the transplanted cells in the stroke lesion, functional integration and replacement of the lost neural circuitry does not appear to be the mechanism of action of intracerebral transplanted MSCs to improve stroke outcome. The results of these studies suggest that the soluble factors secreted by the MSCs are the main actors in improving the functional outcome.^{59, 70, 130, 137} After cerebral transplantation of the MSCs, improved angiogenesis,^{71, 130, 138} increased neuronal activity,⁷¹ reduced loss of periinfarct cells,¹³⁷ and immunomodulatory effects¹⁴² were observed. After transplantation, the local levels of soluble factors such as BDNF, vascular endothelial growth factor (VEGF), bFGF, and angiopoietin-2 were elevated.^{71, 130} In order to increase the regenerative potential of intracerebral administered MSCs, several alternative research approaches were investigated. For example, umbilical cord matrix stem cells were cultured in the presence of conditioned medium obtained from a 5-day culture of rat-derived neural cells. However, this approach did not lead to an additional improvement of stroke outcome.⁷¹ Alternatively, it was shown that hypoxic preconditioning of MSCs promotes the survival, migration, and homing of the transplanted cells toward the ischemic lesion compared to nonpreconditioned cells.¹⁴⁰ Transplantation of these cells also leads to an additional functional improvement after intranasal administration, which is assumed to be mediated by an increase in the expression of migration-related proteins such as CXCR4 and MMPs in the hypoxic preconditioned stem cells.¹⁴⁰

Intracerebral administration of MSCs is an invasive procedure and can lead to iatrogenic damage. Therefore, systemic administration of the transplanted cells via the arterial or venous route was considered. Intraarterial administration (IA) of MSCs in stroke has shown beneficial

1 16 • GERVOIS ET AL.

2 effects in animal stroke models. Interestingly, IA-transplanted cells are able to cross the BBB
3 after ischemic stroke and migrate toward the stroke lesion. Even though MSCs were found
4 in the core and periinfarct zone and expressed both astrocyte and neuronal markers,^{128,131,147}
5 paracrine mechanisms of action of IA-delivered MSCs are thought to be responsible for the
6 enhanced stroke outcome, although integration into the host brain was observed but not
7 functionally confirmed.¹²⁷ At the ischemic boundary zone, MSCs are only present in low
8 numbers but were found to enhance axonal sprouting and remyelination,¹⁵⁰ angiogenesis,
9 and BDNF production while decreasing MMP-9 levels and suppressing microglial activity.¹²⁸
10 Furthermore, BMSC administration into aged rats showed a sustained effect and donor cell
11 survival up to 1 year after transplantation.¹⁴⁷

12 Another route of MSC transplantation is via the venous system. Remarkably, MSCs
13 that were intravenously (IV) delivered, improved the functional outcome after stroke with-
14 out MSCs being observed in the ischemic brain or in lower numbers than in IA or intrac-
15 erebral transplantation.^{60,135,139,148,149,151} Despite being almost absent in the ischemic brain,
16 IV transplantation of MSC induced an increase in periinfarct zone microvasculature density
17 and the expression of proangiogenic factors^{60,135,139,148} and improved oligodendrogenesis and
18 synaptogenesis.^{60,148,152} Remarkably, when BMSCs were transplanted IV in a chronic stroke
19 model 60 days after surgery, the beneficial effects of the graft were attributed to their im-
20 munomodulatory effects on the stroke lesion and on the spleen where the cells preferentially
21 homed toward.¹³⁴ Another successfully applied therapeutic approach was to transduce BM-
22 SCs with an adenoviral vector for PIGF¹³⁶ or a lentiviral vector for CXCR4,¹⁴¹ which further
23 improved the functional outcome after stroke compared to nontransduced BMSCs. The SDF-
24 1/CXCR4 interaction in chemotaxis was found to be stimulated after intravenous administra-
25 tion of MSCs and plays an important role in MSC-homing toward the stroke lesion.^{133,141,146}
26 In accordance with the other routes of administration, the paracrine effects of the transplanted
27 cells appear to be responsible for the posttransplantation effects on stroke outcome. This was
28 supported by a recent study by Doepfner et al., who showed that repeated IV administration
29 of extracellular vesicles produced by BMSC led to a similar improvement as injecting the stem
30 cells themselves.¹²⁹ Moreover, this study also showed reduced poststroke immunosuppression
31 after EV transplantation.

32 This overview shows a variety of studies that were performed using subtypes of MSCs and
33 different administration routes. Although similar results were achieved with different subtypes
34 of MSC, it remains debatable which is the most suitable source for MSC. Moreover, MSCs
35 showed great promise in replacing the neuronal tissue after transplantation due to their in
36 vitro neurogenic differentiation potential. However, cell replacement could not be proven in
37 vivo and it is assumed that paracrine factors are responsible for the beneficial effects after
38 MSC injection, which mainly stimulate angiogenesis and protect the host environment against
39 additional damage without adequately stimulating endogenous neurogenesis or replacing lost
40 neurons. Therefore, additional stem cell sources are considered that are able to differentiate to
41 functional mature neurons in vitro and can potentially improve stroke outcome more effectively
42 than MSCs.

43
44 **3. INDUCED PLURIPOTENT STEM CELLS AS A THERAPY IN STROKE**

45
46 In 2006, Takahashi and Yamanaka successfully transformed murine and in 2007, human fi-
47 broblasts to pluripotent stem cells by retroviral transduction with the transcription factors
48 Oct3/4, Sox2, Klf4, and c-Myc, the so-called Yamanaka factors, allowing the formation of
49 a patient-specific source of pluripotent stem cells.^{153,154} These iPSCs are able to differentiate
50 toward tissues from all three germ layers in vitro and in vivo as proven by teratoma formation
51 upon subcutaneous transplantation of iPSCs.

The ability of iPSCs to differentiate into tissues of all three germ layers offers numerous potential therapeutic approaches. However, the main drawback in using iPSCs for transplantation studies in stroke research is that these cells, such as human ESCs, form teratomas when injected in an undifferentiated pluripotent state, with little to no improvement of the disease outcome.^{155,156} Therefore, iPSCs are irreversibly predifferentiated in vitro in order to minimize tumor formation and improve the functional outcome as only the undifferentiated iPSCs form teratomas. However, a recent study by Choi et al. showed that iPSC-derived NPCs reactivate the silenced exogenous retroviral genes caused by a downregulation of DNA methyltransferases during differentiation and can return toward their pluripotent and thus tumorigenic state.¹⁵⁷ Moreover, nondifferentiated iPSCs can remain present within an iPSC-derived progenitor cell pool, which showed teratoma formation after subcutaneous transplantation. This teratoma formation was not observed when fully committed iPSC-derived cells were transplanted.¹⁵⁶ This study by Liu et al. was supported by Fu et al., who demonstrated that residual nondifferentiated iPSCs could not be eliminated by extended cell differentiation.¹⁵⁸ Therefore, adequate full neuronal commitment monitoring prior to transplantation is advised and studies that used nondifferentiated iPSCs in stroke research will be left out of this overview.

A. In vitro Evidence for the Regenerative and (Neuro)protective Potential of iPSCs on the Brain

The multilineage differentiation potential of iPSCs makes these cells an attractive alternative for NSCs as a source for cell-based therapies in neurological disorders. Neurogenic differentiating iPSCs follow similar developmental principles as hESC-derived neurons, although the neural differentiation efficiency can differ between different iPSC lines.¹⁵⁹ Targeted differentiation of iPSCs toward neuronal subtypes has been achieved by various research groups, using different approaches to generate a variety of neuronal cells including medium spiny neurons,¹⁶⁰ dopaminergic neurons,¹⁶¹ motor neurons,^{159,162} nociceptors,¹⁶³ and pyramidal cortical neurons.^{57,87} Furthermore, it was shown that these neuronally differentiated cells are capable of repeated action potential firing, suggesting advanced maturation.^{57,87,159,161,163} More importantly, this targeted differentiation of iPSCs toward specific neurons is of high importance and provides additional regenerative potential as it has been suggested that damaged brain areas can only be successfully repaired by the neurons corresponding to the damaged area, as discussed earlier.^{83–85} Similar to MSCs, various differentiation protocols were developed to induce neurogenic differentiation of iPSCs. The protocols that were most successful are based on retinoic acid and sonic hedgehog signaling or based on SMAD inhibition using Noggin, which blocks SMAD signaling by the transforming growth factor beta superfamily of signaling proteins.^{87,159,163,164}

In contrast to MSCs, where the paracrine effects of the stem cell secretome have been investigated thoroughly, there is a lack of stroke-related in vitro evidence of the (neuro)protective, regenerative, and angiogenic properties of iPSCs. Among the few studies that investigated the secretome of iPSCs, it was shown that iPSCs locally enhanced the production of proangiogenic factors. However, these studies did not include an in vitro secretome analysis or an in vitro evaluation of the angiogenic properties of the iPSC secretome.^{165,166} Although the paracrine effects of the iPSC secretome on angiogenesis were not evaluated in vitro, several studies have reported endothelial cell differentiation of iPSCs,^{91,167,168} which opened up the possibility that transplanted iPSCs can directly contribute to establish new blood vessel formation in the damaged brain.

Despite the lack of in vitro evidence for paracrine-mediated regeneration, the successful differentiation of iPSCs toward endothelial cells, neuronal precursors, and mature neuronal subtypes encouraged the use of iPSCs in animal models of ischemic stroke.

1 18 • GERVOIS ET AL.

2 **B. Neurogenic Predifferentiated iPSCs in In vivo Stroke Models**

3
4 Contrary to MSCs, iPSCs that were transplanted in animal stroke models were neuronally
5 differentiated prior to engraftment in order to avoid teratogenicity (Table II). Moreover, no
6 reports are available in which iPSCs are administered IV or IA. Information about mechanisms
7 of action of iPSCs in ameliorating stroke outcome is therefore only available for intracranially
8 transplanted neurogenic predifferentiated iPSCs.

9 When considering the overall results of the behavioral tests performed after iPSC trans-
10 plantation after stroke, improvements in general neurological score,¹⁶⁹ motor,^{57,58,164,169,170}
11 sensorimotor,^{57,58,164,169,170} and cognitive function¹⁶⁴ were observed. On the contrary, Jensen
12 et al. did not observe an iPSC-mediated improvement in behavioral function.¹⁷¹

Q6 13 Underlying mechanisms of action of the transplanted iPSCs include migration and local
14 neuronal maturation of the transplanted iPSCs,^{57,58,164,170} synaptic integration into the host
15 brain^{57,58} but also paracrine effects are considered to play a role.¹⁶⁹ The transplanted iPSCs were
16 predifferentiated toward long-term expandable neuroepithelial-like stem cells,^{57,58,170} fated to-
17 ward cortical neurons⁵⁷ or toward neuronal precursor cells.^{164,169,171} Transplanted iPSCs were
18 traced with antibodies directed against human nuclei, cytoplasm or mitochondria,^{57,58,169–172}
19 DiI labeling,¹⁶⁴ or by using GFP positive iPSCs.^{57,58,173} The engrafted cells survived up to 10
20 weeks after transplantation but the percentage of surviving cells varied considerably between the
21 different studies, which can be caused by several factors. For example, the host strain (i.e., nude
22 rats vs. immunocompetent rats) and species^{174–177} can have an influence on the stroke outcome
23 and cell survival rates.⁵⁸ Moreover, the altered stroke microenvironment in aged animals might
24 be less favorable for cell transplantation.^{170,178} Another factor that can influence cell survival
25 and amelioration is the time after stroke onset; the transplantation is performed as cell trans-
26 plantation at later time points exposes the cells to the established immune response,¹⁷⁹ whereas
27 cell transplantation early after stroke onset or too close to the ischemic core can expose the
28 cells to limited blood supply, oxidative stress, and trophic factor deficiency.¹³ Remarkably, both
29 engrafted fated cortical neurons derived from long-term expandable neuroepithelial-like stem
30 cells and their nonfated counterparts, which were transplanted 48 hr after stroke onset, could be
31 detected 2 months posttransplantation in the damaged rat cortex in the study by Tornero et al.⁵⁷
32 The transplanted cells differentiated locally toward neuronal cells as shown by the expression
33 of doublecortin (DCX) at the periphery of the graft. The expression of HuD and NeuN was
34 found at the core of the graft,^{57,58,170} which was increased by differentiating the long-term
35 expandable neuroepithelial-like stem cells toward cortical neurons prior to transplantation.⁵⁷
36 A study by Lam et al. demonstrated that 1-week survival of the iPSC grafts is enhanced—
37 although not significant—in photothrombotic stroke by delivering the cells in a hyaluronic
38 acid hydrogel. However, despite claiming to use iPSC-derived NPCs, the neuronal character-
39 istics were not described, which might have had an impact on the survival rate. Additionally,
40 transplantation of the cells in the hyaluronic acid hydrogel favored DCX-positive neuroblast
41 formation 1 week after transplantation.¹⁷³ The study by Yuan et al. also observed the expres-
42 sion of nonmature neuronal markers of the engrafted cells such as beta tubulin and nestin.¹⁶⁴
43 Moreover, a low fraction of cells differentiated toward astrocytes, as shown by GFAP expres-
44 sion. Nonetheless, this study showed a preferential neuronal differentiation of engrafted iPSC-
45 derived neuronal precursors after transplantation,¹⁶⁴ which is supported by the study of Jensen
46 et al.¹⁷¹

47 To examine whether the transplanted iPSCs contributed to the synaptic network in the
48 host brain, various methods to determine synaptic integration were applied. One method
49 that was used to determine synaptic integration was by retrograde tracing of fluorogold in
50 the ipsilateral globus pallidus, 9 weeks after the injection of iPSCs.⁵⁸ In this study by Oki
51 et al., it was shown that a small fraction of the transplanted iPSCs were fluorogold positive,

1
2 meaning that striatal transplanted iPSCs extended projections toward the globus pallidus. Other
3 strategies to identify axonal projections of transplanted iPSCs used antibodies directed against
4 donor specific cytoplasmic⁵⁷ or surface markers.¹⁶⁹ The study by Polentes et al. observed iPSC-
5 derived axonal projections from the site of engraftment that was in the lesion cavity located
6 in the lateral quadrant of the striatum to the striatum and globus pallidus after 1 month and
7 these projections extended into the substantia nigra 1 month later. A few fibers were found
8 in the corpus callosum.¹⁶⁹ The study by Tornero et al. that compared long-term expandable
9 neuroepithelial-like stem cells and long-term expandable neuroepithelial-like stem cells fated
10 toward cortical neurons observed a higher density of projections extending from the site of
11 engraftment in the cortex over the corpus callosum in fated cells compared to the nonfated
12 cells.⁵⁷ In addition to axonal projections, the engrafted cells were shown to be functionally active
13 up to 6 months after transplantation, as was shown by whole-cell patch-clamp recordings in
14 acute brain slice preparations.^{57,58}

15 In addition to cell-replacement mechanisms and synaptic integration, paracrine-mediated
16 improvement of stroke outcome is another mechanism by which transplanted iPSCs can exert
17 their effect. The study by Oki et al. showed that functional recovery was independent of long-
18 term engraftment, suggesting a beneficial effect early after transplantation.⁵⁸ Furthermore,
19 they showed that VEGF reactivity was upregulated following transplantation in astrocytes and
20 the blood vessel wall of the damaged brain as early as 1 week after transplantation. How-
21 ever, 9 weeks after transplantation, when animals receiving the cell graft showed significant
22 functional improvement, no difference in vessel length density and immunoreactivity for the
23 endothelial marker CD31 was observed between animals that were injected with iPSCs com-
24 pared to vehicle-injected animals. An additional study by this research group detected only
25 weak expression of VEGF in the grafted cells and blood vessel walls of the damaged brain
26 8 weeks posttransplantation. VEGF-expressing astrocytes were not observed.¹⁷⁰ This suggests
27 that VEGF signaling is important in early recovery after stroke and can trigger long-lasting
28 effects in brain plasticity. In addition, it can also be postulated that VEGF signaling alone is not
29 sufficient to clarify the improved functional recovery after iPSC transplantation.¹⁸⁰ Moreover,
30 Tatarishvili et al. observed functional improvement from 1 to 4 weeks after transplantation,
31 making it inconceivable that the behavioral improvement is due to cell-mediated neuronal
32 replacement.¹⁷⁰

33 Another possible mechanism by which transplanted iPSCs can influence functional recovery
34 after stroke is by modulating the immune response. Microglia were not found to be more
35 prominent or more activated in vehicle-treated animals compared to animals that received
36 the iPSC graft 8 weeks after transplantation. However, microglia in the vehicle-treated group
37 showed a more round/amoeboid morphology compared to animals that received the iPSC
38 graft.¹⁷⁰ There is no information on how the number and activation status of the microglia
39 was affected in the early time points after transplantation. Previous studies have shown that
40 NSCs, transplanted in the cortex or striatum after stroke, can reduce the number of microglia
41 in both early as late time points after engraftment.^{54,181} In addition to these results, Chen
42 et al. previously reported an upregulation of antiinflammatory cytokines and a downregula-
43 tion of proinflammatory cytokines after nonpredifferentiated iPSC administration in the stroke
44 brain.¹⁸² More recently, a study by Eckert et al. showed that intrahippocampal transplantation
45 of iPSC-derived NSCs 24 hours after stroke onset reduced the expression of proinflammatory
46 markers, microglial activation, and adhesion molecules while attenuating BBB damage, leading
47 to a significant improvement in motor and sensorimotor function within the first week after
48 transplantation. These data support the influence of early transplantation on the host immune
49 response, leading to an improved stroke outcome.¹⁷²

50 These results show that functional integration of transplanted iPSCs in the host model
51 is achievable. However, the timing of graft-induced improvement in behavioral tests suggests

1 20 • GERVOIS ET AL.

2 that early amelioration of stroke symptoms is mediated by paracrine mechanisms and may be
3 mediated by the early influence of the graft on the host immune system. In addition, functional
4 improvement caused by the integration of graft-derived neuronal cells is expected to take longer
5 than the 8–9 weeks follow-up in the presented studies. The presented studies by Oki et al.⁵⁸ and
6 Tornero et al.⁵⁷ followed the engrafted cells for up to 6 months after transplantation and were
7 able to observe neuronal differentiation and functionality of the cells, but did not examine the
8 underlying mechanisms into further detail.

9 One of the major issues in determining the fate of the transplanted cells and determining
10 the exact onset of functional recovery is that most studies described above used traditional
11 histopathological techniques to determine the fate of the transplanted cells and to visualize
12 possible methods of action of the transplanted cells. Therefore, as will be discussed next, novel
13 techniques are being developed to allow real-time and longitudinal noninvasive imaging and
14 tracking of the transplanted cells. Furthermore, additional imaging methods can be applied in
15 the same animals to get real-time information on revascularization and reinnervation as will be
16 discussed next.

17 18 19 **4. CURRENT CHALLENGES AND FUTURE PROSPECTS OF STEM CELL BASED** 20 **THERAPIES IN STROKE: VISUALIZING DONOR CELLS AND THE INJURED BRAIN**

21
22 Traditional histopathological techniques can only be applied ex vivo. The most frequently
23 used cell trackers include the use of GFP-transduced or BrdU-labeled stem cells in addition to
24 antibodies directed against human-specific epitopes. However, it has been shown by Burns et al.
25 that thymidine analogs such as BrdU may not be a suitable marker to track donor cells due
26 to label transfer to phagocytizing cells.¹⁸³ Similar to cell tracking, morphological changes such
27 as revascularization and reinnervation, evoked by the transplanted cells, are investigated by
28 using tissue-specific antibodies or other ex vivo methods. Fortunately, noninvasive, quantifiable
29 imaging methods have been developed to track the fate of the transplanted cells and evaluate
30 the host environment, which can also be applied in humans. These methods will provide more
31 insight into the exact timing of behavioral improvements following stem cell transplantation by
32 correlating the presence or absence of the engrafted cells with morphological adaptations such
33 as revascularization and reinnervation at the injured site.

34 35 **A. Visualizing Engrafted Cells in the Injured Brain**

36
37 Noninvasive imaging methods of donor cells are based on magnetic resonance imaging (MRI),
38 positron emission tomography (PET), single photon emission computed tomography (SPECT),
39 and optical methods such as bioluminescence imaging (BLI) and fluorescence imaging (FLI).
40 The advantage of MRI compared to the other imaging methodologies is its high spatial reso-
41 lution. However, PET-, SPECT-, BLI-, and FLI-based imaging hold the advantage of a higher
42 sensitivity, although the last two are only sensitive in preclinical research. Prior to injection,
43 donor cells can be labeled either directly or indirectly. For indirect labeling, imaging reporter
44 genes are introduced into the host cells that encode for proteins or molecules that will lead to
45 the accumulation of a specific substrate or ligand within the cells in which the reporter is ex-
46 pressed. Direct labeling involves the (stable) attachment or incorporation of reporter molecules
47 into the cells after in vitro incubation (extensively reviewed in Ref. [184]). Direct pretransplan-
48 tation donor cell labels such as superparamagnetic iron oxide (SPIO) particles^{60, 185–187} and
49 radionuclides^{144, 145} have also been used to track donor cell fate, but as will be discussed below,
50 are also subjected to several disadvantages.¹⁸⁸ The different labeling strategies are illustrated in
51 Fig. 3.

Direct Labeling	Indirect Labeling
<p>+ Minimal influence on stem cell properties Ease of use – readout with FLI/MRI/PET/SPECT Biosafety risks are minimal</p> <p>- Label transfer to phagocytes Persisting signal after cell death when using SPIOs Distribution and related reduction of signal in proliferating cells Imaging window dependent on radionuclide half-life time</p>	<p>+ Readout with MRI/PET/SPECT/BLI/FLI Substrate-dependent control of imaging time-window Longitudinal imaging</p> <p>- Viral Components – biosafety issues Labour-intensive design of reporter gene constructs</p>




Figure 3. Direct and indirect labeling strategies to track stem cell fate in vivo. Direct labeling methods such as SPIOs and radionuclides have the advantage that they have been shown to have minimal effects on stem cell properties. Furthermore, biosafety risks are minimal and the labeled cells can be easily monitored with noninvasive imaging techniques such as FLI, MRI, PET, and SPECT. The disadvantages of using direct labeling strategies are the decreasing signal over time and the possibility of monitoring a nonspecific signal. These disadvantages can be circumvented by using indirect labeling methods in which reporter genes are incorporated into the donor cell genome of which the reporter protein can be visualized in a substrate-dependent manner, allowing spatiotemporal control of cell tracking. The major disadvantage of using this labeling strategy is the labor-intensive design of reporter gene constructs and biosafety issues regarding the use of viral components to transduce the donor cells. Image was created using Servier Medical Art.

SPIO particles are commonly used tracers that allow direct labeling of the donor cells for use with T2-weighted MR images without a significant effect on stem cell biology and differentiation potential of the labeled cells.^{60, 185–187} Despite the promising use of SPIOs in noninvasive cell tracking with MRI, this labeling method has several limitations.¹⁸⁹ It is impossible to discriminate between viable and dead cells as the SPIO particles remain present after the cells have died. Furthermore, macrophages and microglia present at the lesion site may phagocytize the cell fragments of dead SPIO-labeled cells, which can lead to the occurrence of nonspecific signal not originating from transplanted cells. In proliferating cells, the SPIO signal decreases after transplantation, which is aggravated by asymmetrical replication of the donor cells.¹⁸⁹

Another approach to directly label stem cells is with radionuclides that are detectable with the nuclear imaging methods PET or SPECT. These radionuclides can bind to the cell membrane (i.e., hexadecyl-4[¹⁸F]fluorobenzoate)¹⁹⁰ or can be taken up by the cell via passive diffusion through the cell membrane or via ion channels, transporters, and pumps.^{191–193} The most widely used PET-compatible radionuclide is 2-deoxy-2-¹⁸F-fluoro-D-glucose (¹⁸F-FDG). ¹⁸F-FDG has been successfully used to monitor human autologous BMSC and peripheral hematopoietic stem cell homing after myocardial infarction in a human study.^{194, 195} Survival, proliferation, and differentiation of radiolabeled MSCs is maintained after radionuclide labeling, suggesting minimal radiotoxicity in these cells.^{196–198} The two most frequently used SPECT-compatible radionuclides are oxine-bound Indium-111 (¹¹¹In-oxine) and technetium

1 22 • GERVOIS ET AL.

2 bound to hexamethylpropylene amine oxine (^{99m}Tc -HMPAO). ^{111}In -oxine and ^{99m}Tc -HMPAO
3 have a half-life of 2.8 days and 6 hr, respectively, allowing long- and short-term imaging. ^{99m}Tc -
4 HMPAO has been used as an effective tracer in a human study using labeled intraarterial-
5 injected autologous bone marrow mononuclear cells after acute ischemic stroke¹⁹⁹ and shows
6 minimal radiotoxic effects.¹⁴⁵ Although ^{111}In -oxine is an FDA-approved tracer and has been
7 used in animal models of stroke and myocardial infarction^{144,200} and human clinical studies,²⁰¹
8 it has been reported that ^{111}In -oxine binding is reversible, and that cell function and viability
9 are impaired due to radiotoxic effects.^{202,203}

10 Similar to SPIOs, direct labeling of stem cells with radionuclides also has several disad-
11 vantages, such as tracer leakage into the extracellular space that induces labeling of nontarget
12 cells.^{10,188} Furthermore, the tracer leakage to nontarget cells can be homogeneously distributed
13 to these cells that leads only to a diminishable background signal compared to the initially tar-
14 geted cells. More important is that the application window of these tracers is highly dependent
15 on the half-life of the used radionuclide, reducing the use of these radionuclides for longitudinal
16 follow-up of the labeled cells.¹⁰ Radiotoxicity does not appear to be a major hurdle when using
17 low doses of ^{18}F -FDG or ^{99m}Tc -HMPAO, but becomes a problem when ^{111}In -oxine is used as
18 a tracer and is highly dependent on the dose that is used.¹⁹⁷

19 To minimize the problem of label transfer from donor to host cells or to cope with the
20 decreasing signal after transplantation of the donor cells, other noninvasive imaging methods
21 are available. One of the possibilities is using reporter genes of which the proteins are only
22 encoded in viable cells. Well-known examples are green fluorescent protein,²⁰⁴ which can be
23 detected by FLI, and firefly luciferase (*fluc*),²⁰⁵ which catalyzes a light-emitting reaction that
24 can be detected by BLI upon oxidation of exogenously delivered luciferin. BLI-compatible
25 reporter genes have been successfully transduced in MSCs,^{206,207} and iPSCs¹⁵⁶ and have been
26 successfully used to monitor both donor cell fate²⁰⁷ as endogenous NSCs.²⁰⁸ Although these
27 optical imaging methods provide a highly sensitive technique^{209,210} to monitor cell survival and
28 proliferation, these methods are limited by a loss of spatial resolution due to light scattering and
29 by the anatomical depth of the signal-generating engrafted cells²¹¹ making them only useful for
30 preclinical research in small animals.

31 To improve the spatial resolution and to gain additional information of the engrafted cells,
32 MRI-detectable reporter gene methods have been developed, which are listed in Patrick et al.
33 (Table S1 in Ref. [212]) and reviewed in detail by Vandsburger et al.²¹¹ MRI reporter genes are
34 based on intracellular iron accumulation, enzymatic reactions, membrane-bound proteins such
35 as the biotin–streptavidin interaction, and chemical exchange saturation transfer (CEST).²¹¹
36 It should be noted that noninvasive MRI reporter gene methods that are based on visualizing
37 iron accumulation cannot distinguish between living and dead cells, once the transduced cells
38 have bound or accumulated iron.²¹¹ Although MRI reporter genes based on cell membrane
39 interactions and enzymatic reactions were developed, the most promising MRI reporter method
40 is based on CEST. CEST is based on compounds containing exchangeable protons that resonate
41 at a different frequency than bulk water protons. These protons can be selectively saturated with
42 a radiofrequency pulse and are transferred to the bulk water molecules after proton transfer,
43 attenuating the signal of the water signal.²¹³ However, to date, no stem cell tracking experiments
44 have been performed with CEST.

45 In addition to MRI-detectable reporter genes, nuclear reporter genes that can be detected
46 with PET or SPECT have been developed. These highly sensitive techniques allow for repeated
47 visualization of migration and function of donor and host cells.²¹⁴ Imaging reporter genes can
48 be subdivided into three main categories: enzymes, receptors, and transporters.¹⁰

49 The herpes simplex virus type 1 thymidine kinase gene (*HSV1-tk*) encodes the viral pro-
50 tein HSV1-TK, an enzyme that can phosphorylate nucleoside analogs that are subsequently
51

negatively charged and become entrapped in the cells.²¹⁵ HSV1-TK can phosphorylate isotope-labeled pyrimidine analogs that can subsequently be used as reporter probes in PET and SPECT imaging.¹⁹¹ Unfortunately, the use of viral proteins can potentially cause an immune response and more specific for stroke research, none of the HSV1-TK compatible tracers is able to cross the BBB.^{216,217} These disadvantages of HSV1-TK compatible tracers can be overcome by using reporter genes encoding for receptors such as the dopamine D2 receptor (D2R)²¹⁸ and human somatostatin receptor subtype 2 (hSSTr2)²¹⁹ or by reporter genes coding for transporters such as the human sodium iodide symporter (hNIS) that can transport all radioactive forms of I⁻ as well as other isotopes (i.e., Technetium-99m).²²⁰ The advantage of using D2R, hSSTr2, and hNIS is that they can be labeled with tracers able to cross the BBB and, since they are of human origin, are not likely to elicit an immune response.¹⁹¹ However, D2R and hSSTr2 have not been used for longitudinal tracking of engrafted cells. hNIS has already been successfully used to track MSCs²²¹ and iPSCs²²² in vivo but to date, no studies have been performed to trace donor stem cells after transplantation in stroke with hNIS.

In order to combine the high spatial resolution and anatomical precision of MRI with the sensitivity of radionuclide or BLI, dual-modality probes have been developed to exploit the characteristics of each imaging method^{212,223} (reviewed in Ref. [224]). For example, Patrick et al. used *Oatp1a1* as a reporter.²¹² This molecule is able to mediate the cellular uptake of several small molecules, including gadolinium-based contrast agents that enhance T1-weighted images in MRI and the radionuclide ¹¹¹In, which can be detected with SPECT. Dual-modality imaging has also been used in stroke research to track transplanted NSCs^{225–227} and MSCs.¹⁴³ In these studies, donor NSCs were transduced with *fluc* and labeled with SPIO particles to allow BLI and MRI imaging,^{226,227} or were imaged with ¹⁹F MRI after transduction with *fluc*.²²⁵ In a study by Walczak et al., SPIO-labeled MSCs were monitored with MRI combined with laser Doppler flow.¹⁴³

Over the past decade, several advances have been made in molecular imaging to facilitate stem cell tracking after transplantation. As will be discussed next, some of these methods are not only applicable to track the fate of the donor cells but can also be used to acquire additional information on the endogenous repair mechanisms following ischemic brain injury.

B. In vivo Imaging of the Recovering Brain

Similar to tracking the fate of donor cells, noninvasive imaging modalities based on MRI, radionuclide imaging, or optical methods can be applied to monitor the physiological and/or functional properties of the host environment. These methods allow visualization of neurovascular processes and neurological function but can also be used to study endogenous stem cell responses to treatment.

Optical methods are mainly based on FLI or BLI. In vivo two-photon FLI can be used to monitor blood flow, synapse formation, and neuroinflammation.^{228–230} Two-photon imaging requires fluorophores, although label-free imaging has been successfully used to visualize the mouse brain.^{231,232} FLI can also be used to directly monitor neuronal activity by using voltage sensitive dyes and proteins, which change their fluorescent properties in response to changes in transmembrane voltage.^{233,234} Unfortunately, penetration depth is limited and a cranial window is required to visualize subcortical structures and repair processes.^{233,235} BLI-based optical imaging methods are also limited by penetration depth, although it has been shown that the BLI signal can be detected through the intact skull.²⁰⁸ BLI-based methods have been used to track endogenous neuronal stem cells and neurogenesis after stroke.²⁰⁸ In addition, *fluc* under the control of the VEGFR2 receptor promoter has been used to evaluate poststroke angiogenesis with BLI²³⁶ and when put under control of the toll-like receptor 2, the response of microglia could be observed after photothrombotic stroke.²³⁷

1 24 • GERVOIS ET AL.

2 Another approach is to use PET or SPECT compatible radionuclides such as ^{18}F -FDG,
3 which has been used in animal models of stroke to evaluate stroke outcome with PET after
4 transplanting iPSCs and ESCs.²³⁸ This study by Wang et al. was able to correlate the ^{18}F -FDG
5 PET signal with functional improvement as shown by a decrease in mNSS score in animals
6 that received a nonpredifferentiated iPSC or ESC graft. Although the donor cells were not
7 predifferentiated, tumor formation was not observed up to 4 weeks posttransplantation. A
8 similar study was performed by Daadi et al. where NSCs were labeled with SPIO particles
9 and transduced with HSV1-*tk* to allow PET with the reporter probe [^{18}F]-FHBG prior to
10 transplantation in a rat stroke model.²³⁹ In accordance with the study of Wang et al., Daadi et al.
11 demonstrated an increased ^{18}F -FDG PET signal after NSC transplantation and correlated the
12 ^{18}F -FDG PET signal and the presence of NSCs as shown by [^{18}F]-FHBG detection. Increased
13 glucose uptake might be indicative of enhanced neuronal function but caution needs to be taken
14 when interpreting ^{18}F -FDG PET results as it is unclear which mechanisms are responsible for
15 the increased ^{18}F -FDG uptake in the injured brain.²⁴⁰ More recently, a study by Zinnhardt
16 et al. combined a multitracer PET study with MRI to link the spatiotemporal relationship of
17 MMP and microglial activation after transient MCAO providing more detailed information on
18 early and delayed endogenous stroke responses.²⁴¹

19 MR-based methods have the advantage that they can couple physiological information with
20 anatomical characteristics of the region of interest. Therefore, several MR-based modalities are
21 used to gain additional information of the host environment in stroke and will be discussed
22 next.

23 Blood oxygenation level dependent functional MRI (BOLD-fMRI) depends on the hemo-
24 dynamic response to neuronal activity, which are directly related to the energy demand of the
25 studied brain areas.²⁴² In small animal stroke research, BOLD-fMRI after electric forepaw stim-
26 ulation has been successfully used to assess functional recovery and electric brain activity.²⁴³
27 As stated previously, current therapies for ischemic stroke aim to salvage the ischemic penum-
28 bra. Therefore, it is important to have an adequate monitoring tool to evaluate the size of
29 the penumbra before and after therapeutic intervention. By using diffusion- and perfusion-
30 weighted MRI, the size of the ischemic penumbra can be determined based on the area of
31 diffusion/perfusion mismatch.²⁴⁴ While the diffusion/perfusion ratio is mainly used clinically,
32 it has also been successfully applied in a rat model of ischemic stroke.^{245,246}

33 In addition to providing a ways of evaluating the size of the penumbra, also the architectural
34 information of the affected brain region can be visualized with MR modalities. Diffuse tensor
35 imaging (DTI) images the anisotropy of water molecules in different tissues and is mainly used
36 in stroke research to study and visualize white matter tracts.²⁴⁷ Although its use in experimental
37 stroke in animal models is limited, DTI has been used to study the MRI evolution of stroke
38 macaques.²⁴⁸ While DTI can provide architectural information of white matter organization
39 after stroke, manganese-enhanced MRI (MEMRI) can be used to image synaptic connectivity
40 and assess changes in neuroarchitecture, anterograde axonal transport, and demarcates active
41 regions of the brain independent of hemodynamic contrast compounds.^{249,250} MEMRI uses T1
42 contrast-enhancing Mn^{2+} ions that can enter the cell via Ca^{2+} channels. As Ca^{2+} plays a crucial
43 role in neuronal activation Mn^{2+} is transported toward the synaptic cleft, where it can be taken
44 up by other neurons in the circuit.²⁵¹ Neuronal connectivity during stroke and recovery has
45 been evaluated in MCAO models using this technique.^{252,253}

46 Reperfusion plays a key role in repairing the ischemic lesion and can be examined with
47 arterial spin labelling (ASL) scans in order to quantify the absolute amount of tissue perfusion in
48 different brain areas evoked by therapeutic interventions, including stem cell based therapies.²⁵⁴
49 ASL measurements in small animal models of ischemic stroke were able to visualize perfusion
50 in the ischemic brain, allowing the potential longitudinal follow-up of therapeutic interventions
51 that aim to enhance reperfusion.^{255,256}

5. DISCUSSION AND PERSPECTIVES

Because of the promising preclinical results of stem cell based therapies in in vivo models of ischemic stroke, small-scale human trials were performed using IV delivered autologous MSCs.²⁵⁷⁻²⁵⁹ The outcome of these studies showed that MSC transplantation improved the disease outcome but stress that the underlying mechanisms of action need to be determined to provide a more directed approach, although it was reported that the SDF-1 levels in the serum of MSC-transplanted patients were associated with the clinical outcome.²⁵⁹ Therefore, it is important that in the preclinical phase the potential of stem cell based therapies is thoroughly investigated in animal models of ischemic stroke. Each model and route of administration has its own advantages to investigate specific mechanisms of action of the donor cells. IA- and IV-delivered donor cells would be the preferred method of administration in human applications, but this administration route requires a substantial amount of donor cells compared to intracranial delivered stem cells and are hindered by several limitations. These include high morbidity and cells ending up in the spleen after IA delivery^{143,260} and pulmonary obstruction after IV delivery of donor cells.^{145,261} However, several reports are available that state that donor cell migration toward the spleen is a possible mechanism of action of stem cell mediated regeneration following stroke. Acosta et al. demonstrated that IV-delivered BMSCs end up in the spleen, but that BMSC migration to the spleen inversely correlated with a reduced infarct size, periinfarct size, and the number of MHC-II positive activated cells in the striatum.¹³⁴ Although the influence of MSCs themselves on poststroke immunosuppression was not investigated, Doepfner et al. reported that poststroke immunosuppression was attenuated after MSC extracellular vesicles were injected in stroke mice.¹²⁹ Supporting this hypothesis, Vendrame et al. investigated the immunomodulatory effects of the mononuclear cell fraction of umbilical cord blood cells.²⁶² In this study, it was demonstrated that IV transplantation of these cells diminished spleen reduction and rescued CD8⁺ T-cell counts in addition to a reduction in brain damage. Moreover, it was shown that the cell transplant increased Il-10 and interferon gamma mRNA expression and decreased tumor necrosis factor alfa mRNA expression.²⁶² Donor cells applied to the host circulation migrate toward and integrate in low numbers into the brain lesion and ameliorate the disease outcome (See Table I) presumably by neurotrophic effects as the cerebral level of neurotrophins was found to be elevated in some studies.^{132, 135, 139} Moreover, IA or IV delivery of donor cells allows the interaction of the BBB with donor cells to be studied that can contribute to knowledge of the neuroimmunological response after ischemic stroke.¹²⁸ Intracranial delivery of donor cells is the most invasive route of transplantation but a thorough meta-analysis of preclinical data by Vu et al. showed that intracranial delivery of MSCs provided greater clinical benefit, although this mode of administration is less favorable for human applications due to the highly invasive nature of the transplantation procedure.²⁶³ Nonetheless, intracranial transplantation of fetal nigral tissue to treat patients with Parkinson's disease has been performed although the clinical benefit remained debatable.²⁶⁴

Donor cells remain detectable at the site of injury and ameliorate the disease outcome as shown by behavioral results and postmortem tissue analysis. However, functional integration of the engrafted MSCs is based on marker expression instead of electrophysiological recordings. The latter has only been performed for iPSC-derived cortical differentiated long-term expandable neuroepithelial-like stem cells, demonstrating functional integration of the engrafted cells.^{57,58} IA or IV delivery of iPSC-derived neuronal precursor or committed cells has not been performed to date.

When considering the most suitable source of stem cell as a potential therapy in stroke research, comparative studies are needed to highlight differences in therapeutic potential of stem cells from different sources in an analogous experimental setup. For example, studies comparing different subtypes of MSCs showed that ASCs are more suitable as a candidate

1 26 • GERVOIS ET AL.

2 MSC than BMSCs¹³⁹ while another study compared the same subsets of MSC and could not
3 observe any difference.⁶⁰ Moreover, the studies listed in Tables I and II describe both xenogenic
4 transplantation of human-derived MSCs and iPSCs as allogeneic-derived MSCs in animal
5 models for ischemic stroke. Studies that compare xenogenic and allogeneic MSCs in ischemic
6 stroke are limited to the study by Yasuhara et al.¹³⁷ and Balseanu et al.,¹⁴⁹ who compared
7 human and rat BMSCs, and Gutiérrez-Fernández et al. who compared human ASCs with
8 rat-ASCs.¹⁵¹ In the study by Yasuhara et al., it was shown that the allogeneic BMSC graft
9 showed a higher survival rate and a higher number of neurogenic differentiated cells. In both
10 engrafted groups, improvement in locomotor and neurological function and a reduced loss of
11 striatal periinfarct cells was observed.¹³⁷ Although Balseanu et al. used both rat and human
12 BMSC in their study, no direct comparison was made between the xenograft and allograft.¹⁴⁹
13 Preclinical studies with autologous-derived stem cells are scarce and were only described in
14 Jiang et al.'s¹³¹ study who transplanted autologous rat ASCs in ischemic rats, whereas for other
15 subtypes of MSCs, preclinical studies with autologous stem cells are not available. This study
16 by Jiang et al. did not compare the outcome of the transplantation between autologous and
17 allogeneic ASCs, which would provide additional information on extrapolability of their results.
18 Although xenogenic stem cell transplants are not likely to be used in human studies, preclinical
19 studies using xenogenic grafts provide insight into the potential of human stem cell sources in
20 ischemic stroke.

21 To date, no studies have been performed that compare the therapeutic potential of iPSCs
22 with a subtype of MSCs. Although the most thoroughly studied stem cell type in stroke re-
23 search are MSCs, the potential of iPSCs and iPSC-derived neural progenitor cells is currently
24 being intensively investigated with several very promising results when iPSCs are predifferentiated
25 toward neuronal precursor cells or committed cortical neurons.^{57,58} Nonetheless, adequate
26 screening of fully neuronal committed cells preceding engraftment is recommended for several
27 reasons. As stated previously, a recent study by Choi et al. demonstrated the ability of the
28 iPSC-derived NPCs to return toward their pluripotent and thus tumorigenic state by transgene
29 reactivation during differentiation.¹⁵⁷ This statement was supported by Liu et al.¹⁵⁶ and Fu
30 et al.¹⁵⁸ who demonstrated that nondifferentiated iPSCs remain present in an iPSC-derived
31 progenitor pool. Another problem with iPSCs, and more specifically with retroviral trans-
32 duced iPSCs, is the retroviral gene integration in the host, which promotes tumorigenicity.²⁶⁵
33 Therefore, additional approaches have been developed to generate iPSCs with a lower risk for
34 tumorigenicity.^{266,267} Moreover, it has been shown that iPSCs retain an epigenetic memory
35 related to the somatic donor tissue, leading to spontaneous redifferentiation to the cells of the
36 tissue of origin,^{268–270} although it has been shown that this epigenetic memory and redifferentia-
37 tion rate can vary between the somatic cells of origin with different tumorigenic propensities
38 between the somatic donor cells.^{269,270} The tumorigenicity of human iPSCs (and ESCs) was
39 thoroughly reviewed by Ben-David and Benvenisty.²⁷¹ As an alternative to iPSCs in which the
40 Yamanaka factors are transduced, exogenous gene-free iPSCs can be used.²⁷² Another option
41 is to additionally engineer iPSC-derived cells to express suicide genes to eradicate the cells, an
42 approach that was successfully used in ESCs and BMSCs.^{273,274}

43 It should be noted that in addition to various MSC sources and iPSCs, encouraging results
44 have been achieved by using BMMNCs in animal models of ischemic stroke where this subset
45 of cells was found to stimulate endogenous angiogenesis^{61,62} and neurogenesis by improving
46 the NPC-vascular niche⁶³ or modulate the immune system.⁹⁵ Moreover, it was shown that
47 after ischemic stroke, the amount of CD34⁺ blood cells that migrate from the bone marrow
48 to peripheral blood is increased, which is associated with a better clinical outcome and is be-
49 lieved to be mediated by granulocyte colony-stimulating factor (G-CSF).^{275,276} In addition to
50 its effect on BMMNC recruitment, this factor has previously been shown to possess neuropro-
51 tective and neuroregenerative effects²⁷⁷ and is also believed to mobilize BMSCs and possess

Q7

immunomodulatory properties.²⁷⁸ BMMNCs can also be quickly isolated from peripheral blood by Ficoll-Paque density gradient centrifugation just before administration, circumventing the additional cell culture period, which is required for MSC- or iPSC-based therapies.⁶³ As was a hurdle with MSC-based therapy, also the administration route of BMMNCs remains a topic of debate. Although study by Kamiya et al.⁹⁶ showed superior results after IA-delivered BMMNCs over IV-delivered cells, this was contradicted by Yang et al. who showed that IA delivery was not superior to IV-delivered BMMNCs.²⁷⁹ Subsequently, these cells have been used in several clinical trials. IA administration of these cells was safe and showed an improved clinical outcome.^{280,281} IV delivery of these cells also appeared safe,²⁸² although no clinical improvement was observed in the study by Prasad et al.,²⁸³ whereas Savitz et al. were able to show an improved functional outcome.²⁸² Despite these encouraging results, in depth in vitro evidence on the underlying mechanisms of action is largely unknown. The scarce in vitro data on the effect of BMMNCs on regenerative processes showed that BMMNCs exerted protective effects on rat hippocampal brain slices subjected to oxygen and glucose deprivation⁹⁸ and that the BMMNC secretome induced neuronal differentiation of SH-SY5Y neuroblastoma cells.⁹⁷ Nonetheless, additional in vitro data supporting the mechanism of action of BMMNC-based therapy for ischemic stroke are required.

As stroke is a disease that mainly affects the elderly,⁷ it is important to take into account the effect of the aged microenvironment, age-related comorbidities, and the aged immune system on the outcome of stem cell based therapies.^{230,284,285} Although it does not appear from clinical studies that the aged brain microenvironment is detrimental for stem cell based therapies, differences exist between the young and aged brain. For example, the formation of the glial scar is accelerated after stroke that hinders functional repair in aged rats.²⁸⁶ Moreover, comorbidities such as hypertension, hyperlipidemia, and diabetes mellitus appear to play a role in age-related stroke severity.^{287,288} As described previously, angiogenesis is a key concept in establishing brain repair. Buga et al. compared the transcriptome and immunochemistry of young and aged stroke rats and poststroke patients. Remarkably, although the upregulation of proangiogenic genes associated with processes such as vessel sprouting, tube formation, and maturation was delayed in aged rats, angiogenesis in the aged brains was similar to their younger counterparts. In addition, an upregulation of proinflammatory and scar-promoting genes was found in the aged rats compared to the younger brains, supporting the accelerated scar formation and increased neuroinflammation in aged stroke subjects.²⁸⁹ Of the studies listed in Table I, two studies by Shen et al.^{146,147} and studies by Taguchi et al.,¹⁵² Balseanu et al.,¹⁴⁹ and Zhang et al.¹⁴⁸ used aged rats to perform MSC transplantation studies in ischemic stroke. Although these studies did not directly compare the outcome of their transplantation study between young and aged rats, several encouraging results were found in these studies. These included, but are not limited to, enhanced functional recovery,^{146,148,149,152} a reduction of the glial scar thickness,^{146,147} and improved angiogenesis.^{148,149,152} Remarkably, Balseanu et al. who used a G-CSF treatment, which is believed to possess multiple regenerative effects,²⁷⁸ or a combination of G-CSF and a single BM-MSCs dose to improve the functional outcome in aged stroke animals, observed that the functional improvement was not increased in this combination treatment.¹⁴⁹ Similarly, Buga et al. who used a G-CSF and a combined G-CSF-BMMNC therapy to improve the functional outcome after transplantation in aged stroke animals were also unable to observe an increased functional improvement by using the combination treatment.²⁹⁰ The use of iPSCs in aged rats was described by Tatarishvili et al. who showed that almost 50% of the engrafted iPSC-derived long-term expendable neuroepithelial cells survived 8 weeks post-transplantation and caused functional improvement in the aged rats.¹⁷⁰ Moreover, these cells differentiated toward neuroblast-like cells, compared to the BM-MSCs described in Refs. 146 and 147 where the few surviving cells predominantly differentiated toward astrocytes. These studies support the use of aged animals for in vivo stroke studies in which they were able to observe

1 28 • GERVOIS ET AL.

Q9

2 cell-mediated improvements in brain regeneration and functional recovery. Nonetheless, a di-
3 rect comparison of stroke outcome and the molecular effect after cell transplantation between
4 young and aged animals for the various age-related differences in (brain) microenvironment
5 could provide additional information on which processes are mainly responsible for stem cell
6 mediated brain repair. For example, the previously mentioned study by Buga et al., which
7 described the transcriptome in young and aged rats after stroke, provided several new target
8 pathways that can provide additional insight into age-related stroke pathology and therapeutic
9 opportunities.²⁸⁹

10 The evaluation of the disease outcome in animal models for ischemic stroke is based on
11 behavioral testing, for which various tests can be applied to investigate specific aspects of
12 neuronal recovery.¹²⁶ As stroke symptoms are largely dependent on the brain area involved,
13 different behavioral tests are applied to evaluate general, motor, sensorimotor, and cognitive
14 recovery and each test has its strengths and weaknesses. For example, the Bederson test is
15 easy to perform but reliable neurological ratings on this Bederson scale are limited because
16 of their subjective nature, a common feature of all behavioral tests that are based on human
17 observation. An overview with critical comments on the different behavioral tests that are often
18 used in stroke research is provided by Schaar et al.¹²⁶ Nonetheless, functional assessment of
19 stroke outcome should include tests to cover all aspects of the disease outcome.

20 As indicated above, preclinical stroke research is facing several important issues that need
21 to be resolved prior to more elaborate testing of the clinical potential of stem cell based
22 therapies for ischemic stroke. Noninvasive imaging methods can aid in the longitudinal follow-
23 up of stem cell fate and effect after transplantation via various administration routes. However,
24 with the exception of a few studies,^{238,253} most of the studies that used noninvasive imaging
25 to monitor the poststroke microenvironment or stem cell migratory pathways focused on
26 the proof of principle of the imaging technique and did not link their results to functional
27 recovery,^{206,208,236,246,256} which will most likely be the next step to be performed with these
28 highly promising imaging modalities.

31 6. CONCLUSION

32 Although multiple clinical advances have been made to improve the clinical diagnosis and out-
33 come after acute ischemic stroke, beneficial long-term or delayed interventions are currently
34 not available. Stem cell based therapies with MSCs and iPSCs have shown great promise in
35 vivo models of ischemic stroke through various administration routes. Nonetheless, the mech-
36 anisms of action of the transplanted cells remain poorly understood and are highly dependent
37 on administration route, pretreatment, and full neuronal predifferentiation when using iPSCs.
38 Although postmortem cell tracking methods provide detailed spatial information on the donor
39 cell fate and host microenvironment, they are unable to deliver dynamic information on these
40 subjects. A longitudinal follow-up with noninvasive imaging methods allows donor cell fate and
41 changes in host microenvironment to be linked with behavioral and functional improvements,
42 which can lead to additional insight into the mechanisms responsible for functional recovery
43 in stroke after donor cell transplantation.
44
45

48 ACKNOWLEDGEMENTS

49
50 Esther Wolfs, Jessica Ratajczak, Tim Vangansewinkel, Petra Hilken, and Annelies Bronck-
51 aers are funded by Fonds Wetenschappelijk Onderzoek by grants G0A7514N, G089213N,

STEM CELL-BASED THERAPIES FOR ISCHEMIC STROKE • 29

G029112N, 12D8516N, and 1508015N, respectively. Yörg Dillen is funded by Bijzonder Onderzoeksfonds by grant BOF15DOC04.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

1. Nichols M, Townsend N, Luengo-Fernandez R, Leal J, Gray A, Scarborough P, Rayner M. European Cardiovascular Disease Statistics 2012/2012. Q10
2. Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Baha MJ, Dai SF, Ford ES, Fox CS, Franco S, Fullerton HJ, Gillespie C, Hailpern SM, Heit JA, Howard VJ, Huffman MD, Judd SE, Kissela BM, Kittner SJ, Lackland DT, Lichtman JH, Lisabeth LD, Mackey RH, Magid DJ, Marcus GM, Marelli A, Matchar DB, McGuire DK, Mohler ER, Moy CS, Mussolino ME, Neumar RW, Nichol G, Pandey DK, Paynter NP, Reeves MJ, Sorlie PD, Stein J, Towfighi A, Turan TN, Virani SS, Wong ND, Woo D, Turner MB, Comm AHAS, Subcomm SS. Heart Disease and Stroke Statistics 2014 Update: A report from the American Heart Association. *Circulation* 2014;129(3):E28–E292. Q11
3. Hinkle JL, Guanci MM. Acute ischemic stroke review. *J Neurosci Nurs* 2007;39(5):285–293, 310. Q12
4. Dua T, Cumbre M, Mathers C, Saxena S. Global burden of neurological disorders: Estimates and projections. In: Campanini B, Ed. *Neurological Disorders: Public Health Challenges*. Geneva, Switzerland: World Health Organization; 2006. p 27–40.
5. United Nations, Department of Economic and Social Affairs. *Magnitude and Speed of Population Ageing. World Population Ageing 1950-2050*. New York, NY: Population Division, DESA, United Nations; 2002. p 11–13.
6. Norrving B, Kissela B. The global burden of stroke and need for a continuum of care. *Neurology* 2013;80(3 Suppl 2):S5–S12.
7. Feigin VL, Forouzanfar MH, Krishnamurthi R, Mensah GA, Connor M, Bennett DA, Moran AE, Sacco RL, Anderson L, Truelsen T, O'Donnell M, Venketasubramanian N, Barker-Collo S, Lawes CM, Wang W, Shinohara Y, Witt E, Ezzati M, Naghavi M, Murray C. Global burden of diseases I, risk factors S, the GBDSEG. Global and regional burden of stroke during 1990-2010: Findings from the Global Burden of Disease Study 2010. *Lancet* 2014;383(9913):245–254.
8. Lindvall O, Kokaia Z. Stem cells for the treatment of neurological disorders. *Nature* 2006;441(7097):1094–1096.
9. Lindvall O, Kokaia Z. Stem cell research in stroke: How far from the clinic? *Stroke* 2011;42(8):2369–2375.
10. Wolfs E, Verfaillie CM, Van Laere K, Deroose CM. Radiolabeling strategies for radionuclide imaging of stem cells. *Stem Cell Rev* 2015;11(2):254–274.
11. Goldstein M, Barnett HJM, Orgogozo JM, Sartorius N, Symon L, Vereshchagin NV. Stroke-1989: Recommendations on stroke prevention, diagnosis, and therapy. Report of the WHO Task Force on Stroke and other Cerebrovascular Disorders. *Stroke* 1989;20(10):1407–1431.
12. Donnan GA, Fisher M, Macleod M, Davis SM. Stroke. *Lancet* 2008;371(9624):1612–1623.
13. White BC, Sullivan JM, DeGracia DJ, O'Neil BJ, Neumar RW, Grossman LI, Rafols JA, Krause GS. Brain ischemia and reperfusion: Molecular mechanisms of neuronal injury. *J Neurol Sci* 2000;179(S 1-2):1–33.
14. Dirnagl U, Iadecola C, Moskowitz MA. Pathobiology of ischaemic stroke: An integrated view. *Trends Neurosci* 1999;22(9):391–397.
15. Bandera E, Botteri M, Minelli C, Sutton A, Abrams KR, Latronico N. Cerebral blood flow threshold of ischemic penumbra and infarct core in acute ischemic stroke: A systematic review. *Stroke* 2006;37(5):1334–1339.

1 30 • GERVOIS ET AL.

- 2 16. Fisher M. The ischemic penumbra: Identification, evolution and treatment concepts. *Cerebrovasc*
3 *Dis* 2004; 1(17 Suppl):1–6.
- 4 17. Moretti A, Ferrari F, Villa RF. Neuroprotection for ischaemic stroke: Current status and challenges.
5 *Pharmacol Ther* 2015;146:23–34.
- 6 18. Muir KW, Buchan A, von Kummer R, Rother J, Baron JC. Imaging of acute stroke. *Lancet Neurol*
7 2006;5(9):755–768.
- 8 19. Goyal M, Menon BK, Derdeyn CP. Perfusion imaging in acute ischemic stroke: Let us improve the
9 science before changing clinical practice. *Radiology* 2013;266(1):16–21.
- 10 20. Pan J, Konstas AA, Bateman B, Ortolano GA, Pile-Spellman J. Reperfusion injury follow-
11 ing cerebral ischemia: Pathophysiology, MR imaging, and potential therapies. *Neuroradiology*
12 2007;49(2):93–102.
- 13 21. Dominguez C, Delgado P, Vilches A, Martin-Gallan P, Ribo M, Santamarina E, Molina C, Corbetol
14 N, Rodriguez-Sureda V, Rosell A, Alvarez-Sabin J, Montaner J. Oxidative stress after thrombolysis-
15 induced reperfusion in human stroke. *Stroke* 2010;41(4):653–660.
- 16 22. Jickling GC, Liu D, Stamova B, Ander BP, Zhan X, Lu A, Sharp FR. Hemorrhagic transformation
17 after ischemic stroke in animals and humans. *J Cereb Blood Flow Metab* 2014;34(2):185–199.
- 18 23. Warach S, Latour LL. Evidence of reperfusion injury, exacerbated by thrombolytic therapy, in
19 human focal brain ischemia using a novel imaging marker of early blood-brain barrier disruption.
20 *Stroke* 2004;35(11 Suppl 1):2659–2661.
- 21 24. Paciaroni M, Agnelli G, Corea F, Ageno W, Alberti A, Lanari A, Caso V, Micheli S, Bertolani
22 L, Venti M, Palmerini F, Biagini S, Comi G, Previdi P, Silvestrelli G. Early hemorrhagic transfor-
23 mation of brain infarction: Rate, predictive factors, and influence on clinical outcome: Results of a
24 prospective multicenter study. *Stroke* 2008;39(8):2249–2256.
- 25 25. Langhorne P, Collaborati SUT. Organised inpatient (stroke unit) care for stroke. *Cochrane Database*
26 *Syst Rev* 2013;(9).
- 27 26. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. Tissue
28 plasminogen activator for acute ischemic stroke. *N Engl J Med* 1995;333(24):1581–1587.
- 29 27. Hacke W, Kaste M, Fieschi C, Toni D, Lesaffre E, von Kummer R, Boysen G, Bluhmki E, Hoxter
30 G, Mahagne MH, et al. Intravenous thrombolysis with recombinant tissue plasminogen activator
31 for acute hemispheric stroke. The European Cooperative Acute Stroke Study (ECASS). *JAMA*
32 1995;274(13):1017–1025.
- 33 28. Sandercock P, Collins R, Counsell C, Farrell B, Peto R, Slattery J, Warlow C, Anderson S, Bowie
34 A, Boyle J, Brownlie A, Charlton D, Cranswick G, Day L, Dennis M, Dorman P, Fraser H, Kaye
35 M, Lindley R, Liu M, MacDonald C, McCrindle I, Middleton G, Perry D, Scoltock V, Smith B,
36 Taylor H, Waddell F, Wardlaw J, Crowther J, Heineman J, Knight S, Radley A, Ripley R, Richards
37 S, Wilberforce S, Chen ZM, vanGijn J, Harrison M, Wilhelmsen L. The International Stroke Trial
38 (IST): A randomised trial of aspirin, subcutaneous heparin, both, or neither among 19 435 patients
39 with acute ischaemic stroke. *Lancet* 1997;349(9065):1569–1581.
- 40 29. Vahedi K, Hofmeijer J, Juettler E, Vicaut E, George B, Algra A, Amelink GJ, Schmiedeck P,
41 Schwab S, Rothwell PM, Bousser MG, van der Worp HB, Hacke W, Decimal Destiny, Hamlet
42 Investigators. Early decompressive surgery in malignant infarction of the middle cerebral artery: A
43 pooled analysis of three randomised controlled trials. *Lancet Neurol* 2007;6(3):215–222.
- 44 30. National Institute of Neurological Disorders and Stroke rt-PA Study Group. 1995. Tissue plasmino-
45 gen activator for acute ischemic stroke. *N Engl J Med*;333(24):1581–1588.
- 46 31. Agyeman O, Nedeltchev K, Arnold M, Fischer U, Remonda L, Isenegger J, Schroth G, Mattle HP.
47 Time to admission in acute ischemic stroke and transient ischemic attack. *Stroke* 2006;37(4):963–
48 966.
- 49 32. Ropper AH, Samuels MA, Klein JP. Chapter 34: Cerebrovascular diseases. *Adams and Victor's*
50 *Principles of Neurology*. 10th ed. New York: McGraw-Hill; 2014. p 778–884.
- 51 33. Hacke W, Donnan G, Fieschi C, Kaste M, von Kummer R, Broderick JP, Brott T, Frankel M,
Grotta JC, Haley EC, Jr., Kwiatkowski T, Levine SR, Lewandowski C, Lu M, Lyden P, Marler JR,

STEM CELL-BASED THERAPIES FOR ISCHEMIC STROKE • 31

- Patel S, Tilley BC, Albers G, Bluhmki E, Wilhelm M, Hamilton S, Investigators AT, Investigators ET, Investigators Nr-PSG. Association of outcome with early stroke treatment: Pooled analysis of ATLANTIS, ECASS, and NINDS rt-PA stroke trials. *Lancet* 2004;363(9411):768–774.
34. Hacke W, Kaste M, Bluhmki E, Brozman M, Davalos A, Guidetti D, Larrue V, Lees KR, Medeghri Z, Machnig T, Schneider D, von Kummer R, Wahlgren N, Toni D, Investigators E. Thrombolysis with alteplase 3 to 4.5 hours after acute ischemic stroke. *N Engl J Med* 2008;359(13):1317–1329.
35. Frank JI. Large hemispheric infarction, deterioration, and intracranial pressure. *Neurology* 1995;45(7):1286–1290.
36. von Kummer R, Albers GW, Mori E, Committees DS. The Desmoteplase in Acute Ischemic Stroke (DIAS) clinical trial program. *Int J Stroke* 2012;7(7):589–596.
37. Emberson J, Lees KR, Lyden P, Blackwell L, Albers G, Bluhmki E, Brott T, Cohen G, Davis S, Donnan G, Grotta J, Howard G, Kaste M, Koga M, von Kummer R, Lansberg M, Lindley RI, Murray G, Olivot JM, Parsons M, Tilley B, Toni D, Toyoda K, Wahlgren N, Wardlaw J, Whiteley W, Del Zoppo GJ, Baigent C, Sandercock P, Hacke W, Stroke Thrombolysis Trialists' Collaborative G. Effect of treatment delay, age, and stroke severity on the effects of intravenous thrombolysis with alteplase for acute ischaemic stroke: A meta-analysis of individual patient data from randomised trials. *Lancet* 2014;384(9958):1929–1935.
38. Barreto AD, Alexandrov AV, Shen L, Sisson A, Bursaw AW, Sahota P, Peng H, Ardjomand-Hessabi M, Pandurengan R, Rahbar MH, Barlinn K, Indupuru H, Gonzales NR, Savitz SI, Grotta JC. CLOTBUST-Hands Free: Pilot safety study of a novel operator-independent ultrasound device in patients with acute ischemic stroke. *Stroke* 2013;44(12):3376–3381.
39. Smith WS, Sung G, Starkman S, Saver JL, Kidwell CS, Gobin YP, Lutsep HL, Nesbit GM, Grobelny T, Rymer MM, Silverman IE, Higashida RT, Budzik RF, Marks MP, Investigators MT. Safety and efficacy of mechanical embolectomy in acute ischemic stroke: Results of the MERCI trial. *Stroke* 2005;36(7):1432–1438.
40. Asadi H, Dowling R, Yan B, Wong S, Mitchell P. Advances in endovascular treatment of acute ischaemic stroke. *Intern Med J* 2015;45(8):798–805.
41. Dirnagl U, Klehmet J, Braun JS, Harms H, Meisel C, Ziemssen T, Prass K, Meisel A. Stroke-induced immunodepression: Experimental evidence and clinical relevance. *Stroke* 2007;38(2 Suppl):770–773.
42. Chouchani ET, Pell VR, Gaude E, Aksentijevic D, Sundier SY, Robb EL, Logan A, Nadtochiy SM, Ord EN, Smith AC, Eyassu F, Shirley R, Hu CH, Dare AJ, James AM, Rogatti S, Hartley RC, Eaton S, Costa AS, Brookes PS, Davidson SM, Duchon MR, Saeb-Parsy K, Shattock MJ, Robinson AJ, Work LM, Frezza C, Krieg T, Murphy MP. Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS. *Nature* 2014;515(7527):431–435.
43. Ishiguro M, Mishihiro K, Fujiwara Y, Chen H, Izuta H, Tsuruma K, Shimazawa M, Yoshimura S, Satoh M, Iwama T, Hara H. Phosphodiesterase-III inhibitor prevents hemorrhagic transformation induced by focal cerebral ischemia in mice treated with tPA. *PLoS One* 2010;5(12):e15178.
44. Hawkins KE, DeMars KM, Singh J, Yang C, Cho HS, Frankowski JC, Dore S, Candelario-Jalil E. Neurovascular protection by post-ischemic intravenous injections of the lipoxin A4 receptor agonist, BML-111, in a rat model of ischemic stroke. *J Neurochem* 2014;129(1):130–142.
45. van der Worp HB, Macleod MR, Bath PM, Demotes J, Durand-Zaleski I, Gebhardt B, Gluud C, Kollmar R, Krieger DW, Lees KR, Molina C, Montaner J, Roine RO, Petersson J, Staykov D, Szabo I, Wardlaw JM, Schwab S, Euro HYPi. EuroHYP-1: European multicenter, randomized, phase III clinical trial of therapeutic hypothermia plus best medical treatment vs. best medical treatment alone for acute ischemic stroke. *Int J Stroke* 2014;9(5):642–645.
46. Polderman KH. Mechanisms of action, physiological effects, and complications of hypothermia. *Crit Care Med* 2009;37(7 Suppl):S186–S202.
47. Molina CA. Reperfusion therapies for acute ischemic stroke: Current pharmacological and mechanical approaches. *Stroke* 2011;42(1 Suppl):S16–S19.
48. Teixeira-Salmela LF, Olney SJ, Nadeau S, Brouwer B. Muscle strengthening and physical conditioning to reduce impairment and disability in chronic stroke survivors. *Arch Phys Med Rehabil* 1999;80(10):1211–1218.

1 32 • GERVOIS ET AL.

- 2 49. Burns TC, Verfaillie CM, Low WC. Stem cells for ischemic brain injury: A critical review. *J Comp*
3 *Neurol* 2009;515(1):125–144.
- 4 50. Lundberg C, Martinez-Serrano A, Cattaneo E, McKay RD, Bjorklund A. Survival, integration, and
5 differentiation of neural stem cell lines after transplantation to the adult rat striatum. *Exp Neurol*
6 1997;145(2 Pt 1):342–360.
- 7 51. Liu S, Qu Y, Stewart TJ, Howard MJ, Chakraborty S, Holekamp TF, McDonald JW. Embryo-
8 nic stem cells differentiate into oligodendrocytes and myelinate in culture and after spinal cord
9 transplantation. *Proc Natl Acad Sci USA* 2000;97(11):6126–6131.
- 10 52. Johansson CB, Momma S, Clarke DL, Risling M, Lendahl U, Frisen J. Identification of a neural
11 stem cell in the adult mammalian central nervous system. *Cell* 1999;96(1):25–34.
- 12 53. Reynolds BA, Weiss S. Generation of neurons and astrocytes from isolated cells of the adult
13 mammalian central nervous system. *Science* 1992;255(5052):1707–1710.
- 14 54. Mine Y, Tatarishvili J, Oki K, Monni E, Kokaia Z, Lindvall O. Grafted human neural stem cells
15 enhance several steps of endogenous neurogenesis and improve behavioral recovery after middle
16 cerebral artery occlusion in rats. *Neurobiol Dis* 2013;52:191–203.
- 17 55. McLaren A. Ethical and social considerations of stem cell research. *Nature* 2001;414(6859):129–131.
- 18 56. Nunes MC, Roy NS, Keyoung HM, Goodman RR, McKhann G, 2nd, Jiang L, Kang J, Nedergaard
19 M, Goldman SA. Identification and isolation of multipotential neural progenitor cells from the
20 subcortical white matter of the adult human brain. *Nat Med* 2003;9(4):439–447.
- 21 57. Tornero D, Wattananit S, Gronning Madsen M, Koch P, Wood J, Tatarishvili J, Mine Y, Ge R,
22 Monni E, Devaraju K, Hevner RF, Brustle O, Lindvall O, Kokaia Z. Human induced pluripotent
23 stem cell-derived cortical neurons integrate in stroke-injured cortex and improve functional recovery.
24 *Brain* 2013;136(Pt 12):3561–3577.
- 25 58. Oki K, Tatarishvili J, Wood J, Koch P, Wattananit S, Mine Y, Monni E, Tornero D, Ahlenius H,
26 Ladewig J, Brustle O, Lindvall O, Kokaia Z. Human-induced pluripotent stem cells form functional
27 neurons and improve recovery after grafting in stroke-damaged brain. *Stem Cells* 2012;30(6):1120–
28 1133.
- 29 59. Leong WK, Henshall TL, Arthur A, Kremer KL, Lewis MD, Helps SC, Field J, Hamilton-Bruce
30 MA, Warming S, Manavis J, Vink R, Gronthos S, Koblar SA. Human adult dental pulp stem cells
31 enhance poststroke functional recovery through non-neural replacement mechanisms. *Stem Cells*
32 *Transl Med* 2012;1(3):177–187.
- 33 60. Gutierrez-Fernandez M, Rodriguez-Frutos B, Ramos-Cejudo J, Teresa Vallejo-Cremades M,
34 Fuentes B, Cerdan S, Diez-Tejedor E. Effects of intravenous administration of allogenic bone
35 marrow- and adipose tissue-derived mesenchymal stem cells on functional recovery and brain repair
36 markers in experimental ischemic stroke. *Stem Cell Res Ther* 2013;4(1):11.
- 37 61. Shyu WC, Lin SZ, Chiang MF, Su CY, Li H. Intracerebral peripheral blood stem cell (CD34+)
38 implantation induces neuroplasticity by enhancing beta1 integrin-mediated angiogenesis in chronic
39 stroke rats. *J Neurosci* 2006;26(13):3444–3453.
- 40 62. Taguchi A, Soma T, Tanaka H, Kanda T, Nishimura H, Yoshikawa H, Tsukamoto Y, Iso
41 H, Fujimori Y, Stern DM, Naritomi H, Matsuyama T. Administration of CD34+ cells after
42 stroke enhances neurogenesis via angiogenesis in a mouse model. *J Clin Invest* 2004;114(3):330–
43 338.
- 44 63. Nakano-Doi A, Nakagomi T, Fujikawa M, Nakagomi N, Kubo S, Lu S, Yoshikawa H, Soma T,
45 Taguchi A, Matsuyama T. Bone marrow mononuclear cells promote proliferation of endogenous
46 neural stem cells through vascular niches after cerebral infarction. *Stem Cells* 2010;28(7):1292–1302.
- 47 64. Martens W, Wolfs E, Struys T, Politis C, Bronckaers A, Lambrechts I. Expression pattern of basal
48 markers in human dental pulp stem cells and tissue. *Cells Tissues Organs* 2012;196(6):490–500.
- 49 65. Fujimura J, Ogawa R, Mizuno H, Fukunaga Y, Suzuki H. Neural differentiation of adipose-derived
50 stem cells isolated from GFP transgenic mice. *Biochem Biophys Res Commun* 2005;333(1):116–121.
- 51 66. Gervois P, Struys T, Hilkens P, Bronckaers A, Ratajczak J, Politis C, Brone B, Lambrechts I, Martens
W. Neurogenic maturation of human dental pulp stem cells following neurosphere generation

- induces morphological and electrophysiological characteristics of functional neurons. *Stem Cells Dev* 2014.
67. Bederson JB, Pitts LH, Tsuji M, Nishimura MC, Davis RL, Bartkowski H. Rat middle cerebral artery occlusion: Evaluation of the model and development of a neurologic examination. *Stroke* 1986;17(3):472–476.
68. Bacigaluppi M, Comi G, Hermann DM. Animal models of ischemic stroke. Part two: Modeling cerebral ischemia. *Open Neurol J* 2010;4:34–38.
69. Chen ST, Hsu CY, Hogan EL, Maricq H, Balentine JD. A model of focal ischemic stroke in the rat: Reproducible extensive cortical infarction. *Stroke* 1986;17(4):738–743.
70. Zhao LR, Duan WM, Reyes M, Keene CD, Verfaillie CM, Low WC. Human bone marrow stem cells exhibit neural phenotypes and ameliorate neurological deficits after grafting into the ischemic brain of rats. *Exp Neurol* 2002;174(1):11–20.
71. Lin YC, Ko TL, Shih YH, Lin MY, Fu TW, Hsiao HS, Hsu JY, Fu YS. Human umbilical mesenchymal stem cells promote recovery after ischemic stroke. *Stroke* 2011;42(7):2045–2053.
72. Egashira Y, Sugitani S, Suzuki Y, Mishiro K, Tsuruma K, Shimazawa M, Yoshimura S, Iwama T, Hara H. The conditioned medium of murine and human adipose-derived stem cells exerts neuroprotective effects against experimental stroke model. *Brain Res* 2012;1461:87–95.
73. Mead B, Logan A, Berry M, Leadbeater W, Scheven BA. Paracrine-mediated neuroprotection and neuritogenesis of axotomised retinal ganglion cells by human dental pulp stem cells: Comparison with human bone marrow and adipose-derived mesenchymal stem cells. *PLoS One* 2014;9(10):e109305.
74. Scheibe F, Klein O, Klose J, Priller J. Mesenchymal stromal cells rescue cortical neurons from apoptotic cell death in an in vitro model of cerebral ischemia. *Cell Mol Neurobiol* 2012;32(4):567–576.
75. Eriksson PS, Perfilieva E, Bjork-Eriksson T, Alborn AM, Nordborg C, Peterson DA, Gage FH. Neurogenesis in the adult human hippocampus. *Nat Med* 1998;4(11):1313–1317.
76. Reumers V, Deroose CM, Krylyshkina O, Nuyts J, Geraerts M, Mortelmans L, Gijbbers R, Van den Haute C, Debysier Z, Baekelandt V. Noninvasive and quantitative monitoring of adult neuronal stem cell migration in mouse brain using bioluminescence imaging. *Stem Cells* 2008;26(9):2382–2390.
77. Arvidsson A, Collin T, Kirik D, Kokaia Z, Lindvall O. Neuronal replacement from endogenous precursors in the adult brain after stroke. *Nat Med* 2002;8(9):963–970.
78. Thored P, Arvidsson A, Cacci E, Ahlenius H, Kallur T, Darsalia V, Ekdahl CT, Kokaia Z, Lindvall O. Persistent production of neurons from adult brain stem cells during recovery after stroke. *Stem Cells* 2006;24(3):739–747.
79. Li Y, Huang J, He X, Tang G, Tang YH, Liu Y, Lin X, Lu Y, Yang GY, Wang Y. Postacute stromal cell-derived factor-1alpha expression promotes neurovascular recovery in ischemic mice. *Stroke* 2014;45(6):1822–1829.
80. Huang J, Li Y, Tang Y, Tang G, Yang GY, Wang Y. CXCR4 antagonist AMD3100 protects blood-brain barrier integrity and reduces inflammatory response after focal ischemia in mice. *Stroke* 2013;44(1):190–197.
81. Ruscher K, Kuric E, Liu Y, Walter HL, Issazadeh-Navikas S, Englund E, Wieloch T. Inhibition of CXCL12 signaling attenuates the postischemic immune response and improves functional recovery after stroke. *J Cereb Blood Flow Metab* 2013;33(8):1225–1234.
82. Mithal DS, Banisadr G, Miller RJ. CXCL12 signaling in the development of the nervous system. *J Neuroimmune Pharmacol* 2012;7(4):820–834.
83. Michelsen KA, Acosta-Verdugo S, Benoit-Marand M, Espuny-Camacho I, Gaspard N, Saha B, Gaillard A, Vanderhaeghen P. Area-specific reestablishment of damaged circuits in the adult cerebral cortex by cortical neurons derived from mouse embryonic stem cells. *Neuron* 2015;85(5):982–997.
84. Gaillard A, Prestoz L, Dumartin B, Cantereau A, Morel F, Roger M, Jaber M. Reestablishment of damaged adult motor pathways by grafted embryonic cortical neurons. *Nat Neurosci* 2007;10(10):1294–1299.

1 34 • GERVOIS ET AL.

- 2 85. Thompson L, Barraud P, Andersson E, Kirik D, Bjorklund A. Identification of dopaminergic
3 neurons of nigral and ventral tegmental area subtypes in grafts of fetal ventral mesencephalon based
4 on cell morphology, protein expression, and efferent projections. *J Neurosci* 2005;25(27):6467–6477.
- 5 86. Ebrahimi-Gaillard A, Guitet J, Garnier C, Roger M. Topographic distribution of efferent fibers
6 originating from homotopic or heterotopic transplants: Heterotopically transplanted neurons retain
7 some of the developmental characteristics corresponding to their site of origin. *Brain Res Dev Brain*
8 *Res* 1994;77(2):271–283.
- 9 87. Espuny-Camacho I, Michelsen KA, Gall D, Linaro D, Hasche A, Bonnefont J, Bali C, Orduz D,
10 Bilheu A, Herpoel A, Lambert N, Gaspard N, Peron S, Schiffmann SN, Giugliano M, Gaillard A,
11 Vanderhaeghen P. Pyramidal neurons derived from human pluripotent stem cells integrate efficiently
12 into mouse brain circuits in vivo. *Neuron* 2013;77(3):440–456.
- 13 88. Arthur A, Shi S, Zannettino AC, Fujii N, Gronthos S, Koblar SA. Implanted adult human dental
14 pulp stem cells induce endogenous axon guidance. *Stem Cells* 2009;27(9):2229–2237.
- 15 89. Bronckaers A, Hilkens P, Martens W, Gervois P, Ratajczak J, Struys T, Lambrechts I. Mesenchy-
16 mal stem/stromal cells as a pharmacological and therapeutic approach to accelerate angiogenesis.
17 *Pharmacol Ther* 2014;143(2):181–196.
- 18 90. Hilkens P, Fanton Y, Martens W, Gervois P, Struys T, Politis C, Lambrechts I, Bronckaers A.
19 Pro-angiogenic impact of dental stem cells in vitro and in vivo. *Stem Cell Res* 2014;12(3):778–790.
- 20 91. Kusuma S, Shen YI, Hanjaya-Putra D, Mali P, Cheng L, Gerecht S. Self-organized vascular
21 networks from human pluripotent stem cells in a synthetic matrix. *Proc Natl Acad Sci USA*
22 2013;110(31):12601–12606.
- 23 92. Pierdomenico L, Bonsi L, Calvitti M, Rondelli D, Arpinati M, Chirumbolo G, Becchetti E, Mar-
24 chionni C, Alviano F, Fossati V, Staffolani N, Franchina M, Grossi A, Bagnara GP. Multipotent
25 mesenchymal stem cells with immunosuppressive activity can be easily isolated from dental pulp.
26 *Transplantation* 2005;80(6):836–842.
- 27 93. Uccelli A, Moretta L, Pistoia V. Mesenchymal stem cells in health and disease. *Nat Rev Immunol*
28 2008;8(9):726–736.
- 29 94. Scheibe F, Ladhoff J, Huck J, Grohmann M, Blazej K, Oersal A, Baeva N, Seifert M, Priller J.
30 Immune effects of mesenchymal stromal cells in experimental stroke. *J Cereb Blood Flow Metab*
31 2012;32(8):1578–1588.
- 32 95. Brenneman M, Sharma S, Harting M, Strong R, Cox CS, Jr., Aronowski J, Grotta JC, Savitz SI.
33 Autologous bone marrow mononuclear cells enhance recovery after acute ischemic stroke in young
34 and middle-aged rats. *J Cereb Blood Flow Metab* 2010;30(1):140–149.
- 35 96. Kamiya N, Ueda M, Igarashi H, Nishiyama Y, Suda S, Inaba T, Katayama Y. Intra-arterial
36 transplantation of bone marrow mononuclear cells immediately after reperfusion decreases brain
37 injury after focal ischemia in rats. *Life Sci* 2008;83(11–12):433–437.
- 38 97. Phruksaniyom C, Dharmasaroja P, Issaragrisil S. Bone marrow non-mesenchymal mononuclear
39 cells induce functional differentiation of neuroblastoma cells. *Exp Hematol Oncol* 2013;2(1):9.
- 40 98. Wagner DC, Bojko M, Peters M, Lorenz M, Voigt C, Kaminski A, Hasenclever D, Scholz M, Kranz
41 A, Weise G, Boltze J. Impact of age on the efficacy of bone marrow mononuclear cell transplantation
42 in experimental stroke. *Exp Transl Stroke Med* 2012;4(1):17.
- 43 99. Friedenstein AJ, Chailakhjan RK, Lalykina KS. The development of fibroblast colonies in mono-
44 layer cultures of guinea-pig bone marrow and spleen cells. *Cell Tissue Kinet* 1970;3(4):393–403.
- 45 100. Pittenger MF, Mackay AM, Beck SC, Jaiswal RK, Douglas R, Mosca JD, Moorman MA, Simonetti
46 DW, Craig S, Marshak DR. Multilineage potential of adult human mesenchymal stem cells. *Science*
47 1999;284(5411):143–147.
- 48 101. Zuk PA, Zhu M, Ashjian P, De Ugarte DA, Huang JI, Mizuno H, Alfonso ZC, Fraser JK, Ben-
49 haim P, Hedrick MH. Human adipose tissue is a source of multipotent stem cells. *Mol Biol Cell*
50 2002;13(12):4279–4295.
- 51 102. Gronthos S, Mankani M, Brahimi J, Robey PG, Shi S. Postnatal human dental pulp stem cells
(DPSCs) in vitro and in vivo. *Proc Natl Acad Sci USA* 2000;97(25):13625–13630.

103. Weiss ML, Medicetty S, Bledsoe AR, Rachakatla RS, Choi M, Merchav S, Luo Y, Rao MS, Velagaleti G, Troyer D. Human umbilical cord matrix stem cells: Preliminary characterization and effect of transplantation in a rodent model of Parkinson's disease. *Stem Cells* 2006;24(3):781–792.
104. Erices A, Conget P, Minguell JJ. Mesenchymal progenitor cells in human umbilical cord blood. *Br J Haematol* 2000;109(1):235–242.
105. Miura M, Gronthos S, Zhao M, Lu B, Fisher LW, Robey PG, Shi S. SHED: Stem cells from human exfoliated deciduous teeth. *Proc Natl Acad Sci USA* 2003;100(10):5807–5812.
106. Lee KD, Kuo TK, Whang-Peng J, Chung YF, Lin CT, Chou SH, Chen JR, Chen YP, Lee OK. In vitro hepatic differentiation of human mesenchymal stem cells. *Hepatology* 2004;40(6):1275–1284.
107. Toma C, Pittenger MF, Cahill KS, Byrne BJ, Kessler PD. Human mesenchymal stem cells differentiate to a cardiomyocyte phenotype in the adult murine heart. *Circulation* 2002;105(1):93–98.
108. Wislet-Gendebien S, Hans G, Leprince P, Rigo JM, Moonen G, Rogister B. Plasticity of cultured mesenchymal stem cells: Switch from nestin-positive to excitable neuron-like phenotype. *Stem Cells* 2005;23(3):392–402.
109. Hilkens P, Gervois P, Fanton Y, Vanormelingen J, Martens W, Struys T, Politis C, Lambrechts I, Bronckaers A. Effect of isolation methodology on stem cell properties and multilineage differentiation potential of human dental pulp stem cells. *Cell Tissue Res* 2013;353(1):65–78.
110. Dominici M, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini F, Krause D, Deans R, Keating A, Prockop D, Horwitz E. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy* 2006;8(4):315–317.
111. Kaiser S, Hackanson B, Follo M, Mehlhorn A, Geiger K, Ihorst G, Kapp U. BM cells giving rise to MSC in culture have a heterogeneous CD34 and CD45 phenotype. *Cytotherapy* 2007;9(5):439–450.
112. Struys T, Moreels M, Martens W, Donders R, Wolfs E, Lambrechts I. Ultrastructural and immunocytochemical analysis of multilineage differentiated human dental pulp- and umbilical cord-derived mesenchymal stem cells. *Cells Tissues Organs* 2011;193(6):366–378.
113. Anghileri E, Marconi S, Pignatelli A, Cifelli P, Galie M, Sbarbati A, Krampera M, Belluzzi O, Bonetti B. Neuronal differentiation potential of human adipose-derived mesenchymal stem cells. *Stem Cells Dev* 2008;17(5):909–916.
114. Yang H, Xie Z, Wei L, Yang H, Yang S, Zhu Z, Wang P, Zhao C, Bi J. Human umbilical cord mesenchymal stem cell-derived neuron-like cells rescue memory deficits and reduce amyloid-beta deposition in an A β PP/PS1 transgenic mouse model. *Stem Cell Res Ther* 2013;4(4):76.
115. Kiraly M, Porcsalmy B, Pataki A, Kadar K, Jelitai M, Molnar B, Hermann P, Gera I, Grimm WD, Ganss B, Zsembery A, Varga G. Simultaneous PKC and cAMP activation induces differentiation of human dental pulp stem cells into functionally active neurons. *Neurochem Int* 2009;55(5):323–332.
116. Hau S, Reich DM, Scholz M, Naumann W, Emmrich F, Kamprad M, Boltze J. Evidence for neuroprotective properties of human umbilical cord blood cells after neuronal hypoxia in vitro. *BMC Neurosci* 2008;9:30.
117. Liu Y, Zhang Y, Lin L, Lin F, Li T, Du H, Chen R, Zheng W, Liu N. Effects of bone marrow-derived mesenchymal stem cells on the axonal outgrowth through activation of PI3K/AKT signaling in primary cortical neurons followed oxygen-glucose deprivation injury. *PLoS One* 2013;8(11):e78514.
118. Crigler L, Robey RC, Asawachaicharn A, Gaupp D, Phinney DG. Human mesenchymal stem cell subpopulations express a variety of neuro-regulatory molecules and promote neuronal cell survival and neurogenesis. *Exp Neurol* 2006;198(1):54–64.
119. Kingham PJ, Kolar MK, Novikova LN, Novikov LN, Wiberg M. Stimulating the neurotrophic and angiogenic properties of human adipose-derived stem cells enhances nerve repair. *Stem Cells Dev* 2014;23(7):741–754.
120. Wilkins A, Kemp K, Ginty M, Hares K, Mallam E, Scolding N. Human bone marrow-derived mesenchymal stem cells secrete brain-derived neurotrophic factor which promotes neuronal survival in vitro. *Stem Cell Res* 2009;3(1):63–70.

1 36 • GERVOIS ET AL.

- 2 121. Nosrat IV, Smith CA, Mullally P, Olson L, Nosrat CA. Dental pulp cells provide neurotrophic
3 support for dopaminergic neurons and differentiate into neurons in vitro; implications for tissue
4 engineering and repair in the nervous system. *Eur J Neurosci* 2004;19(9):2388–2398.
- 5 122. Martens W, Sanen K, Georgiou M, Struys T, Bronckaers A, Ameloot M, Phillips J, Lambrechts I.
6 Human dental pulp stem cells can differentiate into Schwann cells and promote and guide neurite
7 outgrowth in an aligned tissue-engineered collagen construct in vitro. *FASEB J* 2014;28(4):1634–
8 1643.
- 9 123. Choi M, Lee HS, Naidansaren P, Kim HK, O E, Cha JH, Ahn HY, Yang PI, Shin JC, Joe YA.
10 Proangiogenic features of Wharton's jelly-derived mesenchymal stromal/stem cells and their ability
11 to form functional vessels. *Int J Biochem Cell Biol* 2013;45(3):560–570.
- 12 124. Bronckaers A, Hilkens P, Fanton Y, Struys T, Gervois P, Politis C, Martens W, Lambrechts I.
13 Angiogenic properties of human dental pulp stem cells. *PLoS One* 2013;8(8):e71104.
- 14 125. Hung SC, Pochampally RR, Chen SC, Hsu SC, Prockop DJ. Angiogenic effects of human mul-
15 tipotent stromal cell conditioned medium activate the PI3K-Akt pathway in hypoxic endothelial
16 cells to inhibit apoptosis, increase survival, and stimulate angiogenesis. *Stem Cells* 2007;25(9):2363–
17 2370.
- 18 126. Schaar KL, Breneman MM, Savitz SI. Functional assessments in the rodent stroke model. *Exp*
19 *Transl Stroke Med* 2010;2(1):13.
- 20 127. Toyoshima A, Yasuhara T, Kameda M, Morimoto J, Takeuchi H, Wang F, Sasaki T, Sasada S,
21 Shinko A, Wakamori T, Okazaki M, Kondo A, Agari T, Borlongan CV, Date I. Intra-arterial
22 transplantation of allogeneic mesenchymal stem cells mounts neuroprotective effects in a transient
23 ischemic stroke model in rats: Analyses of therapeutic time window and its mechanisms. *PLoS One*
24 2015;10(6):e0127302.
- 25 128. Ishizaka S, Horie N, Satoh K, Fukuda Y, Nishida N, Nagata I. Intra-arterial cell transplan-
26 tation provides timing-dependent cell distribution and functional recovery after stroke. *Stroke*
27 2013;44(3):720–726.
- 28 129. Doepfner TR, Herz J, Gorgens A, Schlechter J, Ludwig AK, Radtke S, de Miroshedji K, Horn PA,
29 Giebel B, Hermann DM. Extracellular vesicles improve post-stroke neuroregeneration and prevent
30 postischemic immunosuppression. *Stem Cells Transl Med* 2015;4(10):1131–1143.
- 31 130. Liao W, Xie J, Zhong J, Liu Y, Du L, Zhou B, Xu J, Liu P, Yang S, Wang J, Han Z, Han ZC.
32 Therapeutic effect of human umbilical cord multipotent mesenchymal stromal cells in a rat model
33 of stroke. *Transplantation* 2009;87(3):350–359.
- 34 131. Jiang W, Liang G, Li X, Li Z, Gao X, Feng S, Wang X, Liu M, Liu Y. Intracarotid transplantation
35 of autologous adipose-derived mesenchymal stem cells significantly improves neurological deficits
36 in rats after MCAo. *J Mater Sci Mater Med* 2014;25(5):1357–1366.
- 37 132. Leu S, Lin YC, Yuen CM, Yen CH, Kao YH, Sun CK, Yip HK. Adipose-derived mesenchymal
38 stem cells markedly attenuate brain infarct size and improve neurological function in rats. *J Transl*
39 *Med* 2010;8:63.
- 40 133. Lee SH, Jin KS, Bang OY, Kim BJ, Park SJ, Lee NH, Yoo KH, Koo HH, Sung KW. Differential
41 migration of mesenchymal stem cells to ischemic regions after middle cerebral artery occlusion in
42 rats. *PLoS One* 2015;10(8):e0134920.
- 43 134. Acosta SA, Tajiri N, Hoover J, Kaneko Y, Borlongan CV. Intravenous bone marrow stem
44 cell grafts preferentially migrate to spleen and abrogate chronic inflammation in stroke. *Stroke*
45 2015;46(9):2616–2627.
- 46 135. Wakabayashi K, Nagai A, Sheikh AM, Shiota Y, Narantuya D, Watanabe T, Masuda J, Kobayashi
47 S, Kim SU, Yamaguchi S. Transplantation of human mesenchymal stem cells promotes functional
48 improvement and increased expression of neurotrophic factors in a rat focal cerebral ischemia model.
49 *J Neurosci Res* 2010;88(5):1017–1025.
- 50 136. Liu H, Honmou O, Harada K, Nakamura K, Houkin K, Hamada H, Kocsis JD. Neuroprotection
51 by PIGF gene-modified human mesenchymal stem cells after cerebral ischaemia. *Brain* 2006;129(Pt
10):2734–2745.

137. Yasuhara T, Matsukawa N, Hara K, Maki M, Ali MM, Yu SJ, Bae E, Yu G, Xu L, McGrogan M, Bankiewicz K, Case C, Borlongan CV. Notch-induced rat and human bone marrow stromal cell grafts reduce ischemic cell loss and ameliorate behavioral deficits in chronic stroke animals. *Stem Cells Dev* 2009;18(10):1501–1514.
138. Kubis N, Tomita Y, Tran-Dinh A, Planat-Benard V, Andre M, Karaszewski B, Waeckel L, Penicaud L, Silvestre JS, Casteilla L, Seylaz J, Pinard E. Vascular fate of adipose tissue-derived adult stromal cells in the ischemic murine brain: A combined imaging-histological study. *Neuroimage* 2007;34(1):1–11.
139. Ikegame Y, Yamashita K, Hayashi S, Mizuno H, Tawada M, You F, Yamada K, Tanaka Y, Egashira Y, Nakashima S, Yoshimura S, Iwama T. Comparison of mesenchymal stem cells from adipose tissue and bone marrow for ischemic stroke therapy. *Cytotherapy* 2011;13(6):675–685.
140. Wei N, Yu SP, Gu X, Taylor TM, Song D, Liu XF, Wei L. Delayed intranasal delivery of hypoxic-preconditioned bone marrow mesenchymal stem cells enhanced cell homing and therapeutic benefits after ischemic stroke in mice. *Cell Transplant* 2013;22(6):977–991.
141. Yu X, Chen D, Zhang Y, Wu X, Huang Z, Zhou H, Zhang Y, Zhang Z. Overexpression of CXCR4 in mesenchymal stem cells promotes migration, neuroprotection and angiogenesis in a rat model of stroke. *J Neurol Sci* 2012;316(1–2):141–149.
142. Zhou F, Gao S, Wang L, Sun C, Chen L, Yuan P, Zhao H, Yi Y, Qin Y, Dong Z, Cao L, Ren H, Zhu L, Li Q, Lu B, Liang A, Xu GT, Zhu H, Gao Z, Ma J, Xu J, Chen X. Human adipose-derived stem cells partially rescue the stroke syndromes by promoting spatial learning and memory in mouse middle cerebral artery occlusion model. *Stem Cell Res Ther* 2015;6:92.
143. Walczak P, Zhang J, Gilad AA, Kedziorek DA, Ruiz-Cabello J, Young RG, Pittenger MF, van Zijl PC, Huang J, Bulte JW. Dual-modality monitoring of targeted intraarterial delivery of mesenchymal stem cells after transient ischemia. *Stroke* 2008;39(5):1569–1574.
144. Mitkari B, Kerkela E, Nystedt J, Korhonen M, Mikkonen V, Huhtala T, Jolkkonen J. Intra-arterial infusion of human bone marrow-derived mesenchymal stem cells results in transient localization in the brain after cerebral ischemia in rats. *Exp Neurol* 2013;239:158–162.
145. Detante O, Moisan A, Dimastromatteo J, Richard MJ, Riou L, Grillon E, Barbier E, Desruet MD, De Fraipont F, Segebarth C, Jaillard A, Hommel M, Ghezzi C, Remy C. Intravenous administration of ^{99m}Tc-HMPAO-labeled human mesenchymal stem cells after stroke: In vivo imaging and biodistribution. *Cell Transplant* 2009;18(12):1369–1379.
146. Shen LH, Li Y, Chen J, Zacharek A, Gao Q, Kapke A, Lu M, Raginski K, Vanguri P, Smith A, Chopp M. Therapeutic benefit of bone marrow stromal cells administered 1 month after stroke. *J Cereb Blood Flow Metab* 2007;27(1):6–13.
147. Shen LH, Li Y, Chen J, Cui Y, Zhang C, Kapke A, Lu M, Savant-Bhonsale S, Chopp M. One-year follow-up after bone marrow stromal cell treatment in middle-aged female rats with stroke. *Stroke* 2007;38(7):2150–2156.
148. Zhang L, Yi L, Chopp M, Kramer BC, Romanko M, Gosiewska A, Hong K. Intravenous administration of human umbilical tissue-derived cells improves neurological function in aged rats after embolic stroke. *Cell Transplant* 2013;22(9):1569–1576.
149. Balseanu AT, Buga AM, Catalin B, Wagner DC, Boltze J, Zagrean AM, Reymann K, Schaebitz W, Popa-Wagner A. Multimodal approaches for regenerative stroke therapies: Combination of granulocyte colony-stimulating factor with bone marrow mesenchymal stem cells is not superior to G-CSF alone. *Front Aging Neurosci* 2014;6:130.
150. Shen LH, Li Y, Chen J, Zhang J, Vanguri P, Borneman J, Chopp M. Intracarotid transplantation of bone marrow stromal cells increases axon-myelin remodeling after stroke. *Neuroscience* 2006;137(2):393–399.
151. Gutierrez-Fernandez M, Rodriguez-Frutos B, Ramos-Cejudo J, Otero-Ortega L, Fuentes B, Vallejo-Cremades MT, Sanz-Cuesta BE, Diez-Tejedor E. Comparison between xenogeneic and allogeneic adipose mesenchymal stem cells in the treatment of acute cerebral infarct: Proof of concept in rats. *J Transl Med* 2015;13:46.

1 38 • GERVOIS ET AL.

- 2 152. Taguchi A, Zhu P, Cao F, Kikuchi-Taura A, Kasahara Y, Stern DM, Soma T, Matsuyama T, Hata
3 R. Reduced ischemic brain injury by partial rejuvenation of bone marrow cells in aged rats. *J Cereb*
4 *Blood Flow Metab* 2011;31(3):855–867.
- 5 153. Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, Yamanaka S. Induction of
6 pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 2007;131(5):861–872.
- 7 154. Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult
8 fibroblast cultures by defined factors. *Cell* 2006;126(4):663–676.
- 9 155. Seminatore C, Polentes J, Ellman D, Kozubenko N, Itier V, Tine S, Tritschler L, Brenot M, Guidou
10 E, Blondeau J, Lhuillier M, Bugi A, Aubry L, Jendelova P, Sykova E, Perrier AL, Finsen B, Onte-
11 niente B. The postischemic environment differentially impacts teratoma or tumor formation after
12 transplantation of human embryonic stem cell-derived neural progenitors. *Stroke* 2010;41(1):153–
13 159.
- 14 156. Liu Z, Tang Y, Lu S, Zhou J, Du Z, Duan C, Li Z, Wang C. The tumorigenicity of iPS cells and
15 their differentiated derivatives. *J Cell Mol Med* 2013;17(6):782–791.
- 16 157. Choi HW, Kim JS, Choi S, Hong YJ, Kim MJ, Seo HG, Do JT. Neural stem cells differentiated
17 from iPS cells spontaneously regain pluripotency. *Stem Cells* 2014;32(10):2596–2604.
- 18 158. Fu W, Wang SJ, Zhou GD, Liu W, Cao Y, Zhang WJ. Residual undifferentiated cells during
19 differentiation of induced pluripotent stem cells in vitro and in vivo. *Stem Cells Dev* 2012;21(4):521–
20 529.
- 21 159. Hu BY, Weick JP, Yu J, Ma LX, Zhang XQ, Thomson JA, Zhang SC. Neural differentiation of
22 human induced pluripotent stem cells follows developmental principles but with variable potency.
23 *Proc Natl Acad Sci USA* 2010;107(9):4335–4340.
- 24 160. Fink KD, Crane AT, Leveque X, Dues DJ, Huffman LD, Moore AC, Story DT, Dejonge RE,
25 Antcliff A, Starski PA, Lu M, Lescaudron L, Rossignol J, Dunbar GL. Intraatrial transplantation
26 of adenovirus-generated induced pluripotent stem cells for treating neuropathological and functional
27 deficits in a rodent model of Huntington's disease. *Stem Cells Transl Med* 2014;3(5):620–631.
- 28 161. Wernig M, Zhao JP, Pruszak J, Hedlund E, Fu D, Soldner F, Broccoli V, Constantine-Paton M,
29 Isacson O, Jaenisch R. Neurons derived from reprogrammed fibroblasts functionally integrate into
30 the fetal brain and improve symptoms of rats with Parkinson's disease. *Proc Natl Acad Sci USA*
31 2008;105(15):5856–5861.
- 32 162. Raitano S, Ordovas L, De Muynck L, Guo W, Espuny-Camacho I, Geraerts M, Khurana S,
33 Vanuytsel K, Toth BI, Voets T, Vandenberghe R, Cathomen T, Van Den Bosch L, Vanderhaeghen
34 P, Van Damme P, Verfaillie CM. Restoration of progranulin expression rescues cortical neuron
35 generation in an induced pluripotent stem cell model of frontotemporal dementia. *Stem Cell Reports*
36 2015;4(1):16–24.
- 37 163. Chambers SM, Qi Y, Mica Y, Lee G, Zhang XJ, Niu L, Bilsland J, Cao L, Stevens E, Whiting P, Shi
38 SH, Studer L. Combined small-molecule inhibition accelerates developmental timing and converts
39 human pluripotent stem cells into nociceptors. *Nat Biotechnol* 2012;30(7):715–720.
- 40 164. Yuan T, Liao W, Feng NH, Lou YL, Niu X, Zhang AJ, Wang Y, Deng ZF. Human induced
41 pluripotent stem cell-derived neural stem cells survive, migrate, differentiate, and improve neurologic
42 function in a rat model of middle cerebral artery occlusion. *Stem Cell Res Ther* 2013;4(3):73.
- 43 165. Song G, Li X, Shen Y, Qian L, Kong X, Chen M, Cao K, Zhang F. Transplantation of iPSC restores
44 cardiac function by promoting angiogenesis and ameliorating cardiac remodeling in a post-infarcted
45 swine model. *Cell Biochem Biophys* 2014.
- 46 166. Zhou Y, Wang S, Yu Z, Hoyt RF, Jr., Hunt T, Kindzelski B, Shou D, Xie W, Du Y, Liu C, Horvath
47 KA. Induced pluripotent stem cell transplantation in the treatment of porcine chronic myocardial
48 ischemia. *Ann Thorac Surg* 2014;98(6):2130–2137.
- 49 167. Belair DG, Whisler JA, Valdez J, Velazquez J, Molenda JA, Vickerman V, Lewis R, Daigh C, Hansen
50 TD, Mann DA, Thomson JA, Griffith LG, Kamm RD, Schwartz MP, Murphy WL. Human vascular
51 tissue models formed from human induced pluripotent stem cell derived endothelial cells. *Stem Cell*
52 *Rev* 2014.

Q16

STEM CELL-BASED THERAPIES FOR ISCHEMIC STROKE • 39

168. Choi KD, Yu J, Smuga-Otto K, Salvaggio G, Rehauer W, Vodyanik M, Thomson J, Slukvin I. Hematopoietic and endothelial differentiation of human induced pluripotent stem cells. *Stem Cells* 2009;27(3):559–567.
169. Polentes J, Jendelova P, Cailleret M, Braun H, Romanyuk N, Tropel P, Brenot M, Itier V, Seminatore C, Baldauf K, Turnovcova K, Jirak D, Teletin M, Come J, Tournois J, Reymann K, Sykova E, Viville S, Onteniente B. Human induced pluripotent stem cells improve stroke outcome and reduce secondary degeneration in the recipient brain. *Cell Transplant* 2012;21(12):2587–2602.
170. Tatarishvili J, Oki K, Monni E, Koch P, Memanishvili T, Buga AM, Verma V, Popa-Wagner A, Brustle O, Lindvall O, Kokaia Z. Human induced pluripotent stem cells improve recovery in stroke-injured aged rats. *Restor Neurol Neurosci* 2014;32(4):547–558.
171. Jensen MB, Yan H, Krishnaney-Davison R, Al Sawaf A, Zhang SC. Survival and differentiation of transplanted neural stem cells derived from human induced pluripotent stem cells in a rat stroke model. *J Stroke Cerebrovasc Dis* 2013;22(4):304–308.
172. Eckert A, Huang L, Gonzalez R, Kim HS, Hamblin MH, Lee JP. Bystander effect fuels human induced pluripotent stem cell-derived neural stem cells to quickly attenuate early stage neurological deficits after stroke. *Stem Cells Transl Med* 2015;4(7):841–851.
173. Lam J, Lowry WE, Carmichael ST, Segura T. Delivery of iPS-NPCs to the stroke cavity within a hyaluronic acid matrix promotes the differentiation of transplanted cells. *Adv Funct Mater* 2014;24(44):7053–7062.
174. Majid A, He YY, Gidday JM, Kaplan SS, Gonzales ER, Park TS, Fenstermacher JD, Wei L, Choi DW, Hsu CY. Differences in vulnerability to permanent focal cerebral ischemia among 3 common mouse strains. *Stroke* 2000;31(11):2707–2714.
175. Barone FC, Knudsen DJ, Nelson AH, Feuerstein GZ, Willette RN. Mouse strain differences in susceptibility to cerebral ischemia are related to cerebral vascular anatomy. *J Cereb Blood Flow Metab* 1993;13(4):683–692.
176. Sauter A, Rudin M. Strain-dependent drug effects in rat middle cerebral artery occlusion model of stroke. *J Pharmacol Exp Ther* 1995;274(2):1008–1013.
177. Braeuninger S, Kleinschnitz C. Rodent models of focal cerebral ischemia: Procedural pitfalls and translational problems. *Exp Transl Stroke Med* 2009;1:8.
178. Popa-Wagner A, Carmichael ST, Kokaia Z, Kessler C, Walker LC. The response of the aged brain to stroke: Too much, too soon? *Curr Neurovasc Res* 2007;4(3):216–227.
179. Darsalia V, Allison SJ, Cusulin C, Monni E, Kuzdas D, Kallur T, Lindvall O, Kokaia Z. Cell number and timing of transplantation determine survival of human neural stem cell grafts in stroke-damaged rat brain. *J Cereb Blood Flow Metab* 2011;31(1):235–242.
180. Hermann DM, Zechariah A. Implications of vascular endothelial growth factor for postischemic neurovascular remodeling. *J Cereb Blood Flow Metab* 2009;29(10):1620–1643.
181. Horie N, Pereira MP, Niizuma K, Sun G, Keren-Gill H, Encarnacion A, Shamloo M, Hamilton SA, Jiang K, Huhn S, Palmer TD, Bliss TM, Steinberg GK. Transplanted stem cell-secreted vascular endothelial growth factor effects poststroke recovery, inflammation, and vascular repair. *Stem Cells* 2011;29(2):274–285.
182. Chen SJ, Chang CM, Tsai SK, Chang YL, Chou SJ, Huang SS, Tai LK, Chen YC, Ku HH, Li HY, Chiou SH. Functional improvement of focal cerebral ischemia injury by subdural transplantation of induced pluripotent stem cells with fibrin glue. *Stem Cells Dev* 2010;19(11):1757–1767.
183. Burns TC, Ortiz-Gonzalez XR, Gutierrez-Perez M, Keene CD, Sharda R, Demorest ZL, Jiang Y, Nelson-Holte M, Soriano M, Nakagawa Y, Luquin MR, Garcia-Verdugo JM, Prosper F, Low WC, Verfaillie CM. Thymidine analogs are transferred from prelabeled donor to host cells in the central nervous system after transplantation: A word of caution. *Stem Cells* 2006;24(4):1121–1127.
184. Massoud TF, Gambhir SS. Molecular imaging in living subjects: Seeing fundamental biological processes in a new light. *Genes Dev* 2003;17(5):545–580.
185. Arbab AS, Bashaw LA, Miller BR, Jordan EK, Lewis BK, Kalish H, Frank JA. Characterization of biophysical and metabolic properties of cells labeled with superparamagnetic iron ox-

1 40 • GERVOIS ET AL.

- 2 ide nanoparticles and transfection agent for cellular MR imaging. *Radiology* 2003;229(3):838–
3 846.
- 4 186. Struys T, Ketkar-Atre A, Gervois P, Leten C, Hilkens P, Martens W, Bronckaers A, Dresselaers T,
5 Politis C, Lambrichts I, Himmelreich U. Magnetic resonance imaging of human dental pulp stem
6 cells in vitro and in vivo. *Cell Transplant* 2013;22(10):1813–1829.
- 7 187. Hoehn M, Kustermann E, Blunk J, Wiedermann D, Trapp T, Wecker S, Focking M, Arnold H,
8 Hescheler J, Fleischmann BK, Schwandt W, Buhrle C. Monitoring of implanted stem cell migration
9 in vivo: A highly resolved in vivo magnetic resonance imaging investigation of experimental stroke
10 in rat. *Proc Natl Acad Sci USA* 2002;99(25):16267–16272.
- 11 188. Gu E, Chen WY, Gu J, Burrige P, Wu JC. Molecular imaging of stem cells: Tracking survival,
12 biodistribution, tumorigenicity, and immunogenicity. *Theranostics* 2012;2(4):335–345.
- 13 189. Bulte JW. In vivo MRI cell tracking: Clinical studies. *AJR Am J Roentgenol* 2009;193(2):314–325.
- 14 190. Zhang Y, Dasilva JN, Hadizad T, Thorn S, Kuraitis D, Renaud JM, Ahmadi A, Kordos M, Dekemp
15 RA, Beanlands RS, Suuronen EJ, Ruel M. (18)F-FDG cell labeling may underestimate transplanted
16 cell homing: More accurate, efficient, and stable cell labeling with hexadecyl-4-[(18)F]fluorobenzoate
17 for in vivo tracking of transplanted human progenitor cells by positron emission tomography. *Cell*
18 *Transplant* 2012;21(9):1821–1835.
- 19 191. Deroose CM, Reumers V, Debysier Z, Baekelandt V. Seeing genes at work in the living brain with
20 non-invasive molecular imaging. *Curr Gene Ther* 2009;9(3):212–238.
- 21 192. Yaghoubi SS, Campbell DO, Radu CG, Czernin J. Positron emission tomography reporter genes
22 and reporter probes: Gene and cell therapy applications. *Theranostics* 2012;2(4):374–391.
- 23 193. Serganova I, Ponomarev V, Blasberg R. Human reporter genes: Potential use in clinical studies.
24 *Nucl Med Biol* 2007;34(7):791–807.
- 25 194. Hofmann M, Wollert KC, Meyer GP, Menke A, Arseniev L, Hertenstein B, Ganser A, Knapp
26 WH, Drexler H. Monitoring of bone marrow cell homing into the infarcted human myocardium.
27 *Circulation* 2005;111(17):2198–2202.
- 28 195. Kang WJ, Kang HJ, Kim HS, Chung JK, Lee MC, Lee DS. Tissue distribution of 18F-FDG-labeled
29 peripheral hematopoietic stem cells after intracoronary administration in patients with myocardial
30 infarction. *J Nucl Med* 2006;47(8):1295–1301.
- 31 196. Elhami E, Goertzen AL, Xiang B, Deng J, Stillwell C, Mzengeza S, Arora RC, Freed D, Tian G.
32 Viability and proliferation potential of adipose-derived stem cells following labeling with a positron-
33 emitting radiotracer. *Eur J Nucl Med Mol Imaging* 2011;38(7):1323–1334.
- 34 197. Chen MF, Lin CT, Chen WC, Yang CT, Chen CC, Liao SK, Liu JM, Lu CH, Lee KD. The
35 sensitivity of human mesenchymal stem cells to ionizing radiation. *Int J Radiat Oncol Biol Phys*
36 2006;66(1):244–253.
- 37 198. Wolfs E, Struys T, Notelaers T, Roberts SJ, Sohni A, Bormans G, Van Laere K, Luyten FP, Gheysens
38 O, Lambrichts I, Verfaillie CM, Deroose CM. 18F-FDG labeling of mesenchymal stem cells and
39 multipotent adult progenitor cells for PET imaging: Effects on ultrastructure and differentiation
40 capacity. *J Nucl Med* 2013;54(3):447–454.
- 41 199. Correa PL, Mesquita CT, Felix RM, Azevedo JC, Barbirato GB, Falcao CH, Gonzalez C, Men-
42 donca ML, Manfrim A, de Freitas G, Oliveira CC, Silva D, Avila D, Borojevic R, Alves S, Oliveira
43 AC, Jr., Dohmann HF. Assessment of intra-arterial injected autologous bone marrow mononuclear
44 cell distribution by radioactive labeling in acute ischemic stroke. *Clin Nucl Med* 2007;32(11):839–
45 841.
- 46 200. Schachinger V, Aicher A, Dobert N, Rover R, Diener J, Fichtlscherer S, Assmus B, Seeger FH,
47 Menzel C, Brenner W, Dimmeler S, Zeiher AM. Pilot trial on determinants of progenitor cell
48 recruitment to the infarcted human myocardium. *Circulation* 2008;118(14):1425–1432.
- 49 201. Cavelliers V, De Keulenaer G, Everaert H, Van Riet I, Van Camp G, Verheye S, Roland J, Schoors
50 D, Franken PR, Schots R. In vivo visualization of 111In labeled CD133+ peripheral blood stem
51 cells after intracoronary administration in patients with chronic ischemic heart disease. *Q J Nucl*
Med Mol Imaging 2007;51(1):61–66.

202. Brenner W, Aicher A, Eckey T, Massoudi S, Zuhayra M, Koehl U, Heeschen C, Kampen WU, Zeiher AM, Dimmeler S, Henze E. 111In-labeled CD34+ hematopoietic progenitor cells in a rat myocardial infarction model. *J Nucl Med* 2004;45(3):512–518.
203. Nowak B, Weber C, Schober A, Zeiffer U, Liehn EA, von Hundelshausen P, Reinartz P, Schaefer WM, Buell U. Indium-111 oxine labelling affects the cellular integrity of haematopoietic progenitor cells. *Eur J Nucl Med Mol Imaging* 2007;34(5):715–721.
204. Tsien RY. The green fluorescent protein. *Annu Rev Biochem* 1998;67:509–544.
205. de Wet JR, Wood KV, DeLuca M, Helinski DR, Subramani S. Firefly luciferase gene: Structure and expression in mammalian cells. *Mol Cell Biol* 1987;7(2):725–737.
206. Wolfs E, Holvoet B, Gijsbers R, Casteels C, Roberts SJ, Struys T, Maris M, Ibrahim A, Debyser Z, Van Laere K, Verfaillie CM, Deroose CM. Optimization of multimodal imaging of mesenchymal stem cells using the human sodium iodide symporter for PET and Cerenkov luminescence imaging. *PLoS One* 2014;9(4):e94833.
207. Kim DE, Schellingerhout D, Ishii K, Shah K, Weissleder R. Imaging of stem cell recruitment to ischemic infarcts in a murine model. *Stroke* 2004;35(4):952–957.
208. Vandeputte C, Reumers V, Aelvoet SA, Thiry I, De Swaef S, Van den Haute C, Pascual-Brazo J, Farr TD, Vande Velde G, Hoehn M, Himmelreich U, Van Laere K, Debyser Z, Gijsbers R, Baekelandt V. Bioluminescence imaging of stroke-induced endogenous neural stem cell response. *Neurobiol Dis* 2014;69:144–155.
209. Contag CH, Bachmann MH. Advances in in vivo bioluminescence imaging of gene expression. *Annu Rev Biomed Eng* 2002;4:235–260.
210. Shaner NC, Campbell RE, Steinbach PA, Giepmans BN, Palmer AE, Tsien RY. Improved monomeric red, orange and yellow fluorescent proteins derived from *Discosoma* sp. red fluorescent protein. *Nat Biotechnol* 2004;22(12):1567–1572.
211. Vandsburger MH, Radoul M, Cohen B, Neeman M. MRI reporter genes: Applications for imaging of cell survival, proliferation, migration and differentiation. *NMR Biomed* 2013;26(7):872–884.
212. Patrick PS, Hammersley J, Loizou L, Kettunen MI, Rodrigues TB, Hu DE, Tee SS, Hesketh R, Lyons SK, Soloviev D, Lewis DY, Aime S, Fulton SM, Brindle KM. Dual-modality gene reporter for in vivo imaging. *Proc Natl Acad Sci USA* 2014;111(1):415–420.
213. van Zijl PC, Yadav NN. Chemical exchange saturation transfer (CEST): What is in a name and what isn't? *Magn Reson Med* 2011;65(4):927–948.
214. Rueger MA, Backes H, Walberer M, Neumaier B, Ullrich R, Simard ML, Emig B, Fink GR, Hoehn M, Graf R, Schroeter M. Noninvasive imaging of endogenous neural stem cell mobilization in vivo using positron emission tomography. *J Neurosci* 2010;30(18):6454–6460.
215. Tjuvajev JG, Avril N, Oku T, Sasajima T, Miyagawa T, Joshi R, Safer M, Beattie B, DiResta G, Daghigian F, Augensen F, Koutcher J, Zweit J, Humm J, Larson SM, Finn R, Blasberg R. Imaging herpes virus thymidine kinase gene transfer and expression by positron emission tomography. *Cancer Res* 1998;58(19):4333–4341.
216. Jacobs A, Braunlich I, Graf R, Lercher M, Sakaki T, Voges J, Hesselmann V, Brandau W, Wienhard K, Heiss WD. Quantitative kinetics of [124I]FIAU in cat and man. *J Nucl Med* 2001;42(3):467–475.
217. Yaghoubi S, Barrio JR, Dahlbom M, Iyer M, Namavari M, Satyamurthy N, Goldman R, Herschman HR, Phelps ME, Gambhir SS. Human pharmacokinetic and dosimetry studies of [(18)F]FHBG: A reporter probe for imaging herpes simplex virus type-1 thymidine kinase reporter gene expression. *J Nucl Med* 2001;42(8):1225–1234.
218. MacLaren DC, Gambhir SS, Satyamurthy N, Barrio JR, Sharfstein S, Toyokuni T, Wu L, Berk AJ, Cherry SR, Phelps ME, Herschman HR. Repetitive, non-invasive imaging of the dopamine D2 receptor as a reporter gene in living animals. *Gene Ther* 1999;6(5):785–791.
219. Zhang H, Moroz MA, Serganova I, Ku T, Huang R, Vider J, Maecke HR, Larson SM, Blasberg R, Smith-Jones PM. Imaging expression of the human somatostatin receptor subtype-2 reporter gene with 68Ga-DOTATOC. *J Nucl Med* 2011;52(1):123–131.

1 **42** • GERVOIS ET AL.

- 2 220. Penheiter AR, Russell SJ, Carlson SK. The sodium iodide symporter (NIS) as an imaging reporter
3 for gene, viral, and cell-based therapies. *Curr Gene Ther* 2012;12(1):33–47.
- 4 221. Dwyer RM, Ryan J, Havelin RJ, Morris JC, Miller BW, Liu Z, Flavin R, O'Flatharta C, Foley
5 MJ, Barrett HH, Murphy JM, Barry FP, O'Brien T, Kerin MJ. Mesenchymal stem cell-mediated
6 delivery of the sodium iodide symporter supports radionuclide imaging and treatment of breast
7 cancer. *Stem Cells* 2011;29(7):1149–1157.
- 8 222. Templin C, Zweigerdt R, Schwanke K, Olmer R, Ghadri JR, Emmert MY, Muller E, Kuest SM,
9 Cohrs S, Schibli R, Kronen P, Hilbe M, Reinisch A, Strunk D, Haverich A, Hoerstrup S, Luseher
10 TF, Kaufmann PA, Landmesser U, Martin U. Transplantation and tracking of human-induced
11 pluripotent stem cells in a pig model of myocardial infarction: Assessment of cell survival, en-
12 graftment, and distribution by hybrid single photon emission computed tomography/computed
13 tomography of sodium iodide symporter transgene expression. *Circulation* 2012;126(4):430–
14 439.
- 15 223. Wang Y, Miao Z, Ren G, Xu Y, Cheng Z. A novel Affibody bioconjugate for dual-modality imaging
16 of ovarian cancer. *Chem Commun (Camb)* 2014;50(85):12832–12835.
- 17 224. Lee S, Chen X. Dual-modality probes for in vivo molecular imaging. *Mol Imaging* 2009;8(2):87–100.
- 18 225. Boehm-Sturm P, Aswendt M, Minassian A, Michalk S, Mengler L, Adamczak J, Mezzanotte L,
19 Lowik C, Hoehn M. A multi-modality platform to image stem cell graft survival in the naive and
20 stroke-damaged mouse brain. *Biomaterials* 2014;35(7):2218–2226.
- 21 226. Daadi MM, Li Z, Arac A, Grueter BA, Sofilos M, Malenka RC, Wu JC, Steinberg GK. Molecular
22 and magnetic resonance imaging of human embryonic stem cell-derived neural stem cell grafts in
23 ischemic rat brain. *Mol Ther* 2009;17(7):1282–1291.
- 24 227. Pendharkar AV, Chua JY, Andres RH, Wang N, Gaeta X, Wang H, De A, Choi R, Chen S, Rutt
25 BK, Gambhir SS, Guzman R. Biodistribution of neural stem cells after intravascular therapy for
26 hypoxic-ischemia. *Stroke* 2010;41(9):2064–2070.
- 27 228. Sigler A, Murphy TH. In vivo 2-photon imaging of fine structure in the rodent brain: Before, during,
28 and after stroke. *Stroke* 2010;41(10 Suppl):S117–S123.
- 29 229. Svoboda K, Yasuda R. Principles of two-photon excitation microscopy and its applications to
30 neuroscience. *Neuron* 2006;50(6):823–839.
- 31 230. Scheller A, Vivien D, Kirchhoff F, Orset C, Sandu RE, Popa-Wagner A. Imaging neuroinflammation
32 after brain injuries by ultrasensitive MRI and two-photon laser-scanning microscopy. *Rom J*
33 *Morphol Embryol* 2014;55(3):735–743.
- 34 231. Witte S, Negrean A, Lodder JC, de Kock CP, Testa Silva G, Mansvelter HD, Louise Groot M.
35 Label-free live brain imaging and targeted patching with third-harmonic generation microscopy.
36 *Proc Natl Acad Sci USA* 2011;108(15):5970–5975.
- 37 232. Schain AJ, Hill RA, Grutzendler J. Label-free in vivo imaging of myelinated axons in health and
38 disease with spectral confocal reflectance microscopy. *Nat Med* 2014;20(4):443–449.
- 39 233. Brown CE, Aminoltejari K, Erb H, Winship IR, Murphy TH. In vivo voltage-sensitive dye imaging in
40 adult mice reveals that somatosensory maps lost to stroke are replaced over weeks by new structural
41 and functional circuits with prolonged modes of activation within both the peri-infarct zone and
42 distant sites. *J Neurosci* 2009;29(6):1719–1734.
- 43 234. Akemann W, Mutoh H, Perron A, Rossier J, Knopfel T. Imaging brain electric signals with geneti-
44 cally targeted voltage-sensitive fluorescent proteins. *Nat Methods* 2010;7(8):643–649.
- 45 235. Crowe SE, Ellis-Davies GC. Longitudinal in vivo two-photon fluorescence imaging. *J Comp Neurol*
46 2014;522(8):1708–1727.
- 47 236. Adamczak JM, Schneider G, Nelles M, Que I, Suidgeest E, van der Weerd L, Lowik C, Hoehn M.
48 In vivo bioluminescence imaging of vascular remodeling after stroke. *Front Cell Neurosci* 2014;8:
49 274.
- 50 237. Quattromani MJ, Cordeau P, Ruscher K, Kriz J, Wieloch T. Enriched housing down-regulates
51 the Toll-like receptor 2 response in the mouse brain after experimental stroke. *Neurobiol Dis*
2014;66:66–73.

238. Wang J, Chao F, Han F, Zhang G, Xi Q, Li J, Jiang H, Wang J, Yu G, Tian M, Zhang H. PET demonstrates functional recovery after transplantation of induced pluripotent stem cells in a rat model of cerebral ischemic injury. *J Nucl Med* 2013;54(5):785–792.
239. Daadi MM, Hu S, Klausner J, Li Z, Sofilos M, Sun G, Wu JC, Steinberg GK. Imaging neural stem cell graft-induced structural repair in stroke. *Cell Transplant* 2013;22(5):881–892.
240. Cross DJ, Minoshima S. Perspectives on assessment of stem cell therapy in stroke by 18F-FDG PET. *J Nucl Med* 2013;54(5):668–669.
241. Zinnhardt B, Viel T, Wachsmuth L, Vrachimis A, Wagner S, Breyholz HJ, Faust A, Hermann S, Kopka K, Faber C, Dolle F, Pappata S, Planas AM, Tavitian B, Schafers M, Sorokin LM, Kuhlmann MT, Jacobs AH. Multimodal imaging reveals temporal and spatial microglia and matrix metalloproteinase activity after experimental stroke. *J Cereb Blood Flow Metab* 2015;35(11):1711–1721.
242. Nair DG. About being BOLD. *Brain Res Brain Res Rev* 2005;50(2):229–243.
243. Weber R, Ramos-Cabrer P, Justicia C, Wiedermann D, Strecker C, Sprenger C, Hoehn M. Early prediction of functional recovery after experimental stroke: Functional magnetic resonance imaging, electrophysiology, and behavioral testing in rats. *J Neurosci* 2008;28(5):1022–1029.
244. Neumann-Haefelin T, Wittsack HJ, Wenserski F, Siebler M, Seitz RJ, Modder U, Freund HJ. Diffusion- and perfusion-weighted MRI. The DWI/PWI mismatch region in acute stroke. *Stroke* 1999;30(8):1591–1597.
245. Meng X, Fisher M, Shen Q, Sotak CH, Duong TQ. Characterizing the diffusion/perfusion mismatch in experimental focal cerebral ischemia. *Ann Neurol* 2004;55(2):207–212.
246. Reid E, Graham D, Lopez-Gonzalez MR, Holmes WM, Macrae IM, McCabe C. Penumbra detection using PWI/DWI mismatch MRI in a rat stroke model with and without comorbidity: Comparison of methods. *J Cereb Blood Flow Metab* 2012;32(9):1765–1777.
247. Assaf Y, Pasternak O. Diffusion tensor imaging (DTI)-based white matter mapping in brain research: A review. *J Mol Neurosci* 2008;34(1):51–61.
248. Liu Y, D’Arceuil HE, Westmoreland S, He J, Duggan M, Gonzalez RG, Pryor J, de Crespigny AJ. Serial diffusion tensor MRI after transient and permanent cerebral ischemia in nonhuman primates. *Stroke* 2007;38(1):138–145.
249. Aoki I, Wu YJ, Silva AC, Lynch RM, Koretsky AP. In vivo detection of neuroarchitecture in the rodent brain using manganese-enhanced MRI. *Neuroimage* 2004;22(3):1046–1059.
250. Pautler RG, Silva AC, Koretsky AP. In vivo neuronal tract tracing using manganese-enhanced magnetic resonance imaging. *Magn Reson Med* 1998;40(5):740–748.
251. Massaad CA, Pautler RG. Manganese-enhanced magnetic resonance imaging (MEMRI). *Methods Mol Biol* 2011;711:145–174.
252. van der Zijden JP, Wu O, van der Toorn A, Roeling TP, Bleys RL, Dijkhuizen RM. Changes in neuronal connectivity after stroke in rats as studied by serial manganese-enhanced MRI. *Neuroimage* 2007;34(4):1650–1657.
253. van Meer MP, van der Marel K, Otte WM, Berkelbach van der Sprenkel JW, Dijkhuizen RM. Correspondence between altered functional and structural connectivity in the contralesional sensorimotor cortex after unilateral stroke in rats: A combined resting-state functional MRI and manganese-enhanced MRI study. *J Cereb Blood Flow Metab* 2010;30(10):1707–1711.
254. Buxton RB, Frank LR, Wong EC, Siewert B, Warach S, Edelman RR. A general kinetic model for quantitative perfusion imaging with arterial spin labeling. *Magn Reson Med* 1998;40(3):383–396.
255. Thomas DL. Arterial spin labeling in small animals: Methods and applications to experimental cerebral ischemia. *J Magn Reson Imaging* 2005;22(6):741–744.
256. Bratane BT, Walvick RP, Corot C, Lancelot E, Fisher M. Characterization of gadolinium-based dynamic susceptibility contrast perfusion measurements in permanent and transient MCAO models with volumetric based validation by CASL. *J Cereb Blood Flow Metab* 2010;30(2):336–342.
257. Bang OY, Lee JS, Lee PH, Lee G. Autologous mesenchymal stem cell transplantation in stroke patients. *Ann Neurol* 2005;57(6):874–882.

1 44 • GERVOIS ET AL.

- 2 258. Honmou O, Houkin K, Matsunaga T, Niitsu Y, Ishiai S, Onodera R, Waxman SG, Kocsis JD.
3 Intravenous administration of auto serum-expanded autologous mesenchymal stem cells in stroke.
4 *Brain* 2011;134(Pt 6):1790–1807.
- 5 259. Lee JS, Hong JM, Moon GJ, Lee PH, Ahn YH, Bang OY, Starting Collaborators. A long-term
6 follow-up study of intravenous autologous mesenchymal stem cell transplantation in patients with
7 ischemic stroke. *Stem Cells* 2010;28(6):1099–1106.
- 8 260. Li L, Jiang Q, Ding G, Zhang L, Zhang ZG, Li Q, Panda S, Lu M, Ewing JR, Chopp M. Effects of
9 administration route on migration and distribution of neural progenitor cells transplanted into rats
10 with focal cerebral ischemia, an MRI study. *J Cereb Blood Flow Metab* 2010;30(3):653–662.
- 11 261. Fischer UM, Harting MT, Jimenez F, Monzon-Posadas WO, Xue H, Savitz SI, Laine GA, Cox
12 CS, Jr. Pulmonary passage is a major obstacle for intravenous stem cell delivery: The pulmonary
13 first-pass effect. *Stem Cells Dev* 2009;18(5):683–692.
- 14 262. Vendrame M, Gemma C, Pennypacker KR, Bickford PC, Davis Sanberg C, Sanberg PR, Willing
15 AE. Cord blood rescues stroke-induced changes in splenocyte phenotype and function. *Exp Neurol*
16 2006;199(1):191–200.
- 17 263. Vu Q, Xie K, Eckert M, Zhao W, Cramer SC. Meta-analysis of preclinical studies of mesenchymal
18 stromal cells for ischemic stroke. *Neurology* 2014;82(14):1277–1286.
- 19 264. Olanow CW, Goetz CG, Kordower JH, Stoessl AJ, Sossi V, Brin MF, Shannon KM, Nauert
20 GM, Perl DP, Godbold J, Freeman TB. A double-blind controlled trial of bilateral fetal nigral
21 transplantation in Parkinson's disease. *Ann Neurol* 2003;54(3):403–414.
- 22 265. Okita K, Ichisaka T, Yamanaka S. Generation of germline-competent induced pluripotent stem
23 cells. *Nature* 2007;448(7151):313–317.
- 24 266. Nakagawa M, Koyanagi M, Tanabe K, Takahashi K, Ichisaka T, Aoi T, Okita K, Mochiduki Y,
25 Takizawa N, Yamanaka S. Generation of induced pluripotent stem cells without Myc from mouse
26 and human fibroblasts. *Nat Biotechnol* 2008;26(1):101–106.
- 27 267. Fusaki N, Ban H, Nishiyama A, Saeki K, Hasegawa M. Efficient induction of transgene-free human
28 pluripotent stem cells using a vector based on Sendai virus, an RNA virus that does not integrate
29 into the host genome. *Proc Jpn Acad Ser B Phys Biol Sci* 2009;85(8):348–362.
- 30 268. Hu Q, Friedrich AM, Johnson LV, Clegg DO. Memory in induced pluripotent stem cells: Repro-
31 programmed human retinal-pigmented epithelial cells show tendency for spontaneous redifferentiation.
32 *Stem Cells* 2010;28(11):1981–1991.
- 33 269. Ghosh Z, Wilson KD, Wu Y, Hu S, Quertermous T, Wu JC. Persistent donor cell gene expression
34 among human induced pluripotent stem cells contributes to differences with human embryonic stem
35 cells. *PLoS One* 2010;5(2):e8975.
- 36 270. Miura K, Okada Y, Aoi T, Okada A, Takahashi K, Okita K, Nakagawa M, Koyanagi M, Tanabe
37 K, Ohnuki M, Ogawa D, Ikeda E, Okano H, Yamanaka S. Variation in the safety of induced
38 pluripotent stem cell lines. *Nat Biotechnol* 2009;27(8):743–745.
- 39 271. Ben-David U, Benvenisty N. The tumorigenicity of human embryonic and induced pluripotent stem
40 cells. *Nat Rev Cancer* 2011;11(4):268–277.
- 41 272. Wu S, Wu Y, Zhang X, Capecchi MR. Efficient germ-line transmission obtained with transgene-free
42 induced pluripotent stem cells. *Proc Natl Acad Sci USA* 2014;111(29):10678–10683.
- 43 273. Leten C, Roobrouck VD, Struys T, Burns TC, Dresselaers T, Vande Velde G, Santermans J, Lo
44 Nigro A, Ibrahim A, Gijsbers R, Eggermont K, Lambrechts I, Verfaillie CM, Himmelreich U.
45 Controlling and monitoring stem cell safety in vivo in an experimental rodent model. *Stem Cells*
46 2014;32(11):2833–2844.
- 47 274. Schuldiner M, Itskovitz-Eldor J, Benvenisty N. Selective ablation of human embryonic stem cells
48 expressing a “suicide” gene. *Stem Cells* 2003;21(3):257–265.
- 49 275. Paczkowska E, Larysz B, Rzeuski R, Karbicka A, Jalowski R, Kornacewicz-Jach Z, Ratajczak
50 MZ, Machalinski B. Human hematopoietic stem/progenitor-enriched CD34(+) cells are mobilized
51 into peripheral blood during stress related to ischemic stroke or acute myocardial infarction. *Eur J
Haematol* 2005;75(6):461–467.

276. Dunac A, Frelin C, Popolo-Blondeau M, Chatel M, Mahagne MH, Philip PJ. Neurological and functional recovery in human stroke are associated with peripheral blood CD34+ cell mobilization. *J Neurol* 2007;254(3):327–332.
277. Schneider A, Kruger C, Steigleder T, Weber D, Pitzer C, Laage R, Aronowski J, Maurer MH, Gassler N, Mier W, Hasselblatt M, Kollmar R, Schwab S, Sommer C, Bach A, Kuhn HG, Schabitz WR. The hematopoietic factor G-CSF is a neuronal ligand that counteracts programmed cell death and drives neurogenesis. *J Clin Invest* 2005;115(8):2083–2098.
278. Minnerup J, Sevimli S, Schabitz WR. Granulocyte-colony stimulating factor for stroke treatment: Mechanisms of action and efficacy in preclinical studies. *Exp Transl Stroke Med* 2009;1:2.
279. Yang B, Migliati E, Parsha K, Schaar K, Xi X, Aronowski J, Savitz SI. Intra-arterial delivery is not superior to intravenous delivery of autologous bone marrow mononuclear cells in acute ischemic stroke. *Stroke* 2013;44(12):3463–3472.
280. Banerjee S, Bentley P, Hamady M, Marley S, Davis J, Shlebak A, Nicholls J, Williamson DA, Jensen SL, Gordon M, Habib N, Chataway J. Intra-arterial immunoselected CD34+ stem cells for acute ischemic stroke. *Stem Cells Transl Med* 2014;3(11):1322–1330.
281. Moniche F, Montaner J, Gonzalez-Marcos JR, Carmona M, Pinero P, Espigado I, Cayuela A, Escudero I, de la Torre-Laviana FJ, Boada C, Rosell A, Mayol A, Jimenez MD, Gil-Peralta A, Gonzalez A. Intra-arterial bone marrow mononuclear cell transplantation correlates with GM-CSF, PDGF-BB, and MMP-2 serum levels in stroke patients: Results from a clinical trial. *Cell Transplant* 2014;23(Suppl 1):S57–S64.
282. Savitz SI, Misra V, Kasam M, Juneja H, Cox CS, Jr., Alderman S, Aisiku I, Kar S, Gee A, Grotta JC. Intravenous autologous bone marrow mononuclear cells for ischemic stroke. *Ann Neurol* 2011;70(1):59–69.
283. Prasad K, Sharma A, Garg A, Mohanty S, Bhatnagar S, Johri S, Singh KK, Nair V, Sarkar RS, Gorthi SP, Hassan KM, Prabhakar S, Marwaha N, Khandelwal N, Misra UK, Kalita J, Nityanand S, Inve STSG. Intravenous autologous bone marrow mononuclear stem cell therapy for ischemic stroke: A multicentric, randomized trial. *Stroke* 2014;45(12):3618–3624.
284. Popa-Wagner A, Buga AM, Doepfner TR, Hermann DM. Stem cell therapies in preclinical models of stroke associated with aging. *Front Cell Neurosci* 2014;8:347.
285. Popa-Wagner A, Filfan M, Uzoni A, Pourgolafshan P, Buga AM. Poststroke cell therapy of the aged brain. *Neural Plast* 2015;2015:839638.
286. Badan I, Buchhold B, Hamm A, Gratz M, Walker LC, Platt D, Kessler C, Popa-Wagner A. Accelerated glial reactivity to stroke in aged rats correlates with reduced functional recovery. *J Cereb Blood Flow Metab* 2003;23(7):845–854.
287. Zechariah A, ElAli A, Hagemann N, Jin F, Doepfner TR, Helfrich I, Mies G, Hermann DM. Hyperlipidemia attenuates vascular endothelial growth factor-induced angiogenesis, impairs cerebral blood flow, and disturbs stroke recovery via decreased pericyte coverage of brain endothelial cells. *Arterioscler Thromb Vasc Biol* 2013;33(7):1561–1567.
288. Ay H, Koroshetz WJ, Vangel M, Benner T, Melinosky C, Zhu M, Menezes N, Lopez CJ, Sorensen AG. Conversion of ischemic brain tissue into infarction increases with age. *Stroke* 2005;36(12):2632–2636.
289. Buga AM, Margaritescu C, Scholz CJ, Radu E, Zelenak C, Popa-Wagner A. Transcriptomics of post-stroke angiogenesis in the aged brain. *Front Aging Neurosci* 2014;6:44.
290. Buga AM, Scheibe J, Moller K, Ciobanu O, Posel C, Boltze J, Popa-Wagner A. Granulocyte colony-stimulating factor and bone marrow mononuclear cells for stroke treatment in the aged brain. *Curr Neurovasc Res* 2015;12(2):155–162.

1 46 • GERVOIS ET AL.

2 *Belgium. He started as a Ph.D. student and teaching assistant at Hasselt University in September*
3 *2011 at the Department of Morphology. His main research focus is to investigate the neurogenic*
4 *differentiation potential of human dental pulp stem cells and their neuroregenerative effect after*
5 *transplantation in a mouse stroke model.*

6
7 **Esther Wolfs** obtained her master's degree in Biomedical Sciences in 2008 at Hasselt University,
8 Belgium. In 2014, she obtained a Ph.D. degree at the Katholieke Universiteit Leuven at the
9 Department of Nuclear Medicine and Molecular Imaging on radionuclide imaging of stem cells.
10 Currently, she is a postdoctoral researcher at Hasselt University at the Department of Functional
11 Morphology. Her research interests are situated within the field of dental pulp stem cells as
12 treatment vehicles in oncology as well as neurological applications and noninvasive imaging.

13
14 **Jessica Ratajczak** obtained her degree of Master of Biomedical Sciences (2013) at Hasselt
15 University, Belgium. She is currently a Ph.D. student at Hasselt University and focuses on the
16 angiogenic properties of periodontal ligament stem cells after in vitro preconditioning.

17
18 **Yörg Dillen** obtained his degree of Master of Biomedical Sciences (2015) at Hasselt University,
19 Belgium. He started as a predoctoral researcher at Hasselt University in October 2015 at the
20 Department of Morphology. His thesis research project is focused on the neuroprotective and
21 regenerative effects of human dental pulp stem cell transplantation in ischemic stroke.

22
23 **Tim Vangansewinkel** obtained his degree of Master of Biomedical Sciences (2010) at Hasselt
24 University, Belgium. He did his Ph.D. research at Hasselt University where he investigated the
25 effect of mast cell specific proteases on scarring and functional recovery after traumatic spinal
26 cord injury. He is currently working as a scientist in the Morphology Research Group at Hasselt
27 University. His research projects are mainly focused on regenerative therapies for peripheral and
28 central nerve injuries by combining stem cells and different biomaterials for tissue engineering.

29
30 **Petra Hilkens** obtained her master's degree in Biomedical Sciences (2011) at Hasselt University,
31 Belgium. In 2015, she obtained her Ph.D. degree in Biomedical Sciences at Hasselt University.
32 Currently, she is working as a FWO research fellow at Hasselt University, focusing on the cardio-
33 vascular properties of human dental pulp stem cells and their potential application in a wide range
34 of clinical disorders.

35
36 **Annelies Bronckaers** obtained her degree of Master of Biological Sciences (2004) at the
37 Katholieke Universiteit Leuven (K.U. Leuven), Belgium. In 2009, she obtained her Ph.D. at
38 the Rega Institute (K.U. Leuven) where she studied the role of thymidine phosphorylase and
39 thymidine phosphorylase inhibitors in angiogenesis. Since then, she is a postdoctoral researcher
40 at BIOMED, Hasselt University, Belgium, where she investigates the angiogenic capacities of
41 various dental stem cells populations and explores the use of dental pulp stem cells as carriers for
42 gene therapy.

43
44
45 **Ivo Lambrechts, DDS, PhD**, is currently full professor microscopic anatomy at the Faculty of
46 Medicine, Hasselt University, Belgium. He is member of the board of directors of the Belgian-
47 Dutch Society of Oral Biology, and of the board of directors of the Belgian Society of Cell Biology.
48 He is chairman of the Histology-Imaging Group of the BIOMED institute. At present, he is vice-
49 dean of the Faculty of Medicine of Hasselt University and he served in the external quality control
50 of the Flanders dental schools. He is member of the medical and biomedical education management
51 teams of Medicine and Biomedicine of Hasselt University. He contributes to stem cell research,

STEM CELL-BASED THERAPIES FOR ISCHEMIC STROKE • 47

oral biology, and oral imaging research and is involved in tissue banking, tissue reconstruction, and regeneration. He is member of the board of directors of the BIOMED research institute of Hasselt University. He was expert from the EC in the COST action B8 on odontogenesis and member of the management committee of the COST action B23 on oral facial development. He was also member of the central council of the European Association of Dento-Maxillo-Facial Radiology (EADMFR). He is panel member (Med8 and Clinical research) of the Fund for Scientific Research Flanders (FWO). He is referee of the IWT, The Inbev-Latour Award, and review editor of *Frontiers in Craniofacial Biology*. At present, he is vice president of the European Society of Dental and Cranio-Facial Stem Cells (ESCDSC). His research group was partner in a SBO/IWT project "IMAGINE" on nanoparticles, stem cells, and anticancer therapy. Recently, he obtained projects and research grants in the field of cancer research and angiogenesis research from the FWO.

Tom Struys obtained his degree of Master of Biomedical Sciences (2003) at Hasselt University, Belgium. In 2011, he obtained his Ph.D. in biomedical sciences at the Biomedical Research Institute (BIOMED), lab of histology at Hasselt University, Belgium. Following a postdoctoral position at the Molecular Small Animal Imaging Center (MoSAIC) and the biomedical MRI unit at the Katholieke Universiteit Leuven (KU Leuven, Belgium), he received a tenure track position at the Morphology Research Group of the biomedical research institute (BIOMED) at Hasselt University, Belgium. There, he currently acts as assistant professor human anatomy within the faculty of medicine and life sciences. His research projects focus on investigating the neuroprotective effect of stem cell transplantation following ischemic stroke and the use of noninvasive imaging techniques to monitor cell therapy.

Author Query Form

Journal **MED**

Article **med21400**

Dear Author

During the copy-editing of your paper, the following queries arose. Please respond to these by marking up your proofs with the necessary changes/additions. Please write your answers clearly on the query sheet if there is insufficient space on the page proofs. If returning the proof by fax do not write too close to the paper's edge. Please remember that illegible mark-ups may delay publication.

Query No.	Description	Remarks
Q1	Author: Please confirm that given names (red) and surnames/family names (green) have been identified correctly.	
Q2	Author: A running head short title was not supplied; please check if this one is suitable and, if not, please supply a short title that can be used instead.	
Q3	Author: Please define the term "FDA."	
Q4	Author: Please check the usage of term "toward" appearing throughout the article for correctness.	
Q5	Author: Please define terms "ASC" and "UMSC."	
Q6	Author: Sense of the sentence "Underlying mechanisms of . . ." is not clear. Please check.	
Q7	Author: Please check whether edit in the sentence "This statement was . . ." retains the intended meaning.	
Q8	Author: Please check whether edit in the sentence "Although study by Kamiya . . ." retains the intended meaning.	
Q9	Author: Please check whether edit in the sentence "Nonetheless, a direct comparison . . ." retains the intended meaning.	
Q10	Author: Please provide the complete publication details in Ref. 1.	
Q11	Author: Please check the page range as typeset for correctness in Ref. 2.	
Q12	Author: Please check the page range as typeset for correctness in Ref. 3.	
Q13	Author: Please provide the page range in Ref. 25.	
Q14	Author: As per style, please provide the names of all the authors in Ref. 27.	
Q15	Author: Please provide the vol. no. and page range in Ref. 66.	

Q16	Author: Please provide the vol. no. and page range in Refs. 165 and 167.	
-----	--	--