

**Angel O. Rojas Vistorte, Wagner Silva Ribeiro, Denisse Jaen,
Miguel R. Jorge, Sara Evans-Lacko and Jair de Jesus Mari**
**Stigmatizing attitudes of primary care
professionals towards people with mental
disorders: a systematic review**
Article (Accepted version)
(Refereed)

Original citation:

Rojas Vistorte, Angel O. and Silva Ribeiro, Wagner and Jaen, Denisse and Jorge, Miguel R. and Evans-Lacko, Sara and Mari, Jair de Jesus (2018) Stigmatizing attitudes of primary care professionals towards people with mental disorders: a systematic review. *International Journal of Psychiatry in Medicine*, 53 (4). pp. 317-338. ISSN 0091-2174
DOI: [10.1177/0091217418778620](https://doi.org/10.1177/0091217418778620)

© 2018 Sage

This version available at: <http://eprints.lse.ac.uk/89508/>

Available in LSE Research Online: July 2018

LSE has developed LSE Research Online so that users may access research output of the School. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LSE Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain. You may freely distribute the URL (<http://eprints.lse.ac.uk>) of the LSE Research Online website.

This document is the author's final accepted version of the journal article. There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

Stigmatizing attitudes of primary care professionals towards people with mental disorders: a systematic review

Authors: Angel O. Rojas Vistorte 1, Wagner Silva Ribeiro 1,2, Denisse Jaen 1, Miguel R. Jorge 1, Sara Evans-Lacko 2, 3 and Jair de Jesus Mari 1,3

1. Federal University of São Paulo, Psychiatry Department, São Paulo, Brazil.
2. Personal Social Services Research Unit, London School of Economics and Political Science.
3. King's College London, Health Service and Population Research Department, Institute of Psychiatry Psychology & Neuroscience, London, United Kingdom.

Address for Correspondence:

Angel O. Rojas Vistorte

Rua Borges Lagoa No. 570, 7o Andar.

Vila Clementino, São Paulo. Brazil.

CEP: 04038-000

Summary

Objective: To examine stigmatizing attitudes towards people with mental disorders among primary care professionals, and to identify potential factors related to stigmatizing attitudes through a systematic review. **Methods:** A systematic literature search was conducted in Medline, Lilacs, IBECs, Index Psicologia, CUMED, MedCarib, Sec. Est. Saúde SP, WHOLIS, Hanseníase, LIS-Localizador de Informação em Saúde, PAHO, CVSO-Regional and Latindex, through the Virtual Health Library portal (<http://www.bireme.br> website) through to June 2017. The articles included in the review were summarized through a narrative synthesis. **Results:** After applying eligibility criteria, eleven articles, out of 19.109 references identified, were included in the review. Primary care physicians do present stigmatizing attitudes towards patients with mental disorders, and show more negative attitudes towards patients with schizophrenia than towards those with depression. Older and more experience doctors have more stigmatizing attitudes towards people with mental illness compared with younger and less experienced doctors. Health care providers who endorse more stigmatizing attitudes towards mental illness were likely to be more pessimistic about the patient's adherence to treatment. **Conclusions:** Stigmatizing attitudes towards people with mental disorders are common among physicians in primary care settings, particularly among older and more experienced doctors. Stigmatizing attitudes can act as an important barrier for patients to receive the treatment they need. The primary care physicians feel they need better preparation, training and information to deal with and to treat mental illness, such as a user friendly and pragmatic classification system that addresses the high prevalence of mental disorders in primary care and community settings.

Keywords: Stigma, stigmatizing attitudes, primary care, mental disorders.

Introduction

Mental, neurological and substance use disorders (MNSD) exact a high toll, accounting for 13% of the total global burden of disease ⁽¹⁾. The burden of disease attributable to MNSD increased by 41% between 1990 and 2010. MNSD accounts for one in every 10 disability adjusted life years (DALYs) ⁽²⁾ and affects disproportionately low- and middle-income countries (LMIC) ⁽²⁾. Individuals with major depression and schizophrenia have a 40% to 60% greater chance of dying prematurely than the general population, owing to physical health problems that are often left untreated (e.g., cancer, cardiovascular diseases, diabetes and HIV infection) ⁽²⁾.

It is widely known that people with mental disorders commonly present in primary care and that disorders such as depression, alcohol abuse and epilepsy can be treated by primary care professionals ⁽³⁾. For instance, Ansseau et al. (2004) in a survey performed in Belgium found that, in the year prior to the study, 31% of adult patients in primary care services had affective disorders, 19% had some type of anxiety disorder, 18% had some type of somatoform disorder and 10% had disorders related to use of alcohol ⁽⁴⁾, and other studies estimate that 24% of patients who present to primary care physicians have a well-defined ICD-10 mental disorder ⁽⁵⁾. Scientific evidence shows that, worldwide, 69% of patients with mental disorders in primary care settings present to physicians with physical symptoms and that in many of those patients the physical symptoms or physical illness remain undetected ⁽⁵⁾.

It is recommended that common mental disorders be treated in primary care settings. However, a number of obstacles hinder primary care physicians (PCP's) ability to identify and to treat mental disorders. Firstly, it has been shown that PCP have limited ability diagnose a mental disorder because the training required for developing this ability is not routinely available in general medical undergraduate courses and medical training ⁽⁶⁾. Lack of training leads to difficulties in dealing with mental health problems, and to the maintenance of misconceptions about mental disorders ^(6, 7). Another important obstacle is the stigma or prejudice that primary care providers may have in relation to patients with mental disorders ⁽⁸⁻¹³⁾.

Stigma is defined as a process involving labeling, separation, stereotype endorsement, prejudice and discrimination in a context in which social, economic or political power is exercised to the detriment of members of a social group ⁽¹⁴⁾. It comprises three related dimensions, which, when combined are a powerful driver of social exclusion: a) lack of knowledge (ignorance and misinformation); b) negative attitudes (prejudice); and c) excluding or avoiding behavior's (discrimination) ^(12, 15-18). Stigmatizing attitudes may influence how primary care physicians diagnose and treat individuals with mental disorders ⁽¹⁹⁾. Thus, although individuals with mental disorders may seek treatment more frequently from primary care than from specialized services, in part to avoid the labeling related to psychiatric treatment, they may also face stigma in primary care ⁽²⁰⁾.

Health professionals who work in primary health care services and support stigmatizing attitudes toward people with mental disorders are more pessimistic about their adherence to treatment, both for mental illness and physical illness. Stigma can lead professionals to make clinical decisions based on mistaken assumptions or against clinical standards or guidelines ⁽¹³⁾. In this context, unreceptive attitudes of professionals who perpetuate stigmatizing beliefs tend to distance people with mental disorders even more from the health care they need ^(20, 21).

The main aim of this study was examined stigmatizing attitudes towards people with mental disorders among primary care physicians, and to identify potential factors related to stigmatizing attitudes through a systematic review.

Methods and Procedures

We performed a systematic review of the scientific literature through the Biblioteca Virtual em Saúde (Virtual Health Library) – BVS. The BVS automatically runs searches in the following databases: Medline, Lilacs, PubMed, IBECs, Lilacs, Index Psicologia – Periódicos técnicos-científicos, CUMED, CidSaúde-Cidades saudáveis, MedCarib, BDEF-Enfermagem, Sec. Est. Saúde SP, WHOLIS, BBO-Odontologia, Hanseníase, HomeoIndex-Homeopatia, LIS-Localizador de Informação em Saúde, PAHO, CVSO-Regional, Latindex. Additionally, reference lists of included studies and reviews were checked for potentially relevant articles not identified through the

electronic search. We used the following search terms: stigma-related terms AND primary care AND mental health-related terms as follows: “((tw:(stigma*)) OR (tw:(discriminat*)) OR (tw:(prejudice*)) OR (tw:(social distance*)) OR (tw:(stereotyp*)) OR (tw:(attitude*)) OR (tw:(behav*))) AND (tw:(primary care)) AND ((tw:(mental disorder*)) OR (tw:(mental illness*)) OR (tw:(mental disease*)) OR (tw:(mental disabilit*)) OR (tw:(psychiatr* disorder*)) OR (tw:(psychiatry* illness*)) OR (tw:(psychiatry* disease*)) OR (tw:(psychiatry* diagnos*)))”.

Titles and abstracts were screened and full reports of potentially relevant studies were obtained. Two authors (AORV and DJV) independently assessed the reports for eligibility, with discrepancies resolved by discussion with a third author (WSR).

We included quantitative studies which assessed stigmatizing attitudes towards mental health problems, mental health symptoms or mental disorders, among primary care physicians in English, Spanish and Portuguese. Articles were excluded based on the following exclusion criteria: 1) if they referred to non-data-based studies (e.g., editorials, commentaries, opinion papers and review papers); and 2) if stigmatizing attitudes were assessed among non-physician primary care professionals, such as nurses, technicians, social workers and other professionals, among mental health professionals, or in the general population. Data on study design, sample characteristics and findings were extracted independently by two authors (AORV and DJV). Because of heterogeneity between studies, which hindered a statistical synthesis of their results, we summarized evidence from articles included in the review through a narrative synthesis ⁽²²⁾.

Results

The database search identified 19.109 non-duplicate references. After reviewing titles and abstracts, 15 articles were identified as potentially relevant and were assessed against eligibility criteria. Eleven studies fulfilled inclusion criteria (**Figure 1**) and are summarized in **Table 1**.

All the eleven studies included in the review used a cross-sectional design to assess stigmatizing attitudes towards mental disorders through questionnaires developed by the authors in six studies and one study using a standardized scale. Two studies were conducted in Brazil ^(23, 24); two in the USA ^(19, 25), one in Australia ⁽¹⁰⁾, one in

Hong Kong ⁽²⁶⁾, one in Spain ⁽²⁷⁾, one from Finland ⁽²⁸⁾, one from USA and Canada ⁽²⁹⁾, one from Israel ⁽³⁰⁾ and one from Switzerland ⁽³¹⁾.

Three studies included in the review assessed attitudes towards depression and anxiety among general practitioners working in primary care ^(23, 29, 30). All the studies considered that depression is a frequent problem in primary care. One of the studies ⁽²³⁾ found that 42% of physicians considered it “difficult to differentiate whether patients are presenting with unhappiness or a clinical depressive disorder that needs treatment”; that 47% of professionals agreed or strongly agreed that most of depressive symptoms “originates from patients’ recent misfortunes”; that 46% and 27% of physicians, respectively, considered it to be “heavy going” and “not rewarding” to work with depressed patients. However, 57% of respondents reported feeling “comfortable in dealing with the need of depressive patients”. Despite the consideration that depression is a problem that should be treated in primary care, in the study from Israel ⁽³⁰⁾ almost the 50% of the sample stated that the preferential locus for treatment of depression and anxiety are mental health clinics, 80.6% of the physicians agreed that there is under-diagnosis and under-treatment of depression and anxiety in primary care; and 37.3% stated that they have no interest in treating depression and anxiety in primary care. Coinciding with this, in the third study ⁽²⁹⁾ the PCP’s who think their patients feel uncomfortable with their broaching issues about depression see depression as less important in primary care. Moreover, PCP’s are more likely to feel satisfied if their self-efficacy for diagnosing, treating and managing depression is high.

Three articles assessed PCP’s views, knowledge’s, beliefs and attitudes (including stereotypes and moral attributions) towards alcohol consumption and drugs dependence ^(24, 27, 28). Whereas one of the studies ⁽²⁴⁾ found that PCP had the lowest scores, meaning less negative attitudes, towards alcohol addiction and towards marijuana/cocaine dependence, when compared to other health professionals, the other study ⁽²⁷⁾ found that 69.9% of PCP presented skepticism when treating patients with problems caused by alcohol misuse and presented indifference in their work with patients with alcohol-related disorders. The third study ⁽²⁸⁾ found that 87% of PCP’s “have positive attitudes towards discussing alcohol with patients”, and that 81% of them “thought that detection and treating of early phase alcohol abusers was appropriate to their everyday work”.

Two articles ^(25, 31) evaluated stigmatizing attitudes toward schizophrenia, by assessing stigma characteristics, physician's needs and expectations about patients' adherence to treatment, and subsequent health decisions (referral to a specialist and refill pain prescription). The first study ⁽²⁵⁾ showed that physicians who stigmatized mental illness were more likely to be pessimistic about the patient's adherence to treatment ⁽²⁵⁾. In the other study ⁽³¹⁾ 21% of PCP's reported no problems when treating patients with schizophrenia, 56% kept treating schizophrenic patients, even though they considered patients' behavior to be problematic, and 13% preferred to refer patients for considering their behavior to be problematic. The study also showed that the more PCP's considered the patient's behavior to be problematic, the more often they referred them to specialists.

Three studies compared PCP's attitudes towards patients with schizophrenia with their attitudes towards patients with depression ^(10, 19, 26). PCP's stigmatizing attitudes were greater towards patients with schizophrenia than towards patients with depression. PCP were more willing to treat patients with depression than schizophrenia, and they were less likely to feel comfortable to deal with the needs of patients with schizophrenia than of patients with depression.

Four ^(10, 24, 26, 27) of the articles included in the review assessed factors that might be associated with physicians' stigmatizing attitudes towards mental disorders. Two studies ^(24, 26) found a statistically significant association between age and stigmatizing attitudes, meaning that older physicians presented higher moralization towards alcohol-related disorders ⁽²⁴⁾, and lower willingness to work with patients with alcohol problems ⁽²⁷⁾. Another study, however, found results in the opposite direction when the younger physicians presented higher levels of stigma than the older one ⁽¹⁰⁾. Alongside with age, training is another factor that might impact PCP's attitudes and clinical practice. PCP's with more training in depression, for example, felt significantly more comfortable dealing with depressed patient, when compared to PCP's with no or little training ⁽²⁹⁾. Moreover, a major proportion of PCP's (59.7% indicated lack of knowledge in diagnosis and treatment) referred/appointed lack of training as a major barrier for treating patients with depression and anxiety, and, therefore, referred/appointed the need of training for improving their skills to treat patients with mental health problems (83.6% of the PCP's) ⁽³⁰⁾. Also, Stigma was

greater among those providers who were relatively less comfortable with using mental health services themselves ⁽²⁵⁾.

Five ^(23, 27, 29-31) articles including in their results the impact of the training in the attitudes and clinical practice of the PCP's. In one of the studies ⁽³⁰⁾ almost the 60% of the sample indicated the lack of knowledge in diagnosis and treatment as barriers to care people with depression and anxiety. On the other hand, those PCP's with more training in depression experienced significantly less discomfort while addressing in that those with little or no training ⁽²⁹⁾. The need of more training it's a common conclusion by the PCP's in order to improved their skills to treat people with mental disorders ^(23, 27, 31).

When compared the stigmatizing attitudes from PCP's with mental health professionals or general population one study ⁽¹⁰⁾ found that the public rated positive outcomes as more likely and negative outcomes as less likely than did the physicians and psychiatrists for depression patients. Also, regarding patients with schizophrenia, the general population rated positive outcomes as more likely and negative outcomes as less likely than did all three professional groups, and the general population was less likely to believe there would be discrimination against patients than the physicians did.

Discussion

The results of this systematic review show that stigmatizing attitudes towards patients with mental disorders are common among physicians in primary care settings, and that many physicians don't feel comfortable to deal with patients with mental disorders. Stigmatization towards schizophrenia was found to be significantly higher than towards depression ^(10, 26, 32), and primary care physicians' views were more negative about a patient with schizophrenia than an otherwise identical patient with depression ⁽³³⁾. These findings of primary care providers having more negative views of individuals with schizophrenia are consistent with other studies which considered stigmatizing attitudes among others primary care professionals like nurses, social workers and mental health professionals ⁽³⁴⁻³⁷⁾.

Our results also show that physicians have more negative attitudes towards mental illness when compared to other professionals and to the general population. Studies comparing such groups found that physicians had the highest levels of stigmatizing

attitudes, followed by other primary care professionals, mental health professionals and the general population ^(10, 19, 24, 25).

The literature suggests that stigmatizing attitudes among physicians is associated with a lack of adequate training regarding treatment and identification of mental disorders ⁽³⁸⁻⁴⁰⁾. Research has repeatedly identified deficiencies in both the identification and management of depressive illness in general practice ⁽⁴¹⁻⁴⁴⁾.

There is evidence that physicians stigmatizing attitudes towards mental disorders might be an important barrier for people with mental health problems to receive the treatment they need ^(19, 25). Negative attitudes of primary care providers and mental health providers may contribute to disparities in physical healthcare for persons with serious mental illness such as schizophrenia ^(40, 45-49). One of the most important negative attitudes was the erroneous attribution of the signs and symptoms of physical illness to concurrent mental disorders, as it may result in physical ailments being undertreated due to the so-called “medical bias” or “diagnostic overshadowing” related to providers’ negative attitudes ⁽⁵⁰⁾.

As most depressed patients seek treatment in primary care settings, and these patients are frequently undertreated, it is important to identify ways to increase the effectiveness of primary care clinicians in managing depression. Interventions need to consider the context of the primary care setting and consider barriers to treatment including lack of adequate time, training, competing agendas, and lack of adequate reimbursement. Ineffective treatment leads to patient and physician frustration due to lack of progress ⁽⁵¹⁾. Thus, primary health care providers represent a potential target for interventions to reduce disparities in care for individuals with depression, schizophrenia and in clinical care in general ^(26, 27, 52).

The main limitation of the study was the search strategy, which may have led to publication bias as potential sources of gray literature were not included. However, the search was conducted in the main medical databases, using terms which have been previously used in the literature on stigma, and references of the studies selected were carefully screened to find further studies ⁽⁵³⁾. Another limitation was the heterogeneity between studies – particularly regarding the assessment of stigma. Such heterogeneity limits the possibility of combining and synthesizing findings of different articles. Additionally, it was not possible to combine studies’ results in a statistical synthesis (meta-analysis) given the heterogeneity in analysis strategies

and results from individual articles. For this reason, we decided to summarize the results of articles included in the review through a narrative synthesis. Another important limitation is that most of the results in the review are based on a single study, which have important implications when interpreting and generalizing our results.

The main conclusion of this review is that stigmatizing attitudes towards people with mental disorders are common among primary care professionals, and that professionals need more training and adequate tools to deal with mental disorders, such as a user friendly and pragmatic classification system that addresses the high prevalence of mental disorders in primary care and community settings ⁽⁵⁴⁾. The reduction of stigma among health professionals is important to reduce barriers to treatment faced by people with mental disorders and, thus, to increase their access to optimal care, which would lead to improvement in their mental health status and well-being.

Acknowledgments

AORV is receiving a scholarship from CAPES foundation of the Brazilian Ministry of Education for his doctorate training in the Department of Psychiatry, Federal University of São Paulo. DJ is also funded by Capes for her doctorate training. JJM is an I-A Senior Researcher from the National Brazilian Council (CNPq). SE-L and WSR are funded by the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement number [337673].

References

1. Mental health action plan 2013-2020 [Internet]. 2013 [cited 2013]. Available from: http://apps.who.int/iris/bitstream/10665/89966/1/9789241506021_eng.pdf?ua=1.
2. Patel V, Chisholm D, Parikh R, Charlson FJ, Degenhardt L, Dua T, et al. Addressing the burden of mental, neurological, and substance use disorders: key messages from Disease Control Priorities, 3rd edition. The Lancet. 2016;387(10028):1672-85.
3. World Health Organization. Salud Mental: Nuevos conocimientos, nuevas esperanzas. . Ginebra 2001; 2001.
4. Anseau M, Dierick M, Buntinx F, Cnockaert P, De Smedt J, Van Den Haute M, et al. High prevalence of mental disorders in primary care. Journal of affective disorders. PLoS ONE. 2004;78(1):49-55.
5. WHO WHO. Mental Disorders in Primary Care. Department of Mental Health and Substance Abuse. 1998.
6. Reed GM, Mendonca Correia J, Esparza P, Saxena S, Maj M. The WPA-WHO Global Survey of Psychiatrists' Attitudes Towards Mental Disorders Classification. World Psychiatry. 2011 Jun;10(2):118-31. PubMed PMID: 21633689. Pubmed Central PMCID: PMC3105474. Epub 2011/06/03. eng.
7. Lam TP, Goldberg DP, Dowell AC, Fortes S, Mbatia JK, Minhas FA, et al. Proposed new diagnoses of anxious depression and bodily stress syndrome in ICD-11-PHC: an international focus group study. Family practice. 2013 Feb;30(1):76-87. PubMed PMID: 22843638. Epub 2012/07/31. eng.
8. Jorge M. Concepções populares e estigma relacionados às doenças mentais. Nova Perspectiva Sistêmica. 2013;46:8-19.
9. Hodges B, Inch C, Silver I. Improving the psychiatric knowledge, skills, and attitudes of primary care physicians, 1950-2000: a review. The American journal of psychiatry. 2001 Oct;158(10):1579-86. PubMed PMID: 11578983. Epub 2001/10/02. eng.
10. Jorm AF, Korten AE, Jacomb PA, Christensen H, Henderson S. Attitudes Towards People with a Mental Disorder: A Survey of the Australian Public and Health Professionals. Australian and New Zealand Journal of Psychiatry. 1999 February 1, 1999;33(1):77-83.
11. Pinfold V, Huxley P, Thornicroft G, Farmer P, Toulmin H, Graham T. Reducing psychiatric stigma and discrimination--evaluating an educational intervention with the police force in England. Social psychiatry and psychiatric epidemiology. 2003 Jun;38(6):337-44. PubMed PMID: 12799785. Epub 2003/06/12. eng.
12. Thornicroft G, Rose D, Kassam A, Sartorius N. Stigma: ignorance, prejudice or discrimination? . The British Journal of Psychiatry. 2007;190:192-3.
13. Shim R, Rust G. Primary care, behavioral health, and public health: partners in reducing mental health stigma. Am J Public Health. 2013 May;103(5):774-6. PubMed PMID: 23488498. Pubmed Central PMCID: PMC3698818. Epub 2013/03/16. eng.
14. Link B, Phelan J. Conceptualizing stigma. American Review of Sociology. 2001;27:363-85.

15. Thornicroft G. *Shunned: Discrimination against People with Mental Illness*. Oxford University Press. 2006b.
16. Corrigan P. *On the Stigma of Mental Illness*. In: Association AP, editor. Washington, D.C. 2005.
17. Sartorius N, Schulze H. *Reducing the Stigma of Mental Illness. A Report from a Global Programme of the World Psychiatric Association*. Cambridge: 2005.
18. Thornicroft G. *Actions Speak Louder: Tackling Discrimination against People with Mental Illness*. In: Foundation MH, editor. London 2006a.
19. Mittal D, Corrigan P, Sherman M, Chekuri L, Han X, Reaves C, et al. *Healthcare Providers' Attitudes Toward Persons With Schizophrenia*. *Psychiatric Rehabilitation Journal*. 2014;37(4):297-303.
20. Pingitore D, Snowden L, Sansone RA, Klinkman M. *Persons with depressive symptoms and the treatments they receive: a comparison of primary care physicians and psychiatrists*. *Int J Psychiatry Med*. 2001;31(1):41-60. PubMed PMID: 11529390. Epub 2001/09/01. eng.
21. Henderson C, Evans-Lacko S, Thornicroft G. *Mental illness stigma, help seeking, and public health programs*. *Am J Public Health*. 2013 May;103(5):777-80. PubMed PMID: 23488489. Pubmed Central PMCID: PMC3698814. Epub 2013/03/16. Eng.
22. Popay J, Roberts H, Sowden A, Petticrew M, Arai L, Rodgers M, et al. *Guidance on the conduct of narrative synthesis in systematic reviews. A product from the ESRC methods programme Version*. 2006;1:b92.
23. Botega NJ, Silveira GM. *General Practitioners' Attitudes Towards Depression: a Study in Primary Care Setting in Brazil*. *International Journal of Social Psychiatry*. 1996 September 1, 1996;42(3):230-7.
24. Ronzani TM, Higgins-Biddle J, Furtado EF. *Stigmatization of alcohol and other drug users by primary care providers in Southeast Brazil*. *Social science & medicine (1982)*. 2009 Oct;69(7):1080-4. PubMed PMID: 19692163. Epub 2009/08/21. eng.
25. Corrigan PW, Mittal D, Reaves CM, Haynes TF, Han X, Morris S, et al. *Mental health stigma and primary health care decisions*. *Psychiatry research*. 2014 04/18;218(0):35-8. PubMed PMID: PMC4363991.
26. Lam TP, Lam KF, Lam EW, Ku YS. *Attitudes of primary care physicians towards patients with mental illness in Hong Kong*. *Asia-Pacific psychiatry : official journal of the Pacific Rim College of Psychiatrists*. 2013 Mar;5(1):E19-28. PubMed PMID: 23857792. Epub 2013/07/17. eng.
27. Rodríguez Fernández E, Espí Martínez F, Canteras Jordana M, Gómez Moraga A. *Actitud de los médicos de atención primaria sobre los pacientes con consumo excesivo de alcohol*. *Atención Primaria*. 1999;24(8):447-55.
28. Aalto M, Pekuri P, Seppä K. *Primary health care nurses' and physicians' attitudes, knowledge and beliefs regarding brief intervention for heavy drinkers*. *Addiction (Abingdon, England)*. 2001 Feb;96(2):305-11. PubMed PMID: 11182876. Epub 2001/02/22. eng.
29. Main DS, Lutz LJ, Barrett JE, Matthew J, Miller RS. *The role of primary care clinician attitudes, beliefs, and training in the diagnosis and treatment of depression: a report from the Ambulatory Sentinel Practice Network Inc*. *Archives of Family Medicine*. 1993;2(10):1061.
30. Goldfracht M, Shalit C. *Attitudes of Israeli Primary Care Physicians Towards Mental Health Care/Commentary-Attitudes*. *The Israel journal of psychiatry and related sciences*. 2007;44(3):225.
31. Simon AE, Lauber C, Ludewig K, Braun-Scharm H, Umbricht DS. *General practitioners and schizophrenia: results from a Swiss survey*. *The British Journal of Psychiatry*. 2005;187(3):274-81.
32. Mann CE, Himelein MJ. *Factors Associated With Stigmatization of Persons With Mental Illness*. *Psychiatric Services*. 2004;55(2):185-7. PubMed PMID: 14762246.
33. Lawrie SM, Martin K, McNeill G, Drife J, Chrystie P, Reid A, et al. *General practitioners' attitudes to psychiatric and medical illness*. *Psychol Med*. 1998 Nov;28(6):1463-7. PubMed PMID: 9854287. Epub 1998/12/17. eng.

34. Bjorkman T, Angelman T, Jonsson M. Attitudes towards people with mental illness: A cross-sectional study among nursing staff in psychiatric and somatic care. *Scandinavian Journal of Caring Sciences*. 2008;22:170-7.
35. Caldwell T, Jorm A. Mental health nurses' beliefs about likely outcomes for people with schizophrenia or depression: A comparison with the public and other healthcare professionals. *Australian and New Zealand Journal of Mental Health Nursing*. 2001;10:42-54.
36. Hori H, Richards M, Kawamoto Y, Kunugi H. Attitudes toward schizophrenia in the general population, psychiatric staff, physicians, and psychiatrists: A web-based survey in Japan. *Psychiatry Research*. 2011;186(183-189).
37. Hugo M. Mental health professionals' attitudes towards people who have experienced a mental health disorder. *J Psychiatr Ment Health Nurs*. 2001 Oct;8(5):419-25. PubMed PMID: 11882162. Epub 2002/03/08. eng.
38. Mari JdJ, Iacoponi E, Williams P, Simões O, Silva JBT. Detection of psychiatric morbidity in the primary medical care setting in Brazil. *Revista de Saúde Pública*. 1987;21:501-7.
39. Pincus HA. The future of behavioral health and primary care: drowning in the mainstream or left on the bank? *Psychosomatics*. 2003 Jan-Feb;44(1):1-11. PubMed PMID: 12515832. Epub 2003/01/08. eng.
40. M DEH, Correll CU, Bobes J, Cetkovich-Bakmas M, Cohen D, Asai I, et al. Physical illness in patients with severe mental disorders. I. Prevalence, impact of medications and disparities in health care. *World Psychiatry*. 2011 Feb;10(1):52-77. PubMed PMID: 21379357. Pubmed Central PMCID: PMC3048500. Epub 2011/03/08. eng.
41. Williamson J, Stokoe IH, Gray S, Fisher M, Smith A, McGhee A, et al. OLD PEOPLE AT HOME. THEIR UNREPORTED NEEDS. *Lancet*. 1964 May 23;1(7343):1117-20. PubMed PMID: 14132624. Epub 1964/05/23. eng.
42. Goldberg DP, Blackwell B. Psychiatric Illness in General Practice: A Detailed Study Using a New Method of Case Identification. *British Medical Journal*. 1970;2(5707):439-43. PubMed PMID: PMC1700485.
43. JOHNSON DAW. A Study of the Use of Antidepressant Medication in General Practice. *The British Journal of Psychiatry*. 1974 1974-08-01 00:00:00;125(585):186-92.
44. Freeling P, Rao BM, Paykel ES, Sireling LI, Burton RH. Unrecognised depression in general practice. *British Medical Journal (Clinical research ed)*. 1985;290(6485):1880-3. PubMed PMID: PMC1416762.
45. Druss BG, Bradford DW, Rosenheck RA, Radford MJ, Krumholz HM. Mental disorders and use of cardiovascular procedures after myocardial infarction. *Jama*. 2000 Jan 26;283(4):506-11. PubMed PMID: 10659877. Epub 2000/02/05. eng.
46. Koroukian SM, Bakaki PM, Golchin N, Tyler C, Loue S. Mental illness and use of screening mammography among Medicaid beneficiaries. *American journal of preventive medicine*. 2012 Jun;42(6):606-9. PubMed PMID: 22608377. Pubmed Central PMCID: PMC3631560. Epub 2012/05/23. eng.
47. Li Y, Cai X, Cram P. Are patients with serious mental illness more likely to be admitted to nursing homes with more deficiencies in care? *Medical care*. 2011 Apr;49(4):397-405. PubMed PMID: 21368683. Pubmed Central PMCID: PMC3784995. Epub 2011/03/04. eng.
48. Lord O, Malone D, Mitchell AJ. Receipt of preventive medical care and medical screening for patients with mental illness: a comparative analysis. *Gen Hosp Psychiatry*. 2010 Sep-Oct;32(5):519-43. PubMed PMID: 20851274. Epub 2010/09/21. eng.
49. Thornicroft G, Brohan E, Rose D, Sartorius N, Leese M. Global pattern of experienced and anticipated discrimination against people with schizophrenia: a cross-sectional survey. *Lancet*. 2009 Jan 31;373(9661):408-15. PubMed PMID: 19162314. Epub 2009/01/24. eng.
50. Thornicroft G, Rose D, Kassam A. Discrimination in health care against people with mental illness. *International Review of Psychiatry*. 2007;19(2):113-22.

51. Vega WA, Rodriguez MA, Ang A. Addressing stigma of depression in Latino primary care patients. *Gen Hosp Psychiatry*. 2010 Mar-Apr;32(2):182-91. PubMed PMID: 20302993. Epub 2010/03/23. eng.
52. Kapungwe A, Cooper S, Mayeya J, Mwanza J, Mwape L, Sikwese A, et al. Attitudes of primary health care providers towards people with mental illness: evidence from two districts in Zambia. *African journal of psychiatry*. 2011 Sep;14(4):290-7. PubMed PMID: 22038427. Epub 2011/11/01. eng.
53. Clement S, Schauman O, Graham T, Maggioni F, Evans-Lacko S, Bezborodovs N, et al. What is the impact of mental health-related stigma on help-seeking? A systematic review of quantitative and qualitative studies. *Psychological Medicine*. 2015;45(01):11-27.
54. Jacob KS, Patel V. Classification of mental disorders: a global mental health perspective. *The Lancet*. 2014;383(9926):1433-5.