



Published in final edited form as:

Nat Cell Biol. 2007 June ; 9(6): 636–645. doi:10.1038/ncb1590.

STIM1 heteromultimerizes TRPC channels to determine their function as store-operated channels

Joseph P. Yuan^{1,4}, Weizhong Zeng^{1,4}, Guo N. Huang², Paul F. Worley^{2,3,5}, and Shmuel Muallem^{1,5}

¹ Department of Physiology, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA

² Department of Neuroscience, The Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

³ Department of Neurology, The Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA

Abstract

Stromal interacting molecule 1 (STIM1) is a Ca²⁺ sensor that conveys the Ca²⁺ load of the endoplasmic reticulum to store-operated channels (SOCs) at the plasma membrane. Here, we report that STIM1 binds TRPC1, TRPC4 and TRPC5 and determines their function as SOCs. Inhibition of STIM1 function inhibits activation of TRPC5 by receptor stimulation, but not by La³⁺, suggesting that STIM1 is obligatory for activation of TRPC channels by agonists, but STIM1 is not essential for channel function. Through a distinct mechanism, STIM1 also regulates TRPC3 and TRPC6. STIM1 does not bind TRPC3 and TRPC6, and regulates their function indirectly by mediating the heteromultimerization of TRPC3 with TRPC1 and TRPC6 with TRPC4. TRPC7 is not regulated by STIM1. We propose a new definition of SOCs, as channels that are regulated by STIM1 and require the store depletion-mediated clustering of STIM1. By this definition, all TRPC channels, except TRPC7, function as SOCs.

A key component of the receptor-evoked Ca²⁺ signal is activation of a Ca²⁺-influx channel at the plasma membrane in response to depletion of Ca²⁺ from the endoplasmic reticulum, the so-called store-operated Ca²⁺ channels (SOCs)¹. Ca²⁺ influx through SOCs mediates numerous physiological functions^{1,2}. At least two types of SOCs can be distinguished electrophysiologically and now molecularly. The first type of SOCs are the highly Ca²⁺-selective *I_{crac}* currents¹ that recently were shown to be mediated by the Orai family of proteins^{3–8}, and the second type of SOCs are the non-selective, Ca²⁺ permeable TRPC channels^{1,9}.

Very little is known about the OraIs. The three OraIs are four trans-membrane-span proteins^{3,7,8}, with Orai1 the most prominent *I_{crac}* channel^{6,10}. Much more information is available on the TRP family of ion channels (recently reviewed in ref. ¹¹). The TRP superfamily is grouped into seven subfamilies and mediates many cellular functions^{1,9,11}. All

⁵Correspondence should be addressed to S.M. or P.F.W. (e-mail: E-mail: Shmuel.muallem@utsouthwestern.edu; pworley.edu; E-mail: pworley@jhmi.edu).

⁴These authors contributed equally to this work.

Note: Supplementary Information is available on the Nature Cell Biology website.

AUTHOR CONTRIBUTIONS

J.P.Y., W.Z. and G.N.H. performed and analysed the experiments. P.F.W. and S.M. planned and analysed the experiments. All authors contributed to writing the manuscript.

COMPETING FINANCIAL INTERESTS

The authors declare that they have no competing financial interests.

TRPC channels are activated by receptor stimulation and several TRPCs have been suggested to function as SOCs^{1,9,11,12}. Deletion in mice^{13,14} and knockdown by siRNA¹⁵ implicate TRPC1 and TRPC4 as SOCs. TRPC3, TRPC6 and TRPC7 are activated by the lipid diacylglycerol^{16,17}. Several studies reported that TRPC3 behaves as a SOC^{15,18}, whereas others concluded that TRPC3 does not function as a SOC^{19,20}. The same uncertainty exists with respect to TRPC6 (ref. 21, 22), whereas the behaviour of TRPC7 as a SOC seems to depend on its expression level²³.

The molecular mechanism by which the plasma membrane SOCs sense the Ca²⁺ filling of the endoplasmic reticulum was revealed by the recent identification of STIM1 as the endoplasmic reticulum Ca²⁺-content sensor^{24,25}. STIM1 has an amino-terminal EF hand Ca²⁺-binding domain that resides in the endoplasmic reticulum lumen. Depletion of endoplasmic reticulum Ca²⁺ results in rearrangement of STIM1 in punctae underneath the plasma membrane and subsequent activation of Ca²⁺-influx channels^{24–26}. Activation of the SOCs requires the sterile a motif (SAM), coiled-coil and serine–threonine (ST) domains of STIM1 (ref. 27) and is probably aided by multimerization of STIM1 (ref. 28). STIM1 regulates the activity of all known SOCs, including native SOCs^{24,25} and *I_{crac}*^{6,24,29,30}. Moreover, we showed that STIM1 binds TRPC1, TRPC4 and TRPC5, but not TRPC3, TRPC6 and TRPC7 (ref. 31), and the soluble carboxyl-terminus of STIM1 (amino acids 235–685) is sufficient to activate TRPC1 (ref. 31). The EF-hand mutant STIM1^{D76A} that does not bind Ca²⁺ is constitutively active, whereas the ΔERM(Ezrin–radixin–moesin)–STIM1^{D76A} acts as a dominant-negative STIM1 (refs 25, 31). Similarly, STIM1 was shown to regulate native TRPC1 in platelets³². In HSG cells, STIM1 was reported to exist in a complex with TRPC1 and Orai1 (ref. 33). However, very little is known about the mechanism by which STIM1 regulates TRPC and other Ca²⁺ influx channels. Fundamental unanswered questions are: which TRPC channels are regulated by STIM1; is STIM1 a subunit of the Ca²⁺ influx channels and is it obligatory for channel activity; and does STIM1 regulate the channel by affecting their activity at the plasma membrane.

Here, we show that STIM1 regulates directly or indirectly all TRPC channels, except TRPC7. However, although STIM1 directly regulates TRPC1, TRPC4 and TRPC5, the regulation of TRPC3 and TRPC6 by STIM1 is mediated by STIM1-dependent heteromultimerization of TRPC3 with TRPC1 and of TRPC6 with TRPC4. STIM1 is required for activation of all TRPC channels by agonist stimulation, but it is not essential for channel function. These findings clarify fundamental aspect of TRPC channels function by showing that under physiological conditions, STIM1 heteromultimerizes TRPC channels, assembling them into specific complexes to determine their function as SOCs.

RESULTS

Regulation of TRPC3 by STIM1

As STIM1 does not bind TRPC3, TRPC6 and TRPC7 (ref. 31), we expected that STIM1 would similarly regulate TRPC1, TRPC4 and TRPC5, but not regulate TRPC3, TRPC6 and TRPC7. Surprisingly, this was not the case for TRPC3 and TRPC6. Multiple probes were used to determine regulation of TRPC channels by STIM1; knockdown of *STIM1* with siRNA (*STIM1* siRNA) and the dominant negative ΔERM–STIM1^{D76A} (ref. 31) were used to inhibit the action of native STIM1 and the constitutively active STIM1^{D76A} and STIM1^{CT} (amino acids 235–685) were used to activate the channels in a store- and agonist-independent manner. The traces in Fig. 1a and b and the summary in Fig. 1c show that, when expressed at low levels, TRPC3 current is inhibited by *STIM1* siRNA and ΔERM–STIM1^{D76A} and is activated by STIM1^{D76A} and STIM1^{CT}, as was found for TRPC1 (ref. 31).

The results in Fig. 1 indicate that TRPC3 does function as a STIM1-regulated channel when expressed at low levels. Furthermore, inhibition of TRPC3 by Δ ERM-STIM1^{D76A}, which inhibits the translocation of wild-type STIM1 from the endoplasmic reticulum to the plasma membrane³¹, indicates that translocation of STIM1 to the plasma membrane is required for activation of TRPC3. We reasoned that as TRPC3 does not bind STIM1, regulation of TRPC3 by STIM1 may be mediated by one of the TRPC channels that do bind STIM1. The results in Fig. 2 show that this is indeed the case. In these experiments, the ability of native STIM1 to coimmunoprecipitate TRPC3 was examined in cells expressing TRPC3 alone or together with TRPC1, TRPC4 or TRPC5 (Fig. 2a). Remarkably, immunoprecipitation of native STIM1 resulted in coimmunoprecipitation of TRPC3 only when it is coexpressed with TRPC1, but not when it is expressed alone or with TRPC4 or TRPC5. STIM1 is obligatory for the interaction between TRPC3 and TRPC1 (Fig. 2b, c). TRPC3 plus TRPC1 were transfected into cells treated with scrambled siRNA or *STIM1* siRNA. TRPC3 coimmunoprecipitation with TRPC1, and expression of recombinant STIM1 did not further enhance the coimmunoprecipitate, suggesting that native STIM1 is sufficient to mediate the interaction between the channels under the expression conditions used. Significantly, knockdown of native *STIM1* nearly abolished the coimmunoprecipitation of TRPC3 with TRPC1.

An important question is whether the plasma membrane resident or the intracellular STIM1 mediate the TRPC1-TRPC3 heteromultimerization. This was addressed by testing the ability of YFP-STIM1 to rescue the interaction between the channels in cells treated with *STIM1* siRNA. Previous work showed that YFP-STIM1 does not translocate to the plasma membrane⁶ and its mRNA is not recognized by *STIM1* siRNA³⁴. These properties of YFP-STIM1 were confirmed and expression of YFP-STIM1 in cells treated with *STIM1* siRNA rescued the interaction between TRPC1 and TRPC3 (Fig. 2c). These findings suggest that migration of STIM1 to the plasma membrane stabilizes or promotes the formation of the TRPC1-TRPC3 complex. This raises the question of whether formation of the heteromultimer is affected by cell stimulation. Cell stimulation stabilized or enhanced formation of the TRPC1-TRPC3 complex and endogenous STIM1 was sufficient for this effect (Fig. 2d). Importantly, termination of cell stimulation resulted in reduction in the complex to nearly basal level within 3 min (Fig. 2d).

The functional significance of the STIM1-mediated interaction between TRPC1 and TRPC3 is shown (Fig. 3). Further evidence that STIM1 regulates TRPC3 is shown in Fig. 3a, in which YFP-STIM1 similarly rescued the activity of TRPC1 and TRPC3 expressed in *STIM1* siRNA-treated cells. This experiment is the functional complement of the experiment in Fig. 2c. The role of native TRPC1 in the regulation of TRPC3 by STIM1 was then tested. Immunoprecipitation of native TRPC1 was sufficient to coimmunoprecipitate TRPC3 and the coimmunoprecipitate was nearly abolished by knockdown of *STIM1* (Fig. 3b and see Supplementary Information, Fig. S1). In control HEK cells that expressed native TRPC1 (ref. 15), Δ ERM-STIM1^{D76A} inhibited TRPC3 current by about 85%. Knockdown of native *TRPC1* markedly reduced the inhibition of TRPC3 by Δ ERM-STIM1^{D76A}. Importantly, knockdown of the native *TRPC4* had no effect on inhibition of TRPC3 by Δ ERM-STIM1^{D76A}. These findings indicate that STIM1 indirectly regulates TRPC3 and that the regulation of TRPC3 by STIM1 is mediated by TRPC1. Further evidence for indirect regulation of TRPC3 by STIM1 was obtained by examining the effect of knockdown of *STIM1* on TRPC3 expressed at low and high levels. Previous work has shown that the behaviour of TRPC3 as a SOC^{18, 19} and regulation by InsP₃ receptors^{19,35} is observed only when TRPC3 is expressed at low levels. The level of native TRPC3 and when TRPC3 is expressed at low and high levels is shown in Fig. 3c, which also shows that overexpression of TRPC3 partially overcame the inhibition of TRPC3 current by knockdown of *STIM1*. It is likely that at low expression levels, most TRPC3 is in a complex with TRPC1 and STIM1. When it is overexpressed, large fraction of TRPC3 is not complexed and can be activated by agonist in a STIM1-independent manner.

The implication of these experiments is that STIM1 is not obligatory for the function of TRPC3 as a channel. However, when TRPC3 is present in a complex with TRPC1, as found in several cell types^{36,37}, it behaves as a STIM1-regulated SOC.

The significance of the heteromultimerization for channel function was examined further by measuring the rate of TRPC1 and TRPC3 activation on repeated stimulation. The half time for activation of TRPC1 was the same for the first and second stimulation (Fig. 3d). In contrast, the rate of TRPC3 activation increased approximately 2.4-fold in the second compared to the first stimulation. These results raised the possibility of a 'memory' of the assembled STIM1–TRPC1–TRPC3 complex, which may be important for Ca²⁺ oscillations and perhaps reloading of the stores on termination of cell stimulation. Thus, Ca²⁺ oscillations require repeated activation of Ca²⁺ influx in phase with Ca²⁺ release³⁸ and Ca²⁺ influx is required to sustain the Ca²⁺ oscillations^{1,39}. The memory of complex formation and facilitation of SOC activation can function to better coordinate Ca²⁺ release and activation of Ca²⁺ influx.

Regulation of TRPC4 and TRPC6 by STIM1

STIM1 binds TRPC4 through its ERM domain³¹ and activation of TRPC4 by agonist is inhibited by knockdown of *STIM1* and by Δ ERM-STIM1^{D76A} (see Supplementary Information, Fig. S2). These results indicate that STIM1 regulates TRPC4 in a mechanism that is similar to the regulation of TRPC1 by STIM1.

Although TRPC6 did not bind STIM1, STIM1 indirectly regulated activation of TRPC6 by agonist stimulation (Figs 4, 5). When TRPC6 was expressed at low levels, activation of TRPC6 by agonist was inhibited by knockdown of *STIM1* and by Δ ERM–STIM1^{D76A}, and TRPC6 became constitutively active in the presence of STIM1^{D76A} (Fig. 4). When TRPC6 was expressed at high levels, its activation by agonist was no longer regulated by STIM1, indicating that STIM1 is not obligatory for the function of TRPC6 (Fig. 4c), as was found for TRPC3.

Suspecting that regulation of TRPC6 by STIM1 is mediated by one of the TRPC channels that interact with STIM1, we examined which TRPC channel mediated the interaction between TRPC6 and STIM1. Native STIM1 does not coimmunoprecipitate TRPC6 when expressed alone (Fig. 5a). Notably, coexpression of TRPC6 with TRPC4, but not with TRPC1 or TRPC5, resulted in coimmunoprecipitation of STIM1 and TRPC6. In addition, immunoprecipitation of TRPC4 coimmunoprecipitated TRPC6 when both were coexpressed, and knockdown of *STIM1* markedly reduced the coimmunoprecipitation (Fig. 5b).

The functional significance of the STIM1-mediated interaction between TRPC6 and TRPC4 is shown in Fig. 5c. When TRPC6 was expressed at low levels, Δ ERM–STIM1^{D76A} inhibited activation of TRPC6 by about 87%. Knockdown of TRPC4 had no effect on current density or activation of TRPC6. However, knockdown of *TRPC4* markedly reduced inhibition of TRPC6 by Δ ERM–STIM1^{D76A}. As a control, knockdown of *TRPC1* did not prevent inhibition of TRPC6 stimulation by Δ ERM–STIM1^{D76A} (Fig. 5d). This was in contrast with the effect of knockdown of *TRPC1* on the inhibition of agonist stimulation of TRPC3 by Δ ERM–STIM1^{D76A}. Thus, these findings indicate that STIM1 indirectly regulates TRPC6 and that TRPC4 mediates the regulation of TRPC6 by STIM1.

STIM1 is obligatory for activation of TRPC channels by agonists, but not for channel activity

A critical question in understanding regulation of Ca²⁺ influx channels by STIM1 is whether STIM1 is obligatory for ion conductance by the channels that are directly regulated by STIM1, such as TRPC1, TRPC4 and TRPC5. This is highlighted in the finding that Orail activity as an *I_{crac}* channel can be observed only when it is coexpressed with STIM1 (refs 6, 8, 29, 30, 40). To address this question, it was necessary to examine the effect of STIM1 on a channel

that could be activated by receptor stimulation and by ligands in a receptor-independent manner. TRPC5 is such a channel, and can be activated by receptor stimulation or directly by lanthanides⁴¹. Immunoprecipitation of native STIM1 coimmunoprecipitated TRPC5 and the coimmunoprecipitated was not affected by coexpression of TRPC5 with TRPC3 or TRPC6 (Fig. 6a). Selective traces of the typical double rectification current/voltage relationship of TRPC5-mediated current is shown in Fig. 6b. These results indicate that STIM1 is obligatory for activation of TRPC5 by receptor stimulation. Hence, unlike the findings for TRPC3 and TRPC6, activation of TRPC5 by receptor stimulation requires STIM1 independent of the expression level of TRPC5. The second important finding from these experiments is that STIM1 is not obligatory for the function of TRPC5. Thus, knockdown of *STIM1* that largely inhibits activation of TRPC5 by agonist stimulation had no effect on direct activation of TRPC5 by La^{3+} . In addition, knockdown of *STIM1* had no effect on TRPC5 current density or its voltage-dependence. The significance of these findings is that STIM1 is required only for activation of TRPC channels by receptor stimulation, but STIM1 is not obligatory for ion conductance by the channels and does not regulate the voltage-dependence of channel function.

Not all TRPC channels are regulated by STIM1

TRPC7 function depends on its expression level. When expressed at low levels, activation of TRPC7 by agonist required its coupling to InsP_3 receptors, and TRPC7 activity was inhibited by $1 \mu\text{M Gd}^{3+}$. However, when TRPC7 was expressed at high levels, its activation was independent of InsP_3 receptors and TRPC7 activity was inhibited only by $10 \mu\text{M Gd}^{3+}$ (refs 23, 42). Consistent with the finding that TRPC7 does not bind STIM1 (ref. 31)³¹, the activity of TRPC7 was independent of STIM1 whether functioning as an InsP_3 receptor-dependent or -independent channel (Fig. 7). Thus, agonist-mediated activation of TRPC7 was not inhibited by $\Delta\text{ERM-STIM1}^{\text{D76A}}$ when its activity was largely inhibited by $1 \mu\text{M Gd}^{3+}$ (Fig. 7a) or when its inhibition required $10 \mu\text{M Gd}^{3+}$ (Fig. 7b). Similarly, knockdown of *STIM1* did not prevent activation of TRPC7 and did not change its sensitivity to Gd^{3+} (Fig. 7c). Finally, $\text{STIM1}^{\text{D76A}}$ did not activate TRPC7 and did not increase its spontaneous activity.

DISCUSSION

The study examines fundamental aspects of the function of TRPC channels and their regulation by STIM1, and the mechanism by which STIM1 may regulate SOCs. The first major finding is that all TRPC subunits, except TRPC7, are directly or indirectly regulated by STIM1. Indirect regulation is dependent on a STIM1-mediated heteromultimerization with TRPC subunits that can directly bind STIM1. It has long been controversial whether TRPC channels, in particular TRPC3 and TRPC6, function as SOCs^{15,18–21,22}. Regulation of a channel by STIM1 and the requirement of translocation of STIM1 from the endoplasmic reticulum to plasma-membrane domains for channel activation can be considered one form of SOC — that is, an endoplasmic reticulum Ca^{2+} -content sensor (STIM1) regulates a plasma membrane Ca^{2+} channel, and the regulation by the sensor requires rearrangement of STIM1 in response to depletion of the stored Ca^{2+} . We propose this as a new molecular definition of SOCs. Regulation of TRPC channels by STIM1 is concluded from inhibition of channel function by knockdown of *STIM1*. Requirement for translocation of STIM1 is concluded from activation of the channels by the constitutively active $\text{STIM1}^{\text{D76A}}$ and from their inhibition by $\Delta\text{ERM-STIM1}^{\text{D76A}}$, which prevents the rearrangement of STIM1 into punctae in response to store depletion. By these criteria, it is clear that TRPC1, TRPC4 and TRPC5 function as SOCs under all conditions, whereas TRPC3 and TRPC6 function as SOCs when expressed at low, physiological levels. However, the function of TRPC3 and TRPC6 as SOCs is indirect and is mediated by their regulated interaction with other TRPC channels.

A second finding of note is that STIM1 heteromultimerizes TRPC channels to confer their function as SOCs. The specificity of this effect is quite remarkable. STIM1 mediates the interaction between TRPC1–TRPC3 and between TRPC4–TRPC6. Moreover, the TRPC1–TRPC3 and TRPC4–TRPC6 multimerization is required for regulation of TRPC3 and TRPC6 by STIM1 and their function as SOCs. Assembly of the TRPC1–TRPC3 and TRPC4–TRPC6 into stable complexes is enhanced or stabilized by agonist stimulation, and is mediated by intracellular, rather than the plasma membrane, STIM1. This implies that the store depletion-mediated clustering of STIM1 is likely to facilitate interaction between the channels to allow regulation of all heteromeric TRPC channels by STIM1. The STIM1-mediated formation of the heteromultimers not only increases the diversity of SOC channels, but also provides the cells with a mechanism to regulate their function and store Ca^{2+} content.

Another important finding of this study is that STIM1 seems to be obligatory for activation of the TRPC channels by receptor stimulation, but not for channel function. This conclusion is based on the finding that high overexpression of TRPC1, TRPC4 (data not shown) and TRPC5 (Fig. 6) does not result in STIM1-independent function of these channels. On the other hand, high overexpression of TRPC3 and TRPC6 results in STIM1-independent channel function and increased spontaneous activity. The findings that TRPC3 and TRPC6 can function in a STIM1-dependent and STIM1-independent manner indicate that STIM1 is not obligatory for channel function. The strongest evidence for the conclusion that STIM1 is not obligatory for channel function was obtained with TRPC5. Deletion of STIM1 prevents activation of TRPC5 by agonist stimulation, but has no effect on activation of TRPC5 by La^{3+} , the size of the current or its regulation by voltage. These observations lead to the conclusion that STIM1 mediates regulation of TRPC channels by receptor stimulation, but STIM1 is not a channel subunit and is not needed for TRPC channel to function. Instead, TRPC channels can be activated by more than one mechanism, one of which is activation by STIM1 that is used by receptors to activate Ca^{2+} influx. The mechanism by which STIM1 activates and/or gates TRPC channels has yet to be determined.

METHODS

Solutions, reagents and clones

TRPC1, TRPC3 and TRPC4 clones were previously described⁴³, and GFP–TRPC5, TRPC6 and TRPC7 clones were generously provided by Y. Mori (Kyoto University, Kyoto, Japan). The antibodies used were anti-TRPC1 (1:500; Sigma, St Louis, MO), anti-TRPC3 (1:3000; a gift from C. Montell, Johns Hopkins, Baltimore, MD), anti-STIM1 antibody for coimmunoprecipitation (BD Biosciences, Franklin Lakes, NJ), HRP-conjugated anti-Myc (Santa Cruz, Santa Cruz, CA) and HRP-conjugated anti-HA (Covance, Princeton, NJ), and anti-Flag (Sigma). The *STIM1* siRNA sequence used was GGCTCTGGATACAGTGCTC, and the siRNA to knockdown *TRPC1* and *TRPC4* were previously reported¹⁵ and their effectiveness was confirmed in the present work. siRNA transfection of HEK293 cells was performed using the Qiagen TransMessenger Transfection kit. The amount of siRNA used was 0.8 μg per 12-well containing 80–90% HEK cells. Wells were coated with 0.5 mg ml^{-1} poly-L-ornithine in 0.15 M borate buffer at pH 8.6. After 6 h siRNA transfection, 70% of cells were replated into fresh wells so that confluency was 80–90% the following day. Plasmids transfection was performed using Lipofectamine 2000 reagent for 6 h. The total amount of cDNA used per 12-well was 0.5 μg . Thus, for a typical transfection, 0.13 μg TRPC channel was used for low expression levels, together with 0.13 μg M3 receptor, 0.13 μg STIM1 and 0.1 μg GFP. For high expression levels of TRPC channels, 0.4–0.5 μg cDNA was used. For coimmunoprecipitation analysis, transfection was performed in 6-well plates, and each cDNA amount was scaled up to a total of 1 μg . Current was measured or cells were harvested and extracted for coimmunoprecipitation analysis the following day.

Western blot and coimmunoprecipitation

Transfected cells were harvested and lysed using 500 μ l binding buffer (PBS containing 1 mM NaVO₃, 10 mM sodium pyrophosphate, 50 mM NaF at pH 7.4 and 1% Triton X-100). The extracts were sonicated and insoluble material was removed by centrifugation at 30,000g for 20 min. For coimmunoprecipitation, 2 μ l anti-HA antibody, 1 μ g anti-STIM1 antibody or 4 μ g anti-TRPC1 antibody were added to 100 μ l cell extract and incubated for 1 h at 4 °C. Then, 50 μ l of 1:1 slurry of protein G–Sepharose 4B beads were added and incubated continued for an additional hour at 4 °C. Beads were washed three times with binding buffer, proteins were released from the beads with 50 μ l of SDS-loading buffer and analysed by SDS–PAGE.

Current measurement

The current of TRPC channels in transiently transfected HEK cells was measured in whole current recording configuration, as described previously^{31,44}. Briefly, the pipette solution contained 140 mM CsCl, 2 mM MgCl₂, 1 mM ATP, 5 mM EGTA, 1.5 mM CaCl₂ (free Ca²⁺, 70 nM) and 10 mM HEPES at pH 7.2, to eliminate K⁺ current and prevent inhibition of the channels by high cytoplasmic Ca²⁺. The bath solution contained 140 mM NaCl or 140 mM NMDG-Cl, 5 mM KCl, 0.5 mM EGTA and 10 mM HEPES at pH 7.4 with NaOH or NMDG-OH⁻). The current was recorded by stepping the membrane potential to –100 mV for 200 ms every 5 s from a holding potential of 0 mV or by 400 ms rapid alterations of membrane potential (RAMPs) from –100 to +100 mV from a holding potential of 0 mV. The current recorded at –100 mV was used to calculate current density as pA/pF and current recorded in multiple experiments was used to obtain the mean \pm s.e.m. and calculate significance by Student's *t*-test.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Y Mori for GFP–TRPC5, TRPC6 and TRPC7 plasmids. This work was supported in part by grant BGIA 06651924 from the Texas American Heart Association to W.Z., National Institutes of Health Grants DE12309 and DK38938 and the Ruth S. Harrell Professorship in Medical Research to S.M. and by the National Institute on Drug Abuse (NIDA; DA00266, DA10309) and the National Institute of Mental Health (NIMH; MH068830) to P.F.W.

References

1. Parekh AB, Putney JW Jr. Store-operated calcium channels. *Physiol Rev* 2005;85:757–810. [PubMed: 15788710]
2. Berridge MJ, Bootman MD, Roderick HL. Calcium signalling: dynamics, homeostasis and remodelling. *Nature Rev Mol Cell Biol* 2003;4:517–529. [PubMed: 12838335]
3. Vig M, et al. CRACM1 is a plasma membrane protein essential for store-operated Ca²⁺ entry. *Science* 2006;312:1220–1223. [PubMed: 16645049]
4. Smyth JT, et al. Emerging perspectives in store-operated Ca²⁺ entry: roles of Orai, Stim and TRP. *Biochim Biophys Acta* 2006;1763:1147–1160. [PubMed: 17034882]
5. Soboloff J, Spassova MA, Dziadek MA, Gill DL. Calcium signals mediated by STIM and Orai proteins — a new paradigm in inter-organelle communication. *Biochim Biophys Acta* 2006;1763:1161–1168. [PubMed: 17084918]
6. Mercer JC, et al. Large store-operated calcium selective currents due to co-expression of Orai1 or Orai2 with the intracellular calcium sensor, Stim1. *J Biol Chem* 2006;281:24979–24990. [PubMed: 16807233]
7. Feske S, et al. A mutation in Orai1 causes immune deficiency by abrogating CRAC channel function. *Nature* 2006;441:179–185. [PubMed: 16582901]

8. Zhang SL, et al. Genome-wide RNAi screen of Ca²⁺ influx identifies genes that regulate Ca²⁺ release-activated Ca²⁺ channel activity. *Proc Natl Acad Sci USA* 2006;103:9357–9362. [PubMed: 16751269]
9. Pedersen SF, Owsianik G, Nilius B. TRP channels: an overview. *Cell Calcium* 2005;38:233–252. [PubMed: 16098585]
10. Takahashi Y, et al. Essential role of the N-terminus of murine Orai1 in store-operated Ca²⁺ entry. *Biochem Biophys Res Comm* 2007;356:45–52. [PubMed: 17343823]
11. Minke B. TRP channels and Ca²⁺ signaling. *Cell Calcium* 2006;40:261–275. [PubMed: 16806461]
12. Kiselyov K, Kim JY, Zeng W, Muallem S. Protein-protein interaction and function TRPC channels. *Pflugers Arch* 2005;451:116–124. [PubMed: 16044307]
13. Freichel M, et al. Lack of an endothelial store-operated Ca²⁺ current impairs agonist-dependent vasorelaxation in *TRPA4*^{-/-} mice. *Nature Cell Biol* 2001;3:121–127.
14. Dietrich A, et al. Increased vascular smooth muscle contractility in *TRPC6*^{-/-} mice. *Mol Cell Biol* 2005;25:6980–6989. [PubMed: 16055711]
15. Zagranichnaya TK, Wu X, Villereal ML. Endogenous TRPC1, TRPC3, and TRPC7 proteins combine to form native store-operated channels in HEK-293 cells. *J Biol Chem* 2005;280:29559–29569. [PubMed: 15972814]
16. Hofmann T, et al. Direct activation of human TRPC6 and TRPC3 channels by diacylglycerol. *Nature* 1999;397:259–263. [PubMed: 9930701]
17. Okada T, et al. Molecular and functional characterization of a novel mouse transient receptor potential protein homologue TRP7. *J Biol Chem* 1999;274:27359–27370. [PubMed: 10488066]
18. Kiselyov K, et al. receptors and store-operated Functional interaction between InsP₃ Htrp3 channels. *Nature* 1998;396:478–482. [PubMed: 9853757]
19. Vazquez G, Lievremon JP, St JBG, Putney JW Jr. Human Trp3 forms both inositol trisphosphate receptor-dependent and receptor-independent store-operated cation channels in DT40 avian B lymphocytes. *Proc Natl Acad Sci USA* 2001;98:11777–11782. [PubMed: 11553786]
20. Trebak M, Bird GS, McKay RR, Putney JW Jr. Comparison of human TRPC3 channels in receptor-activated and store-operated modes. *J Biol Chem* 2002;277:21617–21623. [PubMed: 11943785]
21. Boulay G. Ca²⁺-calmodulin regulates receptor-operated Ca²⁺ entry activity of TRPC6 in HEK-293 cells. *Cell Calcium* 2002;32:201–207. [PubMed: 12379180]
22. Dietrich A, Chubanov V, Kalwa H, Rost BR, Gudermann T. Cation channels of the transient receptor potential superfamily: Their role in physiological and pathophysiological processes of smooth muscle cells. *Pharmacol Therap* 2006;112:744–760. [PubMed: 16842858]
23. Lievremon JP, Bird GS, Putney JW Jr. Canonical transient receptor potential TRPC7 can function as both a receptor- and store-operated channel in HEK-293 cells. *Am J Physiol Cell Physiol* 2004;287:C1709–C1716. [PubMed: 15342342]
24. Roos J, et al. STIM1, an essential and conserved component of store-operated Ca²⁺ channel function. *J Cell Biol* 2005;169:435–445. [PubMed: 15866891]
25. Liou J, et al. STIM is a Ca²⁺ sensor essential for Ca²⁺-store-depletion-triggered Ca²⁺ influx. *Curr Biol* 2005;15:1235–1241. [PubMed: 16005298]
26. Wu MM, Buchanan J, Luik RM, Lewis RS. Ca²⁺ store depletion causes STIM1 to accumulate in ER regions closely associated with the plasma membrane. *J Cell Biol* 2006;174:803–813. [PubMed: 16966422]
27. Baba Y, et al. Coupling of STIM1 to store-operated Ca²⁺ entry through its constitutive and inducible movement in the endoplasmic reticulum. *Proc Natl Acad Sci USA* 2006;103:16704–16709. [PubMed: 17075073]
28. Stathopoulos PB, Li GY, Plevin MJ, Ames JB, Ikura M. Stored Ca²⁺ depletion-induced oligomerization of stromal interaction molecule 1 (STIM1) via the EF-SAM Region. *J Biol Chem* 2006;281:35855–35862. [PubMed: 17020874]
29. Soboloff J, et al. Orai1 and STIM reconstitute store-operated calcium channel function. *J Biol Chem* 2006;281:20661–20665. [PubMed: 16766533]
30. Peinelt C, et al. Amplification of CRAC current by STIM1 and CRACM1 (Orai1). *Nature Cell Biol* 2006;8:771–773. [PubMed: 16733527]

31. Huang GN, et al. STIM1 carboxyl-terminus activates native SOC, I(crac) and TRPC1 channels. *Nature Cell Biol* 2006;8:1003–1010. [PubMed: 16906149]
32. Lopez JJ, Salido GM, Pariente JA, Rosado JA. Interaction of STIM1 with endogenously expressed human canonical TRP1 upon depletion of intracellular Ca²⁺ stores. *J Biol Chem* 2006;281:28254–28264. [PubMed: 16870612]
33. Ong HL, et al. Dynamic assembly of TRPC1/STIM1/Orai1 ternary complex is involved in store operated calcium influx: Evidence for similarities in SOC and CRAC channel components. *J Biol Chem* 2007;282:9105–9116. [PubMed: 17224452]
34. Wedel B, Boyles RR, Putney JW, Bird GS. Role of the store-operated calcium entry proteins, Stim1 and Orai1, in muscarinic-cholinergic receptor stimulated calcium oscillations in human embryonic kidney cells. *J Physiol* 2007;579:679–689. [PubMed: 17218358]
35. Vazquez G, Wedel BJ, Trebak M, St John Bird G, Putney JW Jr. Expression level of the canonical transient receptor potential 3 (TRPC3) channel determines its mechanism of activation. *J Biol Chem* 2003;278:21649–21654. [PubMed: 12686562]
36. Xu XZ, Li HS, Guggino WB, Montell C. Coassembly of TRP and TRPL produces a distinct store-operated conductance. *Cell* 1997;89:1155–1164. [PubMed: 9215637]
37. Liu X, Bandyopadhyay BC, Singh BB, Groschner K, Ambudkar IS. Molecular analysis of a store-operated and 2-acetyl-sn-glycerol-sensitive non-selective cation channel. *J Biol Chem* 2005;280:21600–21606. [PubMed: 15834157]
38. Loessberg PA, Zhao H, Muallem S. Synchronized oscillation of Ca²⁺ entry and Ca²⁺ release in agonist-stimulated AR42J cells. *J Biol Chem* 1991;266:1363–1366. [PubMed: 1899088]
39. Kiselyov K, Wang X, Shin DM, Zang W, Muallem S. Calcium signaling complexes in microdomains of polarized secretory cells. *Cell Calcium* 2006;40:451–459. [PubMed: 17034849]
40. Yeromin AV, et al. Molecular identification of the CRAC channel by altered ion selectivity in a mutant of Orai. *Nature* 2006;443:226–229. [PubMed: 16921385]
41. Jung S, et al. Lanthanides potentiate TRPC5 currents by an action at extracellular sites close to the pore mouth. *J Biol Chem* 2003;278:3562–3571. [PubMed: 12456670]
42. Vazquez G, Bird GS, Mori Y, Putney JW Jr. Native TRPC7 channel activation by an inositol trisphosphate receptor-dependent mechanism. *J Biol Chem* 2006;281:25250–25258. [PubMed: 16822861]
43. Yuan JP, et al. Homer binds TRPC family channels and is required for gating of TRPC1 by IP3 receptors. *Cell* 2003;114:777–789. [PubMed: 14505576]
44. Kim JY, et al. Homer 1 mediates store- and IP3Rs- dependent translocation and retrieval of TRPC3 to the plasma membrane. *J Biol Chem* 2006;281:32540–32549. [PubMed: 16887806]






Figure 1.

Regulation of TRPC3 by STIM1. HEK cells were transfected with TRPC3 alone or TRPC3 and the indicated constructs of STIM1. **(a)** The experimental protocol involves incubation of cells in media containing 150 mM Na⁺ to measure the monovalent current before and after cell stimulation with 100 μM carbachol. Na⁺ was then replaced with NMDG⁺ to determine the zero current level (dashed line). Whole-cell currents measured in cells transfected with TRPC3 alone or with STIM1^{D76A}, STIM1^{CT} or ΔERM–STIM1^{D76A} are shown. **(b)** The I/V relationships of HEK cells expressing TRPC3 before (grey trace) and after stimulation with carbachol (carb), cells transfected with TRPC3 and STIM1^{D76A} before (blue trace) and after stimulation with carbachol and cells transfected with ΔERM–STIM1^{D76A} and stimulated with carbachol. Note the typical TRPC3 I/V. **(c)** Summary of the results of TRPC3 alone ($n = 8$) and experiments with the indicated conditions ($n = 5$). The spontaneous current density measured before cell stimulation, the current activated by carbachol and the total current are shown. The error bars represent mean \pm s.e.m.

**Figure 2.**

STIM1-mediated interaction of TRPC3 with TRPC1. **(a)** STIM1 specifically coimmunoprecipitates TRPC1 and TRPC3. HEK cells were transfected with Flag-TRPC3 alone (C3), Flag-TRPC3 + HA-TRPC1 (+C1), HA-TRPC4 (+C4) or HA-TRPC5 (+C5) and the extracts were used to immunoprecipitate native STIM1 and probed for coimmunoprecipitation of TRPC3. Note that TRPC1 selectively mediates the coimmunoprecipitation of STIM1 with TRPC3. **(b)** STIM1 is required for TRPC1-TRPC3 interaction. HEK cells treated with scrambled (first three lanes) and *STIM1* siRNA were transfected with Flag-TRPC3 and Myc-STIM1 (C3+STIM1), Flag-TRPC3 + HA-TRPC1 (C3+C1) or Flag-TRPC3 + HA-TRPC1 + Myc-STIM1 (C3+C1+STIM1) and the extracts were used to immunoprecipitate TRPC1 with anti-HA antibody and probe for coimmunoprecipitation of TRPC3. Note that knockdown of *STIM1* almost abolishes the coimmunoprecipitation of TRPC1 and TRPC3. **(c)** Rescue of TRPC1-TRPC3 interaction by cytosolic STIM1. HEK cells treated with scrambled (first two lanes) or *STIM1* siRNA (last two lanes) were transfected with TRPC3 and YFP-STIM1 (Y-S) or C3+ C1+Y-S and the extracts were used to immunoprecipitate TRPC1 and probe for coimmunoprecipitate of TRPC3 and TRPC1. Note that YFP-STIM1, which does not translocate to the plasma membrane, restores the TRPC1-TRPC3 complex. **(d)** Agonist stimulation enhances complex formation. HEK cells were transfected with Flag-TRPC3 alone (C3, first lane control); C3+HA-TRPC1 or C3+C1+Myc-STIM1. The cells were left unstimulated (rest), were stimulated with 100 μ M carbachol for 3 min or were stimulated with carbachol for 3 min and inhibited with atropine (Atr) for 3 min. At the end of the treatments, the cells were harvested to prepare the extracts

that were used to immunoprecipitate TRPC1 and blot for coimmunoprecipitate of TRPC1 and TRPC3. Uncropped images of the blots are shown in the Supplementary Information, Fig. S1.

**Figure 3.**

Regulation of TRPC3 by STIM1 requires TRPC1. (a) Rescue of TRPC3 activity by cytosolic STIM1. HEK cells treated with *STIM1* siRNA were transfected with HA-TRPC1 (C1) alone, C1 and YFP-STIM1 (C1+Y-S), Flag-TRPC3 alone (C3) or C3 and YFP-STIM1 (C3+Y-S) and receptor-stimulated current density was measured. The error bars indicate the mean \pm s.e.m. of four experiments. (b) TRPC1 is required for regulation of TRPC3 by STIM1. The insert shows coimmunoprecipitation of TRPC3 with native TRPC1 from cells treated with scrambled or *STIM1* siRNA and transfected with Flag-TRPC3. The native TRPC1 was immunoprecipitated with the Sigma anti-TRPC1 and probed for coimmunoprecipitation of TRPC3 expressed at low levels. Current density was measured in cells transfected with TRPC3 alone or TRPC3+ Δ ERM-STIM1^{D76A}. Expression was in cells treated with scrambled siRNA (control) and *TRPC1* siRNA or *TRPC4* siRNA. The yellow columns represent the percent residual current recorded for TRPC3+ Δ ERM-STIM1^{D76A} in scrambled siRNA cells versus cells treated with *TRPC1* siRNA. The error bars indicate the mean \pm s.e.m. of five experiments with *TRPC1* siRNA and four experiments with *TRPC4* siRNA. (c) STIM1 is not obligatory for TRPC3 activity. Control and *STIM1* siRNA-treated cells were transfected with low and high TRPC3. The blot shows the level of native and expressed TRPC3. For native and expressed TRPC3 the blot was probed with anti-TRPC3 and developed for 15 s and 5 s, respectively. Cells were stimulated with carbachol to record the TRPC3 current. Note that overexpression

of TRPC3 overcomes the dependence of TRPC3 activation on STIM1. The error bars indicate the mean \pm s.e.m. of five experiments. **(d)** Faster activation of heteromultimeric TRPC3. HEK cells transfected with TRPC1 or TRPC3 were repeatedly stimulated with carbachol. Example traces and the average half time for activation of TRPC1 and TRPC3 by the first and second stimulation are shown. The TRPC3 current is approximately twice as large as the TRPC1 current, but it was rescaled for illustration purposes. The error bars indicate the mean \pm s.e.m. of four experiments. Uncropped images of the blots are shown in the Supplementary Information, Fig. S1.




Figure 4.

Regulation of TRPC6 by STIM1. **(a)** HEK cells were transfected with Flag-TRPC6 alone or Flag-TRPC6 and the indicated constructs of STIM1. The experimental protocol was the same as in Fig. 1. Whole-cell current recording measured in cells transfected with TRPC6 alone or with TRPC6+ΔERM-STIM1^{D76A} is shown. **(b)** The I/V relationships of HEK cells expressing TRPC6 before (grey trace) and after (thick dark trace) stimulation with carbachol, and of cells transfected with TRPC6 and ΔERM-STIM1^{D76A} and stimulated with carbachol (thin dark trace). **(c)** The blot shows the level of Flag-TRPC6 expression under the indicated conditions and the graph summarizes the results of six experiments with low and four experiments with high TRPC6 levels. The spontaneous current density measured before cell stimulation, the current activated by carbachol and the total current are plotted. Cells were transfected with low or high levels of TRPC6 alone or together with ΔERM-STIM1^{D76A} or D76A-STIM1 and in

cells treated with *STIM1* siRNA, as indicated. Cells with high levels of TRPC6 were transfected with three times more cDNA than cells with low level of TRPC6. Note that Δ ERM-*STIM1*^{D76A} and *STIM1* siRNA inhibited TRPC6 and *STIM1*^{D76A} increased the spontaneous activity of TRPC6 only when TRPC6 was expressed at low levels. The error bars indicate the mean \pm s.e.m.




Figure 5. Interaction of TRPC6 with TRPC4 mediates regulation of TRPC6 by STIM1. **(a)** Native STIM1 specifically coimmunoprecipitates TRPC4 and TRPC6. HEK cells were transfected with Flag-TRPC6 alone (C6), or C6 with HA-TRPC1 (C6+C1), HA-TRPC4 (C6+C4) or HA-TRPC5 (C6+C5) and the extracts were used to immunoprecipitate native STIM1 and probed for coimmunoprecipitation of TRPC6. **(b)** STIM1 is required for TRPC4-TRPC6 coimmunoprecipitation. HEK cells treated with scrambled (first and second lanes) or *STIM1* siRNA (third lane) were transfected with Flag-TRPC6 (C6) or C6+HA-TRPC4 (C6+C4); last two lanes) and the extracts were used to immunoprecipitate TRPC4 and probe for coimmunoprecipitation of TRPC6. **(c, d)** TRPC4, but not TRPC1, is required for regulation of TRPC6 by STIM1. Current density was measured in cells transfected with TRPC6 alone or TRPC6 and ΔERM-STIM1^{D76A}. TRPC6 and TRPC6+ΔERM-STIM1^{D76A} were expressed in cells treated with scrambled siRNA (control; **c, d**), *TRPC4* siRNA (**c**) or *TRPC1* siRNA (**d**). The spontaneous, stimulated and total current densities are shown. The error bars indicate the mean ± s.e.m. of five experiments with *TRPC4* siRNA and four experiments with *TRPC1* siRNA.





Figure 6.

STIM1 regulates TRPC5 but it is not obligatory for channel function. **(a)** STIM1 interacts with TRPC5. HEK cells were transfected with GFP-TRPC5 alone (C5); C5+ Flag-TRPC3 (C5+C3) and C5+ Flag-TRPC6 (C5+C6) and the extracts were used to immunoprecipitate native STIM1 and probe for coimmunoprecipitation of TRPC5. Note that STIM1 interacts with TRPC5 and the interaction is not affected by TRPC3 or TRPC6. **(b)** High levels of TRPC5 (four times more cDNA) were transfected in HEK cells treated with scrambled siRNA or with *STIM1* siRNA. The current was measured in response to stimulation with carbachol, and then in response to activation of TRPC5 by La³⁺ in the same cells. Representative I/V curves are shown. **(c)** Histogram indicating the mean \pm s.e.m. of the current density recorded in five

experiments of cells expressing low or high levels of TRPC5 and treated with scrambled (control) or *STIM1* siRNA. The cells were stimulated with carbachol and then activated with La^{3+} . Note that knockdown of *STIM1* inhibits receptor-mediated activation of TRPC5 irrespective of expression levels and that knockdown of *STIM1* has no effect on activation of TRPC5 by La^{3+} in the same cells.

**Figure 7.**

TRPC7 is not regulated by STIM1. **(a, b)** HEK cells were transfected with low **(a)** or high (four times more cDNA; **b**) levels of TRPC7 alone or with Δ ERM–STIM1^{D76A} and stimulated with carbachol. The mode of TRPC7 function was determined by measuring the sensitivity to inhibition by 1 and 10 μ M Gd³⁺. Typical I/V relationships at the indicated conditions and the mean \pm s.e.m. of four experiments with low and five experiments with high levels of SLC26A7 are also shown. **(c)** TRPC7 was transfected into HEK cells treated with scrambled (control) or *STIM1* siRNA and the effect of *STIM1* knockdown on TRPC7 current and its sensitivity to Gd³⁺ were measured. **(d)** TRPC7 was expressed alone or with the constitutively active STIM1^{D76A} and the spontaneous, carbachol-stimulated and total current were measured.

STIM1^{D76A} did not increase spontaneous activity of TRPC7. The error bars in **c** and **d** indicate the mean \pm s.e.m. of four experiments.