

Stimulation of Endothelial Cell Prostacyclin Production by Thrombin, Trypsin, and the Ionophore A 23187

BABETTE B. WEKSLER, CHRISTOPHER W. LEY, and ERIC A. JAFFE, *Division of Hematology-Oncology, Department of Medicine, Cornell University Medical College, New York 10021*

ABSTRACT Prostacyclin (PGI₂) is an unstable prostaglandin which inhibits platelet aggregation and serotonin release and causes vasodilation. The PGI₂ activity produced by monolayers of cultured human endothelial cells and fibroblasts was measured by the ability of their supernates to inhibit platelet aggregation in platelet-rich plasma, or to inhibit thrombin-induced [¹⁴C]serotonin release from aspirin-treated, washed platelet suspensions. Monolayers of cultured human endothelial cells, stimulated with sodium arachidonate, thrombin, the ionophore A 23187, or trypsin, secreted PGI₂ into the supernatant medium. Monolayers of fibroblasts produced PGI₂ activity only when stimulated by arachidonate. "Resting," intact monolayers did not produce detectable PGI₂, nor did monolayers treated with ADP or epinephrine. Production of PGI₂ activity was abolished by treatment of the monolayers with indomethacin, tranylcypromine, or 15-hydroperoxy arachidonic acid. The PGI₂ activity of the supernates was destroyed by boiling or acidification. Inhibition of thrombin with diisopropylfluorophosphate, and of trypsin with soybean trypsin inhibitor, abolished the stimulation of PGI₂ production by these enzymes. Production of thrombin at a site of vascular injury could, by stimulating PGI₂ synthesis by endothelial cells adjacent to the injured area, limit the number of platelets involved in the primary hemostatic response and help to localize thrombus formation.

INTRODUCTION

Prostacyclin (PGI₂)¹ is a labile prostaglandin which inhibits platelet aggregation and serotonin release (1-3)

Eric A. Jaffe is the recipient of National Institutes of Health Research Career Development award 1 KO4 HL 00237 and a Career Scientist award from the Irma T. Hirschl Trust.

Received for publication 10 March 1978 and in revised form 30 May 1978.

¹Abbreviations used in this paper: DFP, diisopropylfluorophosphate; HPAA, 15-hydroperoxy arachidonic acid; PGI₂, prostacyclin; PRP, platelet-rich plasma.

and counteracts the vasoconstrictive effects of platelet-derived thromboxane A₂ by directly inducing vasodilation (4, 5). In tissues and in platelets, PGI₂ produces a rise in intracellular cyclic AMP levels by stimulating adenylate cyclase (6-9). We have previously demonstrated that suspensions of cultured human and bovine endothelial cells produce PGI₂ when stimulated with sodium arachidonate or when incubated with platelets capable of generating prostaglandin endoperoxides (10).

We now report that intact monolayers of human cultured endothelial cells incubated with thrombin, the ionophore A 23187, trypsin, or sodium arachidonate produce PGI₂. However, "resting", undisturbed endothelial cell monolayers as well as monolayers treated with ADP or epinephrine do not produce significant amounts of PGI₂.

METHODS

Endothelial cell and fibroblast monolayer cultures. Human endothelial cells derived from umbilical cord veins were cultured in T-25 flasks in Medium 199 which contained 20% pooled human serum, with techniques previously described (11). Flasks which contained confluent endothelial cell cultures in passages 1-5 were used in these studies; 38 separate lines were evaluated. 10 lines of fibroblasts derived from human embryo skin, human adult skin, and human foreskin were obtained from Doctors Gretchen Darlington and Ted Brown (Cornell University Medical College), the American Type Culture Collection, Rockville, Md. (Registry nos. CCL 110 and CCL 75 (WI-38)), and the Institute for Medical Research, Camden, N. J. (Registry nos. IMR-90, IMR-91, GM-1603, GM-1604, AG-2257, and AG-2258). The fibroblasts were cultured in minimal essential medium (Flow Laboratories, Inc., Rockville, Md.) which contained 10 or 15% fetal calf serum (depending upon the fibroblast line used), and were used in passages 3-37 when confluent. Before use, the culture medium was removed and the cell monolayers were washed twice with gentle rinsing with 5 ml Buffer A (10 mM Hepes, pH 7.55, 150 mM NaCl, 5 mM KCl, 1.8 mM CaCl₂, 1 mM MgCl₂, and 5 mM glucose). Stimuli were added to Buffer A at desired concentrations and then 2.5 ml of the mixture was incubated with the washed monolayers for 30 s-10 min. Stimuli used included arachidonic acid, 3'5'-ADP (both from Sigma Chemical Co., St. Louis, Mo.), ionophore A 23187 (a gift from

Dr. R. L. Hamill, Eli Lilly and Company, Indianapolis, Ind.), epinephrine (Parke, Davis & Company, Detroit, Mich.), 9,11-azoprostanoid III (a gift from Dr. E. J. Corey, Harvard University, Cambridge, Mass.), and purified human thrombin (a gift from J. Fenton, Albany, N. Y.). The supernatant fluid was then removed and tested immediately for PGI₂ by bioassay (*vide infra*) or kept on ice until use. Authentic PGI₂ (a gift from Dr. K. C. Nicolaou, University of Pennsylvania, Philadelphia, Pa.) was used as a standard. After the incubation, the endothelial cells (or fibroblasts) were removed from the culture flasks by treatment with collagenase (0.1%, 10 min, 37°C) and counted in a hemocytometer. Cell counts in all flasks of any one cell type used on any given day varied by <10%. Flasks of endothelial cells contained 0.6–1.7 × 10⁶ cells/flask, and flasks of fibroblasts contained 0.8–1.7 × 10⁶ cells/flask.

Platelet preparations. Blood from normal human donors who had not ingested aspirin-containing drugs for at least 10 days was obtained by venipuncture, with a two-syringe technique, and mixed with 0.1 vol of 3.2% trisodium citrate. Platelet-rich plasma (PRP) was prepared by centrifugation at 150 g for 10 min at room temperature and was kept tightly capped under 5% CO₂; 95% air until use.

Washed platelet suspensions for the thrombin release assay (*vide infra*) were prepared from normal human venous blood drawn into 1/6 vol of acid citrate dextrose following a modification of the procedure of Mustard et al. (12) as previously described (10). In addition, the platelets were incubated with 100 μM aspirin for 20 min in the first washing buffer to prevent generation of endogenous prostaglandin endoperoxide or thromboxane A₂. The platelet suspension was labeled with [2'-¹⁴C]5-hydroxytryptamine creatinine sulfate (sp act 53 mCi/mmol, Amersham Corp., Arlington Heights, Ill.) in the second washing buffer by addition of 0.167–0.555 μCi/ml platelet suspension (to give a serotonin concentration of 3–10.5 μM) for 15 min at 37°C. Approximately 90–95% of the [¹⁴C]serotonin was taken up by the platelets. The washing procedure was then completed and the platelet suspension was adjusted to 500,000 platelets/μl in Tyrode's solution which contained 0.1% apyrase and 0.35% bovine serum albumin.

Platelet aggregation. Platelet aggregation studies were performed with 0.4 ml aliquots of PRP stirred at 1,000 rpm at 37°C in a dual channel Payton aggregation module with a linear recorder (Payton Associates, Buffalo, N. Y.). Final reaction volumes were 0.5 ml. Threshold values of arachidonic acid (usually 0.2–0.5 mM) were determined for each batch of PRP and were used throughout the experiment to induce aggregation. 90–98 μl of Buffer A or endothelial cell (or fibroblast) supernate was added to the cuvette which contained PRP at 37°C and stirred for 1 min. Then the threshold-dose of aggregating agent was added and the response recorded.

Thrombin release assay. The release of [¹⁴C]serotonin from prelabeled, washed, aspirin-treated platelet suspensions by thrombin in the presence or absence of PGI₂ or supernatant fluids which contained PGI₂ was measured by modifications of the method of Baenziger et al. (13). The final suspension of platelets contained 1 μM imipramine to prevent re-uptake of released serotonin during the assay. Endothelial cell or fibroblast supernates were removed from the culture flasks, 320 μl of supernate was mixed with 1.6 ml of platelet suspension for 2 min and 240-μl portions of the treated platelet suspension were then placed into a series of 1.5 ml polypropylene centrifuge tubes in a 37°C dry heating block. At 10-s intervals, 10 μl of different dilutions of thrombin were added to give final thrombin concentrations of 0, 0.05, 0.1, 0.25, 0.5, 1.0, 5.0, and 10.0 U/ml. After the platelet suspensions were incubated at 37°C with thrombin for exactly 2 min, the reaction was stopped by addition of 0.25 ml of

ice-cold 2.4% paraformaldehyde, and the tubes were transferred to an ice bath. The chilled tubes were then centrifuged for 30 s at 8,000 g and 200-μl samples of each supernate were placed in scintillation minivials which contained 2.7 ml of PCS (Amersham Corp.) and 0.1 ml of H₂O, and were counted in a Searle Mark III liquid scintillation counter (Searle Diagnostics Inc., subsid. of G. D. Searle & Co., Des Plaines, Ill.). In some experiments, the platelet suspension was directly layered over the cell monolayers in tissue culture flasks for 2 min and then removed and tested as above.

A release curve was constructed by plotting the percent of serotonin released into the supernate of the platelet suspension against the log of the thrombin concentration. The data were analyzed by linearizing the standard dose:response curve with the logit transformation (14). A linear least-squares regression analysis of logit (B/B₀) versus ln thrombin concentration (in U/ml) was performed on an HP-97 programmable desk calculator (Hewlett-Packard Corp., McMinnville, Oreg.). The correlation coefficient (*r*) in all cases was >0.95, and was usually >0.99. In this analysis, B is the serotonin (as disintegrations per minute) retained in the platelets (i.e., the total serotonin minus the serotonin released) at an experimental point, B₀ is the disintegrations per minute retained at zero thrombin concentration, and N is the disintegrations per minute retained at the concentration of thrombin giving maximal release. Logit (B/B₀) is ln (B – N)/(B₀ – B). The thrombin concentration which yielded 50% of maximum release was calculated (T₅₀). Thus the effects of PGI₂ in monolayer supernates on thrombin-induced serotonin release can be expressed quantitatively as the percent change in the thrombin concentration producing 50% of maximal release (ΔT₅₀) where,

$$\Delta T_{50} = \frac{T_{50}(\text{sample}) - T_{50}(\text{control})}{T_{50}(\text{control})} \times 100.$$

Expressing results in terms of ΔT₅₀ allowed us to compensate for the variation in thrombin sensitivity observed among platelet donors. The results of experiments performed on the same day on replicates from the same cell line usually varied by <12%. The results of this assay were assessed statistically with a one sample *t* test where the null hypothesis was a ΔT₅₀ = 0 (i.e., no effect).

Evaluation for carryover of agents used to stimulate monolayers. The possibility that agents used to stimulate endothelial cell or fibroblast monolayers might be carried over in the monolayer supernate and alter the thrombin-induced serotonin release assay results was evaluated. Thrombin carryover will affect the [¹⁴C]serotonin release assay either by decreasing the T₅₀ or by causing the release of [¹⁴C]serotonin from platelets in control tubes from which thrombin has been omitted. To assay for residual thrombin, PGI₂ was eliminated either by running an empty-dish control or by treating the cell monolayers with indomethacin to block PGI₂ production. We therefore incubated thrombin (0.1 U/ml) in buffer either in an empty dish or with indomethacin-treated endothelial and fibroblast monolayers for 2 min at 37°C and then assayed the thrombin containing buffer in our usual assay procedure. In both models, thrombin at an initial concentration of 0.1 U/ml (which would yield thrombin at 0.02 U/ml after 1:5 dilution in the assay) neither lowered the T₅₀ nor released [¹⁴C]serotonin from platelets in control tubes from which thrombin was omitted. However, thrombin at an initial concentration of 0.5 U/ml in both models produced a decrease of T₅₀ and caused release of [¹⁴C]serotonin from platelets in control tubes from which thrombin was omitted. The release of [¹⁴C]serotonin seen in this case was directly equivalent to that obtained with thrombin (0.1 U/ml) in our assay (because the supernate is diluted 1:5

in the assay). Despite this finding, when nonindomethacin-treated endothelial cells were incubated with thrombin at 0.5 U/ml, the endothelial cell supernate reproducibly increased the T_{50} and did not release [14 C]serotonin from platelets in control tubes which lacked thrombin. On the other hand, supernates from nonindomethacin-treated fibroblasts incubated with thrombin at 0.5 U/ml consistently decreased the T_{50} and produced [14 C]serotonin release in platelets in control tubes which lacked thrombin equivalent to that produced by thrombin at 0.1 U/ml. Thus, in the case of endothelial cells stimulated with thrombin at 0.5 U/ml, a net effect is seen whereby the amount of PGI₂ produced reverses not only the effect of the residual thrombin, but also some of the thrombin added in the assay, thus significantly increasing the T_{50} . It is therefore likely that the increased T_{50} , observed when endothelial cells are stimulated with thrombin at 0.5 U/ml, underestimates the true amount of PGI₂ produced.

Carryover of sodium arachidonate, ADP, trypsin, and the ionophore A 23187 at the doses used did not affect the assay because the platelets had been treated with aspirin and thus were unresponsive to these agents. Epinephrine produced a decrease in the T_{50} at all doses used. Possible production of "hidden" PGI₂ by endothelial cells and fibroblasts stimulated by epinephrine was ruled out by showing that the decrease in T_{50} was the same whether or not the cells had been treated with indomethacin.

Similar experiments were performed to rule out the effects of carryover of agents in monolayer supernates used in platelet aggregation studies.

Inhibition of PGI₂ production. Endothelial cell and fibroblast monolayers were washed free of medium and incubated for 20 min at 37°C with the cyclo-oxygenase inhibitor indomethacin (10 µg/ml) (Sigma Chemical Co.), or with the inhibitors of prostacyclin synthesis (15, 16), 15-hydroperoxyarachidonic acid (HPAA) (30 µg/ml) (generously provided by Dr. A. Marcus, Cornell University Medical College) or tranilcypromine (500 µg/ml) (a gift from Dr. H. Green, Smith, Kline & French, Philadelphia, Pa.). Incubations with inhibitors were carried out in Buffer A. Indomethacin and 15-HPAA were removed before the monolayers were stimulated. Tranilcypromine inhibition was assayed both in the presence of drug and after its removal. While indomethacin is a reversible inhibitor of cyclo-oxygenase (17), its effect was not reversed during the short course of our experiment, which included the removal of the indomethacin, a washing of the monolayer with buffer, and a stimulation of the cell monolayers for 2 min with a variety of agents which included arachidonate (20 µM).

Thrombin was inactivated by incubation with diisopropylfluorophosphate (DFP), 10 mM, for 30 min at room temperature, followed by dialysis overnight against 0.75 M NaCl-0.1 M Tris, pH 8.3, at 4°C to remove the excess DFP. DFP treatment decreased thrombin ability to aggregate platelets by more than 90%. Native thrombin used as a control was similarly dialyzed and retained full platelet aggregating activity.

Inhibition of PGI₂ activity. PGI₂ or monolayer supernates which contained PGI₂ activity were inactivated by boiling for 30 s or by acidification to pH 3 for 10 min followed by back titration to pH 7.5.

RESULTS

Inhibition of platelet aggregation by supernates from cell monolayers. Endothelial cell monolayers were washed free of culture medium and incubated with Buffer A. The release of PGI₂ activity into the overlying buffer was measured by the buffer capacity

to inhibit platelet aggregation when added to PRP (Fig. 1). This method detects 0.1 ng/ml of PGI₂ (data not shown). No inhibition of platelet aggregation occurred in the presence of buffer or 10% (vol/vol) monolayer supernate taken from resting endothelial cells (Fig. 1A, curves 1 and 2). However, the supernatant fluid from endothelial cell monolayers incubated for 2 min with 20 µM sodium arachidonate markedly inhibited platelet aggregation (Fig. 1A, curve 3). Endothelial cell monolayers were compared to fibroblast monolayers for capacity to generate PGI₂ activity in their supernate (Fig. 1B). Fibroblasts derived from embryonic skin, foreskin, or adult skin all yielded the same results and are thus considered together. Resting skin fibroblasts did not produce inhibitory supernates (Fig. 1B, curve 5). Skin fibroblast monolayers often, but not invariably, generated inhibitory activity in the presence of 20 µM arachidonate (Fig. 1B, curve 6). As little as 2 µl of supernate derived from incubation of 20 µM sodium arachidonate with 4 × 10⁵ endothelial cells/ml of Buffer A completely inhibited platelet aggregation in PRP. This was equivalent to inhibition by 0.1 nM PGI₂ or by 90–100 µl of supernate from similarly incubated fibroblast monolayers.

Endothelial cell and fibroblast monolayers were incubated with a number of agents that aggregate platelets (with concentrations, however, which at the same dilution alone did not produce aggregation in test PRP). The supernatant medium was then removed and tested for its effects on platelet aggregation in PRP challenged




FIGURE 1 Inhibition of platelet aggregation in PRP by supernates from monolayers of endothelial cells (A) and fibroblasts (B). Supernates were added at the first arrow and 0.3 mM sodium arachidonate (AA) at the second arrow. (A) (1) Control-Buffer A, (2) Supernate from unstimulated endothelial cells, and (3) supernate from endothelial cells stimulated with 20 µM sodium arachidonate (AA). (B) (4) Control-Buffer A, (5) supernate from unstimulated skin fibroblasts, and (6) supernate from skin fibroblasts stimulated with 20 µM sodium arachidonate (AA).

with sodium arachidonate. Neither the culture medium nor the buffer used to wash the resting monolayers produced any change in the normal aggregation responses of PRP. Endothelial cell monolayers incubated with sodium arachidonate (20 μ M), thrombin (0.1 U/ml), or the ionophore A 23187 (10 μ M) released PGI₂ activity into the supernatant buffer. These supernates inhibited platelet aggregation in PRP stimulated by arachidonate. ADP (1–5 μ M), epinephrine (0.1–1 μ M), and the endoperoxide analogue, azoprostanoic acid (12.6 μ g/ml), did not stimulate production of PGI₂ by endothelial cell monolayers in this test system. Fibroblast monolayers derived from skin produced PGI₂ activity only after incubation with arachidonate (20 μ M). Incubation with ionophore A 23187, thrombin, ADP, epinephrine, or azoprostanoic acid at the same doses as used on endothelial cells did not stimulate production of PGI₂ by fibroblast monolayers in this test system.

Mechanical stimulation of endothelial cell monolayers by a rubber policeman in the absence of any other stimulus led to the rapid appearance of PGI₂ activity in the supernate.

Inhibition of [¹⁴C]serotonin release: quantitation of PGI₂ activity produced by endothelial cell and fibroblast monolayers. Inhibition of thrombin-induced release of [¹⁴C]serotonin from aspirin-treated, washed platelets by synthetic PGI₂ or by supernatant fluids from endothelial cell and fibroblast monolayers was used to measure PGI₂ activity. Treatment of a washed platelet suspension with PGI₂ or with cell monolayer supernates which contained PGI₂ activity made the platelets less sensitive to thrombin, and thus shifted the curve of serotonin release. Examples of a standard curve of serotonin release and the effect of 10 nM of PGI₂ on this curve are shown in Fig. 2A. By means of a logit transformation of the data (Fig. 2B) the release curves are linearized and the T₅₀ (thrombin concentration which produces 50% of the maximal serotonin release) can be calculated. This assay can detect PGI₂ at concentrations as low as 10–100 pM depending upon the platelet donor. The concentration of thrombin required to liberate 50% of maximally releasable [¹⁴C]serotonin from control platelets was 0.25 ± 0.14 U/ml (mean ± SD, n = 36) in this system. While the inter-donor variation in platelet sensitivity to thrombin was quite large, the intra-donor variation was much smaller (i.e., average coefficient of variation = 15.3%).

We used the quantitative [¹⁴C]serotonin-release assay to compare the production of PGI₂ activity by endothelial cell monolayers stimulated by different agents. Because PGI₂ activity shifted the [¹⁴C]serotonin release curve and increased the T₅₀, the PGI₂ effect was expressed as percent change in T₅₀ or Δ T₅₀ (Methods).

Endothelial cell monolayers, incubated with buffer alone, produced no detectable PGI₂ activity when compared to control, untreated platelets (Table I). The ad-




FIGURE 2 Inhibition of thrombin-induced [¹⁴C]serotonin release from aspirin-treated, washed platelets by PGI₂. (A) [¹⁴C]Serotonin release as a function of thrombin concentration in the presence of Buffer A (●) and in the presence of 10 nM PGI₂ (○). (B) Logit plot of data in panel A. 50% release of [¹⁴C]serotonin (T₅₀) occurs at logit = 0.

dition of exogenous arachidonate resulted in significant production of PGI₂ activity. Thrombin treatment of endothelial cell monolayers also resulted in significant PGI₂ production related to the dose of thrombin used. No thrombin carryover effect was noted with thrombin at 0.1 U/ml. However, 0.5 U/ml doses of thrombin produced carryover (Methods). Despite the carryover of thrombin, the endothelial cells produced enough PGI₂ to completely mask the presence of residual thrombin. It is therefore likely that the T₅₀ observed when endothelial cells are stimulated with 0.5 U/ml of thrombin underestimates the true amount of PGI₂ produced. Treatment of endothelial cell monolayers with DFP-inactivated thrombin did not stimulate PGI₂ synthesis. The ionophore A 23187 stimulated production of large amounts of PGI₂ activity in a dose-related fashion. ADP did not lead to significant PGI₂ production. Epinephrine was ineffective in stimulating PGI₂ production. More-

TABLE I
PGI₂ Production in Stimulated Endothelial Cell Monolayers

	ΔT_{50}^*	SEM	n	P
Control platelets†	0	—	36	—
Endothelial cells§				
Buffer A	-1.3	5.3	5	NS
Sodium arachidonate (20 μ M)	271	43.4	18	<0.0001
Thrombin (0.5 U/ml)	171	21.9	7	<0.0001
(0.1 U/ml)	73	21.3	5	<0.05
Ionophore A 23187 (10 μ M)	512	177	6	<0.05
(5 μ M)	433	67.9	4	<0.01
ADP (5 μ M)	10.7	6.6	6	NS
Epinephrine (0.9 μ M)	-33.8	2.9	8	<0.0001

$$* \Delta T_{50} = \frac{T_{50} - T_{50}(\text{control})}{T_{50}(\text{control})} \times 100\%$$

† T_{50} of control platelets = 0.25 ± 0.14 U/ml (mean \pm SD, $n = 36$).

§ Endothelial cell monolayers were incubated with the listed agents for 2 min at 37°C and the supernates removed and tested. All agents were dissolved in Buffer A.

over, supernates from the epinephrine-treated endothelial cells potentiated thrombin-induced serotonin release and lowered the T_{50} because of epinephrine carryover.

Fibroblasts, incubated under identical conditions, produced statistically significant amounts of PGI₂ only when stimulated by arachidonate (Table II). Thrombin, ionophore A 23187, and ADP did not stimulate the production of statistically significant amounts of PGI₂. Epinephrine, as in endothelial cells, significantly potentiated thrombin-induced serotonin release. Fibroblasts as early as passage 3 were unresponsive to thrombin and ionophore, whereas endothelial cells at all passages tested (passages 1–5) produced PGI₂ in response to these stimuli.

Time course of PGI₂ production after stimulation of monolayers. The production of PGI₂ activity after stimulation of endothelial cell monolayers by sodium arachidonate or thrombin was rapid and transient. Maximum production of PGI₂ was observed within 1–2 min, as measured semiquantitatively by the delay in platelet aggregation induced by aliquots of supernate sequentially removed from monolayers stimulated with arachidonate (20 μ M) or thrombin (0.5 U/ml). PGI₂ activity was undetectable 10 min after stimulation. Production of PGI₂ activity by fibroblasts stimulated with sodium arachidonate followed a similar time course.

Once stimulated, an endothelial cell monolayer was less responsive to a second stimulation given 15 min to 1 h later. Thus the response to a second arachidonate stimulus was about 40% less than to the first arachi-

TABLE II
PGI₂ Production in Stimulated Skin Fibroblast Monolayers

	ΔT_{50}^*	SEM	n	P
Control platelets†	0	—	17	—
Fibroblasts§				
Sodium arachidonate (20 μ M)	148.4	51.4	9	<0.02
Thrombin (0.1 U/ml)	21.8	13.9	5	NS
Ionophore A 23187 (10 μ M)	33.3	17.5	4	NS
ADP (5 μ M)	2.2	15.5	3	NS
Epinephrine (0.9 μ M)	-33	7.5	3	<0.05

$$* \Delta T_{50} = \frac{T_{50} - T_{50}(\text{control})}{T_{50}(\text{control})} \times 100.$$

† T_{50} of control platelets = 0.25 ± 0.09 U/ml (mean \pm SD, $n = 17$).

§ Fibroblast monolayers were incubated with the listed agents for 2 min at 37°C and the supernates removed and tested. All agents were dissolved in Buffer A.

donate stimulus. A monolayer stimulated first with thrombin produced PGI₂ in response to a second stimulation by arachidonate; however, a monolayer stimulated first with arachidonate subsequently responded poorly to thrombin. In addition, a monolayer which produced PGI₂ upon thrombin-stimulation failed to respond to a second exposure with thrombin.

Inhibition of PGI₂ production in monolayers of endothelial cells. Indomethacin-treated monolayers of endothelial cells did not produce PGI₂ activity when stimulated with sodium arachidonate (Table III). Production of PGI₂ activity by endothelial cell monolayers stimulated with sodium arachidonate was also prevented by incubation of the monolayers with 15-HPAA (30 μ g/ml) or with tranylcypromine (0.5 mg/ml), both inhibitors of PGI₂ synthesis. Significant inhibition of

TABLE III
Inhibition of PGI₂ Production by Endothelial Cell Monolayers

Inhibitor*	Inhibition†
	%
None (Buffer control)	0
Indomethacin (10 μ g/ml)	94
15-Hydroperoxy arachidonate (30 μ g/ml)	100
Tranylcypromine	
Prior incubation only (500 μ g/ml)	29
Coincubation (500 μ g/ml)	69

* Endothelial cell monolayers (10^6 cells) were treated with the stated inhibitors and then incubated for 2 min with 2.5 ml of Buffer A which contained 20 μ M arachidonate. Supernates were then removed and tested in the [¹⁴C]serotonin release assay.

† Inhibition is expressed as the percent decrease in ΔT_{50} .

PGI₂ production by tranlycypromine, unlike 15-HPAA, required the latter's presence during incubation of endothelial cell monolayers with arachidonate, which indicates that the inhibitory effect of tranlycypromine was reversible. The production of PGI₂ by endothelial cells stimulated by thrombin, trypsin, and ionophore A 23187 were also inhibited by treatment with indomethacin. Indomethacin-treated monolayers could produce PGI₂ when incubated with normal PRP but not when incubated with aspirin-treated PRP. These results are consistent with the inhibition of endothelial cell cyclo-oxygenase by indomethacin. Because normal PRP can generate endoperoxides which act as substrates for PGI₂ formation in the endothelial cells, indomethacin treatment of the monolayers would not be expected to inhibit PGI₂ generation in the presence of normal PRP. Fibroblast production of PGI₂ was similarly inhibited by indomethacin, tranlycypromine, and 15-HPAA (data not shown).

Inactivation of PGI₂ in endothelial cell monolayer supernates. The PGI₂ activity of supernatant fluid removed from endothelial cell (or fibroblast) monolayers stimulated with all of the agents tested was destroyed by brief boiling or acidification to pH 3 for 10 min. Inhibitory activity was also lost when the supernate was incubated at room temperature for 30 min at pH 7.4, or at 37°C for 15 min. Activity was preserved for several hours when the supernate was kept on ice at pH 7.4 or pH 8.6, and for days when it was frozen at -70°C at pH 8.6.

Effects on PGI₂ production of incubating endothelial cell monolayers with trypsin and chymotrypsin. Endothelial cell monolayers were incubated for 2 min with 0.0001-0.0025% trypsin and the culture supernatant fluid then examined for PGI₂ activity. These concentrations of trypsin had no effect on platelet aggregation or on thrombin-induced [¹⁴C]serotonin release from platelets. The cells appeared to contract during trypsin treatment when viewed by phase microscopy, but did not detach from the culture flask. The trypsin-treated monolayers released PGI₂ and significantly shifted the T₅₀. Thus, monolayers treated with 0.001% trypsin gave a $\Delta T_{50} = 225 \pm 32$ (mean \pm S.E.M., $P < 0.01$). Prior treatment of trypsin with a twofold excess of soybean trypsin inhibitor abolished the stimulatory effect of trypsin on PGI₂ production. The trypsin concentration required for this effect on the T₅₀ was, on a molar basis, about 200-fold greater than the thrombin concentration which gave a similar effect. Chymotrypsin at a concentration of 0.001% did not stimulate PGI₂ activity.

Fibroblasts incubated with similar concentrations of trypsin did not produce PGI₂.

DISCUSSION

Monolayers of cultured human endothelial cells produced PGI₂ after brief incubation with sodium arachi-

donate, thrombin, the calcium ionophore A 23187, and trypsin, or after mechanical stimulation. However, resting monolayers, or monolayers incubated with ADP, or epinephrine did not produce PGI₂. PGI₂ activity was assayed by inhibition both of platelet aggregation and of [¹⁴C]serotonin release. Fibroblasts produced PGI₂ only after incubation with arachidonate. Our results with fibroblast monolayers are consistent with the findings of Baenziger et al, who showed that PGI₂ is produced by fibroblasts stimulated by scraping the cells from the culture dish (13) and by intact cells incubated with arachidonate.² Confirmation that the inhibitor produced by the monolayers was a prostaglandin was obtained by blocking its production with indomethacin or aspirin.³ Such indomethacin-treated monolayers, however, could generate PGI₂ activity when they were placed in contact with normal, heparinized PRP which acted as a source of prostaglandin endoperoxides. Specific evidence that the platelet-inhibitory activity was PGI₂ was provided by studies in which 15-HPAA and tranlycypromine, inhibitors of endoperoxide conversion of PGI₂, prevented its synthesis (15, 16). Moreover, heating or acidifying the active cell supernates, conditions known to inactivate PGI₂ but not prostaglandin D₂ (15), abolished the inhibitory activity. These experiments clearly indicated that the inhibition of platelet aggregation and of thrombin-induced [¹⁴C]serotonin release from platelets exposed to the cell monolayer supernates was a result of PGI₂ in the supernates.

Because thrombin stimulated the production of PGI₂ by the endothelial cell monolayers without addition of arachidonate, the substrate for PGI₂ synthesis must have been endogenous arachidonate. We suggest that thrombin bound to a recently described specific receptor (18) on the endothelial cells and then activated phospholipase A₂, which in turn released arachidonic acid from membrane phospholipid for prostaglandin synthesis. DFP-treated thrombin did not stimulate endothelial cell PGI₂ production, although DFP-treated thrombin has been reported to bind endothelial cells as well as native thrombin (18). This suggests that thrombin protease activity is required for the stimulation of PGI₂ synthesis in endothelial cell monolayers. Endothelial cell monolayers restimulated with thrombin produced little or no PGI₂. The poor response to thrombin stimulation after an initial arachidonate stimulus probably represents depletion of endogenous arachidonate recruited during the initial response. The lack of response to the second of two sequential thrombin stimuli may represent depletion of endogenous arachidonate and(or) destruction of a thrombin-sensitive cellular component.

Monolayers of stimulated endothelial cells produced

² Baenziger, N. Personal communication.

³ Jaffe, E. A., C. W. Ley, and B. B. Weksler. Manuscript in preparation.

PGI₂ in a "burst"-like pattern. PGI₂ peaked 2–3 min after stimulation and could no longer be detected at 10 min. The limiting factor appeared to be the availability of arachidonate. Thus the cells again produced PGI₂ when exogenous arachidonate was added. It is possible that in intact blood vessels in vivo, exogenous arachidonate from plasma may continuously replenish the endothelial cells' endogenous arachidonate pool and thus allow the endothelial cells to respond continuously with PGI₂ production to a variety of stimuli, either chemical or mechanical. Because mechanical stimulation of endothelial cell monolayers produced PGI₂, mechanical stimuli in vivo (i.e., shearing forces resulting from blood flow) may also stimulate PGI₂ production.

The ability of trypsin to stimulate PGI₂ production indicates that other enzymes besides thrombin may be capable of stimulating PGI₂ synthesis by endothelial cells. It is likely that proteolytic enzymes activate endothelial cell phospholipase A₂ from a precursor form (19). Because PGI₂ produced by endothelial cells has been shown to inhibit polymorphonuclear leukocyte function (20), the ability of various proteolytic enzymes to induce PGI₂ synthesis may represent a modulating mechanism in the control of inflammatory processes.

The ionophore A 23187 has recently been shown to stimulate prostaglandin and thromboxane synthesis in platelets by mobilizing intracellular calcium and by activating phospholipase A₂, a calcium-dependent enzyme (21, 22). At similar concentrations (5–10 μM) to those giving maximal thromboxane A₂ production in platelets (21), ionophore A 23187 stimulated PGI₂ production in endothelial cell monolayers in the absence of exogenous arachidonate, which suggests that the mobilization of intracellular calcium can also activate phospholipase A₂ in endothelial cells, thus making endogenous arachidonate available for PGI₂ synthesis. The finding that indomethacin treatment of the cells blocked PGI₂ synthesis induced by ionophore A 23187 is consistent with this mechanism, because indomethacin-treated endothelial cells cannot utilize arachidonate for PGI₂ synthesis.

Other agents which stimulate platelets, namely ADP and epinephrine, did not stimulate endothelial cell monolayers to synthesize PGI₂. Assessment of ADP-mediated stimulation of PGI₂ production may be complicated by the fact that endothelial cells possess ADPase activity (23, 24). Thus, ADP added to the endothelial cells would be degraded. It is possible that endothelial cells can respond to locally higher doses of ADP, but these interfere with our assay method. Epinephrine did not stimulate PGI₂ production. In fact, epinephrine sensitized the aspirin-treated platelets and decreased the T₅₀.

In conclusion, intact monolayers of cultured human endothelial cells can synthesize PGI₂ either from endogenous arachidonic acid when the cells are stimulated by thrombin, trypsin, or the ionophore A 23187, or from

exogenous endoperoxides. Production of thrombin at a site of vascular injury could, by stimulating PGI₂ production by endothelial cells adjacent to the injured area, limit the number of platelets involved in the primary hemostatic response and help to localize thrombus formation.

ACKNOWLEDGMENTS

We thank Christine Baranowski and Joyce Knapp for excellent technical assistance and Naomi Nemtsov for typing the manuscript.

This work was supported by The National Institutes of Health through Specialized Center for Thrombosis grant HL 18828, the New York Heart Association, and the Arnold R. Krakower Hematology Foundation.

REFERENCES

1. Moncada, S., R. Gryglewski, S. Bunting, and J. R. Vane. 1976. An enzyme isolated from arteries transforms prostaglandin endoperoxides to an unstable substance that inhibits platelet aggregation. *Nature (Lond.)* **263**: 663–665.
2. Johnson, R. A., D. R. Morton, J. H. Kinner, R. R. Gorman, J. C. McGuire, F. F. Sun, N. Whittaker, S. Bunting, J. Salmon, S. Moncada, and J. R. Vane. 1976. The chemical structure of prostaglandin X (prostacyclin). *Prostaglandins*. **12**: 915–927.
3. Moncada, S., E. A. Higgs, and J. R. Vane. 1977. Human arterial and venous tissues generate prostacyclin (prostaglandin X), a potent inhibitor of platelet aggregation. *Lancet*. **1**: 18–20.
4. Dusting, G. J., S. Moncada, and J. R. Vane. 1977. Prostacyclin (PGX) is the endogenous metabolite responsible for relaxation of coronary arteries induced by arachidonic acid. *Prostaglandins*. **13**: 3–15.
5. Raz, A., P. C. Isakson, M. S. Minkes, and P. Needleman. 1977. Characterization of a novel metabolic pathway of arachidonate in coronary arteries which generates a potent endogenous coronary vasodilator. *J. Biol. Chem.* **252**: 1123–1126.
6. Gorman, R. R., S. Bunting, and O. V. Miller. 1977. Modulation of human platelet adenylate cyclase by prostacyclin (PGX). *Prostaglandins*. **13**: 337–338.
7. Tateson, J. E., S. Moncada, and J. R. Vane. 1977. Effects of prostacyclin (PGX) on cyclic AMP concentrations in human platelets. *Prostaglandins*. **13**: 389–397.
8. Lapetina, E. G., C. J. Schmitges, K. Chandrabose, and P. Cuatrecasas. 1977. Cyclic adenosine 3'5'-monophosphate and prostacyclin inhibit membrane phospholipase activity in platelets. *Biochem Biophys. Res. Commun.* **76**: 828–835.
9. Gerrard, J. M., J. D. Peller, T. P. Krick, and J. G. White. 1977. Cyclic AMP and platelet prostaglandin synthesis. *Prostaglandins*. **14**: 39–50.
10. Weksler, B. B., A. J. Marcus, and E. A. Jaffe. 1977. Synthesis of prostaglandin I₂ (prostacyclin) by cultured human and bovine endothelial cells. *Proc. Natl. Acad. Sci. U. S. A.* **74**: 3922–3926.
11. Jaffe, E. A., R. L. Nachman, C. G. Becker, and C. R. Minick. 1973. Culture of human endothelial cells derived from umbilical veins. *J. Clin. Invest.* **52**: 2745–2756.
12. Mustard, J. F., D. W. Perry, N. G. Ardlie, and M. A. Packham. 1972. Preparation of suspensions of washed platelets from humans. *Br. J. Haematol.* **22**: 193–204.
13. Baenziger, N. L., M. J. Dillender, and P. W. Majerus. 1977. Cultured human skin fibroblasts and arterial cells

- produce a labile platelet-inhibitory prostaglandin. *Biochem. Biophys. Res. Commun.* **78**: 294–301.
14. Rodbard, D., and G. R. Frazier. 1975. Statistical analysis of radioligand assay data. *Methods Enzymol.* **37**: 3–22.
 15. Moncada, S., P. Needleman, S. Bunting, and J. R. Vane. 1976. Prostaglandin endoperoxide and thromboxane generating systems and their selective inhibition. *Prostaglandins.* **12**: 323–333.
 16. Moncada, S., R. J. Gryglewski, S. Bunting, and J. R. Vane. 1976. A lipid peroxide inhibits the enzyme in blood vessel microsomes that generates from prostaglandin endoperoxides the substance (prostaglandin X) which prevents platelet aggregation. *Prostaglandins.* **12**: 715–738.
 17. Stanford, N., G. J. Roth, T. Y. Shen, and P. W. Majerus. 1977. Lack of covalent modification of prostaglandin synthetase (cyclo-oxygenase) by indomethacin. *Prostaglandins.* **13**: 669–675.
 18. Awbrey, B. J., W. G. Owen, J. C. Hoak, and G. L. Fry. 1977. Binding of thrombin to endothelial cells. *Blood.* **50**(Suppl. 1): 257. (Abstr.)
 19. Pickett, W. C., R. L. Jesse, and P. Cohen. 1976. Trypsin-induced phospholipase activity in human platelets. *Biochem. J.* **160**: 405–408.
 20. Weksler, B. B., J. M. Knapp, and E. A. Jaffe. 1977. Prostacyclin (PGI₂) synthesized by cultured endothelial cells modulates polymorphonuclear leukocyte function. *Blood.* **50**(Suppl. 1): 287. (Abstr.)
 21. Knapp, H. R., O. Oelz, L. J. Roberts, B. J. Sweetman, J. O. Oates, and P. W. Reed. 1977. Ionophores stimulate prostaglandin and thromboxane biosynthesis. *Proc. Natl. Acad. Sci. U. S. A.* **74**: 4251–4255.
 22. Pickett, W. C., R. L. Jesse, and P. Cohen. 1977. Initiation of phospholipase A₂ activity in human platelets by the calcium ion ionophore A 23187. *Biochim. Biophys. Acta.* **486**: 209–213.
 23. Heyns, A du P., C. J. Badenhorst, and F. P. Retief. 1977. ADPase activity of normal and atherosclerotic human aorta intima. *Thromb. Haemostasis.* **37**: 429–435.
 24. Glasgow, J. E., and F. A. Pitlick. 1977. Type-specific differences in the ability of cultured human cells to hydrolyze adenine nucleotides and to trigger the platelet release reaction. *Fed. Proc.* **36**: 1082. (Abstr.)