

# Stochastic Block Model and Community Detection in Sparse Graphs: A spectral algorithm with optimal rate of recovery \*

**Peter Chin**  
*Boston University*

SPCHIN@CS.BU.EDU

**Anup Rao**  
*Yale University*

ANUP.RAO@YALE.EDU

**Van Vu**  
*Yale University*

VAN.VU@YALE.EDU

## Abstract

In this paper, we present and analyze a simple and robust spectral algorithm for the stochastic block model with  $k$  blocks, for any  $k$  fixed. Our algorithm works with graphs having constant edge density, under an optimal condition on the gap between the density inside a block and the density between the blocks. As a co-product, we settle an open question posed by Abbe et. al. concerning censor block models.

**Keywords:** Stochastic Block Model, Censor Block Model, Spectral Algorithm

## 1. Introduction

Community detection is an important problem in statistics, theoretical computer science and image processing. A widely studied theoretical model in this area is the stochastic block model. In the simplest case, there are two blocks  $V_1, V_2$  each of size of  $n$ ; one considers a random graph generated from the following distribution: an edge between vertices belonging to the same block appears with probability  $\frac{a}{n}$  and an edge between vertices across different blocks appear with probability  $\frac{b}{n}$ , where  $a > b > 0$ . Given an instance of this graph, we would like to identify the two blocks as correctly as possible. Our paper will deal with the general case of  $k \geq 2$  blocks, but for the sake of simplicity, let us first focus on  $k = 2$ .

For  $k = 2$ , the problem can be seen as a variant of the well known hidden bipartition problem, which has been studied by many researchers in theoretical computer science, starting with the work of [Bui et al. \(1987\)](#); (see [Dyer and Frieze \(1989\)](#) [Boppana \(1987\)](#) [Jerrum and Sorkin \(1993\)](#) [McSherry \(2001\)](#) and the references therein for further developments). In these earlier papers,  $a$  and  $b$  are large (at least  $\log n$ ) and the goal is to recover both blocks completely. It is known that one can efficiently obtain a complete recovery if  $\frac{(a-b)^2}{a+b} \geq C \frac{\log n}{n}$  and  $a, b \geq C \log n$  for some sufficiently large constant  $C$  (see, for instance [Vu \(2014\)](#)).

In the stochastic block model problem, the graph is sparse with  $a$  and  $b$  being constants. Classical results from random graph theory tell us that in this range the graph contains, with high probability, a linear portion of isolated vertices [Bollobas \(2001\)](#). Apparently, there is no way to tell these

---

\* P. Chin is supported by NSF-DMS-1222567 from NSF and FA9550-12-1-0136 from the Air Force Office of Scientific Research. A. Rao is supported by NSF grant CCF-1111257. V. Vu is supported by research grants DMS-0901216 and AFOSAR-FA-9550-09-1-0167 and a grant from Draper Lab.

vertices apart and so a complete recovery is out of question. The goal here is to recover a large portion of each block, namely finding a partition  $V'_1 \cup V'_2$  of  $V = V_1 \cup V_2$  such that  $V_i$  and  $V'_i$  are close to each other. For quantitative purposes, let us introduce a definition

**Definition 1** A collection of subsets  $V'_1, V'_2$  of  $V_1 \cup V_2$  is  $\gamma$ -correct if  $|V_i \cap V'_i| \geq (1 - \gamma)n$ .

In [Coja-Oghlan \(2010\)](#), Coja-Oghlan proved

**Theorem 2** For any constant  $\gamma > 0$  there are constants  $d_0, C > 0$  such that if  $a, b > d_0$  and  $\frac{(a-b)^2}{a+b} > C \log(a+b)$ , one can find a  $\gamma$ -correct partition using a polynomial time algorithm.

Coja-Oghlan proved [Theorem 2](#) as part of a more general problem, and his algorithm was rather involved. Furthermore, the result is not yet sharp and it has been conjectured that the log term is removable<sup>1</sup>. Even when the log term is removed, an important question is to find out the optimal relation between the accuracy  $\gamma$  and the ratio  $\frac{(a-b)^2}{a+b}$ . This is the main goal of this paper.

**Theorem 3** There are constants  $C_0$  and  $C_1$  such that the following holds. For any constants  $a > b > C_0$  and  $\gamma > 0$  satisfying

$$\frac{(a-b)^2}{a+b} \geq C_1 \log \frac{1}{\gamma},$$

we can find a  $\gamma$ -correct partition with probability  $1 - o(1)$  using a simple spectral algorithm.

The constants  $C_0, C_1$  can be computed explicitly via a careful, but rather tedious, book keeping. We try not to optimize these constants to simplify the presentation. The proof of [Theorem 3](#) yields the following corollary

**Corollary 4** There are constants  $C_0$  and  $\epsilon$  such that the following holds. For any constants  $a > b > C_0$  and  $\epsilon > \gamma > 0$  satisfying

$$\frac{(a-b)^2}{a+b} \geq 8.1 \log \frac{2}{\gamma},$$

we can find a  $\gamma$ -correct partition with probability  $1 - o(1)$  using a simple spectral algorithm.

In parallel to our study, [Zhang and Zhou](#), proving a minimax rate result that suggested that there is a constant  $c > 0$

$$\frac{(a-b)^2}{a+b} \leq c \log \frac{1}{\gamma}$$

then one *cannot* recover a  $\gamma$ -correct partition (in expectation), regardless the algorithm.

In order to prove [Theorem 3](#), we design a fast and robust algorithm which obtains a  $\gamma$ -correct partition under the condition  $\frac{(a-b)^2}{a+b} \geq C \log \frac{1}{\gamma}$ . Our algorithm guarantees  $\gamma$ -correctness with high probability.

We can refine the algorithm to handle the (more difficult) general case of having  $k$  blocks, for any fixed number  $k$ . Suppose now there are  $k$  blocks  $V_1, \dots, V_k$  with  $|V_i| = \frac{n}{k}$  with edge probabilities  $\frac{a}{n}$  between vertices within the same block and  $\frac{b}{n}$  between vertices in different blocks. As before, a collection of subsets  $V'_1, V'_2, \dots, V'_k$  of  $V_1 \cup V_2 \cup \dots \cup V_k$  is  $\gamma$ -correct if  $|V_i \cap V'_i| \geq (1 - \gamma)\frac{n}{k}$ .

---

1. We would like to thank E. Abbe for communicating this conjecture.

**Theorem 5** *There exists constants  $C_1, C_2$ , such that if  $k$  is any constant as  $n \rightarrow \infty$  and if*

1.  $a > b \geq C_1$
2.  $(a - b)^2 \geq C_2 k^2 a \log \frac{1}{\gamma}$ ,

*then we can find a  $\gamma$ -correct partition with probability at least  $1 - o(1)$  using a simple spectral algorithm.*

We believe that this result is sharp, up to the values of  $C_1$  and  $C_2$ ; in particular, the requirement  $\frac{(a-b)^2}{a} = \Omega(k^2)$  is optimal.

Our method also works (without significant changes) in the case the blocks are not equal, but have comparable sizes (say  $cn \geq |V_i| \geq n$  for some constant  $c \geq 1$ ). In this case, the constants  $C_1, C_2$  above will also depend on  $c$ . While the emphasis of this paper is on the case  $a, b$  are constants, we would like to point out that this assumption is not required in our theorems, so our algorithms work on denser graphs as well.

Let us now discuss some recent works, which we just learned after posting the first version of this paper on arxiv. Mossel informed us about a recent result in [Mossel et al. \(2013b\)](#) which is similar to Theorem 3 (see [Mossel et al., 2013b](#), Theorem 5.3). They give a polynomial time algorithm and prove that there exists a constant  $C$  such that if  $(a - b)^2 > C(a + b)$  and  $a, b$  are fixed as  $n \rightarrow \infty$ , then the algorithm recovers an optimal fraction of the vertices. The algorithm in [Mossel et al. \(2013b\)](#) is very different from ours, and uses non-back tracking walks. This algorithm does not yet handle the case of more than 2 blocks, and its analysis looks very delicate. Next, Guedon sent us [Guédon and Vershynin \(2014\)](#), in which the authors also proved a result similar to Theorem 3, under a stronger assumption  $(a - b)^2 \geq C \frac{1}{\gamma^2} (a + b)$  (see Theorem 1.1 and Corollary 1.2 of [Guédon and Vershynin \(2014\)](#)). Their approach relies on an entirely different (semi-definite program) algorithm, which, in turn, was based on Grothendick's inequality. This approach seems to extend to the general  $k > 2$  case; however, the formulation of the result in this case, using matrix approximation, is somewhat different from ours (see [Guédon and Vershynin, 2014](#), Theorem 1.3)). Two more closely related papers have been brought to our attention by the reviewers. In [Lelarge et al. \(2013\)](#), authors have worked out the spectral part of the result in this paper. Also, in a paper posted less than a month prior to this paper, authors of [Yun and Proutiere \(2014\)](#) have also achieved a  $\log 1/\gamma$  dependence using similar techniques.

It is remarkable to see so many progresses, using different approaches, on the same problem in such a short span of time. This suggests that the problem is indeed important and rich, and it will be really pedagogical to study the performance of the existing algorithms in practice. We are going to discuss the performance of our algorithm in Sections 3 and 4.

We next present an application of our method to the *Censor Block Model* studied by Abbe et. al. in [Abbe et al. \(2014\)](#). As before, let  $V$  be the union of two blocks  $V_1, V_2$ , each of size  $n$ . Let  $G = (V, E)$  be a random graph with edge probability  $p$  with incidence matrix  $B_G$  and  $\mathbf{x} = (x_1, \dots, x_{2n})$  be the indicator vector of  $V_2$ . Let  $\mathbf{z}$  be a random noise vector whose coordinates  $z_{e_i}$  are i.i.d Bernoulli( $\epsilon$ ) (taking value 1 with probability  $\epsilon$  and 0 otherwise), where  $e_i$  are the edges of  $G$ .

Given a noisy observation

$$\mathbf{y} = B_G \mathbf{x} \oplus \mathbf{z}$$

where  $\oplus$  is the addition in mod 2, one would like identify the blocks. In [Abbe et al. \(2014\)](#), the authors proved that exact recovery ( $\gamma = 0$ ) is possible if and only if  $\frac{np}{\log n} \geq \frac{2}{(1-2\epsilon)^2} + o(\frac{1}{(1-2\epsilon)^2})$  in the limit  $\epsilon \rightarrow 1/2$ . Further, they gave a semidefinite programming based algorithm which succeeds up to twice the threshold. They posed the question of partial recovery ( $\gamma > 0$ ) for sparse graphs. Addressing this question, we show

**Theorem 6** *For any given constants  $\gamma, 1/2 > \epsilon > 0$ , there exists constant  $C_1, C_2$  such that if  $np \geq \frac{C_1}{(1-2\epsilon)^2}$  and  $p \geq \frac{C_2}{n}$ , then we can find a  $\gamma$ -correct partition with probability  $1 - o(1)$ , using a simple spectral algorithm.*

Let us conclude this section by mentioning a related, interesting, problem, where the purpose is just to do better than a random guess (in our terminology, to find a partition which is  $(1/2 + \epsilon)$ -correct). It was conjectured in [Decelle et al. \(2011\)](#) that this is possible if and only if  $(a - b)^2 > (a + b)$ . This conjecture has been settled recently by Mossel et. al. [Mossel et al. \(2012\)](#) [Mossel et al. \(2013a\)](#) and Massoulié [Massoulié \(2013\)](#). Another closely related problem which has been studied in [Abbe et al. \(2014\)](#) [Mossel et al. \(2014\)](#) is about when one can recover at least  $1 - o(1)$  fraction of the vertices.

The rest of the paper is organized as follows. In section 2, we describe our algorithm for Theorem 7 and an overview of the proof. The full proof comes in sections 3. In section 4, we show how to modify the algorithm to handle the  $k$  block case and prove theorem 5. Finally, in section 5, we prove theorem 6.

## 2. Two communitites

We first consider the case  $k = 2$ . Our algorithm will have two steps. First we use a spectral algorithm to recover a partition where the dependence between  $\gamma$  and  $\frac{(a-b)^2}{a+b}$  is sub-optimal.

Let  $A_0$  denote the adjacency matrix of a random graph generated from the distribution as in Theorem 7. Let  $\bar{A}_0 \stackrel{\text{def}}{=} \mathbb{E}A_0$  and  $E_0 \stackrel{\text{def}}{=} A_0 - \bar{A}_0$ . Then  $\bar{A}_0$  is a rank two matrix with the two non zero eigenvalues  $\lambda_1 = a + b$  and  $\lambda_2 = a - b$ . The eigenvector  $\mathbf{u}_1$  corresponding to the eigenvalue  $a + b$  has coordinates

$$\mathbf{u}_1(i) = \frac{1}{\sqrt{2n}}, \text{ for all } i \in V$$

and eigenvector  $\mathbf{u}_2$  corresponding to the eigenvalue  $a - b$  has coordinates

$$\mathbf{u}_2(i) = \begin{cases} \frac{1}{\sqrt{2n}} & \text{if } i \in V_1 \\ \frac{-1}{\sqrt{2n}} & \text{if } i \in V_2. \end{cases}$$

Notice that the second eigenvector of  $\bar{A}_0$  identifies the partition. We would like to use the second eigenvector of  $A_0$  to approximately identify the partition. Since  $A_0 = \bar{A}_0 + E_0$ , perturbation theory tells us that we get a good approximation if  $\|E_0\|$  is sufficiently small. However, with probability  $1 - o(1)$ , the norm of  $E_0$  is rather large (even larger than the norm of the main term). In order to handle this problem, we modify  $E_0$  using the auxiliary deletion, at the cost of losing a few large degree vertices.

**Spectral Partition.**

1. Input the adjacency matrix  $A_0$ ,  $d := a + b$ .
2. Zero out all the rows and columns of  $A_0$  corresponding to vertices whose degree is bigger than  $20d$ , to obtain the matrix  $A$ .
3. Find the eigenspace  $W$  corresponding to the top two eigenvalues of  $A$ .
4. Compute  $v_1$ , the projection of all-ones vector on to  $W$ .
5. Let  $v_2$  be the unit vector in  $W$  perpendicular to  $v_1$ .
6. Sort the vertices according to their values in  $v_2$ , and let  $V'_1 \subset V$  be the top  $n$  vertices, and  $V'_2 \subset V$  be the remaining  $n$  vertices.
7. Output  $(V'_1, V'_2)$ .

Figure 1: Spectral Partition

Let  $\bar{A}$ ,  $A$ ,  $E$  be the matrices obtained from  $\bar{A}_0$ ,  $A_0$ ,  $E_0$  after the deletion, respectively. Let  $\Delta \stackrel{\text{def}}{=} \bar{A} - \bar{A}_0$ ; we have

$$\begin{aligned} A &= \bar{A} + E \\ &= \bar{A}_0 + \Delta + E. \end{aligned}$$

The key observation is that  $\|E\|$  is significantly smaller than  $\|E_0\|$ . In the next section we will show that  $\|E\| = \mathcal{O}(\sqrt{d})$ , with probability  $1 - o(1)$ , while  $\|E_0\|$  is  $\Theta(\sqrt{\frac{\log n}{\log \log n}})$ , with probability  $1 - o(1)$ . Furthermore, we could show that  $\|\Delta\|$  is only  $\mathcal{O}(1)$  with probability  $1 - o(1)$ . Therefore, if the second eigenvalue gap for the matrix  $A_0$  is greater than  $C\sqrt{d}$ , for some large enough constant  $C$ , then Davis-Kahan sin  $\Theta$  theorem would allow us to bound the angle between the second eigenvector of  $\bar{A}_0$  and  $A$  by an arbitrarily small constant. This will, in turn, enable us to recover a large portion of the blocks, proving the following statement

**Theorem 7** *There are constants  $C_0$  and  $C_1$  such that the following holds. For any constants  $a > b > C_0$  and  $\gamma > 0$  satisfying  $\frac{(a-b)^2}{a+b} \geq C_1 \frac{1}{\gamma^2}$ , then with probability  $1 - o(1)$ , **Spectral Partition** outputs a  $\gamma$ -correct partition.*

**Remark 8** *The parameter  $d := a + b$  can be estimated very efficiently from the adjacency matrix  $A$ . We take this as input for a simpler exposition.*

Step 2 is a further correction that gives us the optimal (logarithmic) dependence between  $\gamma$  and  $\frac{(a-b)^2}{a+b}$ . The idea here is to use the degree sequence to correct the mislabeled vertices. Consider a mislabeled vertex  $u \in V'_1 \cap V_2$ . As  $u \in V_2$ , we expect  $u$  to have  $b$  neighbors in  $V_1$  and  $a$  neighbors in  $V_2$ . Assume that **Spectral Partition** output  $V'_1, V'_2$  where  $|V_1 \setminus V'_1| \leq 0.1n$ , we expect  $u$  to have at most  $0.9b + 0.1a$  neighbors in  $V'_1$  and at least  $0.1b + 0.9a$  neighbors in  $V'_2$ . As

**Partition**

1. Input the adjacency matrix  $A_0, d := a + b$ .
2. Randomly color the edges with Red and Blue with equal probability.
3. Run **Spectral Partition** on Red graph, outputting  $V'_1, V'_2$ .
4. Run **Correction** on the Blue graph.
5. Output the corrected sets  $V'_1, V'_2$ .

Figure 2: Partition

$0.1b + 0.9a > \frac{a+b}{2} > 0.9b + 0.1a$ , we can correctly reclassify  $u$  by thresholding. There are, however, few problems with this argument. First, everything is in expectation. This turns out to be a minor problem; we can use a large deviation result to show that a majority of mislabeled vertices can be detected this way. As a matter of fact, the desired logarithmic dependence is achieved at this step, thanks to the exponential probability bound in the large deviation result.

The more serious problem is the lack of independence. Once **Spectral Partition** has run, the neighbors of  $u$  are no longer random. We can avoid this problem using a splitting trick as given in **Partition**. We sample randomly half of the edges of the input graph and used the graph formed by them in **Spectral Partition**. After receiving the first partition, we use the other (random) half of the edges for correction. This doesn't make the two steps completely independent, but we can still prove the stated result.

The sub-routine **Correction** is as follows:

**Correction.**

1. Input: a partition  $V'_1, V'_2$  and a Blue graph on  $V'_1 \cup V'_2$ .
2. For any  $u \in V'_1$ , label  $u$  *bad* if the number of neighbors of  $u$  in  $V'_2$  is at least  $\frac{a+b}{4}$  and *good* otherwise.
3. Do the same for any  $v \in V'_2$ .
4. Correct  $V'_i$  by deleting its bad vertices and adding the bad vertices from  $V'_{3-i}$ .

Figure 3: Correction

Figure 4 is the density plot of the matrix before and after clustering according to the algorithm described above. We can prove

**Lemma 9** *Given a 0.1-correct partition  $V'_1, V'_2$  and a Blue graph on  $V'_1 \cup V'_2$  as input to the sub-routine **Correction** given in figure 3, we get a  $\gamma$ -correct partition with  $\gamma = 2 \exp(-0.072 \frac{(a-b)^2}{a+b})$ .*




Figure 4: On the left is the density plot of the input (unclustered) matrix with parameters  $n = 7500, a = 10, b = 3$  and on the right is the density plot of the permuted matrix after running the algorithm described above. This took less than 3secs in Matlab running on a 2009 MacPro.

### 3. First step: Proof of Theorem 7

We now turn to the details of the proof. Using the notation in the previous section, we let  $W$  be the two dimensional eigenspace corresponding to the top two eigenvalues of  $A$  and  $\bar{W}$  be the corresponding space of  $\bar{A}$ . For any two vector subspaces  $W_1, W_2$  of same dimension, we use the usual convention  $\sin \angle(W_1, W_2) := \|P_{W_1} - P_{W_2}\|$ , where  $P_{W_i}$  is the orthogonal projection onto  $W_i$ . The proof has two main steps:

1. *Bounding the angle:* We show that  $\sin \angle(W, \bar{W})$  is small, under the conditions of the theorem.
2. *Recovering the partition:* If  $\sin \angle(W, \bar{W})$  is small, we find an approximate partition which can then improved to find an optimal one.

#### 3.1. Bounding the angle

For the first part, recall that  $A = \bar{A}_0 + \Delta + E$ . We first prove that  $\|\Delta\|$  and  $\|E\|$  are small with probability  $1 - o(1)$ . Bounding  $\|\Delta\|$  is easy as it will be sufficient to bound the number of vertices of high degrees. We need the following

**Lemma 10** *There exist a constant  $d_0$  such that if  $d := a + b \geq d_0$ , then with probability  $1 - \exp(-\Omega(a^{-2}n))$  not more than  $a^{-3}n$  vertices have degree  $\geq 20d$ .*

Note that the proof of the above lemma and other missing proofs in this subsection appear in appendix A.1. If there are at most  $a^{-3}n$  vertices with degree  $\geq 20d$ , then by definition,  $\Delta$  has at most  $2a^{-3}n^2$  non-zero entries, and the magnitude of each entry is bounded by  $\frac{a}{n}$ . Therefore, its Hilbert-Schmidt norm is bounded by  $\|\Delta\|_{HS} \leq \sqrt{2}a^{-1/2}$ .

**Corollary 11** For  $d_0$  sufficiently large, with probability  $1 - \exp(-\Omega(a^{-3}n))$ ,  $\|\Delta\| \leq 1$ .

Now we address the harder task of bounding  $\|E\|$ . Here is the key lemma

**Lemma 12** Suppose  $M$  is random symmetric matrix with zero on the diagonal whose entries above the diagonal are independent with the following distribution

$$M_{ij} = \begin{cases} 1 - p_{ij} & \text{w.p. } p_{ij} \\ -p_{ij} & \text{w.p. } 1 - p_{ij} \end{cases}.$$

Let  $\sigma$  be a quantity such that  $p_{ij} \leq \sigma^2$  and  $M_1$  be the matrix obtained from  $M$  by zeroing out all the rows and columns having more than  $20\sigma^2 n$  positive entries. Then with probability  $1 - o(1)$ ,  $\|M_1\| \leq C\sigma\sqrt{n}$  for some constant  $C > 0$ .

Lemma 12 implies

**Corollary 13** There exist constants  $C_0, C$  such that if  $a > b \geq C_0$ , and  $E$  is obtained as described before, then we have,

$$\|E\| \leq C\sqrt{d}$$

with probability  $1 - o(1)$ .

Now, let  $\bar{v}_1, \bar{v}_2$  be eigenvectors of  $\bar{A}_0$  corresponding to the largest two eigenvalues  $\lambda_1 \geq \lambda_2$   $v_1, v_2$  be eigenvectors of  $A = \bar{A}_0 + \Delta + E$  corresponding to the largest two eigenvalues. Further,  $\bar{W} := \text{Span}\{\bar{v}_1, \bar{v}_2\}$  and  $W := \text{Span}\{v_1, v_2\}$ .

**Lemma 14** For any constant  $c < 1$ , we can choose constants  $C_2$  and  $C_3$  such that such that if  $a - b \geq C_2\sqrt{a+b} = C_2\sqrt{d}$  and  $a \geq C_3$  then,  $\sin(\angle \bar{W}, W) \leq c < 1$  with probability  $1 - o(1)$ .

**Proof of Lemma 14:** Let  $C_3$  be a constant such that if  $a \geq C_3$ , then theorem 11 holds giving us  $\|\Delta\| \leq 1$ . From lemma 13 we have that  $\|E\| \leq C\sqrt{d}$ . The lemma then follows from the Davis-Kahan Davis (1963) Bhatia (1997) bound for matrices  $\bar{A}_0$  and  $A$ , which gives  $\sin(\angle W, \bar{W}) \leq \frac{\|E+\Delta\|}{\lambda_2}$ . Therefore, the lemma follows by choosing  $C_1$  big enough.

### 3.2. Recovery

Given a subspace  $W$  satisfying  $\sin(\angle \bar{W}, W) \leq c < 1/16$ , we can recover a big portion of the vertices. We prove (in appendix A.2) that

**Lemma 15** Given a subspace  $W$  satisfying  $\sin(\angle \bar{W}, W) \leq c < 1/16$ , we can recover a  $8c/3$ -correct partition.

Once we have an approximate partition, we can use the Blue edges to boost it in the **Correction** step. We prove (in appendix A.3)

**Lemma 16** Given a 0.1 correct partition  $V'_1, V'_2$  as input to the **Correction** routine in figure 3, the algorithm outputs a  $\gamma$  correction partition with  $\gamma = 2 \exp(-0.072 \frac{(a-b)^2}{a+b})$ .



## 4. Multiple communities

### 4.1. Overview

Let us start with the algorithm, which (compared to the algorithm for the case of 2 blocks) has an additional step of random splitting. This additional step is needed in order to recover the partitions. We will start by computing an approximation of the space spanned by the first  $k$  eigenvectors of the hidden matrix. However, when  $k > 2$ , it is not obvious how to approximate the eigenvectors themselves. To handle this problem, we need a new argument that requires this extra step.




Figure 5: Partition

Since we use different set of edges for each step, we have independence across the steps.

### 4.2. Details

Step 1 is a spectral algorithm on a portion of the adjacency matrix  $A_0$  as given in figure 6. This will enable us to recover a large portion of the blocks  $Z \cap V_1, \dots, Z \cap V_k$ . We will prove the following statement (appendix B.1)

**Theorem 17** *There exists constants  $C_1, C_2$  such that for any fixed integer  $k$  the following holds.*

1.  $a > b \geq C_1$
2.  $\frac{(a-b)^2}{a} \geq C_2 k^2 \frac{1}{\gamma}$ , and

*then we can find a  $\gamma$ -correct partition  $U'_1, \dots, U'_k$  of  $Z$  with high probability using a simple spectral algorithm.*

**Spectral Partition.**

1. Input  $B$  (a matrix of dimension  $|Z| \times |Y|$ ),  $a$ ,  $b$  and  $k$ .
2. Let  $Y_1$  be a random subset of  $Y$  by selecting each element with probability  $\frac{1}{2}$  independently and let  $A_1, A_2$  be the sub matrix of  $B$  formed by the columns indexed by  $Y_1, Y_2 := Y \setminus Y_1$ , respectively.
3. Let  $d := a + (k - 1)b$ . Zero out all the rows and columns of  $A_1$  corresponding to vertices whose degree is bigger than  $20d$ , to obtain the matrix  $A$ .
4. Find the space spanned by  $k$  left singular vectors of  $A$ , say  $W$
5. Let  $\mathbf{a}_1, \dots, \mathbf{a}_m$  be some  $m = 2 \log n$  random columns of  $A_2$ . For each  $i$ , project  $\mathbf{a}_i - \mathbf{a}$  onto  $W$ , where  $\mathbf{a}(j) = \frac{a+b}{2n}$  for all  $j$  is a constant vector.
6. For each projected vector, identify the top (in value)  $n/2k$  coordinates. Of the  $2 \log n$  sets so obtained, discard half of the sets with the lowest Blue edge density in them.
7. Of the remaining subsets, identify some  $k$  subsets  $U'_1, \dots, U'_k$  such that  $|U'_i \cap U'_j| < 0.2n/2k$ , for  $i \neq j$ .
8. Output  $U'_1, \dots, U'_k$ .

Figure 6: Spectral Partition

Step 2 (figure 7) is a further correction that gives us the optimal (logarithmic) dependence between  $\gamma$  and  $\frac{(a-b)^2}{a+b}$ . The idea here is to use the degree sequence to correct the mislabeled vertices in  $Z$ . Consider a mislabeled vertex  $u \in Z \cap V_1$ . As  $u \in Z \cap V_1$ , we expect  $u$  to have  $a/4$  Red neighbors in  $Z \cap V_1$  and  $b/4$  Red neighbors in  $Z \cap V_i$  for all  $i \neq 1$ . Assume that **Spectral Partition** output  $U'_1, \dots, U'_k$  where  $|U_1 \setminus U'_1| \leq .1n/2k$ , we expect  $u$  to have at most  $0.9b/4k + 0.1a/4k$  Red neighbors in  $U'_i$  and at least  $0.1b/4k + 0.9a/4k$  Red neighbors in  $U'_1$ . As

$$0.1b/8k + 0.9a/8k > \frac{a+b}{8k} > 0.9b/8k + 0.1a/8k$$

we can correctly reclassify  $u$  by thresholding. We can prove (appendix B.2)

**Lemma 18** *Given a 0.1 correction partition of  $Z = (Z \cap V_1) \cup \dots \cup (Z \cap V_k)$  and the Red graph over  $Z$ , the sub-routine **Correction** given in figure 7 computes a  $\gamma$  correct partition with  $\gamma = 2k \exp(-0.04 \frac{(a-b)^2}{k(a+b)})$ .*

Step 3 is to use the clustering information of vertices in  $Z$  to label the vertices in  $Y$ , and is similar to step 2. We prove (appendix B.3)

**Lemma 19** *Given a 0.1 correction partition of  $Z = (Z \cap V_1) \cup \dots \cup (Z \cap V_k)$  and the Blue graph over  $Y \cup Z$ , the sub-routine **Merge** is given in (figure 8) computes a  $\gamma$  correct partition with  $\gamma = 2k \exp(-0.0324 \frac{(a-b)^2}{k(a+b)})$ .*

Combining lemmas 18, 19, we get the stated result.

**Correction.**

1. Input: A collection of subsets  $U'_1, \dots, U'_k \subset Z$  and a graph on  $Z$ .
2. For every  $u \in Z$ , if  $i \in \{1, 2, \dots, k\}$  is such that  $u$  has maximum neighbors in  $U'_i$ , then add  $u$  to  $U''_i$ . Break ties arbitrarily.
3. Output  $U''_1, \dots, U''_k$ .

Figure 7: Correction

**Merging.**

1. Input: A partition  $U'_1, \dots, U'_k$  of  $(Z \cap V_1) \cup (Z \cap V_2) \cup \dots \cup (Z \cap V_k)$  and a graph between vertices  $Y$  and  $Z$ .
2. For all  $u \in Y$ , label  $u$  with ‘i’ if the number of neighbors of  $u$  in  $U'_i$  is at least  $\frac{a+b}{8}$ . Label the conflicts arbitrarily.
3. Output the label classes as the clusters  $V'_1, \dots, V'_k$ .

Figure 8: Merge

## 5. Censor Block Model

We first introduce some notations so as to write this problem in a way similar to the other problems in this paper. To simplify the analysis, we make the following assumptions. We assume that there are  $|V| = 2n$  vertices, with exactly  $n$  of them labeled 1, and the rest labeled 0. As in [Abbe et al. \(2014\)](#), we assume that  $G \in G_{2n,p}$  is a graph generated from the Erdos-Renyi model with edge probability  $p$ . Since any edge  $(i, j)$  appears with probability  $p$ , and that  $z_e \sim \text{Bernoulli}(\epsilon)$ , we have

$$y_{i,j} = \begin{cases} x_i \oplus x_j & w.p. \quad p(1 - \epsilon) \\ x_i \oplus x_j \oplus 1 & w.p. \quad p\epsilon \\ 0 & w.p. \quad 1 - p \end{cases}.$$

For any  $i, j \in V$ , let us write  $w_{ij} := x_i \oplus x_j$ , and  $W := (w_{ij})_{ij}$  the associated  $2n \times 2n$  matrix.

We note that  $\bar{y}_{i,j} := \mathbf{E}(y_{i,j}) = p\epsilon + p(1 - 2\epsilon)w_{i,j}$ . Therefore, we can write  $y_{i,j} = \bar{y}_{i,j} + \zeta_{i,j}$ , where  $\zeta_{i,j}$ s are mean zero random variables satisfying  $\text{Var}(\zeta_{i,j}) \leq p$ . First we note that we can recover the two communities from the eigenvectors of the  $2n \times 2n$  matrix  $\bar{Y} := (\bar{y}_{i,j}) = p\epsilon I + p(1 - 2\epsilon)W$ .  $\bar{Y}$  is a rank 2 matrix with eigenvalues  $pn$  and  $p(1 - 2\epsilon)n$ , with the corresponding eigenvectors  $\mathbf{v}_1 = (1, 1, \dots, 1)$  and  $\mathbf{v}_2 = (1, \dots, 1, -1, \dots, -1)$ . if we can find  $\mathbf{v}_2$ , we can identify the two blocks. Let  $Y = (y_{i,j})$  and  $E = (\zeta_{i,j})$  be  $2n \times 2n$  matrices. Algorithm 10 (which is essentially same as algorithm 2) which takes as input the adjacency matrix  $Y$  and the edge probability  $p$  achieves this when  $np \geq \frac{C_2}{(1-2\epsilon)^2}$ . More detail appears in appendix C.




Figure 9: On the left is the density plot of the input (unclustered) matrix with parameters  $n = 3000$ ,  $a = 22$ ,  $b = 2$  and on the right is the density plot of the permuted matrix after running the algorithm described above. This took less than 1sec in Matlab running on a 2009 MacPro.

### Spectral Partition II.

1. Input the adjacency matrix  $Y$ ,  $p$ .
2. Zero out all the rows and columns of  $Y$  corresponding to vertices whose degree is bigger than  $20pn$ , to obtain the matrix  $Y_0$ .
3. Find the eigenspace  $U$  corresponding to the top two eigenvalues of  $Y_0$ .
4. Compute  $v_1$ , the projection of all-ones vector on to  $U$
5. Let  $v_2$  be the unit vector in  $W$  perpendicular to  $v_1$ .
6. Sort the vertices according to their values in  $v_2$ , and let  $V'_1 \subset V$  be the top  $n$  vertices, and  $V'_2 \subset V$  be the remaining  $n$  vertices
7. Output  $(V'_1, V'_2)$ .

Figure 10: Algorithm 3

### References

- E. Abbe, A. S. Bandeira, and G. Hall. Exact Recovery in the Stochastic Block Model. *ArXiv e-prints*, 1405.3267, May 2014.
- Emmanuel Abbe, Afonso S. Bandeira, Annina Bracher, and Amit Singer. Decoding binary node labels from censored edge measurements: Phase transition and efficient recovery. *CoRR*, abs/1404.4749, 2014.

- N. Alon and J.H. Spencer. *The Probabilistic Method*. Wiley-Interscience series in discrete mathematics and optimization. Wiley, 2004. ISBN 9780471653981. URL <http://books.google.com/books?id=q3lUjheWiMoC>.
- Noga Alon and Nabil Kahale. A spectral technique for coloring random 3-colorable graphs. In *SIAM Journal on Computing*, pages 346–355, 1994.
- Rajendra Bhatia. *Matrix Analysis*, volume 169. Springer, 1997. ISBN 0387948465.
- B. Bollobas. *Random Graphs*. Cambridge University Press, 2001.
- Ravi B. Boppana. Eigenvalues and graph bisection: An average-case analysis. In *Proceedings of the 28th Annual Symposium on Foundations of Computer Science*, SFCS '87, pages 280–285, Washington, DC, USA, 1987. IEEE Computer Society. ISBN 0-8186-0807-2. doi: 10.1109/SFCS.1987.22. URL <http://dx.doi.org/10.1109/SFCS.1987.22>.
- S. Boucheron, G. Lugosi, and P. Massart. Concentration inequalities. A nonasymptotic theory of independence, 2013. URL [halv3-preprod.archives-ouvertes.fr/hal-00794821](http://halv3-preprod.archives-ouvertes.fr/hal-00794821).
- T.N. Bui, S. Chaudhuri, F.T. Leighton, and M. Sipser. Graph bisection algorithms with good average case behavior. *Combinatorica*, 7(2):171–191, 1987. ISSN 0209-9683. doi: 10.1007/BF02579448. URL <http://dx.doi.org/10.1007/BF02579448>.
- Amin Coja-Oghlan. Graph partitioning via adaptive spectral techniques. *Combinatorics, Probability and Computing*, 19:227–284, 3 2010. ISSN 1469-2163. doi: 10.1017/S0963548309990514. URL [http://journals.cambridge.org/article\\_S0963548309990514](http://journals.cambridge.org/article_S0963548309990514).
- Chandler Davis. The rotation of eigenvectors by a perturbation. *Journal of Mathematical Analysis and Applications*, 6(2):159 – 173, 1963. ISSN 0022-247X. doi: [http://dx.doi.org/10.1016/0022-247X\(63\)90001-5](http://dx.doi.org/10.1016/0022-247X(63)90001-5). URL <http://www.sciencedirect.com/science/article/pii/0022247X63900015>.
- Aurelien Decelle, Florent Krzakala, Cristopher Moore, and Lenka Zdeborová. Asymptotic analysis of the stochastic block model for modular networks and its algorithmic applications. *Phys. Rev. E*, 84:066106, Dec 2011. doi: 10.1103/PhysRevE.84.066106. URL <http://link.aps.org/doi/10.1103/PhysRevE.84.066106>.
- M. E. Dyer and A. M. Frieze. The solution of some random np-hard problems in polynomial expected time. *J. Algorithms*, 10(4):451–489, December 1989. ISSN 0196-6774. doi: 10.1016/0196-6774(89)90001-1. URL [http://dx.doi.org/10.1016/0196-6774\(89\)90001-1](http://dx.doi.org/10.1016/0196-6774(89)90001-1).
- Uriel Feige and Eran Ofek. Spectral techniques applied to sparse random graphs. *Random Structures & Algorithms*, 27(2):251–275, 2005. ISSN 1098-2418. doi: 10.1002/rsa.20089. URL <http://dx.doi.org/10.1002/rsa.20089>.
- J. Friedman, J. Kahn, and E. Szemerédi. On the second eigenvalue of random regular graphs. In *Proceedings of the Twenty-first Annual ACM Symposium on Theory of Computing*, STOC '89, pages 587–598, New York, NY, USA, 1989. ACM. ISBN 0-89791-307-8. doi: 10.1145/73007.73063. URL <http://doi.acm.org/10.1145/73007.73063>.

- O. Guédon and R. Vershynin. Community detection in sparse networks via Grothendieck’s inequality. *ArXiv e-prints*, 1411.4686, November 2014.
- Mark Jerrum and G.B. Sorkin. Simulated annealing for graph bisection. In *Foundations of Computer Science, 1993. Proceedings., 34th Annual Symposium on*, pages 94–103, Nov 1993. doi: 10.1109/SFCS.1993.366878.
- M. Lelarge, L. Massoulié, and Jiaming Xu. Reconstruction in the labeled stochastic block model. In *Information Theory Workshop (ITW), 2013 IEEE*, pages 1–5, Sept 2013. doi: 10.1109/ITW.2013.6691264.
- Laurent Massoulié. Community detection thresholds and the weak ramanujan property. *CoRR*, abs/1311.3085, 2013.
- F. McSherry. Spectral partitioning of random graphs. In *Foundations of Computer Science, 2001. Proceedings. 42nd IEEE Symposium on*, pages 529–537, Oct 2001. doi: 10.1109/SFCS.2001.959929.
- E. Mossel, J. Neeman, and A. Sly. Stochastic Block Models and Reconstruction. *ArXiv e-prints*, 1202.1499, February 2012.
- E. Mossel, J. Neeman, and A. Sly. A Proof Of The Block Model Threshold Conjecture. *ArXiv e-prints*, 1311.4115, November 2013a.
- E. Mossel, J. Neeman, and A. Sly. Belief Propagation, Robust Reconstruction, and Optimal Recovery of Block Models. *ArXiv e-prints*, 1309.1380, September 2013b.
- E. Mossel, J. Neeman, and A. Sly. Consistency Thresholds for Binary Symmetric Block Models. *ArXiv e-prints*, 1407.1591, July 2014.
- Joel A. Tropp. User-friendly tail bounds for sums of random matrices. *Foundations of Computational Mathematics*, 12(4):389–434, 2012. ISSN 1615-3375. doi: 10.1007/s10208-011-9099-z. URL <http://dx.doi.org/10.1007/s10208-011-9099-z>.
- V. Vu. A simple SVD algorithm for finding hidden partitions. *ArXiv e-prints*, 1404.3918, April 2014.
- Se-Young Yun and Alexandre Proutiere. Accurate community detection in the stochastic block model via spectral algorithms. *CoRR*, abs/1412.7335, 2014. URL <http://arxiv.org/abs/1412.7335>.
- Y. Zhang and H. Zhou. Minimax rates of community detection in stochastic block model.

## Appendix A. Two communities

### A.1. Bounding $\|\Delta\|$ and $\|E\|$

**Proof of Lemma 10:** One can prove Lemma 10 using a standard argument from random graph theory. Consider a set of vertices  $X \subset V$  of size  $|X| = cn$ , where  $c < 1$  is a constant. We first bound the probability that all the vertices in this set have degree greater than  $20d$ .

Let us denote the set of edges on  $X$  by  $E(X)$  and the set of edges with exactly one end point in  $X$  by  $E(X, X^c)$ . If each degree in  $X$  is at least  $20d$ , then a quick consideration reveals that either  $|E(X)| \geq 2cnd$  or  $|E(X, X^c)| \geq 8cnd$ . The expected number of edges  $\mu_{E(X)} := \mathbf{E}(|E(X)|)$  satisfies

$$0.25(cn)^2 \frac{a}{n} \leq \mu_{E(X)} \leq 0.5(cn)^2 \frac{a}{n}.$$

Let  $\delta_1 := \frac{2}{c} \leq \frac{2cnd}{\mu_{E(X)}}$ , then Chernoff bound (see [Alon and Spencer \(2004\)](#) for example) gives

$$\begin{aligned} \mathbb{P}(|E(X)| \geq cnd) &\leq \left( \frac{\exp(\delta_1 - 1)}{\delta_1^{\delta_1}} \right)^{\mu_{E(X)}} \\ &\leq \exp \left( \left( \frac{2}{c} - 1 - \frac{2}{c} \log \left( \frac{2}{c} \right) \right) 0.25(cn)^2 \frac{a}{n} \right) \\ &\leq \exp \left( -\frac{1}{c} \log \left( \frac{1}{c} \right) 0.25(cn)^2 \frac{a}{n} \right) \text{ (for small enough } c) \\ &= \exp \left( -0.25 \log \left( \frac{1}{c} \right) acn \right). \end{aligned}$$

Similarly, the expected number of edges  $\mu_{E(X, X^c)}$  in  $E(X, X^c)$  satisfies

$$c(1-c)n^2 \frac{a}{n} \leq \mu_{E(X, X^c)} \leq c(2-c)n^2 \frac{a}{n}.$$

Let  $\delta_2 := 4 \leq \frac{8cnd}{\mu_{E(X, X^c)}}$ , then by Chernoff bound

$$\begin{aligned} \mathbb{P}(|E(X, X^c)| \geq 8cnd) &\leq \left( \frac{\exp(\delta_2 - 1)}{\delta_2^{\delta_2}} \right)^{\mu_{E(X, X^c)}} \\ &\leq \exp(-c(2-c)an). \end{aligned}$$

Now, if we substitute  $c = a^{-3}$  in the above bounds, we get

$$\begin{aligned} \mathbb{P}(|E(X)| \geq 2cnd) &\leq \exp(-0.75 \log(a) a^{-2} n) \\ \mathbb{P}(|E(X, X^c)| \geq 8cnd) &\leq \exp(-a^{-2} n). \end{aligned}$$

There are at most

$$\binom{2n}{cn} \leq \exp \left( -c \left( \log \left( \frac{c}{2} \right) - 1 \right) n \right)$$

subsets  $X$  of size  $|X| = cn$ . Substituting  $c = a^{-3}$  again, we get

$$\binom{2n}{cn} \leq \exp(4a^{-3} \log(a)n).$$

The claim follows from the union bound.

**Proof of Lemma 12:** We start by proving a simpler result.

**Lemma 20** *Let  $M$  be random symmetric matrix of size  $n$  with zero diagonal whose entries above the diagonal are independent with the following distribution*

$$M_{ij} = \begin{cases} 1 - p_{ij} & \text{w.p. } p_{ij} \\ -p_{ij} & \text{w.p. } 1 - p_{ij} \end{cases}.$$

*Let  $\sigma^2 \geq C_1 \frac{\log n}{n}$  be a quantity such that  $p_{ij} \leq \sigma^2$  for all  $i, j$ , where  $C_1$  is a constant. Then with probability  $1 - o(1)$ ,  $\|M\| \leq C_2 \sigma \sqrt{n}$  for some constant  $C_2 > 0$ .*

Let us address Lemma 20. A weaker bound  $C\sigma\sqrt{n\log n}$  follows easily from Alshwede-Winter type matrix concentration results (see Tropp (2012)). To prove the claimed bound, we need to be more careful and follow the  $\epsilon$ -net approach by Kahn and Szemerédi for random regular graphs in Friedman et al. (1989) (see also Alon and Kahale (1994); Feige and Ofek (2005)).

Consider a  $\frac{1}{2}$ -net  $\mathcal{N}$  of the unit sphere  $\mathcal{S}^n$ . We can assume  $|\mathcal{N}| \leq 5^n$ . It suffices to prove that there exists a constant  $C'_2$  such that with probability  $1 - o(1)$ ,  $|\mathbf{x}^T M \mathbf{y}| \leq C'_2 \sigma \sqrt{n}$  for all  $\mathbf{x}, \mathbf{y} \in \mathcal{N}$ .

For two vectors  $\mathbf{x}, \mathbf{y} \in \mathcal{N}$ , we follow an argument of Kahn and Szemerédi Friedman et al. (1989) and call all pairs  $(i, j)$  such that  $|x_i y_j| \leq \frac{\sigma}{\sqrt{n}}$  *light* and all remaining pairs *heavy* and denote these two classes by  $L$  and  $H$  respectively. We have

$$\mathbf{x}^T M \mathbf{y} = \sum_{i,j} x_i M_{ij} y_j = \sum_L x_i M_{i,j} y_j + \sum_H x_i M_{i,j} y_j.$$

We now show that with probability  $1 - o(1)$ , the last two summands are small in absolute value.

First, let us consider the contribution of light couples. We rewrite  $X := \sum_L x_i M_{i,j} y_j$  as  $\sum_{(i,j) \in L, i > j} M_{i,j} a_{i,j}$ , where

$$a_{i,j} = \begin{cases} x_i y_j + x_j y_i & \text{if } (i, j), (j, i) \in L \\ x_i y_j & \text{if } (i, j) \in L \\ x_j y_i & \text{if } (j, i) \in L \end{cases}$$

By the definition light pairs,  $|a_{i,j}| \leq 2\frac{\sigma}{\sqrt{n}}$ . Also, since  $\mathbf{x}$  and  $\mathbf{y}$  are unit vectors,  $\sum_{i,j} a_{i,j}^2 \leq 4$ . Therefore, by Bernstein's bound (see page 36 in Boucheron et al. (2013) for e.g.)

$$\mathbb{P}(X > t) \leq \exp\left(\frac{-\frac{1}{2}t^2}{4\sigma^2 + \frac{1}{3}2\frac{\sigma}{\sqrt{n}}t}\right).$$

Set  $t = 10\sigma\sqrt{n}$  and use the union bound (combining with the fact that the net has at most  $5^n$  vectors, we can conclude that with probability at least  $1 - \exp(-3n)$ ,  $|\sum_L x_i M_{i,j} y_j| \leq 10\sigma$ .

Next we handle the heavy pairs in  $H$ . Since  $1 \geq \sum_H x_i^2 y_j^2$ , the definition of *heavy* implies that  $\sum_H |x_i y_j| \leq \frac{\sqrt{n}}{\sigma}$ .

Let  $A_{i,j} := M_{i,j} + p_{i,j}$ , then

$$\sum_H x_i M_{i,j} y_j = \sum_H x_i A_{i,j} y_j - \sum_H p_{i,j} x_i y_j.$$

Note that  $A$  defines a graph, say  $G_A$ , such that  $A$  is its adjacency matrix. As  $p_{ij} \leq \sigma^2$ , we have  $\sum_H p_{ij} |x_i y_j| \leq \sigma^2 \frac{\sqrt{n}}{\sigma} = \sigma \sqrt{n}$ . We use the following lemma to bound the first term.



**Lemma 21** Let  $\tilde{G} = (\tilde{V}, \tilde{E})$  be any graph whose adjacency matrix is denoted by  $\tilde{A}$ , and  $\mathbf{x}, \mathbf{y}$  be any two unit vectors. Let  $\tilde{d}$  be such that the maximum degree  $\leq c_1 \tilde{d}$ . Further, let  $\tilde{d}$  satisfy the property that for any two subsets of vertices  $S, T \subset \tilde{V}$  one of the following holds for some constants  $c_2$  and  $c_3$ :

$$\frac{e(S, T)}{|S||T|\frac{\tilde{d}}{n}} \leq c_2 \quad (1)$$

$$e(S, T) \log \left( \frac{e(S, T)}{|S||T|\frac{\tilde{d}}{n}} \right) \leq c_3 |T| \log \frac{n}{|T|} \quad (2)$$

then  $\sum_H x_i \tilde{A}_{i,j} y_j \leq \max(16, 8c_1, 32c_2, 32c_3) \sqrt{\tilde{d}}$ . Here  $H := \{(i, j) \mid |x_i y_j| \geq \sqrt{\tilde{d}/n}\}$ .

The proof appears in appendix D.

**Lemma 22** Let  $\tilde{d} := \sigma^2 n$ . Then with probability  $1 - o(1)$ , the maximum degree in the graph  $G_A$  is  $\leq 20\tilde{d}$  and for any  $S, T \subset V$  one of the conditions (1) or (2) holds.

The two lemmas above guarantee that with probability  $1 - o(1)$ ,  $|\sum_H x_i A_{i,j} y_j| \leq C' \sigma \sqrt{n}$  for some constant  $C'$ .

**Proof** The bound on the maximum degree follows from the Chernoff bound. We have that

$$A_{ij} = \begin{cases} 1 & \text{w.p. } p_{ij} \\ 0 & \text{w.p. } 1 - p_{ij} \end{cases}.$$

Consider a particular vertex  $k$  and let  $X = \sum_i A_{ik}$  be the random variable denoting the number of edges incident on it. We have that

$$\mu = \mathbf{E}X = \sum_i p_{ik} \leq \sigma^2 n.$$

For any  $l \geq 4$ , Chernoff bound (see Alon and Spencer (2004)) implies that

$$\begin{aligned} \mathbb{P}(X > l\sigma^2 n) &\leq \exp\left(-\frac{\sigma^2 n l \ln l}{3}\right) \\ &\leq \exp\left(-\frac{l \log n}{3}\right). \end{aligned}$$

Applying this with  $l = 20$ , and taking a union bound over all the vertices, we can bound the maximum degree by  $20\sigma^2 n$ . Now let  $S, T \subset V$  be any two subsets. Let  $X := e(S, T)$  be the number of edges going between  $S$  and  $T$ . We have  $\mathbf{E}X \leq \sigma^2 |S||T|$ . If  $|T| \geq \frac{n}{e}$ , then since the maximum degree is  $\leq 20\sigma^2 n$ , we have  $e(S, T) \leq |S|20\sigma^2 n \leq 20e\sigma^2 |S||T|$ , giving us 1 in this case. Therefore, we can assume  $|T| \leq \frac{n}{e}$ . By Chernoff bound, it follows that for any  $l \geq 4$ ,

$$\mathbb{P}(e(S, T) > l\sigma^2 |S||T|) \leq \exp\left(-\frac{l \ln(l) \sigma^2 |S||T|}{3}\right).$$

Let  $l'$  be the smallest number such that  $l' \ln(l') \geq \frac{21|T|}{\sigma^2|S||T|} \log\left(\frac{n}{|T|}\right)$ . As in Feige and Ofek (2005), if we choose  $l = \max(l', 4)$ , we can bound the above probability by  $\exp\left(-\frac{l \ln(l) \sigma^2 |S| |T|}{3}\right) \binom{n}{|S|} \binom{n}{|T|} \leq \frac{1}{n^3}$ . Therefore, by the union bound we get that with probability  $1 - o(1)$  for all subsets  $S, T$ , and

$$e(S, T) \leq \max(l', 4) \sigma^2 |S| |T|.$$

This implies that one of the conditions 1 or 2 holds with probability  $1 - o(1)$ . ■

**Proof of Lemma 12:** Now we are ready to prove Lemma 12 by modifying the previous proof. We again handle the light couples and the heavy couples separately, but need to make a modification to the argument for the light couples.

Since we zero out some rows and columns of  $M$  to obtain  $M_1$ , we first bound the norm of the matrix  $M_0$ , obtained from  $M$  by zeroing out a set  $S$  of rows and the corresponding columns. Next, we take a union bound over all choices of  $S$ . For a fixed  $S$ , lemma 20 implies that with probability at least  $1 - \exp(-3n)$ , for all  $\mathbf{x}, \mathbf{y} \in \mathcal{N}_{1/2}$ ,  $|\sum_L x_i (M_0)_{ij} y_j| \leq 10\sigma\sqrt{n}$ . Since there are at most  $2^n = \exp(n \ln 2)$  choices for  $S$ , we can apply a union bound to show that with probability at least  $1 - \exp(-(3 - \ln 2)n)$ ,  $|\sum_L x_i (M_1)_{ij} y_j| \leq 10\sigma\sqrt{n}$ .

The proof for the heavy couples goes through without any modifications. We just have to verify that the conditions of lemma 21 are met. Firstly, the adjacency matrix  $A_1$  obtained from  $M_1$  has bounded degree property by the definition of  $M_1$ . Now we note that only for the case of  $|S| \leq |T| \geq \frac{n}{e}$  did we need that the maximum degree was bounded. So for any  $|S| \leq |T| < \frac{n}{e}$ , the discrepancy properties (1) or (2) holds for  $A_1$ , since zeroing out rows and columns can only decrease the edge count across sets of vertices. In the case  $|T| \geq \frac{n}{e}$ , like before we can show that (1) holds for  $A_1$  since the degrees are bounded. ■

Now, to bound the norm of matrix  $E$ , we just appeal to 12. Suppose  $a > b \geq C_0$ , for a large enough constant  $C_0$  to be determined later. Since  $A = \bar{A}_0 + \Delta + E$  and we have bounded  $\Delta$ , it remains to bound  $\|E\|$ . Note that

$$(E_0)_{ij} = \begin{cases} 1 - \frac{a}{n} & \text{w.p. } \frac{a}{n} \\ -\frac{a}{n} & \text{w.p. } 1 - \frac{a}{n} \end{cases}$$

if  $i, j$  belongs to the same community and

$$(E_0)_{ij} = \begin{cases} 1 - \frac{b}{n} & \text{w.p. } \frac{b}{n} \\ -\frac{b}{n} & \text{w.p. } 1 - \frac{b}{n} \end{cases}$$

if  $i, j$  belongs to different communities. Since  $a > b$ , for all  $i, j$  we have that

$$\text{Var}((E_0)_{ij}) \leq \frac{a}{n} \left(1 - \frac{a}{n}\right) \leq \frac{d}{n}.$$

## A.2. Recovery

Now we focus on the second step in the proof, namely the recovery of the blocks once the angle condition is satisfied.

**Lemma 23** *If  $\sin(\angle \bar{W}, W) \leq c \leq \frac{1}{4}$ , then we can find a vector  $\mathbf{v} \in W$  such that  $\sin(\angle \mathbf{v}, \bar{\mathbf{v}}_2) \leq 2\sqrt{c}$ .*

**Proof** Let  $P_{\bar{W}}, P_W$  be the orthogonal projection operators on to the subspaces  $\bar{W}, W$  respectively. From the angle bound for the subspaces, we have that

$$\|P_{\bar{W}} - P_W\|_2 \leq c.$$

The vector we want is obtained as follows. We first project  $\bar{\mathbf{v}}_1$  on to  $W$ , and then find the unit vector orthogonal to the projection in  $W$ . We will now prove that the vector so obtained satisfies the bound stated in the lemma. Since  $\bar{\mathbf{v}}_1, \bar{\mathbf{v}}_2 \in \bar{W}$ , we have that  $\|P_W \bar{\mathbf{v}}_i - \bar{\mathbf{v}}_i\|_2 \leq c$  for  $i = 1, 2$ . Let us define  $\mathbf{u}_i := P_W \bar{\mathbf{v}}_i$  and  $\mathbf{x}_i := \mathbf{u}_i - \bar{\mathbf{v}}_i$  (note that  $\|\mathbf{x}_i\| \leq c$ ) for  $i = 1, 2$ . We will now show that the vector  $\mathbf{v} \in W$  perpendicular to  $\mathbf{u}_1$  is close to  $\bar{\mathbf{v}}_2$ . Let  $\mathbf{u}_\perp = \mathbf{u}_2 - \frac{\mathbf{u}_1^T \mathbf{u}_2}{\|\mathbf{u}_1\|^2} \mathbf{u}_1$ , it is then clear that  $\|\mathbf{u}_\perp\| \leq 1$ . Note that  $|\mathbf{u}_1^T \mathbf{u}_2| = |\bar{\mathbf{v}}_1^T \mathbf{x}_2 + \bar{\mathbf{v}}_2^T \mathbf{x}_1 + \mathbf{x}_1^T \mathbf{x}_2| \leq 2c + c^2$ . We have,

$$\begin{aligned} \mathbf{u}_\perp^T \bar{\mathbf{v}}_2 &= \mathbf{u}_2^T \bar{\mathbf{v}}_2 - \frac{(\mathbf{u}_1^T \mathbf{u}_2)(\bar{\mathbf{v}}_2^T \mathbf{u}_1)}{\|\mathbf{u}_1\|^2} \\ |\mathbf{u}_\perp^T \bar{\mathbf{v}}_2| &\geq 1 - c - \frac{(2c + c^2)c}{(1 - c)^2} \\ &\geq 1 - 2c. \end{aligned}$$

The last inequality holds when  $c \leq \frac{1}{4}$ . Therefore, it holds that for a unit vector  $\mathbf{v} \perp \mathbf{u}_1$ ,

$$|\mathbf{v}^T \bar{\mathbf{v}}_2| \geq |\mathbf{u}_\perp^T \bar{\mathbf{v}}_2| \geq 1 - 2c.$$

This gives  $\sin(\angle \mathbf{v}, \bar{\mathbf{v}}_2) \leq \sqrt{1 - (1 - 2c)^2} \leq 2\sqrt{c}$ . ■

Lemmas 14 and 23 together give

**Corollary 24** *For any constant  $c < 1$ , we can choose constants  $C_2$  and  $C_3$  in lemma 14 and find a vector  $\mathbf{v}$  such that  $\sin(\angle \bar{\mathbf{v}}_2, \mathbf{v}) \leq c < 1$  with probability  $1 - o(1)$ .*

We now can conclude the proof of our theorem using the following deterministic fact.

**Lemma 25** *If  $\sin(\angle \bar{\mathbf{v}}_2, \mathbf{v}) < c \leq 0.5$ , then we can identify at least a  $(1 - \frac{4}{3}c^2)$  fraction of vertices from each block correctly.*

**Proof of Lemma 25:** Let us define two sets of vertices,  $V'_1 = \{i | \mathbf{v}(i) > 0\}$  and  $V'_2 = \{i | \mathbf{v}(i) < 0\}$ . One of the sets will have less than or equal to  $\frac{n}{2}$  vertices, let us assume without loss of generality that  $|V'_1| \leq \frac{n}{2}$ . Writing  $\mathbf{v} = c_1 \bar{\mathbf{v}}_2 + \mathbf{err}$ , for a vector  $\mathbf{err}$  perpendicular to  $\bar{\mathbf{v}}_2$  and  $\|\mathbf{err}\| < c$ . We also have  $c_1 > \sqrt{1 - c^2}$ . Since  $\|\mathbf{err}\| < c$ , not more than  $\frac{c^2}{1 - c^2}n$  coordinates of  $\mathbf{err}$  can be bigger than  $\frac{\sqrt{1 - c^2}}{\sqrt{n}} < \frac{c_1}{\sqrt{n}}$ . Since  $\mathbf{v} = c_1 \bar{\mathbf{v}}_2 + \mathbf{err}$  at least  $1 - \frac{c^2}{1 - c^2} > 1 - \frac{4}{3}c^2$  (since  $c \leq 0.5$ ) fraction of vertices with  $\bar{\mathbf{v}}_2(i) = \frac{1}{\sqrt{n}}$  will have  $\mathbf{v}(i) > 0$ . Therefore, we get that there are at least  $(1 - \frac{4}{3}c^2)n$  vertices belonging to the first block. ■

### A.3. Proof of lemma 9

We will use the following large deviation result (see page 36 in [Boucheron et al. \(2013\)](#) for e.g.) repeatedly

**Lemma 26** (Chernoff) *If  $X$  is a sum of  $n$  iid indicator random variables with mean at most  $\rho \leq 1/2$ , then for any  $t > 0$*

$$\max\{\mathbb{P}(X \geq \mathbf{E}X + t), \mathbb{P}(X \leq \mathbf{E}X - t)\} \leq \exp\left(-\frac{t^2}{2\mathbf{Var}X + t}\right) \leq \exp\left(-\frac{t^2}{2n\rho + t}\right).$$

In the Red graph, the edge densities are  $a/2n$  and  $b/2n$ , respectively. By Theorem 7, there is a constant  $C$  such that if  $\frac{(a-b)^2}{a+b} \geq C$  then by running **Spectral Partition** on the Red graph, we obtain, with probability  $1 - o(1)$  two sets  $V'_1$  and  $V'_2$ , where

$$|V_i \setminus V'_i| \leq .1n.$$

In the rest, we condition on this event, and the event that the maximum Red degree of a vertex is at most  $\log^2 n$ , which occurs with probability  $1 - o(1)$ .

Now we use the Blue edges. Consider  $e = (u, v)$ . If  $e$  is not a red edge, and  $u \in V_i, v \in V_{3-i}$ , then  $e$  is a Blue edge with probability

$$\mu := \frac{b/2n}{1 - \frac{b}{2n}}.$$

Similarly, if  $e$  is not a Red edge, and  $u, v \in V_i$ , then  $e$  is a Blue edge with probability

$$\tau := \frac{a/2n}{1 - \frac{a}{2n}}.$$

Thus, for any  $u \in V'_i \cap V_i$ , the number of its Blue neighbors in  $V'_{3-i}$  is at most

$$S(u) := \sum_{i=1}^{.9n} \xi_i^u + \sum_{j=1}^{.1n} \zeta_j^u$$

where  $\xi_i^u$  are iid indicator variables with mean  $\mu$  and  $\zeta_j^u$  are iid indicator variables with mean  $\tau$ .

Similarly, for any  $u \in V'_1 \cap V_2$ , the number of its Blue neighbors in  $V'_2$  is at least

$$S'(u) := \sum_{i=1}^{.9n-d(u)} \zeta_i^u + \sum_{j=1}^{.1n} \xi_j^u,$$

where  $d(u) = \log^2 n$  is the Red degree of  $u$ .

After the correction sub-routine, a vertex  $u$  in the (corrected) set  $V'_1$  is misclassified if

- $u \in V'_1 \cap V_1$  and  $S_u \geq \frac{a+b}{4}$ .
- $u \in V'_1 \cap V_2$  and  $S'_u \leq \frac{a+b}{4}$

Let  $\rho_1, \rho_2$  be the probability of the above events. Then the number of misclassified vertices in the (corrected) set  $V_1'$  is at most

$$M := \sum_{k=1}^n \Gamma_k + \sum_{l=1}^{0.1n} \Lambda_l$$

where  $\Gamma_k$  are iid indicator random variables with mean  $\rho_1$  and  $\Lambda_l$  are iid indicator random variables with mean  $\rho_2$ .

The rest is a simple computation. First we use Chernoff bound to estimate  $\rho_1, \rho_2$ . Consider

$$\rho_1 := \mathbb{P}\left(S(u) \geq \frac{a+b}{4}\right).$$

By definition, we have

$$\begin{aligned} \mathbf{E}S(u) &= 0.9n\mu + 0.1n\tau \\ &= 0.9n\left(\frac{b/2n}{1 - \frac{b}{2n}}\right) + 0.1n\left(\frac{a/2n}{1 - \frac{a}{2n}}\right) \\ &= 0.9\frac{b}{2} + 0.1\frac{a}{2} + 0.9\frac{b}{2}\left(\frac{1}{1 - b/2n} - 1\right) + 0.1\frac{a}{2}\left(\frac{1}{1 - a/2n} - 1\right). \end{aligned} \tag{3}$$

Set

$$t := \frac{a+b}{4} - \mathbf{E}S(u),$$

we have

$$t = 0.2(a-b) - 0.9\frac{b}{2}\left(\frac{1}{1 - b/2n} - 1\right) - 0.1\frac{a}{2}\left(\frac{1}{1 - a/2n} - 1\right) \geq 0.2(a-b) - 0.9\frac{b}{2n} - 0.1\frac{a}{2n} \geq 0.19(a-b),$$

for any sufficiently large  $n$ .

Applying Chernoff's bound, we obtain

$$\rho_1 \leq \exp\left(-\frac{(0.19(a-b))^2}{2(0.9n\mu + .1n\tau) + 0.19(a-b)}\right).$$

By (4), one can show that  $2(0.9n\mu + .1n\tau) + 0.19(a-b) = 0.71b + 0.29a + o(1) \leq \frac{a+b}{2}$ . It follows that

$$\rho_1 \leq \exp\left(-0.072\frac{(a-b)^2}{a+b}\right).$$

By a similar argument, we obtain the same estimate for  $\rho_2$  (the contribution of the term  $d(u) \leq \log^2 n$  is negligible). Thus, we can conclude that

$$\mathbf{E}M \leq 1.1n \exp\left(-0.072\frac{(a-b)^2}{a+b}\right).$$

Applying Chernoff's with  $t := 0.9n \exp(-0.072 \frac{(a-b)^2}{a+b})$ , we conclude that with probability  $1 - o(1)$

$$M \leq \mathbf{E}M + t = 2n \exp(-0.072 \frac{(a-b)^2}{a+b}).$$

This implies that with probability  $1 - o(1)$ ,

$$|V'_1 \setminus V_1| \leq 2n \exp(-0.072 \frac{(a-b)^2}{a+b}).$$

By symmetry, the same conclusion holds for  $|V'_2 \setminus V_2|$ .

Set

$$\gamma := 2 \exp(-0.072 \frac{(a-b)^2}{a+b}),$$

we have, for  $i = 1, 2$

$$|V_i \cap V'_i| = n - |V_i \cap V'_{3-i}| = n - |V'_{3-i} \setminus V_{3-i}| \geq n(1 - \gamma).$$

This shows that the output  $V'_1, V'_2$  form a  $\gamma$ -correct partition, with  $\gamma$  satisfying

$$\frac{(a-b)^2}{a+b} = \frac{1}{0.072} \log \frac{2}{\gamma} \approx 13.89 \log \frac{2}{\gamma},$$

proving our claim.

**Proof of Corollary 4:** Notice that in the analysis of **Spectral Partition**, we only require  $\frac{(a-b)^2}{a+b} \geq C$  for a sufficiently large constant  $C$  (so  $\gamma$  does not appear in the bound). In the analysis of **Correction**, we require  $\frac{(a-b)^2}{a+b} \geq 13.89 \log \frac{2}{\gamma}$ , as shown above. If  $\gamma < \epsilon$  for a sufficiently small  $\epsilon$ , this assumption implies the first. Thus, Corollary holds with assumption  $\frac{(a-b)^2}{a+b} \geq 13.89 \log \frac{2}{\gamma}$ .

The constant 13.89 comes from the fact that the partition obtained from **Spectral Partition** is .1-correct. If one improves upon .1, one improves 13.89. In particular, there is a constant  $\delta$  such that if the first partition is  $\delta$ -correct, then one can improve 13.89 to 8.1 (or any constant larger than 8—which is the limit of the method, for that matter).

## Appendix B. Multiple communities

We say the splitting is ‘perfect’ if we have  $|Y_1 \cap V_i| = \frac{n}{4k} = |Y_2 \cap V_i|$  for  $i = 1, \dots, k$ . We will assume the splittings are perfect in the proofs for a simpler exposition. Though the splitting will almost always not be perfect, and there will just be a  $o(1)$  error term that we have to carry throughout to be precise. The bounds we give will all be still be essentially the same.

### B.1. Proof of theorem 17

To analyze this algorithm, we use the machinery developed so far combined with some ideas from [Vu \(2014\)](#). We consider the stochastic block model with  $k$  blocks of size  $n$ , where  $k$  is a fixed

constant as  $n$  grows. This is a graph  $V = V_1 \cup V_2 \cup \dots \cup V_k$  where each  $|V_i| = n/k$  and for  $u \in V_i, v \in V_j$ :

$$\mathbb{P}[(u, v) \in E] = \begin{cases} a/n & \text{if } i = j \\ b/n & \text{if } i \neq j \end{cases}$$

We can write, as before

$$\begin{aligned} A &= \bar{A} + E \\ &= \bar{A}_1 + \Delta + E, \end{aligned}$$

where  $\bar{A}, \bar{A}_1$  are the expected matrices, and  $\Delta$  is matrix containing the deleted rows and columns. Let  $\bar{W}$  be the span of the  $k$  left singular vectors of  $\bar{A}_1$ . We can bound  $\|\Delta\| \leq 1$  by bounding the number of high degree vertices as we did before.  $E$  is given by

$$E_{u,v} = \begin{cases} 1 - \frac{a}{n} & \text{w.p. } \frac{a}{n} \\ -\frac{a}{n} & \text{w.p. } 1 - \frac{a}{n} \end{cases}$$

if  $u, v \in V_i \cap Y_1$  for some  $i \in 1, \dots, k$  and

$$E_{u,v} = \begin{cases} 1 - \frac{b}{n} & \text{w.p. } \frac{b}{n} \\ -\frac{b}{n} & \text{w.p. } 1 - \frac{b}{n} \end{cases}$$

if  $u \in V_i \cap Y_1$  and  $v \in V_j \cap Y_1$  for  $i \neq j$ . Since  $\sigma^2 := \frac{a}{n} \geq \text{Var}(E_{u,v})$ , corollary 12 applied to  $\bar{A}_1 - A$  gives the following result.

**Lemma 27** *There exists a constant  $C$  such that  $\|E\| \leq C\sqrt{a+b}$  with probability  $1 - o(1)$ .*

It is not hard to show that the rank of the matrix  $\bar{A}_1$  is  $k$ , and its least non-trivial singular value is  $\sigma_k(\bar{A}_1) = \frac{a-b}{k}$ . This fact, combined with lemma 27 and an application of Davis-Kahan bound gives

**Lemma 28** *For any  $c > 0$ , there exists constants  $C_1, C_2$  such that if  $(a-b) > C_1 k^2 a$  and  $a > b \geq C_2$ , then  $\sin \angle(\bar{W}, W) \leq c$  with probability  $1 - o(1)$ .*

We pick  $m = 2 \log n$  indices uniformly randomly from  $Y_2$  and project the corresponding columns from the matrix  $B$ . Let  $\bar{a}_{i_1}, \dots, \bar{a}_{i_m}$  and  $e_{i_1}, \dots, e_{i_m}$  be the corresponding columns of  $\bar{A}_1$  and  $E$ , respectively. For a subspace  $W_0$ , let  $P_{W_0}$  be the projection on to the space  $W_0$ . Note that if vertex  $i \in V_{n_i} \cap Y_2$ , then

$$\bar{a}_i(j) = \begin{cases} \frac{a}{n} & \text{if } j \in V_{n_i} \cap Z \\ \frac{b}{n} & \text{otherwise} \end{cases}.$$

We let the vector  $\bar{a}$  be

$$\bar{a}(j) := \begin{cases} \frac{a+b}{2n} & \text{if } j \in Z \end{cases}.$$

and  $b_i = \bar{a}_i - \bar{a}$ . We therefore have

$$b_i(j) = \begin{cases} \frac{a-b}{2n} & \text{if } j \in V_{n_i} \cap Z \\ \frac{b-a}{2n} & \text{otherwise} \end{cases}.$$

Since both  $\bar{a}_i, \bar{a}$  are in the column span of  $\bar{A}_1$ , we have for all  $i$

$$\mathbf{b}_i = P_{\bar{W}} \mathbf{b}_i.$$

We also note that  $\|\mathbf{b}_i\| = \frac{(a-b)}{2\sqrt{2n}}$ . Therefore, if we can recover  $\mathbf{b}_i$ , we can identify the set  $V_{n_i} \cap Z$ . We now argue that we can recover  $\mathbf{b}_i$  approximately. Since  $\mathbf{a}_i - \bar{\mathbf{a}} = \mathbf{b}_i + \mathbf{e}_i$ , we have

$$\begin{aligned} P_W(\mathbf{a}_i - \bar{\mathbf{a}}) &= P_W \mathbf{b}_i + P_W \mathbf{e}_i \\ &= P_{\bar{W}} \mathbf{b}_i + P_W \mathbf{e}_i + \mathbf{err}_i \\ &= \mathbf{b}_i + P_W \mathbf{e}_i + \mathbf{err}_i, \end{aligned}$$

where  $\mathbf{err}_i = (P_W - P_{\bar{W}}) \mathbf{b}_i$ . Since  $\sin \angle(\bar{W}, W) \leq \delta_1$ , we have for any unit vector  $\mathbf{v}$ ,  $\|P_W \mathbf{v} - P_{\bar{W}} \mathbf{v}\| \leq \delta_1$ , which in turn implies for all  $i$

$$\|\mathbf{err}_i\| \leq \delta_1 \|\mathbf{b}_i\|.$$

Therefore, it is enough to bound  $\|P_W \mathbf{e}_i\|$ . We recall that  $k$  is a constant that does not depend on  $n$ .  $W$  is  $k$  dimensional space giving  $\mathbb{E} \|P_W \mathbf{e}_i\|^2 \leq k\sigma^2$ . By Markov's inequality, it follows that

$$\mathbb{P}(\|P_W \mathbf{e}_i\| > 2\sigma k^{1/2}) \leq \frac{1}{4}.$$

By a simple application of Chernoff bound, we have

**Lemma 29** *With probability at least  $1 - o(1)$ , at least  $m/2$  of the vectors  $\mathbf{e}_{i_1}, \dots, \mathbf{e}_{i_m}$  satisfy*

$$\|P_W \mathbf{e}_{i_j}\| < 2\sigma k^{1/2}.$$

Let  $m_1 \geq m/2$  denote the number of such vectors, hence referred to as good vectors. To avoid introducing extra notation, let us say  $\mathbf{e}_{i_1}, \dots, \mathbf{e}_{i_{m_1}}$  are the good vectors and the corresponding indices as good indices. Note that  $\sigma \leq \frac{\sqrt{a}}{\sqrt{n}}$ . For any  $\delta_2 > 0$ , there exists a big enough constant  $C_1$  such that if  $(a - b) > C_1 \sqrt{ka}$ , we have that  $2\sigma k^{1/2} \leq \delta_2 \|\mathbf{b}_{i_j}\|$  whenever  $i_j$  is good. Therefore

**Lemma 30** *Given any  $\delta > 0$ , there exists constants  $C_1, C_2$  such that the following holds. If  $(a - b) > C_1 \sqrt{ka}$  and  $a \geq b \geq C_2$ , then for all good indices  $i_j$ , it holds that  $\|P_W(\mathbf{a}_{i_j} - \bar{\mathbf{a}}) - \mathbf{b}_{i_j}\| \leq \delta \|\mathbf{b}_{i_j}\|$ .*

Let  $U'_{i_j}$  be the top  $n/2k$  coordinates of the projected vector  $\|P_W(\mathbf{a}_{i_j} - \bar{\mathbf{a}})\|$ . If we choose the constants  $C_1, C_2$  appropriately, then for every good index  $i_j$ ,  $U'_{i_j}$  contains 0.95 fraction of the vertices in  $V_{n_{i_j}} \cap Z$ .

Lemma 31 then implies that when we throw away half of the sets  $U'_1, \dots, U'_m$  with the least Blue edge densities, then each of the remaining sets intersects some  $V_{n_i} \cap Z$  in 0.9 fraction of the vertices.

**Lemma 31** *There exists a constant  $c > 0$  such that the following holds. Suppose we are given a set  $X \subset Z$  of size  $|X| = n/2k$ . If for all  $i \in 1, \dots, k$*

$$|X \cap V_i| \leq 0.9|X|,$$



then with probability at least  $1 - e^{-cn}$  the number of Blue edges in the graph induced by  $X$  is at most  $an/16k - 0.09(a-b)n/16k$ . Conversely, if

$$|X \cap V_i| \geq 0.95|X|$$

for some  $i \in 1, \dots, k$ , then with probability at least  $1 - e^{-cn}$  the number of Blue edges in the graph induced by  $X$  is at least  $an/16k - 0.09(a-b)n/16k$ .

**Proof**

Let  $e(X)$  denote the number of Blue edges in the graph induced by vertices in  $X$ . Suppose  $|X \cap V_i| \leq 0.9|X|$  for all  $i \in 1, \dots, k$ . Then

$$\mathbf{E}e(X) \leq an/16k - 0.09(a-b)n/8k.$$

To bound the probability that  $\mathbf{E}e(X) \geq an/16k - .045(a-b)n/8k$ , we can use Chernoff bound.

$$\text{Let } \delta = \frac{0.045(a-b)n/8k}{an/16k - 0.09(a-b)n/8k}.$$

$$\mathbb{P}(\mathbf{E}e(X) \geq an/16k - .045(a-b)n/8k) \leq \exp\left(-\frac{(.045(a-b)n/8k)^2}{2an/16k + 0.045(a-b)n/8k}\right).$$

Similarly, suppose  $|X \cap V_i| \geq 0.95|X|$  for some  $i \in 1, \dots, k$ . Then

$$\mathbf{E}e(X) \geq an/16k - 0.05(a-b)n/16k.$$

To bound the probability that  $\mathbf{E}e(X) \leq an/16k - .045(a-b)n/8k$ , we can use Chernoff bound.

$$\text{Let } \delta = \frac{0.04(a-b)n/16k}{an/16k - 0.05(a-b)n/16k}.$$

$$\mathbb{P}(\mathbf{E}e(X) \leq an/16k - .045(a-b)n/8k) \leq \exp\left(-\frac{(.04(a-b)n/16k)^2}{2an/16k + 0.04(a-b)n/16k}\right).$$

**B.2. Proof of lemma 18**

For notional convenience, let  $U_i := Z \cap V_i$ . We will use the following large deviation result (see page 36 in [Boucheron et al. \(2013\)](#) for e.g.) repeatedly

**Lemma 32** (Chernoff) *If  $X$  is a sum of  $n$  iid indicator random variables with mean at most  $\rho \leq 1/2$ , then for any  $t > 0$*

$$\max\{\mathbb{P}(X \geq \mathbf{E}X + t), \mathbb{P}(X \leq \mathbf{E}X - t)\} \leq \exp\left(-\frac{t^2}{2\mathbf{Var}X + t}\right) \leq \exp\left(-\frac{t^2}{2n\rho + t}\right).$$

By Theorem step1, there is a constant  $C$  such that if  $\frac{(a-b)^2}{a+b} \geq C$  then by running **Spectral Partition** on the Red graph, we obtain, with probability  $1 - o(1)$ , sets  $U'_1, \dots, U'_k$ , where

$$|U'_i \setminus U_i| \leq 0.1n/2k.$$

In the rest, we condition on this event. The probability we will talk about in this section is based on the edges that go between vertices in  $Z$ .

Now we use the edges that go between vertices in  $Z$ . Consider  $e = (u, v)$ . If  $u \in U_i, v \in U_j$  with  $i \neq j$ , then  $e$  is a Red edge with probability

$$\mu := b/2n.$$

Similarly, if  $u, v \in U_i$ , then  $e$  is a Red edge with probability

$$\tau := a/2n.$$

For any  $u \in U_1$ , the number of its neighbors in  $U'_j$  is at most

$$S_{1i}(u) := \sum_{i=1}^{.9n/2k} \xi_i^u + \sum_{j=1}^{.1n/2k} \zeta_j^u$$

Similarly, for any  $u \in U_1$ , the number of its neighbors in  $U'_1$  is at least

$$S'_{11}(u) := \sum_{i=1}^{.9n/2k} \zeta_i^u + \sum_{j=1}^{.1n/2k} \xi_j^u.$$

After the correction sub-routine, if a vertex  $u \in U_1$  is mislabeled then one of the following holds

- $S'_{1j} \geq \frac{a+b}{8}$  for some  $j \neq 1$
- $S_{11} \leq \frac{a+b}{8}$ .

By an application of Chernoff bound, probability that  $S_{11} \leq \frac{a+b}{8}$  can be bounded by  $\rho_1 = \exp(-0.04 \frac{(a-b)^2}{k(a+b)})$ . Similarly, for any fixed  $j \neq 1$ ,  $S'_{1j} \geq \frac{a+b}{8}$  is bounded by  $\rho_1$ . Therefore, the probability that any of these happens is bounded by  $k\rho_1$ . Therefore, number of vertices in  $U_1$  that will be misclassified after the correction step is at most

$$M := \sum_{k=1}^{n/2k} \Gamma_k$$

where  $\Gamma_k$  are iid indicator random variables with mean  $\rho_1$ .

$$\mathbf{E}M \leq \frac{n}{2k} k \exp(-0.04 \frac{(a-b)^2}{k(a+b)}).$$

Applying Chernoff's with  $t := \frac{n}{2} \exp(-0.04 \frac{(a-b)^2}{k(a+b)})$ , we conclude that with probability  $1 - o(1)$

$$M \leq \mathbf{E}M + t = n \exp(-0.04 \frac{(a-b)^2}{k(a+b)}).$$

This implies that with probability  $1 - o(1)$ , number of mislabeled vertices in  $U_1$  is

$$\leq n \exp(-0.04 \frac{(a-b)^2}{k(a+b)}).$$

Set

$$\gamma := 2k \exp(-0.04 \frac{(a-b)^2}{k(a+b)}).$$

Therefore, by a union bound over all  $i$ , we have that with probability  $1-o(1)$  the output  $U'_1, U'_2, \dots, U'_k$  after the correction step form a  $\gamma$ -correct partition, with  $\gamma$  satisfying

$$\frac{(a-b)^2}{k(a+b)} = \frac{1}{0.04} \log \frac{2k}{\gamma} = 25 \log \frac{2k}{\gamma},$$

proving our claim.

### B.3. Proof of lemma 19

In this section, we show how we can merge  $V_i \cap Y$  with  $V_i \cap Z$  based on the Blue edges that go in between vertices in  $Y$  and  $Z$ . We can assume that that we are given a  $\gamma$  correct partition  $U'_1, \dots, U'_k$  of  $U_1, \dots, U_k$ . Now we label the vertices in  $Y$  according to their degrees to  $U'_i$  as given in the **Merge** routine. Let us assume  $\gamma \leq 0.1$ . In the rest, we condition on this event, and the event that the maximum Red degree of a vertex is at most  $\log^2 n$ , which occurs with probability  $1 - o(1)$ .

Now we use the Blue edges. Consider  $e = (u, v)$ . If  $e$  is not a red edge, and  $u \in V_i \cap Y, v \in V_j \cap Z$ , then  $e$  is a Blue edge with probability

$$\mu := \frac{b/2n}{1 - \frac{b}{2n}}.$$

Similarly, if  $e$  is not a Red edge, and  $u \in V_i \cap Z, v \in V_i \cap Z$ , then  $e$  is a Blue edge with probability

$$\tau := \frac{a/2n}{1 - \frac{a}{2n}}.$$

Thus, for any  $u \in Y \cap V_i$ , the number of Blue neighbors in  $U'_j$  is at most

$$S_j := \sum_{i=1}^{.9n/2k} \xi_i^u + \sum_{j=1}^{.1n/2k} \zeta_j^u$$

where  $\xi_i^u$  are iid indicator variables with mean  $\mu$  and  $\zeta_j^u$  are iid indicator variables with mean  $\tau$ .

Similarly, for any  $u \in Y \cap V_i$ , the number of Blue neighbors in  $U'_i$  is at least

$$S'_i := \sum_{i=1}^{.9n/2k-d(u)} \zeta_i^u + \sum_{j=1}^{.1n/2k} \xi_j^u.$$

After the correction sub-routine, if a vertex  $u$  in  $Y \cap V_i$  is misclassified then one of the following holds

- $S_j \geq \frac{a+b}{8k}$
- $S'_i \leq \frac{a+b}{8k}$ .

Let  $\rho$  be the probability that at least one of the above events happens. Then the number of mislabeled vertices in the  $Y_2$  is at most

$$M := \sum_{k=1}^{n/2} \Gamma_k$$

where  $\Gamma_k$  are iid indicator random variables with mean  $\rho$ . First we use Chernoff bound to estimate  $\rho$ . Consider

$$\rho_1 := \mathbb{P}\left(S_j \geq \frac{a+b}{8k}\right).$$

By definition, we have

$$\begin{aligned} \mathbf{E}S(u) &= \frac{0.9n\mu/k + 0.1n\tau/k}{2} \\ &= 0.9n\left(\frac{b/4kn}{1 - \frac{b}{2n}}\right) + 0.1n\left(\frac{a/4kn}{1 - \frac{a}{2n}}\right) \\ &= 0.9\frac{b}{4k} + 0.1\frac{a}{4k} + 0.9\frac{b}{4k}\left(\frac{1}{1 - b/2n} - 1\right) + 0.1\frac{a}{4k}\left(\frac{1}{1 - a/2n} - 1\right). \end{aligned} \tag{4}$$

Set

$$t := \frac{a+b}{8k} - \mathbf{E}S_j,$$

we have

$$t = 0.1\frac{a-b}{k} - 0.9\frac{b}{4k}\left(\frac{1}{1 - b/2n} - 1\right) - 0.1\frac{a}{4k}\left(\frac{1}{1 - a/2n} - 1\right) \geq 0.1\frac{a-b}{k} - 0.9\frac{b}{4k}\frac{b}{n} - 0.1\frac{a}{4k}\frac{a}{n} \geq 0.09\frac{a-b}{k},$$

for any sufficiently large  $n$ .

Applying Chernoff's bound, we obtain

$$\begin{aligned} \rho_1 &\leq \exp\left(-\frac{(0.09(a-b))^2}{k(0.9b/2 + .1a/2) + 0.09k(a-b)}\right) \\ &\leq \exp\left(-\frac{0.0324(a-b)^2}{k(a+b)}\right). \end{aligned}$$

By a similar argument, we get the same bound for

$$\rho_2 := \mathbb{P}\left(S'_i \leq \frac{a+b}{8k}\right).$$

Therefore, by a union bound, we have that

$$\rho \leq k \exp\left(-\frac{0.0324(a-b)^2}{k(a+b)}\right).$$

Thus, we can conclude that

$$\mathbf{EM} \leq \frac{n}{2}k \exp\left(-0.0324 \frac{(a-b)^2}{k(a+b)}\right).$$

Applying Chernoff's with  $t := \frac{n}{2}k \exp\left(-0.0324 \frac{(a-b)^2}{k(a+b)}\right)$ , we conclude that with probability  $1 - o(1)$

$$M \leq \mathbf{EM} + t = nk \exp\left(-0.0324 \frac{(a-b)^2}{k(a+b)}\right).$$

This implies that with probability  $1 - o(1)$ , the number of mislabeled vertices in  $Y$  is bounded by

$$nk \exp\left(-0.0324 \frac{(a-b)^2}{k(a+b)}\right).$$

Set

$$\gamma := 2k \exp\left(-0.0324 \frac{(a-b)^2}{k(a+b)}\right).$$

We have, with probability  $1 - o(1)$ ,  $\gamma$  correct partition of the vertices in  $Y$ , with  $\gamma$  satisfying

$$\frac{(a-b)^2}{k(a+b)} = \frac{1}{0.0324} \log \frac{2k}{\gamma} \leq 31 \log \frac{2k}{\gamma},$$

proving our claim.

### Appendix C. Censor Block Model

All we have to do now is to bound  $\|E\|$ . Let  $\sigma^2 := p \geq \text{Var}(\zeta_{i,j})$  for all  $(i, j)$ .  $Y_0$  is obtained by zeroing out rows and columns of  $Y$  of high degree. We then have the following lemma. The proof is essentially the same as corollary 13, so we skip the details.

**Lemma 33**  $0 < \epsilon_0 \leq \epsilon < \frac{1}{2}$ . Then there exist constants  $C, C_1$  such that if  $p \geq \frac{C}{n}$ , then with probability  $1 - o(1)$ ,  $\|Y_0 - \bar{Y}\| \leq C_1 \sigma \sqrt{n} = C_1 \sqrt{np}$ .

Since the second eigenvalue of  $\bar{Y}$  is  $p(1 - 2\epsilon)n$ , to make the angle between the eigenspace spanned by the two eigenvectors corresponding to the top two eigenvalues small, we need to assume

$$\frac{p(1 - 2\epsilon)n}{\sqrt{np}}$$

is sufficiently large. The assumption

$$np \geq \frac{C_2}{(1 - 2\epsilon)^2}$$

in theorem 6 is precisely this.

## Appendix D. Proof of Lemma 21

This proof is essentially same as that in [Feige and Ofek \(2005\)](#). Let us first define the following sets. For  $\gamma_k := 2^k$ ,

$$S_k := \left\{ i : \frac{\gamma_{k-1}}{\sqrt{n}} < x_i \leq \frac{\gamma_k}{\sqrt{n}} \right\}, s_k := |S_k|, k = \lfloor \log \frac{\sqrt{d}}{n} \rfloor, \dots, 0, 1, 2, \dots, \lceil \log \sqrt{n} \rceil$$

and

$$T_k := \left\{ i : \frac{\gamma_{k-1}}{\sqrt{n}} < y_i \leq \frac{\gamma_k}{\sqrt{n}} \right\}, t_k := |T_k|, k = \lfloor \log \frac{\sqrt{d}}{n} \rfloor, \dots, 0, 1, 2, \dots, \lceil \log \sqrt{n} \rceil.$$

Further, we use the notation  $\mu_{i,j} := s_i t_j \frac{d}{n}$  and  $\lambda_{i,j} := e(S_i, T_j) / \mu_{i,j}$ . We then have

$$\begin{aligned} \sum_H x_i \tilde{A}_{i,j} y_j &\leq \sum_{i,j:\gamma_i \gamma_j \geq \sqrt{d}} s_i t_j \frac{d}{n} \lambda_{i,j} \frac{\gamma_i}{\sqrt{n}} \frac{\gamma_j}{\sqrt{n}} \\ &= \sqrt{d} \sum_{i,j:\gamma_i \gamma_j \geq \sqrt{d}} s_i \frac{\gamma_i^2}{n} t_j \frac{\gamma_j^2}{n} \frac{\lambda_{i,j} \sqrt{d}}{\gamma_i \gamma_j} \\ &= \sqrt{d} \sum_{i,j:\gamma_i \gamma_j \geq \sqrt{d}} \alpha_i \beta_j \sigma_{i,j}. \end{aligned}$$

In the last line, we have used the following notation  $\alpha_i := s_i \frac{\gamma_i^2}{n}$ ,  $\beta_j := t_j \frac{\gamma_j^2}{n}$ ,  $\sigma_{i,j} := \frac{\lambda_{i,j} \sqrt{d}}{\gamma_i \gamma_j}$ . In this notation, we can write 2 as follows:

$$\sigma_{i,j} \alpha_i \log \lambda_{i,j} \leq c_3 \frac{\gamma_i}{\gamma_j \sqrt{d}} \left[ 2 \log \gamma_j + \log \frac{1}{\beta_j} \right]. \quad (5)$$

Now we bound  $\sum_{i,j:\gamma_i \gamma_j \geq \sqrt{d}} \alpha_i \beta_j \sigma_{i,j}$  by a constant. We note that  $\sum_i \alpha_i \leq 4$  and  $\sum_i \beta_i \leq 4$ . We now consider 6 cases.

1.  $\sigma_{i,j} \leq 1$ :

$$\begin{aligned} \sqrt{d} \sum_{i,j:\gamma_i \gamma_j \geq \sqrt{d}} \alpha_i \beta_j \sigma_{i,j} &\leq \sum_{i,j:\gamma_i \gamma_j \geq \sqrt{d}} \alpha_i \beta_j \\ &\leq \sqrt{d} \left( \sum_i \alpha_i \right) \left( \sum_i \beta_i \right) \\ &\leq 16 \sqrt{d}. \end{aligned}$$

2.  $\lambda_{i,j} \leq c_2$ : Since  $\gamma_i \gamma_j \geq \sqrt{d}$  we have in this case  $\sigma_{i,j} \leq c_2$ . Therefore,

$$\begin{aligned} \sqrt{d} \sum_{i,j:\gamma_i \gamma_j \geq \sqrt{d}} \alpha_i \beta_j \sigma_{i,j} &\leq \sum_{i,j:\gamma_i \gamma_j \geq \sqrt{d}} \alpha_i \beta_j c_2 \\ &\leq c_2 \sqrt{d} \left( \sum_i \alpha_i \right) \left( \sum_i \beta_i \right) \\ &\leq 16 c_2 \sqrt{d}. \end{aligned}$$

3.  $\gamma_i > \sqrt{d}\gamma_j$  : Since the maximum degree is  $\leq c_1 d$ , we have that  $\lambda_{i,j} \leq c_1 n/t_j$ . Therefore,

$$\begin{aligned}
 \sqrt{d} \sum_{i,j:\gamma_i\gamma_j \geq \sqrt{d}} \alpha_i \beta_j \sigma_{i,j} &= \sqrt{d} \sum_i \left( \alpha_i \sum_{j:\gamma_i\gamma_j \geq \sqrt{d}} \beta_j \frac{\lambda_{i,j}\sqrt{d}}{\gamma_i\gamma_j} \right) \\
 &\leq \sqrt{d} \sum_i \left( \alpha_i \sum_{j:\gamma_i\gamma_j \geq \sqrt{d}} b_j \frac{\gamma_j^2 (c_1 n/b_j)\sqrt{d}}{n \gamma_i\gamma_j} \right) \\
 &= \sqrt{d} \sum_i \left( \alpha_i \sum_{j:\gamma_i\gamma_j \geq \sqrt{d}} c_1 \sqrt{d} \frac{\gamma_j}{\gamma_i} \right) \\
 &\leq \sqrt{d} \sum_i (\alpha_i c_1 \times 2) \\
 &\leq 2c_1 \sqrt{d} \sum_i \alpha_i \\
 &\leq 8c_1 \sqrt{d}.
 \end{aligned}$$

4. We now assume that we are **not** in cases 1 – 3. Therefore, we can assume that 5 holds. We consider the following sub cases.

(a)  $\log \lambda_{i,j} > (1/4)[2 \log \gamma_j + \log(1/\beta_j)]$  : 5 implies that  $\sigma_{i,j} \alpha_i \leq 4c_3 (\gamma_i/\gamma_j \sqrt{d})$ . Therefore,

$$\begin{aligned}
 \sqrt{d} \sum_{i,j:\gamma_i\gamma_j \geq \sqrt{d}} \alpha_i \beta_j \sigma_{i,j} &= \sqrt{d} \sum_j \beta_j \sum_{i:\gamma_i\gamma_j \geq \sqrt{d}} \alpha_i \sigma_{i,j} \\
 &\leq \sqrt{d} \sum_j \beta_j \sum_{i:\gamma_i\gamma_j \geq \sqrt{d}} 4c_3 \frac{\gamma_i}{\gamma_j \sqrt{d}} \\
 &\leq \sqrt{d} \sum_j \beta_j \times 8c_3 \\
 &\leq 32c_3 \sqrt{d}.
 \end{aligned}$$

Above we made use of the fact that we are not in case 3, and that  $\sum_{i:\gamma_i\gamma_j \geq \sqrt{d}} 4c_3 \frac{\gamma_i}{\gamma_j \sqrt{d}}$  is a geometric sum.

(b)  $2 \log \gamma_j \geq \log(1/\beta_j)$  : We can assume we are not in case (a), and hence  $\lambda_{i,j} \leq \gamma_j$ . Combined with the fact that we are not in case 1, we have that  $\gamma_i \leq \sqrt{d}$ . Since we are not in case 2, we can assume that  $\log \lambda_{i,j} \geq 1$  and hence  $\sigma_{i,j} \alpha_i \leq c_3 \frac{\gamma_i}{\gamma_j \sqrt{d}} 4 \log \gamma_j$ . Therefore,

$$\begin{aligned}
 \sqrt{d} \sum_{i,j:\gamma_i\gamma_j \geq \sqrt{d}} \alpha_i \beta_j \sigma_{i,j} &= \sqrt{d} \sum_j \left( \beta_j \sum_{i:\gamma_i\gamma_j \geq \sqrt{d}} \alpha_i \frac{\lambda_{i,j}\sqrt{d}}{\gamma_i\gamma_j} \right) \\
 &\leq \sqrt{d} \sum_j \left( \beta_j \sum_{i:\gamma_i\gamma_j \geq \sqrt{d}} 4c_3 \frac{\gamma_i}{\sqrt{d}\gamma_j} \log \gamma_j \right)
 \end{aligned}$$

$$\begin{aligned}
 &\leq \sqrt{d} \left( \sum_j \beta_j 4c_3 \sum_{i:\gamma_i\gamma_j \geq \sqrt{d}} \frac{\gamma_i}{\sqrt{d}} \right) \\
 &\leq \sqrt{d} \sum_j \beta_j 4c_3 \times 2 \\
 &\leq 32c_3 \sqrt{d}.
 \end{aligned}$$

(c)  $2 \log \gamma_j \leq \log 1/\beta_j$  : Since we are not in (a) we have  $\log \lambda_{i,j} \leq \log \frac{1}{\beta_j}$ . It follows that

$$\sigma_{i,j} = \frac{\lambda_{i,j} \sqrt{d}}{\gamma_i \gamma_j} \leq \frac{1}{\beta_j} \frac{\sqrt{d}}{\gamma_i \gamma_j}.$$

Therefore:

$$\begin{aligned}
 \sqrt{d} \sum_{i,j:\gamma_i\gamma_j \geq \sqrt{d}} \alpha_i \beta_j \sigma_{i,j} &= \sqrt{d} \sum_i \left( \alpha_i \sum_{j:\gamma_i\gamma_j \geq \sqrt{d}} \beta_j \sigma_{i,j} \right) \\
 &= \sqrt{d} \sum_i \left( \alpha_i \sum_{j:\gamma_i\gamma_j \geq \sqrt{d}} \beta_j \frac{\lambda_{i,j} \sqrt{d}}{\gamma_i \gamma_j} \right)
 \end{aligned}$$



$$\begin{aligned} &\leq \sqrt{d} \sum_i \left( \alpha_i \sum_{j: \gamma_i \gamma_j \geq \sqrt{d}} \frac{\sqrt{d}}{\gamma_i \gamma_j} \right) \\ &\leq \sqrt{d} \sum_i (\alpha_i \times 2) \\ &\leq 8\sqrt{d}. \end{aligned}$$

■