

# Stochastic differential equation with jumps for multi-type continuous state and continuous time branching processes with immigration

Mátyás Barczy, Zenghu Li and Gyula Pap

Faculty of Informatics, University of Debrecen  
Pf. 12, H-4010 Debrecen, Hungary.

*E-mail address:* [barczy.matyas@inf.unideb.hu](mailto:barczy.matyas@inf.unideb.hu)

*URL:* [http://www.inf.unideb.hu/valseg/dolgozok/barczy/barczy\\_angol.html](http://www.inf.unideb.hu/valseg/dolgozok/barczy/barczy_angol.html)

School of Mathematical Sciences, Beijing Normal University  
Beijing 100875, People's Republic of China.

*E-mail address:* [lizh@bnu.edu.cn](mailto:lizh@bnu.edu.cn)

*URL:* <http://math.bnu.edu.cn/~lizh/engpage.htm>

Bolyai Institute, University of Szeged  
Aradi vértanúk tere 1, H-6720 Szeged, Hungary.

*E-mail address:* [papgy@math.u-szeged.hu](mailto:papgy@math.u-szeged.hu)

*URL:* [http://www.math.u-szeged.hu/~papgy/index\\_en.html](http://www.math.u-szeged.hu/~papgy/index_en.html)

**Abstract.** A multi-type continuous state and continuous time branching process with immigration satisfying some moment conditions is identified as a pathwise unique strong solution of certain stochastic differential equation with jumps.

## 1. Introduction

Continuous state and continuous time branching processes with immigration (CBI processes) arise as high density limits of Galton–Watson branching processes with immigration, see, e.g., [Li \(2011, Theorem 3.43\)](#) without immigration and [Li \(2006\)](#) with immigration. A single-type continuous state and continuous time branching process (CB process) is a non-negative Markov process with a branching property. This class of processes has been first introduced by [Jiřina \(1958\)](#) both in discrete and continuous times. As a generalization of CB processes, [Kawazu and](#)

---

*Received by the editors March 2, 2014; accepted January 12, 2015.*

*2010 Mathematics Subject Classification.* 60H10, 60J80.

*Key words and phrases.* Multi-type continuous state and continuous time branching process with immigration.

The research of M. Barczy and G. Pap was realized in the frames of TÁMOP 4.2.4. A/2-11-1-2012-0001 „National Excellence Program – Elaborating and operating an inland student and researcher personal support system”. The project was subsidized by the European Union and co-financed by the European Social Fund. Z. Li has been partially supported by NSFC under Grant No. 11131003 and 973 Program under Grant No. 2011CB808001.

Watanabe (1971) introduced the more general class of CBI processes, where immigrants may come from outer sources. They defined a single-type CBI process as an  $[0, \infty]$ -valued Markov process with  $\infty$  as a trap in terms of Laplace transforms, see Kawazu and Watanabe (1971, Definition 1.1). An analytic characterization of CBI processes was also presented by giving the explicit form of the corresponding non-negative strongly continuous contraction semigroup, see Kawazu and Watanabe (1971, Theorem 1.1'). Further, limit theorems for Galton-Watson branching processes with immigration towards CBI processes were also investigated, see Kawazu and Watanabe (1971, Section 2). Dawson and Li (2006, Theorems 5.1 and 5.2) proved that a general single-type CBI process is the pathwise unique strong solution of a stochastic differential equation (SDE) with jumps driven by Wiener processes and Poisson random measures. Watanabe (1969, Definition 1.1) introduced two-type CB processes as  $[0, \infty)^2$ -valued Markov processes satisfying a branching property. He characterized them in an analytic way by giving the explicit form of the infinitesimal generator of the corresponding non-negative strongly continuous contraction semigroup, see Watanabe (1969, Theorem 1). Fittipaldi and Fontbona (2012, Theorem 2.1) represented a (sub)-critical continuous time and continuous state branching process conditioned to never be extinct as a pathwise unique strong solution of an appropriate SDE with jumps. It was also shown that a two-type *diffusion* CB process can be obtained as a pathwise unique strong solution of an SDE (without jumps), see Watanabe (1969, Theorem 3). Recently, for a *special* two-type (not necessarily diffusion) CBI process (with a *special* immigration mechanism), an SDE with jumps (a special case of the SDE (3.2) given later on) has already been presented by Ma (2013, Theorem 2.1) together with the existence of a pathwise unique  $[0, \infty)^2$ -valued strong solution of this SDE. For a comparison of our results with those of Ma (2013), see Section 5.

The aim of the present paper is to derive and study an SDE with jumps for a general multi-type CBI process. Next, we give an overview of the structure of the paper by explaining some of its technical merits and including some sort of preview of the types of results which are proved.

In Section 2 we recall some facts about CBI processes (e.g., set of admissible parameters, infinitesimal generator) with special emphasis on their identification (under some moment conditions) as special immigration superprocesses. This identification turns out to be very important since it is the starting point for deriving a formula for the expectation and an SDE with jumps for a general multi-type CBI process (see the proofs of Lemma 3.4 and Theorem 3.7).

In Section 3 we formulate an SDE with jumps and, under the same moment conditions, we prove that this SDE admits an  $[0, \infty)^d$ -valued weak solution which is unique in the sense of probability law among  $[0, \infty)^d$ -valued weak solutions. The idea behind of deriving such an SDE goes back to a result of Li (2011, Theorem 9.18) that an immigration superprocess can be represented as a sum of a continuous local martingale, a purely discontinuous local martingale and a drift term. In our special case, this purely discontinuous local martingale takes the form  $\int_0^t \int_{[0, \infty)^d \setminus \{\mathbf{0}\}} \mathbf{z} \tilde{N}_0(ds, d\mathbf{z})$ ,  $t \geq 0$ , with some (not necessarily Poisson) random measure  $N_0(ds, d\mathbf{z})$  on  $(0, \infty) \times ([0, \infty)^d \setminus \{\mathbf{0}\})$ , where  $\tilde{N}_0(ds, d\mathbf{z})$  denotes the compensation of  $N_0(ds, d\mathbf{z})$ . The next key step is that the integral  $\int_0^t \int_{[0, \infty)^d \setminus \{\mathbf{0}\}} \mathbf{z} \tilde{N}_0(ds, d\mathbf{z})$  can be rewritten as an appropriate sum of integrals with respect to a Poisson and compensated Poisson random measures, and some

additional drift term, due to a representation theorem of right continuous martingales, see, e.g., Ikeda and Watanabe (1989, Chapter II, Definition 1.3 and Lemma 1.2). We also prove that any  $[0, \infty)^d$ -valued weak solution of this SDE is a CBI process, see Theorem 3.7. For the proof of Theorem 3.7, we need a formula for the first moment of a CBI process, see Lemma 3.4. The proof of Lemma 3.4 is based on a formula for expectation of immigration superprocesses, see Li (2011, Proposition 9.11).

In Section 4 we prove that, under the same moment conditions, there is a pathwise unique  $[0, \infty)^d$ -valued strong solution to the SDE (3.2) and the solution is a CBI process, see Theorem 4.6. For the proof, we need a comparison theorem for the SDE (3.2) (see, Lemma 4.2), which, in particular, yields that pathwise uniqueness holds for the SDE (3.2) among  $[0, \infty)^d$ -valued weak solutions. The ideas of the proof of Lemma 4.2 follow those of Theorem 3.1 of Ma (2013), which are adaptations of those of Theorem 5.5 of Fu and Li (2010). More precisely, we derive an upper bound for an appropriate deterministic function of the difference of two  $[0, \infty)^d$ -valued weak solutions of the SDE (3.2) and then apply Gronwall's inequality.

In Section 5 we specialize our SDE (3.2) to dimension 1 and 2, respectively, which enables us to compare our results with those of Dawson and Li (2006, Theorems 5.1 and 5.2) (single-type) and Ma (2013, Theorem 2.1) (two-type), respectively. Moreover, we discuss a special case of the SDE (3.2) with  $\nu = 0$ ,  $\mu_i = 0$ ,  $i \in \{1, \dots, d\}$ , i.e., without integrals with respect to (compensated) Poisson random measures (corresponding to the so-called multi-factor Cox-Ingersoll-Ross process if  $\mathbf{B}$  is diagonal, see, e.g., Jagannathan et al. (2003)), and another special case with  $\mathbf{c} = \mathbf{0}$ , i.e., without integral with respect to a Wiener process.

In Appendix A we present some facts about extensions of probability spaces.

Finally, we mention that our work goes beyond that of Ma (2013) in the sense that we consider general multi-type CBI processes with arbitrary branching and immigration mechanisms instead of two-type CBI processes with a special immigration mechanism, and we carefully present some missing details in the proofs of Ma (2013) for the general multi-type case such as the application of Theorem 9.18 in Li (2011) and of Theorem 7.4 in Chapter II in Ikeda and Watanabe (1989). Further, in a companion paper we established Yamada-Watanabe type results for SDEs with jumps that are needed in the proof of Theorem 4.6 (existence of pathwise unique strong solution of the SDE (3.2)). We point out that Ma (2013) implicitly used these results without proving or referring to them.

## 2. Multi-type CBI processes

Let  $\mathbb{Z}_+$ ,  $\mathbb{N}$ ,  $\mathbb{R}$ ,  $\mathbb{R}_+$  and  $\mathbb{R}_{++}$  denote the set of non-negative integers, positive integers, real numbers, non-negative real numbers and positive real numbers, respectively. For  $x, y \in \mathbb{R}$ , we will use the notations  $x \wedge y := \min\{x, y\}$  and  $x^+ := \max\{0, x\}$ . By  $\|\mathbf{x}\|$  and  $\|\mathbf{A}\|$ , we denote the Euclidean norm of a vector  $\mathbf{x} \in \mathbb{R}^d$  and the induced matrix norm of a matrix  $\mathbf{A} \in \mathbb{R}^{d \times d}$ , respectively. The natural basis in  $\mathbb{R}^d$  and the Borel  $\sigma$ -algebras on  $\mathbb{R}^d$  and on  $\mathbb{R}_+^d$  will be denoted by  $\mathbf{e}_1, \dots, \mathbf{e}_d$ , and by  $\mathcal{B}(\mathbb{R}^d)$  and  $\mathcal{B}(\mathbb{R}_+^d)$ , respectively. The  $d$ -dimensional unit matrix is denoted by  $\mathbf{I}_d$ . For  $\mathbf{x} = (x_i)_{i \in \{1, \dots, d\}} \in \mathbb{R}^d$  and  $\mathbf{y} = (y_i)_{i \in \{1, \dots, d\}} \in \mathbb{R}^d$ , we will use the notation  $\mathbf{x} \leq \mathbf{y}$  indicating that  $x_i \leq y_i$  for all  $i \in \{1, \dots, d\}$ .

By  $C_c^2(\mathbb{R}_+^d, \mathbb{R})$  we denote the set of twice continuously differentiable real-valued functions on  $\mathbb{R}_+^d$  with compact support. Throughout this paper, we make the conventions  $\int_a^b := \int_{(a,b]}$  and  $\int_a^\infty := \int_{(a,\infty)}$  for any  $a, b \in \mathbb{R}$  with  $a < b$ .

**Definition 2.1.** A matrix  $\mathbf{A} = (a_{i,j})_{i,j \in \{1, \dots, d\}} \in \mathbb{R}^{d \times d}$  is called essentially non-negative if  $a_{i,j} \in \mathbb{R}_+$  whenever  $i, j \in \{1, \dots, d\}$  with  $i \neq j$ , i.e., if  $\mathbf{A}$  has non-negative off-diagonal entries. The set of essentially non-negative  $d \times d$  matrices will be denoted by  $\mathbb{R}_{(+)}^{d \times d}$ .

**Definition 2.2.** A tuple  $(d, \mathbf{c}, \boldsymbol{\beta}, \mathbf{B}, \nu, \boldsymbol{\mu})$  is called a set of admissible parameters if

- (i)  $d \in \mathbb{N}$ ,
- (ii)  $\mathbf{c} = (c_i)_{i \in \{1, \dots, d\}} \in \mathbb{R}_+^d$ ,
- (iii)  $\boldsymbol{\beta} = (\beta_i)_{i \in \{1, \dots, d\}} \in \mathbb{R}_+^d$ ,
- (iv)  $\mathbf{B} = (b_{i,j})_{i,j \in \{1, \dots, d\}} \in \mathbb{R}_{(+)}^{d \times d}$ ,
- (v)  $\nu$  is a Borel measure on  $U_d := \mathbb{R}_+^d \setminus \{\mathbf{0}\}$  satisfying  $\int_{U_d} (1 \wedge \|\mathbf{z}\|) \nu(d\mathbf{z}) < \infty$ ,
- (vi)  $\boldsymbol{\mu} = (\mu_1, \dots, \mu_d)$ , where, for each  $i \in \{1, \dots, d\}$ ,  $\mu_i$  is a Borel measure on  $U_d$  satisfying

$$\int_{U_d} \left[ \|\mathbf{z}\| \wedge \|\mathbf{z}\|^2 + \sum_{j \in \{1, \dots, d\} \setminus \{i\}} z_j \right] \mu_i(d\mathbf{z}) < \infty. \quad (2.1)$$

*Remark 2.3.* Our Definition 2.2 of the set of admissible parameters is a special case of Definition 2.6 in Duffie et al. (2003), which is suitable for all affine processes. Namely, one should take  $m = d$ ,  $n = 0$  and zero killing rate in Definition 2.6 in Duffie et al. (2003) noting also that part (v) of our Definition 2.2 is equivalent to the corresponding one  $\int_{U_d} \sum_{i=1}^d (1 \wedge z_i) \nu(d\mathbf{z}) < \infty$  in Definition 2.6 in Duffie et al. (2003). Indeed,

$$1 \wedge \|\mathbf{z}\| \leq 1 \wedge \left( \sum_{i=1}^d z_i \right) \leq \sum_{i=1}^d (1 \wedge z_i) \leq d(1 \wedge \|\mathbf{z}\|)$$

for all  $\mathbf{z} = (z_1, \dots, z_d) \in \mathbb{R}_+^d$ . Further, for all  $i \in \{1, \dots, d\}$ , the condition (2.1) is equivalent to

$$\int_{U_d} \left[ (1 \wedge z_i)^2 + \sum_{j \in \{1, \dots, d\} \setminus \{i\}} (1 \wedge z_j) \right] \mu_i(d\mathbf{z}) < \infty \quad (2.2)$$

and  $\int_{U_d} \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_i(d\mathbf{z}) < \infty.$

Indeed, if (2.1) holds, then for all  $i \in \{1, \dots, d\}$ , we have  $\int_{U_d} \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_i(d\mathbf{z}) = \int_{U_d} (\|\mathbf{z}\| \wedge \|\mathbf{z}\|^2) \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_i(d\mathbf{z}) < \infty$ , and using that  $z_i \leq \|\mathbf{z}\|$  and  $(1 \wedge z_i)^2 = (1 \wedge z_i)^2 \mathbb{1}_{\{\|\mathbf{z}\| \leq 1\}} + (1 \wedge z_i)^2 \mathbb{1}_{\{\|\mathbf{z}\| > 1\}} \leq \|\mathbf{z}\|^2 \mathbb{1}_{\{\|\mathbf{z}\| \leq 1\}} + \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| > 1\}} = \|\mathbf{z}\| \wedge \|\mathbf{z}\|^2$ ,  $i \in \{1, \dots, d\}$ , we have (2.2). If (2.2) holds, then, using again  $z_j \leq \|\mathbf{z}\|$ ,  $j \in$

$\{1, \dots, d\}$ , we have

$$\begin{aligned}
& \int_{U_d} \left[ \|\mathbf{z}\| \wedge \|\mathbf{z}\|^2 + \sum_{j \in \{1, \dots, d\} \setminus \{i\}} z_j \right] \mu_i(d\mathbf{z}) \\
&= \int_{U_d} \left[ \|\mathbf{z}\|^2 + \sum_{j \in \{1, \dots, d\} \setminus \{i\}} z_j \right] \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \mu_i(d\mathbf{z}) \\
&\quad + \int_{U_d} \left[ \|\mathbf{z}\| + \sum_{j \in \{1, \dots, d\} \setminus \{i\}} z_j \right] \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_i(d\mathbf{z}) \\
&\leq \int_{U_d} \left[ z_i^2 + 2 \sum_{j \in \{1, \dots, d\} \setminus \{i\}} z_j \right] \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \mu_i(d\mathbf{z}) + \int_{U_d} \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_i(d\mathbf{z}) \\
&\quad + \sum_{j \in \{1, \dots, d\} \setminus \{i\}} \int_{U_d} \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_i(d\mathbf{z}) < \infty, \quad i \in \{1, \dots, d\},
\end{aligned}$$

yielding (2.1). Note that, here the finiteness of the first integral in (2.2) is nothing else but condition (2.11) in Definition 2.6 in Duffie et al. (2003), and the finiteness of the second integral in (2.2) is an additional condition that we assume compared to Duffie et al. (2003), its role is explained in Remark 2.5.  $\square$

**Theorem 2.4.** *Let  $(d, \mathbf{c}, \boldsymbol{\beta}, \mathbf{B}, \nu, \boldsymbol{\mu})$  be a set of admissible parameters in the sense of Definition 2.2. Then there exists a unique conservative transition semigroup  $(P_t)_{t \in \mathbb{R}_+}$  acting on the Banach space (endowed with the supremum norm) of real-valued bounded Borel-measurable functions on the state space  $\mathbb{R}_+^d$  such that its infinitesimal generator is*

$$\begin{aligned}
(\mathcal{A}f)(\mathbf{x}) &= \sum_{i=1}^d c_i x_i f''_{i,i}(\mathbf{x}) + \langle \boldsymbol{\beta} + \mathbf{B}\mathbf{x}, \mathbf{f}'(\mathbf{x}) \rangle + \int_{U_d} (f(\mathbf{x} + \mathbf{z}) - f(\mathbf{x})) \nu(d\mathbf{z}) \\
&\quad + \sum_{i=1}^d x_i \int_{U_d} (f(\mathbf{x} + \mathbf{z}) - f(\mathbf{x}) - f'_i(\mathbf{x})(1 \wedge z_i)) \mu_i(d\mathbf{z})
\end{aligned} \tag{2.3}$$

for  $f \in C_c^2(\mathbb{R}_+^d, \mathbb{R})$  and  $\mathbf{x} \in \mathbb{R}_+^d$ , where  $f'_i$  and  $f''_{i,i}$ ,  $i \in \{1, \dots, d\}$ , denote the first and second order partial derivatives of  $f$  with respect to its  $i$ -th variable, respectively, and  $\mathbf{f}'(\mathbf{x}) := (f'_1(\mathbf{x}), \dots, f'_d(\mathbf{x}))^\top$ . Moreover, the Laplace transform of the transition semigroup  $(P_t)_{t \in \mathbb{R}_+}$  has a representation

$$\int_{\mathbb{R}_+^d} e^{-\langle \boldsymbol{\lambda}, \mathbf{y} \rangle} P_t(\mathbf{x}, d\mathbf{y}) = e^{-\langle \mathbf{x}, \mathbf{v}(t, \boldsymbol{\lambda}) \rangle - \int_0^t \psi(\mathbf{v}(s, \boldsymbol{\lambda})) ds}, \quad \mathbf{x} \in \mathbb{R}_+^d, \quad \boldsymbol{\lambda} \in \mathbb{R}_+^d, \quad t \in \mathbb{R}_+,$$

where, for any  $\boldsymbol{\lambda} \in \mathbb{R}_+^d$ , the continuously differentiable function  $\mathbb{R}_+ \ni t \mapsto \mathbf{v}(t, \boldsymbol{\lambda}) = (v_1(t, \boldsymbol{\lambda}), \dots, v_d(t, \boldsymbol{\lambda}))^\top \in \mathbb{R}_+^d$  is the unique locally bounded solution to the system of differential equations

$$\partial_t v_i(t, \boldsymbol{\lambda}) = -\varphi_i(\mathbf{v}(t, \boldsymbol{\lambda})), \quad v_i(0, \boldsymbol{\lambda}) = \lambda_i, \quad i \in \{1, \dots, d\}, \tag{2.4}$$

with

$$\varphi_i(\boldsymbol{\lambda}) := c_i \lambda_i^2 - \langle \mathbf{B}\mathbf{e}_i, \boldsymbol{\lambda} \rangle + \int_{U_d} (e^{-\langle \boldsymbol{\lambda}, \mathbf{z} \rangle} - 1 + \lambda_i(1 \wedge z_i)) \mu_i(d\mathbf{z})$$

for  $\lambda \in \mathbb{R}_+^d$  and  $i \in \{1, \dots, d\}$ , and

$$\psi(\lambda) := \langle \beta, \lambda \rangle - \int_{U_d} (e^{-\langle \lambda, z \rangle} - 1) \nu(dz), \quad \lambda \in \mathbb{R}_+^d.$$

Further, the function  $\mathbb{R}_+ \times \mathbb{R}_+^d \ni (t, \lambda) \mapsto \mathbf{v}(t, \lambda)$  is continuous.

*Remark 2.5.* This theorem is a special case of Theorem 2.7 of Duffie et al. (2003) with  $m = d$ ,  $n = 0$  and zero killing rate. The unique existence of a locally bounded solution to the system of differential equations (2.4) is proved by Li (2011, page 45). Here, we point out that the moment condition given in part (vi) in Definition 2.2 (which is stronger than the one (2.11) in Definition 2.6 in Duffie et al. (2003)) ensures that the semigroup  $(P_t)_{t \in \mathbb{R}_+}$  is conservative (we do not need the one-point compactification of  $\mathbb{R}_+^d$ ), see Duffie et al. (2003, Lemma 9.2) and Li (2011, page 45). For the continuity of the function  $\mathbb{R}_+ \times \mathbb{R}_+^d \ni (t, \lambda) \mapsto \mathbf{v}(t, \lambda)$ , see Duffie et al. (2003, Proposition 6.4). Finally, we note that the infinitesimal generator (2.3) can be rewritten in another equivalent form, see formula (2.14) in Lemma 2.11.  $\square$

**Definition 2.6.** A conservative Markov process with state space  $\mathbb{R}_+^d$  and with transition semigroup  $(P_t)_{t \in \mathbb{R}_+}$  given in Theorem 2.4 is called a multi-type CBI process with parameters  $(d, c, \beta, \mathbf{B}, \nu, \mu)$ .

In what follows, we will identify a multi-type CBI process  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  with parameters  $(d, c, \beta, \mathbf{B}, \nu, \mu)$  under a moment condition as a special immigration superprocess. First we parametrize the family of immigration superprocesses for which Theorem 9.18 in Li (2011) is valid. We will use some notations of the book of Li (2011). For a locally compact separable metric space  $E$ , let us introduce the following function spaces:

- $B(E)$  is the space of bounded real-valued Borel functions on  $E$ ,
- $B(E)^+$  is the space of bounded non-negative real-valued Borel functions on  $E$ ,
- $C(E)$  is the space of bounded continuous real-valued functions on  $E$ ,
- $C(E)^+$  is the space of bounded continuous non-negative real-valued functions on  $E$ ,
- $C_0(E)$  is the space of continuous real-valued functions on  $E$  vanishing at infinity.

Let  $M(E)$  denote the space of finite Borel measures on  $E$ . We write  $\mu(f) := \int_E f(x) \mu(dx)$  for the integral of a function  $f : E \rightarrow \mathbb{R}$  with respect to a measure  $\mu \in M(E)$  if the integral exists.

**Definition 2.7.** A tuple  $(E, (R_t)_{t \in \mathbb{R}_+}, c, \beta, b, B, H_1, H_2)$  is called a set of admissible parameters if

- (i)  $E$  is a locally compact separable metric space,
- (ii)  $(R_t)_{t \in \mathbb{R}_+}$  is the transition semigroup of a Hunt process

$$\xi = (\Omega, \mathcal{G}, (\mathcal{G}_t)_{t \in \mathbb{R}_+}, (\xi_t)_{t \in \mathbb{R}_+}, (\theta_t)_{t \in \mathbb{R}_+}, (\mathbb{P}_x)_{x \in E})$$

with values in  $E$  (see, e.g., Li (2011, page 314)) such that  $(R_t)_{t \in \mathbb{R}_+}$  preserves  $C_0(E)$ , and  $\mathbb{R}_+ \ni t \mapsto R_t f \in C_0(E)$  is continuous in the supremum norm for every  $f \in C_0(E)$ ,

- (iii)  $c \in C(E)^+$ ,
- (iv)  $\beta \in M(E)$ ,

- (v)  $b \in C(E)$ ,
- (vi)  $H_1$  is a finite measure on  $M(E)^\circ := M(E) \setminus \{0\}$  (where  $0$  denotes the null measure) satisfying  $\int_{M(E)^\circ} \kappa(1) H_1(d\kappa) < \infty$ ,
- (vii)  $B(x, dy)$  is a bounded kernel on  $E$  (i.e., from  $E$  to  $E$ ) and  $H_2(x, d\kappa)$  is a  $\sigma$ -finite kernel from  $E$  to  $M(E)^\circ$  such that  $E \ni x \mapsto (\kappa(1) \wedge \kappa(1)^2) H_2(x, d\kappa)$  is continuous with respect to the topology of weak convergence in  $M(E)^\circ$ , and the operators

$$f \mapsto \int_{M(E)^\circ} (\kappa(f) \wedge \kappa(f)^2) H_2(\cdot, d\kappa) \quad \text{and} \quad f \mapsto \gamma(\cdot, f)$$

preserve  $C_0(E)^+$ , where the kernel  $\gamma(x, dy)$  on  $E$  is defined by

$$\gamma(x, dy) := B(x, dy) + \int_{M(E)^\circ} \kappa_x(dy) H_2(x, d\kappa),$$

where  $\kappa_x(dy)$  denotes the restriction of  $\kappa(dy)$  to  $E \setminus \{x\}$ , and by  $\gamma(\cdot, f)$  we mean the function  $E \ni x \mapsto \gamma(x, f) := \int_E f(y) \gamma(x, dy)$ .

*Remark 2.8.* Note that Condition (2.25) in Li (2011) readily follows from (vii) of Definition 2.7, since a function in  $C_0(E)$  is bounded, hence

$$\begin{aligned} \sup_{x \in E} \int_{M(E)^\circ} [\kappa(1) \wedge \kappa(1)^2] H_2(x, d\kappa) &< \infty, \\ \sup_{x \in E} \int_{M(E)^\circ} \kappa_x(1) H_2(x, d\kappa) &\leq \sup_{x \in E} \gamma(x, 1) < \infty, \end{aligned}$$

where we used that  $B(x, 1) \in \mathbb{R}_+$  for all  $x \in E$ .  $\square$

**Theorem 2.9.** *Let  $(E, (R_t)_{t \in \mathbb{R}_+}, c, \beta, b, B, H_1, H_2)$  be a set of admissible parameters in the sense of Definition 2.7. Then there exists a unique transition semigroup  $(Q_t)_{t \in \mathbb{R}_+}$  acting on the Banach space (endowed with the supremum norm) of real-valued bounded Borel-measurable functions on the state space  $M(E)$  such that its infinitesimal generator is*

$$\begin{aligned} (\mathcal{A}F)(\mu) &= \int_E c(x) F''(\mu; x) \mu(dx) \\ &+ \int_E (AF'(\mu; x) + \gamma(x, F'(\mu; \cdot)) - b(x) F'(\mu; x)) \mu(dx) \\ &+ \int_E F'(\mu; x) \beta(dx) + \int_{M(E)^\circ} (F(\mu + \kappa) - F(\mu)) H_1(d\kappa) \\ &+ \int_E \left( \int_{M(E)^\circ} (F(\mu + \kappa) - F(\mu) - \kappa(F'(\mu; \cdot))) H_2(x, d\kappa) \right) \mu(dx), \end{aligned} \tag{2.5}$$

for  $\mu \in M(E)$  and functions  $F : M(E) \rightarrow \mathbb{R}$ ,  $F(\mu) = G(\mu(f_1), \dots, \mu(f_n))$ , where  $n \in \mathbb{N}$ ,  $G \in C^2(\mathbb{R}^n, \mathbb{R})$ , and  $f_1, \dots, f_n \in D_0(A)$ , where  $A$  denotes the strong generator of  $(R_t)_{t \in \mathbb{R}_+}$  defined by

$$Af(x) := \lim_{t \downarrow 0} \frac{R_t f(x) - f(x)}{t}, \quad x \in E,$$

where the limit is taken in the supremum norm, and the domain  $D_0(A)$  of  $A$  is the totality of functions  $f \in C_0(E)$  for which the above limit exists,

$$F'(\mu; x) := \lim_{\varepsilon \downarrow 0} \frac{F(\mu + \varepsilon \delta_x) - F(\mu)}{\varepsilon}, \quad \mu \in M(E), \quad x \in E,$$

and  $F''(\mu; x)$  is defined by the limit with  $F(\cdot)$  replaced by  $F'(\cdot; x)$ .

Moreover, the Laplace transform of the transition semigroup  $(Q_t)_{t \in \mathbb{R}_+}$  has a representation

$$\int_{M(E)} e^{-\kappa(f)} Q_t(\mu, d\kappa) = e^{-\mu(V_t f) - \int_0^t I(V_s f) ds}, \quad \mu \in M(E), \quad f \in B(E)^+, \quad t \in \mathbb{R}_+, \quad (2.6)$$

where, for any  $x \in E$  and  $f \in B(E)^+$ , the continuously differentiable function  $\mathbb{R}_+ \ni t \mapsto V_t f(x) \in \mathbb{R}_+$  is the unique locally bounded solution to the integral evolution equation

$$V_t f(x) = R_t f(x) - \int_0^t \left( \int_E \phi(y, V_s f) R_{t-s}(x, dy) \right) ds, \quad t \in \mathbb{R}_+,$$

with

$$\begin{aligned} \phi(x, f) &:= c(x)f(x)^2 + b(x)f(x) - \int_E f(y) B(x, dy) \\ &\quad + \int_{M(E)^\circ} (e^{-\kappa(f)} - 1 + \kappa(\{x\})f(x)) H_2(x, d\kappa) \end{aligned}$$

for  $x \in E$  and  $f \in B(E)^+$ , and

$$I(f) := \beta(f) + \int_{M(E)^\circ} (1 - e^{-\kappa(f)}) H_1(d\kappa), \quad f \in B(E)^+.$$

*Proof:* Formula (2.6), which is, in fact, formula (9.18) in Li (2011), defines a transition semigroup of an immigration superprocess corresponding to the skew convolution semigroup given by (9.7) in Li (2011). Theorem 9.18 in Li (2011) yields that the infinitesimal generator of the immigration superprocess in question has the form given in (2.5), and the unicity of the transition semigroup.  $\square$

**Definition 2.10.** A Markov process with state space  $M(E)$  and with transition semigroup  $(Q_t)_{t \in \mathbb{R}_+}$  given in Theorem 2.9 is called an immigration superprocess with state space  $M(E)$  with parameters  $(E, (R_t)_{t \in \mathbb{R}_+}, c, \beta, b, B, H_1, H_2)$ .

In what follows, we identify a multi-type CBI process  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  with parameters  $(d, \mathbf{c}, \boldsymbol{\beta}, \mathbf{B}, \nu, \boldsymbol{\mu})$  under the moment condition

$$\int_{U_d} \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \nu(d\mathbf{z}) < \infty, \quad (2.7)$$

as a special immigration superprocess.

First, let us introduce the modified parameters  $\tilde{\boldsymbol{\beta}} := (\tilde{\beta}_i)_{i \in \{1, \dots, d\}}$ ,  $\tilde{\mathbf{B}} := (\tilde{b}_{i,j})_{i,j \in \{1, \dots, d\}}$  and  $\mathbf{D} := (d_{i,j})_{i,j \in \{1, \dots, d\}}$  given by

$$\tilde{\boldsymbol{\beta}} := \boldsymbol{\beta} + \int_{U_d} \mathbf{z} \nu(d\mathbf{z}), \quad \tilde{b}_{i,j} := b_{i,j} + \int_{U_d} (z_i - \delta_{i,j})^+ \mu_j(d\mathbf{z}), \quad (2.8)$$

$$d_{i,j} := \tilde{b}_{i,j} - \int_{U_d} z_i \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_j(d\mathbf{z}), \quad (2.9)$$



with  $\delta_{i,j} := 1$  if  $i = j$ , and  $\delta_{i,j} := 0$  if  $i \neq j$ . The moment condition (2.7) together with the fact that  $\nu$  and  $\mu$  satisfy Definition 2.2 imply  $\tilde{\beta} \in \mathbb{R}_+^d$ ,  $\tilde{B} \in \mathbb{R}_{(+)}^{d \times d}$  and  $D \in \mathbb{R}_{(+)}^{d \times d}$ . Indeed,

$$\int_{U_d} \|\mathbf{z}\| \nu(d\mathbf{z}) = \int_{U_d} (1 \wedge \|\mathbf{z}\|) \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \nu(d\mathbf{z}) + \int_{U_d} \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \nu(d\mathbf{z}) < \infty \quad (2.10)$$

by part (v) of Definition 2.2 and (2.7). Moreover, for all  $i \in \{1, \dots, d\}$ ,

$$\int_{U_d} (z_i - 1)^+ \mu_i(d\mathbf{z}) \leq \int_{U_d} z_i \mathbb{1}_{\{z_i \geq 1\}} \mu_i(d\mathbf{z}) \leq \int_{U_d} \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_i(d\mathbf{z}) < \infty \quad (2.11)$$

by  $z_i \leq \|\mathbf{z}\|$ ,  $\mathbf{z} \in \mathbb{R}_+^d$ , and (2.2). Further, for all  $i, j \in \{1, \dots, d\}$ ,  $i \neq j$ ,

$$\begin{aligned} \int_{U_d} z_i \mu_j(d\mathbf{z}) &= \int_{U_d} z_i \mathbb{1}_{\{z_i < 1\}} \mu_j(d\mathbf{z}) + \int_{U_d} z_i \mathbb{1}_{\{z_i \geq 1\}} \mu_j(d\mathbf{z}) \\ &\leq \int_{U_d} (1 \wedge z_i) \mu_j(d\mathbf{z}) + \int_{U_d} \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_j(d\mathbf{z}) < \infty \end{aligned} \quad (2.12)$$

by  $z_i \leq \|\mathbf{z}\|$ ,  $\mathbf{z} \in \mathbb{R}_+^d$ , part (vi) of Definition 2.2 and (2.2). Finally,  $d_{i,j}$  is well-defined for all  $i, j \in \{1, \dots, d\}$  because of (2.2), and, for all  $i, j \in \{1, \dots, d\}$ ,  $i \neq j$ ,

$$\begin{aligned} d_{i,j} &= b_{i,j} + \int_{U_d} z_i \mu_j(d\mathbf{z}) - \int_{U_d} z_i \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_j(d\mathbf{z}) \\ &= b_{i,j} + \int_{U_d} z_i \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \mu_j(d\mathbf{z}) \in \mathbb{R}_+. \end{aligned}$$

Note also that for all  $j \in \{1, \dots, d\}$ ,

$$\int_{U_d} \|\mathbf{z}\|^2 \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \mu_j(d\mathbf{z}) \leq \int_{U_d} \left( z_j^2 + \sum_{k \in \{1, \dots, d\} \setminus \{j\}} z_k \right) \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \mu_j(d\mathbf{z}) < \infty \quad (2.13)$$

by  $z_i \leq \|\mathbf{z}\|$ ,  $\mathbf{z} \in \mathbb{R}_+^d$ , part (vi) of Definition 2.2 and (2.2).

For the discrete metric space  $E := \{1, \dots, d\}$ , we have the following identifications:

- $B(E)$ ,  $C(E)$  and  $C_0(E)$  can be identified with  $\mathbb{R}^d$ , since a function  $f : E \rightarrow \mathbb{R}$  can be identified with the vector  $(f(1), \dots, f(d))^\top \in \mathbb{R}^d$ ,
- $B(E)^+$  and  $C(E)^+$  can be identified with  $\mathbb{R}_+^d$ ,
- $M(E)$  can be identified with  $\mathbb{R}_+^d$ , since a finite Borel measure  $\mu$  on  $E$  can be identified with the vector  $(\mu(\{1\}), \dots, \mu(\{d\}))^\top \in \mathbb{R}_+^d$ ,
- for  $\mu \in M(E)$  and  $f \in B(E)$ , the integral  $\mu(f) = \int_E f(x) \mu(dx) = \sum_{i=1}^d f(i) \mu(\{i\})$  can be identified with the usual Euclidean inner product  $\langle \mu, f \rangle$  in  $\mathbb{R}^d$ ,
- $M(E)^\circ$  can be identified with  $U_d$ .

If  $(\Omega, \mathcal{F}, \mathbb{P})$  is a probability space, then, by  $\mathbb{P}$ -null sets from a sub  $\sigma$ -algebra  $\mathcal{H} \subset \mathcal{F}$ , we mean the elements of the set

$$\{A \subset \Omega : \exists B \in \mathcal{H} \text{ such that } A \subset B \text{ and } \mathbb{P}(B) = 0\}.$$

A filtered probability space  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P})$  is said to satisfy the usual hypotheses if  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$  is right continuous and  $\mathcal{F}_0$  contains all the  $\mathbb{P}$ -null sets in  $\mathcal{F}$ .

**Lemma 2.11.** *Let  $(d, \mathbf{c}, \beta, \mathbf{B}, \nu, \boldsymbol{\mu})$  be a set of admissible parameters in the sense of Definition 2.2 satisfying the moment condition (2.7). Then*

$$(E, (R_t)_{t \in \mathbb{R}_+}, c, \beta, b, B, H_1, H_2)$$

is a set of admissible parameters in the sense of Definition 2.7, where

- (i)  $E := \{1, \dots, d\}$  with the discrete metric,
- (ii)  $(R_t)_{t \in \mathbb{R}_+}$  is the transition semigroup given by  $R_t f := f$ ,  $f \in B(E)$ ,  $t \in \mathbb{R}_+$ ,
- (iii)  $c \in B(E)^+$  is given by  $c(i) := c_i$ ,  $i \in E$ ,
- (iv)  $\beta \in M(E)$  is given by  $\beta(\{i\}) := \beta_i$ ,  $i \in E$ ,
- (v)  $b \in B(E)$ , is given by  $b(i) := -\tilde{b}_{i,i}$ ,  $i \in E$ ,
- (vi)  $B(x, dy)$  is the kernel on  $E$  given by  $B(i, \{i\}) := 0$  for  $i \in \{1, \dots, d\}$  and  $B(i, \{j\}) := b_{j,i}$  for  $i, j \in \{1, \dots, d\}$  with  $i \neq j$ ,
- (vii)  $H_1$  is the measure on  $M(E)^\circ$  identified with the measure  $\nu$  on  $U_d$ ,
- (viii)  $H_2(x, d\kappa)$  is the kernel from  $E$  to  $M(E)^\circ$  such that the measure  $H_2(i, \cdot)$  on  $M(E)^\circ$  is identified with the measure  $\mu_i$  on  $U_d$  for each  $i \in \{1, \dots, d\}$ .

If  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P})$  is a filtered probability space satisfying the usual hypotheses and  $(Y_t)_{t \in \mathbb{R}_+}$  is an  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted càdlàg immigration superprocess with parameters  $(E, (R_t)_{t \in \mathbb{R}_+}, c, \beta, b, B, H_1, H_2)$  satisfying  $\mathbb{E}(Y_0(1)) < \infty$ , then  $\mathbf{X}_t := (Y_t(\{1\}), \dots, Y_t(\{d\}))^\top$ ,  $t \in \mathbb{R}_+$ , is a multi-type CBI process with parameters  $(d, \mathbf{c}, \beta, \mathbf{B}, \nu, \boldsymbol{\mu})$  satisfying  $\mathbb{E}(\|\mathbf{X}_0\|) < \infty$ . The infinitesimal generator (2.3) of  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  can also be written in the form

$$\begin{aligned} (\mathcal{A}_{\mathbf{X}} f)(\mathbf{x}) &= \sum_{i=1}^d c_i x_i f''_{i,i}(x) + \sum_{i=1}^d x_i \int_{U_d} (f(\mathbf{x} + \mathbf{z}) - f(\mathbf{x}) - \langle \mathbf{z}, \mathbf{f}'(\mathbf{x}) \rangle) \mu_i(d\mathbf{z}) \\ &\quad + \langle \beta + \tilde{\mathbf{B}}\mathbf{x}, \mathbf{f}'(\mathbf{x}) \rangle + \int_{U_d} (f(\mathbf{x} + \mathbf{z}) - f(\mathbf{x})) \nu(d\mathbf{z}) \end{aligned} \tag{2.14}$$

for  $f \in C_c^2(\mathbb{R}_+^d, \mathbb{R})$  and  $\mathbf{x} \in \mathbb{R}_+^d$ .

*Proof:* The discrete metric space  $\{1, \dots, d\}$  is trivially a locally compact separable metric space. Clearly,  $R_t f := f$ ,  $f \in B(E)$ ,  $t \in \mathbb{R}_+$ , is the transition semigroup of the Hunt process

$$\xi = (\Omega, \mathcal{G}, (\mathcal{G}_t)_{t \in \mathbb{R}_+}, (\xi_t)_{t \in \mathbb{R}_+}, (\theta_t)_{t \in \mathbb{R}_+}, (\mathbb{P}_x)_{x \in E})$$

with  $\Omega = \{1, \dots, d\}$ ,  $\mathcal{G} = \mathcal{G}_t = 2^\Omega$ ,  $t \in \mathbb{R}_+$ ,  $\xi_t(\omega) = \theta_t(\omega) = \omega$ ,  $\omega \in \Omega$ ,  $t \in \mathbb{R}_+$ ,  $\mathbb{P}_x = \delta_x$ ,  $x \in \{1, \dots, d\}$ . Moreover,  $(R_t)_{t \in \mathbb{R}_+}$  trivially satisfies (ii) of Definition 2.7, and (iii), (iv) and (v) of Definition 2.7 trivially hold. Further (vi) of Definition 2.7 also holds, since  $\int_{U_d} \left( \sum_{i=1}^d z_i \right) \nu(d\mathbf{z}) < \infty$  follows from (2.10) by  $z_i \leq \|\mathbf{z}\|$ ,  $\mathbf{z} \in \mathbb{R}_+^d$ ,  $i \in \{1, \dots, d\}$ . The kernel  $B(x, dy)$  on  $E$  is bounded, since  $\sup_{x \in E} B(x, E) = \max_{i \in \{1, \dots, d\}} \sum_{j \in \{1, \dots, d\} \setminus \{i\}} b_{j,i} < \infty$ . On the discrete metric space  $\{1, \dots, d\}$  every function is continuous, hence  $E \ni$

$x \mapsto (\kappa(1) \wedge \kappa(1)^2)H_2(x, d\kappa)$  is continuous with respect to the topology of weak convergence in  $M(E)^\circ$ . In order to show that the operator

$$f \mapsto \int_{M(E)^\circ} (\kappa(f) \wedge \kappa(f)^2)H_2(\cdot, d\kappa)$$

preserve  $C_0(E)^+$ , it suffices to observe that for each  $\boldsymbol{\lambda} \in \mathbb{R}_+^d$  and  $i \in \{1, \dots, d\}$ , we have

$$\int_{U_d} (\langle \boldsymbol{\lambda}, \mathbf{z} \rangle \wedge \langle \boldsymbol{\lambda}, \mathbf{z} \rangle^2) \mu_i(d\mathbf{z}) \in \mathbb{R}_+,$$

which follows from the estimate

$$\begin{aligned} \int_{U_d} (\langle \boldsymbol{\lambda}, \mathbf{z} \rangle \wedge \langle \boldsymbol{\lambda}, \mathbf{z} \rangle^2) \mu_i(d\mathbf{z}) &\leq \int_{U_d} [(\|\boldsymbol{\lambda}\| \|\mathbf{z}\|) \wedge (\|\boldsymbol{\lambda}\| \|\mathbf{z}\|)^2] \mu_i(d\mathbf{z}) \\ &\leq c_\lambda \int_{U_d} (\|\mathbf{z}\| \wedge \|\mathbf{z}\|^2) \mu_i(d\mathbf{z}) \\ &= c_\lambda \int_{U_d} (\|\mathbf{z}\| \wedge \|\mathbf{z}\|^2) \mathbb{1}_{\{\|\mathbf{z}\| \leq 1\}} \mu_i(d\mathbf{z}) + c_\lambda \int_{U_d} (\|\mathbf{z}\| \wedge \|\mathbf{z}\|^2) \mathbb{1}_{\{\|\mathbf{z}\| > 1\}} \mu_i(d\mathbf{z}) \\ &= c_\lambda \int_{U_d} \|\mathbf{z}\|^2 \mathbb{1}_{\{\|\mathbf{z}\| \leq 1\}} \mu_i(d\mathbf{z}) + c_\lambda \int_{U_d} \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| > 1\}} \mu_i(d\mathbf{z}) < \infty \end{aligned}$$

with  $c_\lambda := \max\{\|\boldsymbol{\lambda}\|, \|\boldsymbol{\lambda}\|^2\}$  by (2.13) and (2.2). In order to show that the operator  $f \mapsto \gamma(\cdot, f)$  preserves  $C_0(E)^+$ , it suffices to observe that for each  $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_d)^\top \in \mathbb{R}_+^d$  and  $i \in \{1, \dots, d\}$ , we have

$$\sum_{j=1}^d \lambda_j B(i, \{j\}) + \sum_{j \in \{1, \dots, d\} \setminus \{i\}} \lambda_j \int_{U_d} z_j \mu_j(d\mathbf{z}) \in \mathbb{R}_+,$$

which follows from (2.12). Consequently,  $(E, (R_t)_{t \in \mathbb{R}_+}, c, \beta, b, B, H_1, H_2)$  is a set of admissible parameters in the sense of Definition 2.7.

By Theorem 2.9, we have

$$\mathbb{E}(e^{-Y_t(f)} | Y_0 = \mu) = \int_{M(E)^\circ} e^{-\kappa(f)} Q_t(\mu, d\kappa) = e^{-\mu(V_t f) - \int_0^t I(V_s f) ds}$$

for  $\mu \in M(E)$ ,  $f \in B(E)^+$  and  $t \in \mathbb{R}_+$ , hence we obtain

$$\mathbb{E}(e^{-\langle \boldsymbol{\lambda}, \mathbf{X}_t \rangle} | \mathbf{X}_0 = \mathbf{x}) = e^{-\langle \mathbf{x}, \mathbf{v}(t, \boldsymbol{\lambda}) \rangle - \int_0^t \psi(\mathbf{v}(s, \boldsymbol{\lambda})) ds}, \quad \mathbf{x}, \boldsymbol{\lambda} \in \mathbb{R}_+^d, \quad t \in \mathbb{R}_+,$$

where, for any  $i \in \{1, \dots, d\}$  and  $\boldsymbol{\lambda} \in \mathbb{R}_+^d$ , the function  $\mathbb{R}_+ \ni t \mapsto \mathbf{v}(t, \boldsymbol{\lambda}) = (v_1(t, \boldsymbol{\lambda}), \dots, v_d(t, \boldsymbol{\lambda}))$  is the unique locally bounded solution to the integral evolution equation

$$v_i(t, \boldsymbol{\lambda}) = \lambda_i - \int_0^t \varphi_i(\mathbf{v}(s, \boldsymbol{\lambda})) ds, \quad t \in \mathbb{R}_+, \quad i \in \{1, \dots, d\},$$

with

$$\varphi_i(\boldsymbol{\lambda}) := c_i \lambda_i^2 - \tilde{b}_{i,i} \lambda_i - \sum_{j \in \{1, \dots, d\} \setminus \{i\}} \lambda_j b_{j,i} + \int_{U_d} (e^{-\langle \boldsymbol{\lambda}, \mathbf{z} \rangle} - 1 + \lambda_i z_i) \mu_i(d\mathbf{z})$$

for  $\boldsymbol{\lambda} \in \mathbb{R}_+^d$  and  $i \in \{1, \dots, d\}$ , and

$$\psi(\boldsymbol{\lambda}) := \langle \boldsymbol{\beta}, \boldsymbol{\lambda} \rangle + \int_{U_d} (1 - e^{-\langle \boldsymbol{\lambda}, \mathbf{z} \rangle}) \nu(d\mathbf{z}), \quad \boldsymbol{\lambda} \in \mathbb{R}_+^d.$$

We have

$$\varphi_i(\boldsymbol{\lambda}) = c_i \lambda_i^2 - \langle \tilde{\mathbf{B}} \mathbf{e}_i, \boldsymbol{\lambda} \rangle + \int_{U_d} (e^{-\langle \boldsymbol{\lambda}, \mathbf{z} \rangle} - 1 + \langle \boldsymbol{\lambda}, \mathbf{z} \rangle) \mu_i(d\mathbf{z}), \quad (2.15)$$

since, by (2.12),

$$\begin{aligned} & \varphi_i(\boldsymbol{\lambda}) - c_i \lambda_i^2 + \langle \tilde{\mathbf{B}} \mathbf{e}_i, \boldsymbol{\lambda} \rangle - \int_{U_d} (e^{-\langle \boldsymbol{\lambda}, \mathbf{z} \rangle} - 1 + \langle \boldsymbol{\lambda}, \mathbf{z} \rangle) \mu_i(d\mathbf{z}) \\ &= -\tilde{b}_{i,i} \lambda_i - \sum_{j \in \{1, \dots, d\} \setminus \{i\}} \lambda_j b_{j,i} + \sum_{j=1}^d \lambda_j \tilde{b}_{j,i} - \sum_{j \in \{1, \dots, d\} \setminus \{i\}} \lambda_j \int_{U_d} z_j \mu_i(d\mathbf{z}) = 0. \end{aligned}$$

Moreover, we can write the functions  $\varphi_i$ ,  $i \in \{1, \dots, d\}$ , in the form

$$\varphi_i(\boldsymbol{\lambda}) = c_i \lambda_i^2 - \langle \mathbf{B} \mathbf{e}_i, \boldsymbol{\lambda} \rangle + \int_{U_d} (e^{-\langle \boldsymbol{\lambda}, \mathbf{z} \rangle} - 1 + \lambda_i (1 \wedge z_i)) \mu_i(d\mathbf{z})$$

for  $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_d)^\top \in \mathbb{R}_+^d$  and  $i \in \{1, \dots, d\}$ . Indeed, by (2.11) and (2.12),

$$\begin{aligned} & \varphi_i(\boldsymbol{\lambda}) - c_i \lambda_i^2 + \langle \mathbf{B} \mathbf{e}_i, \boldsymbol{\lambda} \rangle - \int_{U_d} (e^{-\langle \boldsymbol{\lambda}, \mathbf{z} \rangle} - 1 + \lambda_i (1 \wedge z_i)) \mu_i(d\mathbf{z}) \\ &= \langle (\mathbf{B} - \tilde{\mathbf{B}}) \mathbf{e}_i, \boldsymbol{\lambda} \rangle - \int_{U_d} (\lambda_i (1 \wedge z_i) - \langle \boldsymbol{\lambda}, \mathbf{z} \rangle) \mu_i(d\mathbf{z}) \\ &= -\lambda_i \int_{U_d} (z_i - 1)^+ \mu_i(d\mathbf{z}) - \int_{U_d} (\lambda_i (1 \wedge z_i) - \lambda_i z_i) \mu_i(d\mathbf{z}) = 0. \end{aligned}$$

By Theorem 2.4,  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  is a multi-type CBI with parameters  $(d, \mathbf{c}, \boldsymbol{\beta}, \mathbf{B}, \nu, \boldsymbol{\mu})$  satisfying  $\mathbb{E}(\|\mathbf{X}_0\|) < \infty$ .

Finally, (2.14) follows from

$$\begin{aligned} & (\mathcal{A}_{\mathbf{X}} f)(\mathbf{x}) - \sum_{i=1}^d c_i x_i f''_{i,i}(\mathbf{x}) - \sum_{i=1}^d x_i \int_{U_d} (f(\mathbf{x} + \mathbf{z}) - f(\mathbf{x}) - \langle \mathbf{z}, \mathbf{f}'(\mathbf{x}) \rangle) \mu_i(d\mathbf{z}) \\ & - \langle \boldsymbol{\beta} + \tilde{\mathbf{B}} \mathbf{x}, \mathbf{f}'(\mathbf{x}) \rangle - \int_{U_d} (f(\mathbf{x} + \mathbf{z}) - f(\mathbf{x})) \nu(d\mathbf{z}) \\ &= \sum_{i=1}^d x_i \int_{U_d} (\langle \mathbf{z}, \mathbf{f}'(\mathbf{x}) \rangle - f'_i(\mathbf{x}) (1 \wedge z_i)) \mu_i(d\mathbf{z}) - \langle (\tilde{\mathbf{B}} - \mathbf{B}) \mathbf{x}, \mathbf{f}'(\mathbf{x}) \rangle \\ &= \sum_{i=1}^d x_i \int_{U_d} \left( f'_i(\mathbf{x}) (z_i - (1 \wedge z_i)) + \sum_{j \in \{1, \dots, d\} \setminus \{i\}} z_j f'_j(\mathbf{x}) \right) \mu_i(d\mathbf{z}) \\ & - \sum_{i=1}^d \sum_{j=1}^d x_j f'_i(\mathbf{x}) \int_{U_d} (z_i - \delta_{i,j})^+ \mu_j(d\mathbf{z}) = 0. \end{aligned}$$

using (2.10), (2.11) and (2.12).  $\square$

### 3. Multi-type CBI process as a weak solution of an SDE

Let  $\mathcal{R} := \bigcup_{j=0}^d \mathcal{R}_j$ , where  $\mathcal{R}_j$ ,  $j \in \{0, 1, \dots, d\}$ , are disjoint sets given by

$$\mathcal{R}_0 := U_d \times \{(\mathbf{0}, \mathbf{0})\}^d \subset \mathbb{R}_+^d \times (\mathbb{R}_+^d \times \mathbb{R}_+)^d,$$

and

$$\mathcal{R}_j := \{\mathbf{0}\} \times H_{j,1} \times \cdots \times H_{j,d} \subset \mathbb{R}_+^d \times (\mathbb{R}_+^d \times \mathbb{R}_+)^d, \quad j \in \{1, \dots, d\},$$

where

$$H_{j,i} := \begin{cases} U_d \times U_1 & \text{if } i = j, \\ \{(\mathbf{0}, 0)\} & \text{if } i \neq j. \end{cases}$$

(Recall that  $U_1 = \mathbb{R}_{++}$ .) Let  $m$  be the uniquely defined measure on  $V := \mathbb{R}_+^d \times (\mathbb{R}_+^d \times \mathbb{R}_+)^d$  such that  $m(V \setminus \mathcal{R}) = 0$  and its restrictions on  $\mathcal{R}_j$ ,  $j \in \{0, 1, \dots, d\}$ , are

$$m|_{\mathcal{R}_0}(\mathrm{d}\mathbf{r}) = \nu(\mathrm{d}\mathbf{r}), \quad m|_{\mathcal{R}_j}(\mathrm{d}\mathbf{z}, \mathrm{d}u) = \mu_j(\mathrm{d}\mathbf{z}) \mathrm{d}u, \quad j \in \{1, \dots, d\}, \quad (3.1)$$

where we identify  $\mathcal{R}_0$  with  $U_d$  and  $\mathcal{R}_1, \dots, \mathcal{R}_d$  with  $U_d \times U_1$  in a natural way. Using again this identification, let  $h : \mathbb{R}^d \times V \rightarrow \mathbb{R}_+^d$  be defined by

$$h(\mathbf{x}, \mathbf{r}) := \begin{cases} \mathbf{r}, & \text{if } \mathbf{x} \in \mathbb{R}_+^d, \mathbf{r} \in \mathcal{R}_0, \\ \mathbf{z} \mathbb{1}_{\{u \leq x_j\}}, & \text{if } \mathbf{x} = (x_1, \dots, x_d)^\top \in \mathbb{R}_+^d, \mathbf{r} = (\mathbf{z}, u) \in \mathcal{R}_j, j \in \{1, \dots, d\}, \\ \mathbf{0}, & \text{otherwise.} \end{cases}$$

Consider the decomposition  $\mathcal{R} = V_0 \cup V_1$ , where  $V_0 := \bigcup_{j=1}^d \mathcal{R}_{j,0}$  and  $V_1 := \mathcal{R}_0 \cup (\bigcup_{j=1}^d \mathcal{R}_{j,1})$  with  $\mathcal{R}_{j,k} := \{\mathbf{0}\} \times H_{j,1,k} \times \cdots \times H_{j,d,k}$ ,  $j \in \{1, \dots, d\}$ ,  $k \in \{0, 1\}$ , and

$$H_{j,i,k} := \begin{cases} U_{d,k} \times U_1 & \text{if } i = j, \\ \{(\mathbf{0}, 0)\} & \text{if } i \neq j, \end{cases} \quad U_{d,k} := \begin{cases} \{\mathbf{z} \in U_d : \|\mathbf{z}\| < 1\} & \text{if } k = 0, \\ \{\mathbf{z} \in U_d : \|\mathbf{z}\| \geq 1\} & \text{if } k = 1. \end{cases}$$

Then the sets  $V_0$  and  $V_1$  are disjoint, and the function  $h$  can be decomposed in the form  $h = f + g$  with

$$f(\mathbf{x}, \mathbf{r}) := h(\mathbf{x}, \mathbf{r}) \mathbb{1}_{V_0}(\mathbf{r}), \quad g(\mathbf{x}, \mathbf{r}) := h(\mathbf{x}, \mathbf{r}) \mathbb{1}_{V_1}(\mathbf{r}), \quad (\mathbf{x}, \mathbf{r}) \in \mathbb{R}^d \times V.$$

Let  $(d, \mathbf{c}, \boldsymbol{\beta}, \mathbf{B}, \nu, \boldsymbol{\mu})$  be a set of admissible parameters in the sense of Definition 2.2 such that the moment condition (2.7) holds. Let us consider the  $d$ -dimensional SDE

$$\begin{aligned} \mathbf{X}_t &= \mathbf{X}_0 + \int_0^t b(\mathbf{X}_s) \mathrm{d}s + \int_0^t \sigma(\mathbf{X}_s) \mathrm{d}\mathbf{W}_s \\ &\quad + \int_0^t \int_{V_0} f(\mathbf{X}_{s-}, \mathbf{r}) \tilde{N}(\mathrm{d}s, \mathrm{d}\mathbf{r}) + \int_0^t \int_{V_1} g(\mathbf{X}_{s-}, \mathbf{r}) N(\mathrm{d}s, \mathrm{d}\mathbf{r}), \quad t \in \mathbb{R}_+, \end{aligned} \quad (3.2)$$

where the functions  $b : \mathbb{R}^d \rightarrow \mathbb{R}^d$  and  $\sigma : \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d}$  are defined by

$$b(\mathbf{x}) := \boldsymbol{\beta} + \mathbf{D}\mathbf{x}, \quad \sigma(\mathbf{x}) := \sum_{i=1}^d \sqrt{2c_i x_i^+} \mathbf{e}_i \mathbf{e}_i^\top, \quad \mathbf{x} \in \mathbb{R}^d,$$

$\mathbf{D}$  is defined in (2.9),  $(\mathbf{W}_t)_{t \in \mathbb{R}_+}$  is a  $d$ -dimensional standard Brownian motion,  $N(\mathrm{d}s, \mathrm{d}\mathbf{r})$  is a Poisson random measure on  $\mathbb{R}_{++} \times V$  with intensity measure  $\mathrm{d}s m(\mathrm{d}\mathbf{r})$ , and  $\tilde{N}(\mathrm{d}s, \mathrm{d}\mathbf{r}) := N(\mathrm{d}s, \mathrm{d}\mathbf{r}) - \mathrm{d}s m(\mathrm{d}\mathbf{r})$ . For a short review on point measures and point processes needed for this paper, see, e.g., [Barczy et al. \(2015, Section 2\)](#).

**Definition 3.1.** Let  $n$  be a probability measure on  $(\mathbb{R}_+^d, \mathcal{B}(\mathbb{R}_+^d))$ . An  $\mathbb{R}_+^d$ -valued weak solution of the SDE (3.2) with initial distribution  $n$  is a tuple  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P}, \mathbf{W}, p, \mathbf{X})$ , where

- (D1)  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P})$  is a filtered probability space satisfying the usual hypotheses;
- (D2)  $(\mathbf{W}_t)_{t \in \mathbb{R}_+}$  is a  $d$ -dimensional standard  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -Brownian motion;
- (D3)  $p$  is a stationary  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -Poisson point process on  $V$  with characteristic measure  $m$  given in (3.1);
- (D4)  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  is an  $\mathbb{R}_+^d$ -valued  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted càdlàg process such that
  - (a) the distribution of  $\mathbf{X}_0$  is  $n$ ,
  - (b)  $\mathbb{P}(\int_0^t (\|b(\mathbf{X}_s)\| + \|\sigma(\mathbf{X}_s)\|^2) ds < \infty) = 1$  for all  $t \in \mathbb{R}_+$ ,
  - (c)  $\mathbb{P}(\int_0^t \int_{V_0} \|f(\mathbf{X}_s, \mathbf{r})\|^2 ds m(d\mathbf{r}) < \infty) = 1$  for all  $t \in \mathbb{R}_+$ ,
  - (d)  $\mathbb{P}(\int_0^t \int_{V_1} \|g(\mathbf{X}_{s-}, \mathbf{r})\| N(ds, d\mathbf{r}) < \infty) = 1$  for all  $t \in \mathbb{R}_+$ , where  $N(ds, d\mathbf{r})$  is the counting measure of  $p$  on  $\mathbb{R}_{++} \times V$ ,
  - (e) equation (3.2) holds  $\mathbb{P}$ -a.s., where  $\tilde{N}(ds, d\mathbf{r}) := N(ds, d\mathbf{r}) - ds m(d\mathbf{r})$ .

For the definitions of an  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -Brownian motion and an  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -Poisson point process, see, e.g., Ikeda and Watanabe (1989, Chapter I, Definition 7.2 and Chapter II, Definition 3.2).

*Remark 3.2.* If conditions (D1)–(D3) and (D4)(b)–(d) are satisfied, then the mappings  $\mathbb{R}_+ \times V_0 \times \Omega \ni (s, \mathbf{r}, \omega) \mapsto f(\mathbf{X}_{s-}(\omega), \mathbf{r}) \in \mathbb{R}^d$  and  $\mathbb{R}_+ \times V_1 \times \Omega \ni (s, \mathbf{r}, \omega) \mapsto g(\mathbf{X}_{s-}(\omega), \mathbf{r}) \in \mathbb{R}^d$  are in the (multidimensional versions of the) classes  $\mathbf{F}_p^{2,loc}$  and  $\mathbf{F}_p$ , respectively, defined in Ikeda and Watanabe (1989, pages 61, 62), the integrals in (3.2) are well-defined and have càdlàg modifications as functions of  $t$ , see, e.g., Barczy et al. (2015, Remark 3.2).

Moreover, if  $\mathbb{E}(\int_0^t \|\mathbf{X}_s\| ds) < \infty$  for all  $t \in \mathbb{R}_+$ , and the moment condition (2.7) holds, then conditions (D4)(b)–(d) are satisfied, and the mappings  $\mathbb{R}_+ \times V_0 \times \Omega \ni (s, \mathbf{r}, \omega) \mapsto f(\mathbf{X}_{s-}(\omega), \mathbf{r}) \in \mathbb{R}^d$  and  $\mathbb{R}_+ \times V_1 \times \Omega \ni (s, \mathbf{r}, \omega) \mapsto g(\mathbf{X}_{s-}(\omega), \mathbf{r}) \in \mathbb{R}^d$  are in the (multidimensional versions of the) smaller classes  $\mathbf{F}_p^2$  and  $\mathbf{F}_p^1$ , respectively, defined in Ikeda and Watanabe (1989, page 62). Indeed, with the notation  $\mathbf{X}_s = (X_{s,1}, \dots, X_{s,d})^\top$ ,  $s \in \mathbb{R}_+$ ,

$$\begin{aligned} & \mathbb{E} \left( \int_0^t \int_{V_0} \|f(\mathbf{X}_s, \mathbf{r})\|^2 ds m(d\mathbf{r}) \right) \\ &= \sum_{j=1}^d \mathbb{E} \left( \int_0^t \int_{U_d} \int_{U_1} \|z\|^2 \mathbb{1}_{\{\|z\| < 1\}} \mathbb{1}_{\{u \leq X_{s,j}\}} ds \mu_j(dz) du \right) \\ &= \sum_{j=1}^d \mathbb{E} \left( \int_0^t X_{s,j} ds \right) \int_{U_d} \|z\|^2 \mathbb{1}_{\{\|z\| < 1\}} \mu_j(dz) < \infty \end{aligned}$$

by (2.13), and

$$\begin{aligned} & \mathbb{E} \left( \int_0^t \int_{V_1} \|g(\mathbf{X}_s, \mathbf{r})\| \, ds \, m(d\mathbf{r}) \right) \\ &= \int_0^t \int_{U_d} \|\mathbf{r}\| \, ds \, \nu(d\mathbf{r}) + \sum_{j=1}^d \mathbb{E} \left( \int_0^t \int_{U_d} \int_{U_1} \|z\| \mathbb{1}_{\{\|z\| \geq 1\}} \mathbb{1}_{\{u \leq X_{s,j}\}} \, ds \, \mu_j(dz) \, du \right) \\ &= t \int_{U_d} \|\mathbf{r}\| \, \nu(d\mathbf{r}) + \sum_{j=1}^d \mathbb{E} \left( \int_0^t X_{s,j} \, ds \right) \int_{U_d} \|z\| \mathbb{1}_{\{\|z\| \geq 1\}} \, \mu_j(dz) < \infty \end{aligned}$$

by (2.10) and (2.2). Note that if  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  is a CBI process with  $\mathbb{E}(\|\mathbf{X}_0\|) < \infty$  satisfying the moment condition (2.7), then  $\mathbb{E}(\int_0^t \|\mathbf{X}_s\| \, ds) < \infty$  for all  $t \in \mathbb{R}_+$ , see Lemma 3.4.  $\square$

*Remark 3.3.* Note that if conditions (D1)–(D3) are satisfied, then  $\mathbf{W}$  and  $p$  are automatically independent according to Theorem 6.3 in Chapter II in Ikeda and Watanabe (1989), since the intensity measure  $ds \, m(d\mathbf{r})$  of  $p$  is deterministic. Moreover, if  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P}, \mathbf{W}, p, \mathbf{X})$  is an  $\mathbb{R}_+^d$ -valued weak solution of the SDE (3.2), then  $\mathcal{F}_0$ ,  $\mathbf{W}$  and  $p$  are mutually independent, and hence  $\mathbf{X}_0$ ,  $\mathbf{W}$  and  $p$  are mutually independent as well, see, e.g., Barczy et al. (2015, Remark 3.4).  $\square$

**Lemma 3.4.** *Let  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  be a CBI process with parameters  $(d, \mathbf{c}, \beta, \mathbf{B}, \nu, \mu)$  and with initial distribution  $n$  satisfying  $\int_{\mathbb{R}_+^d} \|z\| \, n(dz) < \infty$ . Suppose that the moment condition (2.7) holds. Then*

$$\mathbb{E}(\mathbf{X}_t) = e^{t\tilde{\mathbf{B}}} \mathbb{E}(\mathbf{X}_0) + \left( \int_0^t e^{u\tilde{\mathbf{B}}} \, du \right) \tilde{\beta}, \quad t \in \mathbb{R}_+,$$

where  $\tilde{\mathbf{B}} \in \mathbb{R}_{(+)}^{d \times d}$  and  $\tilde{\beta} \in \mathbb{R}_+^d$  are defined in (2.8). In particular,  $\int_0^t \mathbb{E}(\|\mathbf{X}_s\|) \, ds < \infty$  for all  $t \in \mathbb{R}_+$ .

*Proof:* By the tower rule for conditional expectations, it suffices to show

$$\mathbb{E}(\mathbf{X}_t | \mathbf{X}_0) = e^{t\tilde{\mathbf{B}}} \mathbf{X}_0 + \left( \int_0^t e^{u\tilde{\mathbf{B}}} \, du \right) \tilde{\beta}, \quad t \in \mathbb{R}_+, \quad (3.3)$$

where the conditional expectation  $\mathbb{E}(\mathbf{X}_t | \mathbf{X}_0) \in [0, \infty]^d$  is meant in the generalized sense, see, e.g., Stroock (2011, Theorem 5.1.6). In order to show (3.3), it is enough to check that for a CBI process  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  with initial value  $\mathbf{X}_0 = \mathbf{x} \in \mathbb{R}_+^d$ , we have

$$\mathbb{E}(\mathbf{X}_t) = e^{t\tilde{\mathbf{B}}} \mathbf{x} + \left( \int_0^t e^{u\tilde{\mathbf{B}}} \, du \right) \tilde{\beta}, \quad t \in \mathbb{R}_+, \quad \mathbf{x} \in \mathbb{R}_+^d. \quad (3.4)$$

Indeed, let  $\phi_n : \mathbb{R}_+^d \rightarrow \mathbb{R}_+^d$ ,  $n \in \mathbb{N}$ , be simple functions such that  $\phi_n(\mathbf{y}) \uparrow \mathbf{y}$  as  $n \rightarrow \infty$  for all  $\mathbf{y} \in \mathbb{R}_+^d$ . Then, by the (multidimensional version of the) monotone convergence theorem for (generalized) conditional expectations, see, e.g., Stroock (2011, Theorem 5.1.6), we obtain  $\mathbb{E}(\phi_n(\mathbf{X}_t) | \mathbf{X}_0) \uparrow \mathbb{E}(\mathbf{X}_t | \mathbf{X}_0)$  as  $n \rightarrow \infty$   $\mathbb{P}$ -almost surely. For each  $B \in \mathcal{B}(\mathbb{R}^d)$ , we have

$$\mathbb{E}(\mathbb{1}_B(\mathbf{X}_t) | \mathbf{X}_0) = \mathbb{P}(\mathbf{X}_t \in B | \mathbf{X}_0) = \int_{\mathbb{R}_+^d} \mathbb{1}_B(\mathbf{y}) \, P_t(\mathbf{X}_0, d\mathbf{y}),$$

hence  $\mathbb{E}(\phi_n(\mathbf{X}_t) | \mathbf{X}_0) = \int_{\mathbb{R}_+^d} \phi_n(\mathbf{y}) P_t(\mathbf{X}_0, d\mathbf{y})$ . By the (multidimensional version of the) monotone convergence theorem,  $\int_{\mathbb{R}_+^d} \phi_n(\mathbf{y}) P_t(\mathbf{X}_0, d\mathbf{y}) \uparrow \int_{\mathbb{R}_+^d} \mathbf{y} P_t(\mathbf{X}_0, d\mathbf{y})$  as  $n \rightarrow \infty$ . By (3.4), we get

$$\mathbb{E}(\mathbf{X}_t | \mathbf{X}_0) = \int_{\mathbb{R}_+^d} \mathbf{y} P_t(\mathbf{X}_0, d\mathbf{y}) = e^{t\tilde{\mathbf{B}}} \mathbf{X}_0 + \left( \int_0^t e^{u\tilde{\mathbf{B}}} du \right) \tilde{\boldsymbol{\beta}},$$

hence we conclude (3.3).

In order to show (3.4), we are going to apply Proposition 9.11 of Li (2011) for the immigration superprocess given in Lemma 2.11. For each  $f \in B(E)$  and  $i \in E$ , the function  $\mathbb{R}_+ \ni t \mapsto \pi_t f(i)$  is the unique locally bounded solution to the linear evolution equation (2.35) in Li (2011) taking the form

$$\begin{aligned} \pi_t f(i) &= f(i) + \int_0^t \gamma(i, \pi_s f) ds - \int_0^t b(i) \pi_s f(i) ds \\ &= f(i) + \int_0^t \left( \sum_{j=1}^d \pi_s f(j) \gamma(i, \{j\}) \right) ds - \int_0^t b(i) \pi_s f(i) ds \\ &= f(i) + \int_0^t \left( \sum_{j=1}^d \pi_s f(j) \tilde{b}_{j,i} \right) ds, \end{aligned}$$

where we used  $R_t f = f$  for  $f \in B(E)$  and  $t \in \mathbb{R}_+$ ,  $b(i) = -\tilde{b}_{i,i}$  and  $\gamma(i, \{i\}) = B(i, \{i\}) = 0$  for  $i \in \{1, \dots, d\}$ , and

$$\gamma(i, \{j\}) = B(i, \{j\}) + \int_{U_d} z_j \mu_i(dz) = b_{j,i} + \int_{U_d} z_j \mu_i(dz) = \tilde{b}_{j,i} \quad (3.5)$$

for  $i, j \in \{1, \dots, d\}$  with  $i \neq j$ . The functions  $\mathbb{R}_+ \ni t \mapsto \pi_t f(i)$ ,  $f \in B(E)$ ,  $i \in \{1, \dots, d\}$ , can be identified with the functions  $\mathbb{R}_+ \ni t \mapsto \pi_i(t, \boldsymbol{\lambda})$ ,  $\boldsymbol{\lambda} \in \mathbb{R}^d$ ,  $i \in \{1, \dots, d\}$ , which are the unique locally bounded solution to the linear evolution equations

$$\pi_i(t, \boldsymbol{\lambda}) = \lambda_i + \int_0^t \langle \tilde{\mathbf{B}} \mathbf{e}_i, \pi(s, \boldsymbol{\lambda}) \rangle ds, \quad t \in \mathbb{R}_+, \quad i \in \{1, \dots, d\}, \quad \boldsymbol{\lambda} \in \mathbb{R}^d.$$

Consequently, the functions  $\mathbb{R}_+ \ni t \mapsto \boldsymbol{\pi}(t, \boldsymbol{\lambda}) := (\pi_1(t, \boldsymbol{\lambda}), \dots, \pi_d(t, \boldsymbol{\lambda}))$ ,  $\boldsymbol{\lambda} \in \mathbb{R}^d$ , satisfies

$$\boldsymbol{\pi}(t, \boldsymbol{\lambda}) = \boldsymbol{\lambda} + \int_0^t \tilde{\mathbf{B}}^\top \boldsymbol{\pi}(s, \boldsymbol{\lambda}) ds, \quad t \in \mathbb{R}_+, \quad \boldsymbol{\lambda} \in \mathbb{R}^d,$$

and hence

$$\boldsymbol{\pi}(t, \boldsymbol{\lambda}) = e^{t\tilde{\mathbf{B}}^\top} \boldsymbol{\lambda}, \quad t \in \mathbb{R}_+, \quad \boldsymbol{\lambda} \in \mathbb{R}^d.$$

The functional  $B(E) \ni f \mapsto \Gamma(f) = \eta(f) + \int_{M(E)^\circ} \kappa(f) H_1(d\kappa)$  of Li (2011, formula (9.20)) can be identified with the functional  $\mathbb{R}^d \ni \mathbf{x} \mapsto \mathbf{x}^\top \boldsymbol{\beta} + \int_{U_d} \mathbf{x}^\top z \nu(dz) = \mathbf{x}^\top \tilde{\boldsymbol{\beta}}$ . Hence Proposition 9.11 of Li (2011) implies

$$\langle \boldsymbol{\lambda}, \mathbb{E}(\mathbf{X}_t) \rangle = \langle e^{t\tilde{\mathbf{B}}^\top} \boldsymbol{\lambda}, \mathbf{x} \rangle + \left( \int_0^t (e^{s\tilde{\mathbf{B}}^\top} \boldsymbol{\lambda})^\top ds \right) \tilde{\boldsymbol{\beta}} = \left\langle \boldsymbol{\lambda}, e^{t\tilde{\mathbf{B}}} \mathbf{x} + \left( \int_0^t e^{s\tilde{\mathbf{B}}} ds \right) \tilde{\boldsymbol{\beta}} \right\rangle$$

for  $t \in \mathbb{R}_+$  and  $\boldsymbol{\lambda} \in \mathbb{R}^d$ , which yields (3.4).  $\square$



*Remark 3.5.* We call the attention that in the proof of the forthcoming Theorem 3.7, which states existence of an  $\mathbb{R}_+^d$ -valued weak solution of the SDE (3.2), we will extensively use that for a CBI process  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  with parameters  $(d, \mathbf{c}, \boldsymbol{\beta}, \mathbf{B}, \nu, \boldsymbol{\mu})$  satisfying  $\mathbb{E}(\|\mathbf{X}_0\|) < \infty$  and the moment condition (2.7), we have  $\int_0^t \mathbb{E}(\|\mathbf{X}_s\|) ds < \infty$ ,  $t \in \mathbb{R}_+$ , proved in Lemma 3.4. We point out that in the proof of Lemma 3.4 we can not use the SDE (3.2), since at that point it has not yet been proved that a CBI process is a solution of this SDE. This drives us back to Definition 2.6 of CBI processes in the proof of Lemma 3.4. Having proved that a CBI process is a solution of the SDE (3.2), one could give another proof of Lemma 3.4 (roughly speaking by taking expectations via localization argument).  $\square$

**Definition 3.6.** We say that uniqueness in the sense of probability law holds for the SDE (3.2) among  $\mathbb{R}_+^d$ -valued weak solutions if whenever  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P}, \mathbf{W}, p, \mathbf{X})$  and  $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{F}}_t)_{t \in \mathbb{R}_+}, \tilde{\mathbb{P}}, \tilde{\mathbf{W}}, \tilde{p}, \tilde{\mathbf{X}})$  are  $\mathbb{R}_+^d$ -valued weak solutions of the SDE (3.2) such that  $\mathbb{P}(\mathbf{X}_0 \in B) = \tilde{\mathbb{P}}(\tilde{\mathbf{X}}_0 \in B)$  for all  $B \in \mathcal{B}(\mathbb{R}^d)$ , then  $\mathbb{P}(\mathbf{X} \in C) = \tilde{\mathbb{P}}(\tilde{\mathbf{X}} \in C)$  for all  $C \in \mathcal{D}(\mathbb{R}_+, \mathbb{R}^d)$ .

**Theorem 3.7.** *Let  $(d, \mathbf{c}, \boldsymbol{\beta}, \mathbf{B}, \nu, \boldsymbol{\mu})$  be a set of admissible parameters in the sense of Definition 2.2 such that the moment condition (2.7) holds. Then for any probability measure  $n$  on  $(\mathbb{R}_+^d, \mathcal{B}(\mathbb{R}_+^d))$  with  $\int_{\mathbb{R}_+^d} \|\mathbf{z}\| n(d\mathbf{z}) < \infty$ , the SDE (3.2) admits an  $\mathbb{R}_+^d$ -valued weak solution with initial distribution  $n$  which is unique in the sense of probability law among  $\mathbb{R}_+^d$ -valued weak solutions. Moreover, any  $\mathbb{R}_+^d$ -valued weak solution is a CBI process with parameters  $(d, \mathbf{c}, \boldsymbol{\beta}, \mathbf{B}, \nu, \boldsymbol{\mu})$ .*

*Proof:* Suppose that  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  is a càdlàg realization of a CBI process with parameters  $(d, \mathbf{c}, \boldsymbol{\beta}, \mathbf{B}, \nu, \boldsymbol{\mu})$  on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  having initial distribution  $n$ , i.e.,  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  is a time homogeneous Markov process having càdlàg trajectories and the same finite dimensional distributions as a CBI process with parameters  $(d, \mathbf{c}, \boldsymbol{\beta}, \mathbf{B}, \nu, \boldsymbol{\mu})$  having initial distribution  $n$  (such a realization exists due to Theorem 9.15 in Li (2011)). Let

$$\mathcal{F}_t := \bigcap_{\varepsilon > 0} \sigma(\mathcal{F}_{t+\varepsilon}^{\mathbf{X}} \cup \mathcal{N}), \quad t \in \mathbb{R}_+,$$

where  $\mathcal{N}$  denotes the collection of null sets under the probability measure  $\mathbb{P}$ , and  $(\mathcal{F}_t^{\mathbf{X}})_{t \in \mathbb{R}_+}$  stands for the natural filtration generated by the process  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$ , hence the filtered probability space  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P})$  satisfies the usual hypotheses.

By the equivalence of parts (3) and (4) of Theorem 9.18 of Li (2011) applied to the immigration superprocess given in Lemma 2.11, we conclude that the process  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  has no negative jumps, the (not necessarily Poisson) random measure

$$N_0(ds, d\mathbf{z}) := \sum_{u \in \mathbb{R}_{++}} \mathbb{1}_{\{\mathbf{X}_u \neq \mathbf{X}_{u-}\}} \delta_{(u, \mathbf{X}_u - \mathbf{X}_{u-})}(ds, d\mathbf{z})$$

on  $\mathbb{R}_{++} \times U_d$  has predictable compensator

$$\hat{N}_0(ds, d\mathbf{z}) := \sum_{j=1}^d X_{s-,j} ds \mu_j(d\mathbf{z}) + ds \nu(d\mathbf{z}),$$

and

$$\mathbf{X}_t - \mathbf{X}_0 - \int_0^t (\tilde{\boldsymbol{\beta}} + \tilde{\mathbf{B}}\mathbf{X}_s) ds - \int_0^t \int_{U_d} \mathbf{z} \tilde{N}_0(ds, d\mathbf{z}), \quad t \in \mathbb{R}_+,$$

is a continuous locally square integrable martingale starting from  $\mathbf{0} \in \mathbb{R}^d$  with quadratic variation process

$$\left( 2\delta_{i,j} c_i \int_0^t X_{s,i} ds \right)_{i,j \in \{1, \dots, d\}}, \quad t \in \mathbb{R}_+,$$

where  $\tilde{N}_0(ds, d\mathbf{z}) := N_0(ds, d\mathbf{z}) - \hat{N}_0(ds, d\mathbf{z})$ . Indeed, first, note that  $R_t f = f$ ,  $t \in \mathbb{R}_+$ ,  $f \in B(E)$ , yields that the strong generator of  $(R_t)_{t \in \mathbb{R}_+}$  is identically 0, i.e.,  $A = 0$ , see Li (2011, (7.1)). Using  $b(i) = -\tilde{b}_{i,i}$  and  $\gamma(i, \{i\}) = B(i, \{i\}) = 0$  for  $i \in \{1, \dots, d\}$  and  $\gamma(i, \{j\}) = \tilde{b}_{j,i}$  for  $i, j \in \{1, \dots, d\}$  with  $i \neq j$  (see, (3.5)), the function  $B(E) \ni f \mapsto Af + \gamma f - bf$  of Li (2011, page 218) can be identified with the function

$$E \ni i \mapsto \sum_{j=1}^d f(j)\gamma(i, \{j\}) - b(i)f(i) = \sum_{j=1}^d \tilde{b}_{j,i}f(j). \quad (3.6)$$

Recalling that the functional  $B(E) \ni f \mapsto \Gamma(f) = \eta(f) + \int_{M(E)^\circ} \kappa(f) H(d\kappa)$  is identified with the functional  $\mathbb{R}^d \ni \mathbf{x} \mapsto \mathbf{x}^\top \tilde{\boldsymbol{\beta}}$  (see, the end of the proof of Lemma 3.4), Theorem 9.18 of Li (2011) yields that for each  $\mathbf{w} = (w_1, \dots, w_d)^\top \in \mathbb{R}^d$ , the process  $(\mathbf{w}^\top \mathbf{X}_t)_{t \in \mathbb{R}_+}$  has no negative jumps, and

$$\mathbf{w}^\top \mathbf{X}_t - \mathbf{w}^\top \mathbf{X}_0 - \int_0^t (\mathbf{w}^\top \tilde{\boldsymbol{\beta}} + \mathbf{w}^\top \tilde{\mathbf{B}}\mathbf{X}_s) ds - \int_0^t \int_{U_d} \mathbf{w}^\top \mathbf{z} \tilde{N}_0(ds, d\mathbf{z}), \quad t \in \mathbb{R}_+,$$

is a continuous locally square integrable martingale starting from  $0 \in \mathbb{R}$  with quadratic variation process

$$\langle \mathbf{w}^\top \mathbf{X} \rangle_t = 2 \sum_{i=1}^d c_i w_i^2 \int_0^t X_{s,i} ds, \quad t \in \mathbb{R}_+.$$

Further, by polarization identity, for all  $\mathbf{w}, \tilde{\mathbf{w}} \in \mathbb{R}^d$ , the cross quadratic variation process of  $(\mathbf{w}^\top \mathbf{X}_t)_{t \in \mathbb{R}_+}$  and  $(\tilde{\mathbf{w}}^\top \mathbf{X}_t)_{t \in \mathbb{R}_+}$  takes the form

$$\begin{aligned} \langle \mathbf{w}^\top \mathbf{X}, \tilde{\mathbf{w}}^\top \mathbf{X} \rangle_t &= \frac{1}{4} (\langle (\mathbf{w} + \tilde{\mathbf{w}})^\top \mathbf{X} \rangle_t - \langle (\mathbf{w} - \tilde{\mathbf{w}})^\top \mathbf{X} \rangle_t) \\ &= \frac{1}{4} \left( 2 \sum_{i=1}^d c_i (w_i + \tilde{w}_i)^2 \int_0^t X_{s,i} ds - 2 \sum_{i=1}^d c_i (w_i - \tilde{w}_i)^2 \int_0^t X_{s,i} ds \right) \\ &= 2 \sum_{i=1}^d c_i w_i \tilde{w}_i \int_0^t X_{s,i} ds, \quad t \in \mathbb{R}_+. \end{aligned}$$

We note that the integral  $\int_0^t \int_{U_d} \mathbf{z} \tilde{N}_0(ds, d\mathbf{z})$  is well-defined, since  $\mathbf{z} = \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} + \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}}$ ,  $\mathbf{z} \in U_d$ , and the functions  $\mathbb{R}_+ \times U_d \times \Omega \ni (s, \mathbf{z}, \omega) \mapsto \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| < 1\}}$  and  $\mathbb{R}_+ \times U_d \times \Omega \ni (s, \mathbf{z}, \omega) \mapsto \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}}$  belong to the classes  $\mathbf{F}_{p_0}^2$  and  $\mathbf{F}_{p_0}^1$ , respectively, where  $p_0$  denotes the point process on  $U_d$  with counting measure  $N_0(ds, d\mathbf{z})$ , i.e.,  $p_0(u) := \mathbf{X}_u - \mathbf{X}_{u-}$  for  $u \in D(p_0)$  with

$D(p_0) := \{u \in \mathbb{R}_{++} : \mathbf{X}_u \neq \mathbf{X}_{u-}\}$ . Indeed,

$$\begin{aligned} & \mathbb{E} \left( \int_0^t \int_{U_d} \|\mathbf{z}\|^2 \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \widehat{N}_0(ds, d\mathbf{z}) \right) \\ &= \int_0^t \int_{U_d} \|\mathbf{z}\|^2 \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} ds \nu(d\mathbf{z}) + \sum_{j=1}^d \int_0^t \int_{U_d} \|\mathbf{z}\|^2 \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \mathbb{E}(X_{s,j}) ds \mu_j(d\mathbf{z}) \\ &\leq t \int_{U_d} \|\mathbf{z}\| \nu(d\mathbf{z}) + \sum_{j=1}^d \int_0^t \mathbb{E}(X_{s,j}) ds \int_{U_d} \|\mathbf{z}\|^2 \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \mu_j(d\mathbf{z}) < \infty \end{aligned}$$

by Lemma 3.4 and the inequalities (2.10) and (2.13), and

$$\begin{aligned} & \mathbb{E} \left( \int_0^t \int_{U_d} \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \widehat{N}_0(ds, d\mathbf{z}) \right) \\ &= \int_0^t \int_{U_d} \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} ds \nu(d\mathbf{z}) + \sum_{j=1}^d \int_0^t \int_{U_d} \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mathbb{E}(X_{s,j}) ds \mu_j(d\mathbf{z}) \\ &\leq t \int_{U_d} \|\mathbf{z}\| \nu(d\mathbf{z}) + \sum_{j=1}^d \int_0^t \mathbb{E}(X_{s,j}) ds \int_{U_d} \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_j(d\mathbf{z}) < \infty \end{aligned}$$

by Lemma 3.4 and the inequalities (2.10) and (2.2).

Using that  $\mathbb{P}(\int_0^t X_{s,i} ds < \infty) = 1$ ,  $i \in \{1, \dots, d\}$  (since  $\mathbf{X}$  has càdlàg trajectories almost surely), by choosing  $\mathbf{w} = \mathbf{e}_j$ ,  $j \in \{1, \dots, d\}$ , a representation theorem for continuous locally square integrable martingales (see, e.g., Ikeda and Watanabe (1989, Chapter II, Theorem 7.1')) yields

$$\mathbf{X}_t = \mathbf{X}_0 + \int_0^t (\tilde{\boldsymbol{\beta}} + \tilde{\mathbf{B}}\mathbf{X}_s) ds + \sum_{i=1}^d \mathbf{e}_i \int_0^t \sqrt{2c_i X_{s,i}} dW_{s,i} + \int_0^t \int_{U_d} \mathbf{z} \tilde{N}_0(ds, d\mathbf{z})$$

for all  $t \in \mathbb{R}_+$ ,  $\tilde{\mathbb{P}}$ -almost surely on an extension  $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{F}}_t)_{t \in \mathbb{R}_+}, \tilde{\mathbb{P}})$  of the filtered probability space  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P})$  (see Definition A.1), and  $(W_{t,1}, \dots, W_{t,d})_{t \in \mathbb{R}_+}$  is a  $d$ -dimensional  $(\tilde{\mathcal{F}}_t)_{t \in \mathbb{R}_+}$ -Brownian motion. We note that, with a little abuse of notation, the extended random variables on the extension  $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{F}}_t)_{t \in \mathbb{R}_+}, \tilde{\mathbb{P}})$  are denoted in the same way as the original ones. Let

$$\tilde{\mathcal{G}}_t := \bigcap_{\varepsilon > 0} \sigma(\tilde{\mathcal{F}}_{t+\varepsilon} \cup \tilde{\mathcal{N}}), \quad t \in \mathbb{R}_+,$$

where  $\tilde{\mathcal{N}}$  denotes the collection of null sets under the probability measure  $\tilde{\mathbb{P}}$ . Then the filtered probability space  $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{G}}_t)_{t \in \mathbb{R}_+}, \tilde{\mathbb{P}})$  satisfies the usual hypotheses, and by Lemma A.4,  $(W_{t,1}, \dots, W_{t,d})_{t \in \mathbb{R}_+}$  is a  $d$ -dimensional  $(\tilde{\mathcal{G}}_t)_{t \in \mathbb{R}_+}$ -Brownian motion.

The aim of the following discussion is to show, by the representation theorem of Ikeda and Watanabe (1989, Chapter II, Theorem 7.4), that the SDE (3.2) holds on an extension of the original probability space. The predictable compensator of the random measure  $N_0(ds, d\mathbf{z})$  can be written in the form  $\widehat{N}_0(ds, d\mathbf{z}) = ds q(s, d\mathbf{z})$ , where

$$q(s, d\mathbf{z}) := \sum_{j=1}^d X_{s-,j} \mu_j(d\mathbf{z}) + \nu(d\mathbf{z}).$$

Let  $\Theta : \mathbb{R}_+ \times V \times \tilde{\Omega} \rightarrow U_d \cup \{\mathbf{0}\} = \mathbb{R}_+^d$  be defined by

$$\Theta(s, \mathbf{r}, \tilde{\omega}) := h(\mathbf{X}_{s-}(\tilde{\omega}), \mathbf{r}), \quad (s, \mathbf{r}, \tilde{\omega}) \in \mathbb{R}_+ \times V \times \tilde{\Omega}.$$

(Note, that  $\Delta = \mathbf{0}$  in the notation of Ikeda and Watanabe (1989, Chapter II, Theorem 7.4).) Then condition (7.26) on page 93 in Ikeda and Watanabe (1989) holds, since for all  $s \in \mathbb{R}_+$ ,  $\tilde{\omega} \in \tilde{\Omega}$ , and  $B \in \mathcal{B}(U_d)$ , we have

$$\begin{aligned} m(\{\mathbf{r} \in V : \Theta(s, \mathbf{r}, \tilde{\omega}) \in B\}) &= \sum_{i=0}^d m(\{\mathbf{r} \in \mathcal{R}_i : \Theta(s, \mathbf{r}, \tilde{\omega}) \in B\}) \\ &= \sum_{i=1}^d (\mu_i \times \ell)(\{(z, u) \in \mathcal{R}_i : z \mathbb{1}_{\{u \leq X_{s-,i}(\tilde{\omega})\}} \in B\}) + \nu(\{\mathbf{r} \in \mathcal{R}_0 : \mathbf{r} \in B\}) \\ &= \sum_{i=1}^d X_{s-,i}(\tilde{\omega}) \mu_i(B) + \nu(B) = q(s, B)(\tilde{\omega}), \end{aligned}$$

where  $\ell$  denotes the Lebesgue measure on  $\mathbb{R}_{++}$ , and we used that  $\mathbf{0} \notin B$ . By Theorem II.7.4 in Ikeda and Watanabe (1989), on an extension  $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{F}}_t)_{t \in \mathbb{R}_+}, \tilde{\mathbb{P}})$  of  $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{G}}_t)_{t \in \mathbb{R}_+}, \tilde{\mathbb{P}})$ , there is a stationary  $(\tilde{\mathcal{F}}_t)_{t \in \mathbb{R}_+}$ -Poisson point process  $p$  on  $V$  with characteristic measure  $m$  such that

$$\begin{aligned} N_0((0, t] \times B) &= \int_0^t \int_V \mathbb{1}_B(\Theta(s, \mathbf{r})) N(ds, d\mathbf{r}) \\ &= \#\{s \in D(p) : s \in (0, t], \Theta(s, p(s)) \in B\} \quad \tilde{\mathbb{P}}\text{-a.s.} \end{aligned}$$

for all  $B \in \mathcal{B}(U_d)$ , where  $N(ds, d\mathbf{r})$  denotes the counting measure of  $p$ , and  $D(p)$  is the domain of  $p$  being a countable subset of  $\mathbb{R}_{++}$  such that  $\{s \in D(p) : s \in (0, t], p(s) \in B\}$  is finite for all  $t \in \mathbb{R}_+$  and compact subsets  $B \in \mathcal{B}(U_d)$ .

Then, by Lemma A.3,  $(W_{t,1}, \dots, W_{t,d})_{t \in \mathbb{R}_+}$  is a  $d$ -dimensional  $(\tilde{\mathcal{F}}_t)_{t \in \mathbb{R}_+}$ -Brownian motion. Let

$$\tilde{\mathcal{G}}_t := \bigcap_{\varepsilon > 0} \sigma\left(\tilde{\mathcal{F}}_{t+\varepsilon} \cup \tilde{\mathcal{N}}\right), \quad t \in \mathbb{R}_+,$$

where  $\tilde{\mathcal{N}}$  denotes the collection of null sets under the probability measure  $\tilde{\mathbb{P}}$ . Then the filtered probability space  $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{G}}_t)_{t \in \mathbb{R}_+}, \tilde{\mathbb{P}})$  satisfies the usual hypotheses. By Lemma A.4,  $(W_{t,1}, \dots, W_{t,d})_{t \in \mathbb{R}_+}$  is a  $d$ -dimensional  $(\tilde{\mathcal{G}}_t)_{t \in \mathbb{R}_+}$ -Brownian motion, and  $p$  is a stationary  $(\tilde{\mathcal{G}}_t)_{t \in \mathbb{R}_+}$ -Poisson point process on  $V$  with characteristic measure  $m$ . Consequently,

$$\#\{s \in D(p_0) : s \in (0, t], p_0(s) \in B\} = \#\{s \in D(p) : s \in (0, t], h(\mathbf{X}_{s-}, p(s)) \in B\} \quad (3.7)$$

for all  $B \in \mathcal{B}(U_d)$ . Using this representation, we will calculate  $\int_0^t \int_{U_d} z \tilde{N}_0(ds, dz)$ ,  $t \in \mathbb{R}_+$ . First observe that

$$\int_0^t \int_{U_d} z \tilde{N}_0(ds, dz) = \int_0^t \int_{U_d} z \mathbb{1}_{\{\|z\| \geq 1\}} \tilde{N}_0(ds, dz) + \int_0^t \int_{U_d} z \mathbb{1}_{\{\|z\| < 1\}} \tilde{N}_0(ds, dz).$$

Since the function  $\mathbb{R}_+ \times U_d \times \Omega \ni (s, \mathbf{z}, \omega) \mapsto \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}}$  belongs to the class  $\mathbf{F}_{p_0}^1$ , by [Ikeda and Watanabe \(1989, Chapter II, \(3.8\)\)](#), we obtain

$$\begin{aligned} & \int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \tilde{N}_0(ds, d\mathbf{z}) \\ &= \int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} N_0(ds, d\mathbf{z}) - \int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \hat{N}_0(ds, d\mathbf{z}). \end{aligned}$$

Applying [\(3.7\)](#), we obtain

$$\begin{aligned} & \int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} N_0(ds, d\mathbf{z}) = \sum_{s \in D(p_0) \cap (0, t]} p_0(s) \mathbb{1}_{\{p_0(s) \geq 1\}} \\ &= \sum_{s \in D(p) \cap (0, t]} h(\mathbf{X}_{s-}, p(s)) \mathbb{1}_{\{\|h(\mathbf{X}_{s-}, p(s))\| \geq 1\}} \\ &= \int_0^t \int_V h(\mathbf{X}_{s-}, \mathbf{r}) \mathbb{1}_{\{\|h(\mathbf{X}_{s-}, \mathbf{r})\| \geq 1\}} N(ds, d\mathbf{r}) \\ &= \int_0^t \int_{\mathcal{R}_0} \mathbf{r} \mathbb{1}_{\{\|\mathbf{r}\| \geq 1\}} N(ds, d\mathbf{r}) + \sum_{j=1}^d \int_0^t \int_{\mathcal{R}_j} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mathbb{1}_{\{u \leq X_{s-, j}\}} N(ds, d\mathbf{r}) \\ &= \int_0^t \int_{V_1} g(\mathbf{X}_{s-}, \mathbf{r}) N(ds, d\mathbf{r}) - \int_0^t \int_{\mathcal{R}_0} \mathbf{r} \mathbb{1}_{\{\|\mathbf{r}\| < 1\}} N(ds, d\mathbf{r}). \end{aligned}$$

Here we used that the function  $\mathbb{R}_+ \times U_d \times \tilde{\Omega} \ni (s, \mathbf{z}, \tilde{\omega}) \mapsto \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}}$  belongs to the class  $\mathbf{F}_{p_0}^1$ , thus the function  $\mathbb{R}_+ \times V \times \tilde{\Omega} \ni (s, \mathbf{r}, \tilde{\omega}) \mapsto h(\mathbf{X}_{s-}(\tilde{\omega}), \mathbf{r}) \mathbb{1}_{\{\|h(\mathbf{X}_{s-}(\tilde{\omega}), \mathbf{r})\| \geq 1\}}$  belongs to the class  $\mathbf{F}_p^1$ , and function  $\mathbb{R}_+ \times V \times \tilde{\Omega} \ni (s, \mathbf{r}, \tilde{\omega}) \mapsto \mathbf{r} \mathbb{1}_{\{\|\mathbf{r}\| < 1\}} \mathbb{1}_{\mathcal{R}_0}(\mathbf{r})$  also belongs to the class  $\mathbf{F}_p^1$  (due to [\(2.10\)](#)), thus the function  $\mathbb{R}_+ \times V \times \tilde{\Omega} \ni (s, \mathbf{r}, \tilde{\omega}) \mapsto g(\mathbf{X}_{s-}(\tilde{\omega}), \mathbf{r})$  belongs to the class  $\mathbf{F}_p^1$  as well. Moreover,

$$\begin{aligned} & \int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \hat{N}_0(ds, d\mathbf{z}) \\ &= \int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} ds \nu(d\mathbf{z}) + \sum_{j=1}^d \int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} X_{s, j} ds \mu_j(d\mathbf{z}) \\ &= \int_0^t \int_{U_d} \mathbf{r} \mathbb{1}_{\{\|\mathbf{r}\| \geq 1\}} ds \nu(d\mathbf{r}) + \sum_{j=1}^d \int_0^t X_{s, j} ds \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_j(d\mathbf{z}). \end{aligned}$$

Let  $\mathcal{M}_2$  denote the complete metric space of square integrable right continuous  $d$ -dimensional martingales on  $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$  with respect to  $(\tilde{\mathcal{F}}_t)_{t \in \mathbb{R}_+}$  starting from  $\mathbf{0}$ , see, e.g., [Ikeda and Watanabe \(1989, Chapter II, Definition 1.3 and Lemma 1.2\)](#). The function  $\mathbb{R}_+ \times U_d \times \tilde{\Omega} \ni (s, \mathbf{z}, \tilde{\omega}) \mapsto \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| < 1\}}$  belongs to the class  $\mathbf{F}_{p_0}^2$ , hence, by [Ikeda and Watanabe \(1989, Chapter II, \(3.9\)\)](#), the process  $(\int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \tilde{N}_0(ds, d\mathbf{z}))_{t \in \mathbb{R}_+}$  belongs to the space  $\mathcal{M}_2$ . Moreover, by [Ikeda and Watanabe \(1989, page 63\)](#),  $\int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \tilde{N}_0(ds, d\mathbf{z})$  is the limit of the sequence  $\int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\frac{1}{n} \leq \|\mathbf{z}\| < 1\}} \tilde{N}_0(ds, d\mathbf{z})$ ,  $n \in \mathbb{N}$ , in  $\mathcal{M}_2$  as  $n \rightarrow \infty$ . For all

$n \in \mathbb{N}$ , the mapping  $\mathbb{R}_+ \times U_d \times \tilde{\Omega} \ni (s, \mathbf{z}, \tilde{\omega}) \mapsto \mathbf{z} \mathbb{1}_{\{\frac{1}{n} \leq \|\mathbf{z}\| < 1\}}$  belongs to the class  $\mathbf{F}_{p_0}^1 \cap \mathbf{F}_{p_0}^2$ , hence we obtain

$$\begin{aligned} & \int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\frac{1}{n} \leq \|\mathbf{z}\| < 1\}} \tilde{N}_0(ds, d\mathbf{z}) \\ &= \int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\frac{1}{n} \leq \|\mathbf{z}\| < 1\}} N_0(ds, d\mathbf{z}) - \int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\frac{1}{n} \leq \|\mathbf{z}\| < 1\}} \hat{N}_0(ds, d\mathbf{z}). \end{aligned}$$

Similarly as above,

$$\begin{aligned} & \int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\frac{1}{n} \leq \|\mathbf{z}\| < 1\}} N_0(ds, d\mathbf{z}) \\ &= \int_0^t \int_{\mathcal{R}_0} \mathbf{r} \mathbb{1}_{\{\frac{1}{n} \leq \|\mathbf{r}\| < 1\}} N(ds, d\mathbf{r}) + \sum_{j=1}^d \int_0^t \int_{\mathcal{R}_j} \mathbf{z} \mathbb{1}_{\{\frac{1}{n} \leq \|\mathbf{z}\| < 1\}} \mathbb{1}_{\{u \leq X_{s-,j}\}} N(ds, d\mathbf{r}) \end{aligned}$$

and

$$\begin{aligned} \int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\frac{1}{n} \leq \|\mathbf{z}\| < 1\}} \hat{N}_0(ds, d\mathbf{z}) &= \int_0^t \int_{U_d} \mathbf{r} \mathbb{1}_{\{\frac{1}{n} \leq \|\mathbf{r}\| < 1\}} ds \nu(d\mathbf{r}) \\ &+ \sum_{j=1}^d \int_0^t \int_{U_d} \int_{U_1} \mathbf{z} \mathbb{1}_{\{\frac{1}{n} \leq \|\mathbf{z}\| < 1\}} \mathbb{1}_{\{u \leq X_{s-,j}\}} ds \mu_j(d\mathbf{z}) du. \end{aligned}$$

Consequently,

$$\begin{aligned} & \int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\frac{1}{n} \leq \|\mathbf{z}\| < 1\}} \tilde{N}_0(ds, d\mathbf{z}) \\ &= \int_0^t \int_{\mathcal{R}_0} \mathbf{r} \mathbb{1}_{\{\frac{1}{n} \leq \|\mathbf{r}\| < 1\}} \tilde{N}(ds, d\mathbf{r}) + \sum_{j=1}^d \int_0^t \int_{\mathcal{R}_j} \mathbf{z} \mathbb{1}_{\{\frac{1}{n} \leq \|\mathbf{z}\| < 1\}} \mathbb{1}_{\{u \leq X_{s-,j}\}} \tilde{N}(ds, d\mathbf{r}). \end{aligned}$$

Taking the limit in  $\mathcal{M}_2$  as  $n \rightarrow \infty$ , we conclude

$$\begin{aligned} & \int_0^t \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \tilde{N}_0(ds, d\mathbf{z}) \\ &= \int_0^t \int_{\mathcal{R}_0} \mathbf{r} \mathbb{1}_{\{\|\mathbf{r}\| < 1\}} \tilde{N}(ds, d\mathbf{r}) + \sum_{j=1}^d \int_0^t \int_{\mathcal{R}_j} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \mathbb{1}_{\{u \leq X_{s-,j}\}} \tilde{N}(ds, d\mathbf{r}) \\ &= \int_0^t \int_{\mathcal{R}_0} \mathbf{r} \mathbb{1}_{\{\|\mathbf{r}\| < 1\}} \tilde{N}(ds, d\mathbf{r}) + \int_0^t \int_{V_0} f(\mathbf{X}_{s-}, \mathbf{r}) \tilde{N}(ds, d\mathbf{r}). \end{aligned}$$

Summarizing, we conclude

$$\begin{aligned}
& \int_0^t \int_{U_d} \mathbf{z} \tilde{N}_0(ds, d\mathbf{z}) \\
&= \int_0^t \int_{V_1} g(\mathbf{X}_{s-}, \mathbf{r}) N(ds, d\mathbf{r}) - \int_0^t \int_{\mathcal{R}_0} \mathbf{r} \mathbb{1}_{\{\|\mathbf{r}\| < 1\}} N(ds, d\mathbf{r}) \\
&\quad - \int_0^t \int_{U_d} \mathbf{r} \mathbb{1}_{\{\|\mathbf{r}\| \geq 1\}} ds \nu(d\mathbf{r}) - \sum_{j=1}^d \int_0^t X_{s,j} ds \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_j(d\mathbf{z}) \\
&\quad + \int_0^t \int_{\mathcal{R}_0} \mathbf{r} \mathbb{1}_{\{\|\mathbf{r}\| < 1\}} \tilde{N}(ds, d\mathbf{r}) + \int_0^t \int_{V_0} f(\mathbf{X}_{s-}, \mathbf{r}) \tilde{N}(ds, d\mathbf{r}) \\
&= \int_0^t \int_{V_0} f(\mathbf{X}_{s-}, \mathbf{r}) \tilde{N}(ds, d\mathbf{r}) + \int_0^t \int_{V_1} g(\mathbf{X}_{s-}, \mathbf{r}) N(ds, d\mathbf{r}) \\
&\quad - \int_0^t \int_{U_d} \mathbf{r} ds \nu(d\mathbf{r}) - \sum_{j=1}^d \int_0^t X_{s,j} ds \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_j(d\mathbf{z}).
\end{aligned}$$

This proves that the SDE (3.2) holds  $\tilde{\mathbb{P}}$ -almost surely, since

$$\begin{aligned}
& \int_0^t (\tilde{\boldsymbol{\beta}} + \tilde{\mathbf{B}} \mathbf{X}_s) ds - \int_0^t \int_{U_d} \mathbf{r} ds \nu(d\mathbf{r}) - \sum_{j=1}^d \int_0^t X_{s,j} ds \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_j(d\mathbf{z}) \\
&= \tilde{\boldsymbol{\beta}} t + \tilde{\mathbf{B}} \int_0^t \mathbf{X}_s ds - t \int_{U_d} \mathbf{r} \nu(d\mathbf{r}) - \sum_{j=1}^d \int_0^t X_{s,j} ds \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_j(d\mathbf{z}) \\
&= \left( \boldsymbol{\beta} + \int_{U_d} \mathbf{r} \nu(d\mathbf{r}) \right) t + \mathbf{D} \int_0^t \mathbf{X}_s ds + \sum_{j=1}^d \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_j(d\mathbf{z}) \int_0^t X_{s,j} ds \\
&\quad - t \int_{U_d} \mathbf{r} \nu(d\mathbf{r}) - \sum_{j=1}^d \int_0^t X_{s,j} ds \int_{U_d} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_j(d\mathbf{z}) = \int_0^t (\boldsymbol{\beta} + \mathbf{D} \mathbf{X}_s) ds.
\end{aligned}$$

The aim of the following discussion is to show that  $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{G}}_t)_{t \in \mathbb{R}_+}, \tilde{\mathbb{P}}, \mathbf{W}, p, \mathbf{X})$  is an  $\mathbb{R}_+^d$ -valued weak solution to the SDE (3.2). Recall that the filtered probability space  $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{G}}_t)_{t \in \mathbb{R}_+}, \tilde{\mathbb{P}})$  satisfies the usual hypotheses, and by Lemma A.4,  $(W_{t,1}, \dots, W_{t,d})_{t \in \mathbb{R}_+}$  is a  $d$ -dimensional  $(\tilde{\mathcal{G}}_t)_{t \in \mathbb{R}_+}$ -Brownian motion, and  $p$  is a stationary  $(\tilde{\mathcal{G}}_t)_{t \in \mathbb{R}_+}$ -Poisson point process on  $V$  with characteristic measure  $m$ . Since  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  is  $\mathbb{R}_+^d$ -valued and has càdlàg trajectories on the original probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , by the definition of an extension of a probability space (see Definition A.1), the extended process (which is denoted by  $\mathbf{X}$  as well) on the extended probability space is  $\mathbb{R}_+^d$ -valued and admits càdlàg trajectories as well. By Remark A.2, the process  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  is  $(\tilde{\mathcal{G}}_t)_{t \in \mathbb{R}_+}$ -adapted, and clearly, the distribution of  $\mathbf{X}_0$  is  $n$ . Since  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  has càdlàg trajectories, (D4)(b) holds. Since the process  $(\int_0^t \int_{V_0} f(\mathbf{X}_{s-}, \mathbf{r}) \tilde{N}(ds, d\mathbf{r}))_{t \in \mathbb{R}_+}$  belongs to the space  $\mathcal{M}_2$ , we

have

$$\begin{aligned} & \tilde{\mathbb{E}} \left( \left\| \int_0^t \int_{V_0} f(\mathbf{X}_{s-}, \mathbf{r}) \tilde{N}(ds, d\mathbf{r}) \right\|^2 \right) \\ &= \sum_{j=1}^d \tilde{\mathbb{E}} \left( \int_0^t \int_{U_d} \int_{U_1} \|z\|^2 \mathbb{1}_{\{\|z\| < 1\}} \mathbb{1}_{\{u \leq X_{s,j}\}} ds \mu_j(dz) du \right) < \infty \end{aligned}$$

by [Ikeda and Watanabe \(1989, Chapter II, \(3.9\)\)](#), which yields (D4)(c). We have already checked that (D4)(d) and (D4)(e) are satisfied.

Now we turn to prove the uniqueness in the sense of probability law for the SDE (3.2) among  $\mathbb{R}_+^d$ -valued weak solutions. If  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P}, \mathbf{W}, p, \mathbf{X})$  is an  $\mathbb{R}_+^d$ -valued weak solution to the SDE (3.2), then for each  $G \in C_c^2(\mathbb{R}, \mathbb{R})$  and  $\mathbf{w} = (w_1, \dots, w_d)^\top \in \mathbb{R}^d$ , by Itô's formula for  $F(\mathbf{x}) := G(\mathbf{w}^\top \mathbf{x})$ ,  $\mathbf{x} = (x_1, \dots, x_d)^\top \in \mathbb{R}^d$ , with  $\partial_{x_k} F(\mathbf{x}) = G'(\mathbf{w}^\top \mathbf{x}) w_k$ ,  $\partial_{x_k} \partial_{x_\ell} F(\mathbf{x}) = G''(\mathbf{w}^\top \mathbf{x}) w_k w_\ell$ ,  $k, \ell \in \{1, \dots, d\}$ , we have

$$G(\mathbf{w}^\top \mathbf{X}_t) = G(\mathbf{w}^\top \mathbf{X}_0) + \sum_{\ell=1}^6 I_\ell(t), \quad t \in \mathbb{R}_+,$$

where

$$\begin{aligned} I_1(t) &:= \int_0^t G'(\mathbf{w}^\top \mathbf{X}_s) \mathbf{w}^\top (\boldsymbol{\beta} + \mathbf{D} \mathbf{X}_s) ds, \\ I_2(t) &:= \sum_{j=1}^d \int_0^t w_j G'(\mathbf{w}^\top \mathbf{X}_s) \sqrt{2c_j X_{s,j}} dW_{s,j}, \\ I_3(t) &:= \sum_{j=1}^d \int_0^t w_j^2 G''(\mathbf{w}^\top \mathbf{X}_s) c_j X_{s,j} ds, \\ I_4(t) &:= \int_0^t \int_{V_0} [G(\mathbf{w}^\top \mathbf{X}_{s-} + \mathbf{w}^\top f(\mathbf{X}_{s-}, \mathbf{r})) - G(\mathbf{w}^\top \mathbf{X}_{s-})] \tilde{N}(ds, d\mathbf{r}), \\ I_5(t) &:= \int_0^t \int_{V_0} [G(\mathbf{w}^\top \mathbf{X}_s + \mathbf{w}^\top f(\mathbf{X}_{s-}, \mathbf{r})) - G(\mathbf{w}^\top \mathbf{X}_s) \\ &\quad - G'(\mathbf{w}^\top \mathbf{X}_s) \mathbf{w}^\top f(\mathbf{X}_{s-}, \mathbf{r})] ds m(d\mathbf{r}), \\ I_6(t) &:= \int_0^t \int_{V_1} [G(\mathbf{w}^\top \mathbf{X}_{s-} + \mathbf{w}^\top g(\mathbf{X}_{s-}, \mathbf{r})) - G(\mathbf{w}^\top \mathbf{X}_{s-})] N(ds, d\mathbf{r}). \end{aligned}$$

The last integral can be written as  $I_6(t) = I_{6,1}(t) + I_{6,2}(t)$ , where

$$\begin{aligned} I_{6,1}(t) &:= \int_0^t \int_{V_1} [G(\mathbf{w}^\top \mathbf{X}_{s-} + \mathbf{w}^\top g(\mathbf{X}_{s-}, \mathbf{r})) - G(\mathbf{w}^\top \mathbf{X}_{s-})] \tilde{N}(ds, d\mathbf{r}), \\ I_{6,2}(t) &:= \int_0^t \int_{V_1} [G(\mathbf{w}^\top \mathbf{X}_s + \mathbf{w}^\top g(\mathbf{X}_s, \mathbf{r})) - G(\mathbf{w}^\top \mathbf{X}_s)] ds m(d\mathbf{r}), \end{aligned}$$



since

$$\begin{aligned}
& \mathbb{E} \left( \int_0^t \int_{V_1} |G(\mathbf{w}^\top \mathbf{X}_{s-} + \mathbf{w}^\top g(\mathbf{X}_{s-}, \mathbf{r})) - G(\mathbf{w}^\top \mathbf{X}_{s-})| \, ds \, m(d\mathbf{r}) \right) \\
&= \mathbb{E} \left( \int_0^t \int_{U_d} |G(\mathbf{w}^\top \mathbf{X}_s + \mathbf{w}^\top \mathbf{r}) - G(\mathbf{w}^\top \mathbf{X}_s)| \, ds \, \nu(d\mathbf{r}) \right) \\
&+ \sum_{j=1}^d \mathbb{E} \left( \int_0^t \int_{U_d} \int_{U_1} |G(\mathbf{w}^\top \mathbf{X}_s + \mathbf{w}^\top \mathbf{z} \mathbb{1}_{\{u \leq X_{s,j}\}}) \right. \\
&\quad \left. - G(\mathbf{w}^\top \mathbf{X}_s)| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \, ds \, \mu_j(d\mathbf{z}) \, du \right) < \infty,
\end{aligned} \tag{3.8}$$

i.e., for all  $\mathbf{w} \in \mathbb{R}^d$ , the function  $\mathbb{R}_+ \times V_1 \times \Omega \ni (s, \mathbf{r}, \omega) \mapsto G(\mathbf{w}^\top \mathbf{X}_{s-}(\omega) + \mathbf{w}^\top g(\mathbf{X}_{s-}(\omega), \mathbf{r})) - G(\mathbf{w}^\top \mathbf{X}_{s-}(\omega))$  belongs to the class  $\mathbf{F}_p^1$ . Indeed, by mean value theorem and (2.10), there exists some  $\theta_0 = \theta_0(\mathbf{w}, \mathbf{X}_s, \mathbf{r}) \in [0, 1]$  such that

$$\begin{aligned}
& \mathbb{E} \left( \int_0^t \int_{U_d} |G(\mathbf{w}^\top \mathbf{X}_s + \mathbf{w}^\top \mathbf{r}) - G(\mathbf{w}^\top \mathbf{X}_s)| \, ds \, \nu(d\mathbf{r}) \right) \\
&= \mathbb{E} \left( \int_0^t \int_{U_d} |G'(\mathbf{w}^\top \mathbf{X}_s + \theta_0 \mathbf{w}^\top \mathbf{r})| |\mathbf{w}^\top \mathbf{r}| \, ds \, \nu(d\mathbf{r}) \right) \\
&\leq \|\mathbf{w}\| \sup_{x \in \mathbb{R}} |G'(x)| \int_{U_d} \|\mathbf{r}\| \, \nu(d\mathbf{r}) < \infty
\end{aligned}$$

due to that  $G'$  is bounded. In a similar way, there exists some  $\theta = \theta(\mathbf{w}, \mathbf{X}_s, \mathbf{z}) \in [0, 1]$  such that for each  $j \in \{1, \dots, d\}$ ,

$$\begin{aligned}
& \mathbb{E} \left( \int_0^t \int_{U_d} \int_{U_1} |G(\mathbf{w}^\top \mathbf{X}_s + \mathbf{w}^\top \mathbf{z} \mathbb{1}_{\{u \leq X_{s,j}\}}) - G(\mathbf{w}^\top \mathbf{X}_s)| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \, ds \, \mu_j(d\mathbf{z}) \, du \right) \\
&= \mathbb{E} \left( \int_0^t \int_{U_d} \int_{U_1} |G(\mathbf{w}^\top \mathbf{X}_s + \mathbf{w}^\top \mathbf{z}) - G(\mathbf{w}^\top \mathbf{X}_s)| \mathbb{1}_{\{u \leq X_{s,j}\}} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \, ds \, \mu_j(d\mathbf{z}) \, du \right) \\
&= \mathbb{E} \left( \int_0^t \int_{U_d} \int_{U_1} |G'(\mathbf{w}^\top \mathbf{X}_s + \theta \mathbf{w}^\top \mathbf{z})| |\mathbf{w}^\top \mathbf{z}| \mathbb{1}_{\{u \leq X_{s,j}\}} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \, ds \, \mu_j(d\mathbf{r}) \, du \right) \\
&\leq \|\mathbf{w}\| \sup_{x \in \mathbb{R}} |G'(x)| \int_0^t \mathbb{E}(X_{s,j}) \, ds \int_{U_d} \|\mathbf{z}\| \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \, \mu_j(d\mathbf{z}) < \infty
\end{aligned}$$

due to that  $G'$  is bounded, Lemma 3.4 (which can be applied since  $\int_{\mathbb{R}_+^d} \|\mathbf{z}\| \, n(d\mathbf{z}) < \infty$ ) and the moment condition (2.2).

In what follows, we identify some of these integrals with some terms in part (5) of Theorem 9.18 of Li (2011). We have

$$\begin{aligned}
I_1(t) &= \int_0^t G'(\mathbf{w}^\top \mathbf{X}_s) \mathbf{w}^\top \boldsymbol{\beta} \, ds + \int_0^t G'(\mathbf{w}^\top \mathbf{X}_s) \mathbf{w}^\top \tilde{\mathbf{B}} \mathbf{X}_s \, ds \\
&\quad - \sum_{i=1}^d \sum_{j=1}^d \int_0^t G'(\mathbf{w}^\top \mathbf{X}_s) w_i X_{s,j} \, ds \int_{U_d} z_i \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \, \mu_j(d\mathbf{z}),
\end{aligned}$$

where, by (3.6), the first two terms on the right hand side can be identified with  $\int_0^t G'(Y_s(f)) \eta(f) \, ds$  and  $\int_0^t G'(Y_s(f)) Y_s (Af + \gamma f - bf) \, ds$ . The sum of the third

term on the right hand side and  $I_{6,2}(t) + I_5(t)$  can be written in the form

$$\begin{aligned}
& - \sum_{j=1}^d \int_0^t \int_{U_d} \int_{U_1} G'(\mathbf{w}^\top \mathbf{X}_s) \mathbf{w}^\top \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mathbb{1}_{\{u \leq X_{s,j}\}} ds \mu_j(d\mathbf{z}) du \\
& + \int_0^t \int_{U_d} [G(\mathbf{w}^\top \mathbf{X}_s + \mathbf{w}^\top \mathbf{r}) - G(\mathbf{w}^\top \mathbf{X}_s)] ds \nu(d\mathbf{r}) \\
& + \sum_{j=1}^d \int_0^t \int_{U_d} \int_{U_1} [G(\mathbf{w}^\top \mathbf{X}_s + \mathbf{w}^\top \mathbf{z} \mathbb{1}_{\{u \leq X_{s,j}\}}) - G(\mathbf{w}^\top \mathbf{X}_s)] \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} ds \mu_j(d\mathbf{z}) du \\
& + \sum_{j=1}^d \int_0^t \int_{U_d} \int_{U_1} [G(\mathbf{w}^\top \mathbf{X}_s + \mathbf{w}^\top \mathbf{z} \mathbb{1}_{\{u \leq X_{s,j}\}}) - G(\mathbf{w}^\top \mathbf{X}_s) \\
& \quad - G'(\mathbf{w}^\top \mathbf{X}_s) \mathbf{w}^\top \mathbf{z} \mathbb{1}_{\{u \leq X_{s,j}\}}] \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} ds \mu_j(d\mathbf{z}) du \\
& = \int_0^t \int_{U_d} [G(\mathbf{w}^\top \mathbf{X}_s + \mathbf{w}^\top \mathbf{r}) - G(\mathbf{w}^\top \mathbf{X}_s)] ds \nu(d\mathbf{r}) \\
& \quad + \sum_{j=1}^d \int_0^t \int_{U_d} \int_{U_1} [G(\mathbf{w}^\top \mathbf{X}_s + \mathbf{w}^\top \mathbf{z} \mathbb{1}_{\{u \leq X_{s,j}\}}) - G(\mathbf{w}^\top \mathbf{X}_s) \\
& \quad \quad - G'(\mathbf{w}^\top \mathbf{X}_s) \mathbf{w}^\top \mathbf{z} \mathbb{1}_{\{u \leq X_{s,j}\}}] ds \mu_j(d\mathbf{z}) du,
\end{aligned}$$

which can be identified with

$$\begin{aligned}
& \int_0^t \int_{M(E)^\circ} [G(Y_s(f) + \kappa(f)) - G(Y_s(f))] H(d\kappa) ds \\
& + \int_0^t \int_E Y_s(dx) \int_{M(E)^\circ} [G(Y_s(f) + \kappa(f)) - G(Y_s(f)) - \kappa(f) G'(Y_s(f))] H(x, d\kappa) ds.
\end{aligned}$$

The integral  $I_3(t)$  can be identified with  $\int_0^t G''(Y_s(f)) Y_s(c f^2) ds$ .

Next we show that the process  $(I_2(t) + I_4(t) + I_{6,1}(t))_{t \in \mathbb{R}_+}$  is a continuous local martingale. Since  $G'$  is bounded and  $\mathbf{X}$  has càdlàg trajectories, we have  $\mathbb{P}(\int_0^t w_j^2 G'(\mathbf{w}^\top \mathbf{X}_s)^2 2c_j X_{s,j} ds < \infty) = 1$  for all  $t \in \mathbb{R}_+$  and  $j \in \{1, \dots, d\}$ , hence  $(I_2(t))_{t \in \mathbb{R}_+}$  is a continuous local martingale (see, e.g., [Karatzas and Shreve \(1991, Definition 3.2.23\)](#)). In order to prove that  $(I_4(t))_{t \in \mathbb{R}_+}$  is a martingale, by page 62 in [Ikeda and Watanabe \(1989\)](#), it is enough to check that

$$\mathbb{E} \left( \int_0^t \int_{V_0} |G(\mathbf{w}^\top \mathbf{X}_s + \mathbf{w}^\top f(\mathbf{X}_s, \mathbf{r})) - G(\mathbf{w}^\top \mathbf{X}_s)|^2 ds m(d\mathbf{r}) \right) < \infty.$$

By mean value theorem, there exists some  $\vartheta_0 = \vartheta_0(\mathbf{w}, \mathbf{X}_s, \mathbf{z}) \in [0, 1]$  such that for each  $j \in \{1, \dots, d\}$ ,

$$\begin{aligned} & \mathbb{E} \left( \int_0^t \int_{U_d} \int_{U_1} |G(\mathbf{w}^\top \mathbf{X}_s + \mathbf{w}^\top \mathbf{z} \mathbb{1}_{\{u \leq X_{s,j}\}}) - G(\mathbf{w}^\top \mathbf{X}_s)|^2 \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} ds \mu_j(d\mathbf{z}) du \right) \\ &= \mathbb{E} \left( \int_0^t \int_{U_d} \int_{U_1} |G'(\mathbf{w}^\top \mathbf{X}_s + \vartheta_0 \mathbf{w}^\top \mathbf{z})|^2 (\mathbf{w}^\top \mathbf{z})^2 \mathbb{1}_{\{u \leq X_{s,j}\}} \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} ds \mu_j(d\mathbf{z}) du \right) \\ &\leq \|\mathbf{w}\|^2 \sup_{x \in \mathbb{R}} |G'(x)|^2 \int_0^t \mathbb{E}(X_{s,j}) ds \int_{U_d} \|\mathbf{z}\|^2 \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \mu_j(d\mathbf{z}) < \infty \end{aligned}$$

due to that  $G'$  is bounded, Lemma 3.4 and (2.13). Hence  $(I_4(t))_{t \in \mathbb{R}_+}$  is a martingale. Further, by (3.8) and page 62 in Ikeda and Watanabe (1989), we get  $(I_{6,1}(t))_{t \in \mathbb{R}_+}$  is a martingale. Consequently, by Theorem 9.18 of Li (2011),  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  is a CBI process with parameters  $(d, \mathbf{c}, \boldsymbol{\beta}, \mathbf{B}, \nu, \boldsymbol{\mu})$ . This yields the uniqueness in the sense of probability law for the SDE (3.2) among  $\mathbb{R}_+^d$ -valued weak solutions, and that any  $\mathbb{R}_+^d$ -valued weak solution is a CBI process with parameters  $(d, \mathbf{c}, \boldsymbol{\beta}, \mathbf{B}, \nu, \boldsymbol{\mu})$  as well.  $\square$

#### 4. Multi-type CBI process as a strong solution of an SDE

**Definition 4.1.** We say that pathwise uniqueness holds for the SDE (3.2) among  $\mathbb{R}_+^d$ -valued weak solutions if whenever

$$(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P}, \mathbf{W}, p, \mathbf{X}) \quad \text{and} \quad (\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P}, \mathbf{W}, p, \widetilde{\mathbf{X}})$$

are  $\mathbb{R}_+^d$ -valued weak solutions of the SDE (3.2) such that  $\mathbb{P}(\mathbf{X}_0 = \widetilde{\mathbf{X}}_0) = 1$ , then  $\mathbb{P}(\mathbf{X}_t = \widetilde{\mathbf{X}}_t \text{ for all } t \in \mathbb{R}_+) = 1$ .

Next we prove a comparison theorem for the SDE (3.2) in  $\boldsymbol{\beta}$ .

**Lemma 4.2.** *Let  $(d, \mathbf{c}, \boldsymbol{\beta}, \mathbf{B}, \nu, \boldsymbol{\mu})$  be a set of admissible parameters in the sense of Definition 2.2 such that the moment condition (2.7) holds. Suppose that  $\boldsymbol{\beta}' \in \mathbb{R}_+^d$  with  $\boldsymbol{\beta} \leq \boldsymbol{\beta}'$ . Let  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P}, \mathbf{W}, p, \mathbf{X})$  and  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P}, \mathbf{W}, p, \mathbf{X}')$  be  $\mathbb{R}_+^d$ -valued weak solutions of the SDE (3.2) with  $\boldsymbol{\beta}$  and  $\boldsymbol{\beta}'$ , respectively. Then  $\mathbb{P}(\mathbf{X}_0 \leq \mathbf{X}'_0) = 1$  implies  $\mathbb{P}(\mathbf{X}_t \leq \mathbf{X}'_t \text{ for all } t \in \mathbb{R}_+) = 1$ . Particularly, pathwise uniqueness holds for the SDE (3.2) among  $\mathbb{R}_+^d$ -valued weak solutions.*

*Proof:* We follow the ideas of the proof of Theorem 3.1 of Ma (2013), which is an adaptation of that of Theorem 5.5 of Fu and Li (2010). There is a sequence  $\phi_k : \mathbb{R} \rightarrow \mathbb{R}_+$ ,  $k \in \mathbb{N}$ , of twice continuously differentiable functions such that

- (i)  $\phi_k(z) \uparrow z^+$  as  $k \rightarrow \infty$ ;
- (ii)  $\phi'_k(z) \in [0, 1]$  for all  $z \in \mathbb{R}_+$  and  $k \in \mathbb{N}$ ;
- (iii)  $\phi'_k(z) = \phi_k(z) = 0$  whenever  $-z \in \mathbb{R}_+$  and  $k \in \mathbb{N}$ ;
- (iv)  $\phi''_k(x - y)(\sqrt{x} - \sqrt{y})^2 \leq 2/k$  for all  $x, y \in \mathbb{R}_+$  and  $k \in \mathbb{N}$ .

For a construction of such functions, see, e.g., the proof of Theorem 3.1 of Ma (2013). Let  $\mathbf{Y}_t := \mathbf{X}_t - \mathbf{X}'_t$  for all  $t \in \mathbb{R}_+$ . By (3.2), and using that

$$\int_0^t \int_{\mathcal{R}_0} g(\mathbf{X}_{s-}, \mathbf{r}) N(ds, d\mathbf{r}) = \int_0^t \int_{\mathcal{R}_0} \mathbf{r} N(ds, d\mathbf{r}) = \int_0^t \int_{\mathcal{R}_0} g(\mathbf{X}'_{s-}, \mathbf{r}) N(ds, d\mathbf{r}),$$

we have

$$\begin{aligned} Y_{t,i} &= Y_{0,i} + \int_0^t \left( \beta_i - \beta'_i + \mathbf{e}_i^\top \mathbf{D} \mathbf{Y}_s \right) ds + \int_0^t \sqrt{2c_i} \left( \sqrt{X_{s,i}} - \sqrt{X'_{s,i}} \right) dW_{s,i} \\ &\quad + \sum_{j=1}^d \int_0^t \int_{\mathcal{R}_{j,0}} \left( \mathbb{1}_{\{u \leq X_{s-,j}\}} - \mathbb{1}_{\{u \leq X'_{s-,j}\}} \right) z_i \mathbb{1}_{\{\|z\| < 1\}} \tilde{N}(ds, d\mathbf{r}) \\ &\quad + \sum_{j=1}^d \int_0^t \int_{\mathcal{R}_{j,1}} \left( \mathbb{1}_{\{u \leq X_{s-,j}\}} - \mathbb{1}_{\{u \leq X'_{s-,j}\}} \right) z_i \mathbb{1}_{\{\|z\| \geq 1\}} N(ds, d\mathbf{r}) \end{aligned}$$

for all  $t \in \mathbb{R}_+$  and  $i \in \{1, \dots, d\}$ . For each  $m \in \mathbb{N}$ , put

$$\tau_m := \inf \left\{ t \in \mathbb{R}_+ : \max_{i \in \{1, \dots, d\}} \max \{ X_{t,i}, X'_{t,i} \} \geq m \right\}.$$

By Itô's formula (which can be used since  $\mathbf{X}$  and  $\mathbf{X}'$  are adapted to the same filtration  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ ), we obtain

$$\phi_k(Y_{t \wedge \tau_m, i}) = \phi_k(Y_{0,i}) + \sum_{\ell=1}^6 I_{i,m,k,\ell}(t)$$

for all  $t \in \mathbb{R}_+$ ,  $i \in \{1, \dots, d\}$  and  $k, m \in \mathbb{N}$ , where

$$\begin{aligned} I_{i,m,k,1}(t) &:= \int_0^{t \wedge \tau_m} \phi'_k(Y_{s,i}) \left( \beta_i - \beta'_i + \mathbf{e}_i^\top \mathbf{D} \mathbf{Y}_s \right) ds, \\ I_{i,m,k,2}(t) &:= \int_0^{t \wedge \tau_m} \phi'_k(Y_{s,i}) \sqrt{2c_i} \left( \sqrt{X_{s,i}} - \sqrt{X'_{s,i}} \right) dW_{s,i}, \\ I_{i,m,k,3}(t) &:= \frac{1}{2} \int_0^{t \wedge \tau_m} \phi''_k(Y_{s,i}) 2c_i \left( \sqrt{X_{s,i}} - \sqrt{X'_{s,i}} \right)^2 ds, \\ I_{i,m,k,4}(t) &:= \sum_{j=1}^d \int_0^{t \wedge \tau_m} \int_{\mathcal{R}_{j,0}} \left[ \phi_k(Y_{s-,i} + (\mathbb{1}_{\{u \leq X_{s-,j}\}} - \mathbb{1}_{\{u \leq X'_{s-,j}\}}) z_i) \right. \\ &\quad \left. - \phi_k(Y_{s-,i}) \right] \mathbb{1}_{\{\|z\| < 1\}} \tilde{N}(ds, d\mathbf{r}) \\ &= \sum_{j=1}^d \int_0^{t \wedge \tau_m} \int_{\mathcal{R}_{j,0}} \left[ \phi_k(Y_{s-,i} + z_i) - \phi_k(Y_{s-,i}) \right] \mathbb{1}_{\{\|z\| < 1\}} \mathbb{1}_{\{X'_{s-,j} < u \leq X_{s-,j}\}} \\ &\quad \times \mathbb{1}_{\{Y_{s-,j} > 0\}} \tilde{N}(ds, d\mathbf{r}) \\ &\quad + \sum_{j=1}^d \int_0^{t \wedge \tau_m} \int_{\mathcal{R}_{j,0}} \left[ \phi_k(Y_{s-,i} - z_i) - \phi_k(Y_{s-,i}) \right] \mathbb{1}_{\{\|z\| < 1\}} \mathbb{1}_{\{X_{s-,j} < u \leq X'_{s-,j}\}} \\ &\quad \times \mathbb{1}_{\{Y_{s-,j} < 0\}} \tilde{N}(ds, d\mathbf{r}), \end{aligned}$$

$$\begin{aligned}
I_{i,m,k,5}(t) &:= \sum_{j=1}^d \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} \left[ \phi_k(Y_{s-,i} + (\mathbb{1}_{\{u \leq X_{s-,j}\}} - \mathbb{1}_{\{u \leq X'_{s-,j}\}})z_i) \right. \\
&\quad \left. - \phi_k(Y_{s-,i}) - \phi'_k(Y_{s-,i})(\mathbb{1}_{\{u \leq X_{s-,j}\}} - \mathbb{1}_{\{u \leq X'_{s-,j}\}})z_i \right] \mathbb{1}_{\{\|z\| < 1\}} ds \mu_j(dz) du \\
&= \sum_{j=1}^d \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} \left[ \phi_k(Y_{s-,i} + z_i) - \phi_k(Y_{s-,i}) - \phi'_k(Y_{s-,i})z_i \right] \\
&\quad \times \mathbb{1}_{\{\|z\| < 1\}} \mathbb{1}_{\{X'_{s-,j} < u \leq X_{s-,j}\}} \mathbb{1}_{\{Y_{s-,j} > 0\}} ds \mu_j(dz) du \\
&\quad + \sum_{j=1}^d \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} \left[ \phi_k(Y_{s-,i} - z_i) - \phi_k(Y_{s-,i}) + \phi'_k(Y_{s-,i})z_i \right] \\
&\quad \times \mathbb{1}_{\{\|z\| < 1\}} \mathbb{1}_{\{X_{s-,j} < u \leq X'_{s-,j}\}} \mathbb{1}_{\{Y_{s-,j} < 0\}} ds \mu_j(dz) du, \\
I_{i,m,k,6}(t) &:= \sum_{j=1}^d \int_0^{t \wedge \tau_m} \int_{\mathcal{R}_{j,1}} \left[ \phi_k(Y_{s-,i} + (\mathbb{1}_{\{u \leq X_{s-,j}\}} - \mathbb{1}_{\{u \leq X'_{s-,j}\}})z_i) \right. \\
&\quad \left. - \phi_k(Y_{s-,i}) \right] \mathbb{1}_{\{\|z\| \geq 1\}} N(ds, d\mathbf{r}),
\end{aligned}$$

where we used that

$$\mathbb{1}_{\{u \leq X_{s-,j}\}} - \mathbb{1}_{\{u \leq X'_{s-,j}\}} = \begin{cases} 1 & \text{if } Y_{s-,j} > 0 \text{ and } X'_{s-,j} < u \leq X_{s-,j}, \\ -1 & \text{if } Y_{s-,j} < 0 \text{ and } X_{s-,j} < u \leq X'_{s-,j}, \\ 0 & \text{otherwise.} \end{cases} \quad (4.1)$$

Using formula (3.8) in Chapter II in [Ikeda and Watanabe \(1989\)](#), the last integral can be written as  $I_{i,m,k,6}(t) = I_{i,m,k,6,1}(t) + I_{i,m,k,6,2}(t)$ , where

$$\begin{aligned}
I_{i,m,k,6,1}(t) &:= \sum_{j=1}^d \int_0^{t \wedge \tau_m} \int_{\mathcal{R}_{j,1}} \left[ \phi_k(Y_{s-,i} + (\mathbb{1}_{\{u \leq X_{s-,j}\}} - \mathbb{1}_{\{u \leq X'_{s-,j}\}})z_i) \right. \\
&\quad \left. - \phi_k(Y_{s-,i}) \right] \mathbb{1}_{\{\|z\| \geq 1\}} \tilde{N}(ds, d\mathbf{r}) \\
I_{i,m,k,6,2}(t) &:= \sum_{j=1}^d \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} \left[ \phi_k(Y_{s-,i} + (\mathbb{1}_{\{u \leq X_{s-,j}\}} - \mathbb{1}_{\{u \leq X'_{s-,j}\}})z_i) \right. \\
&\quad \left. - \phi_k(Y_{s-,i}) \right] \mathbb{1}_{\{\|z\| \geq 1\}} ds \mu_j(dz) du,
\end{aligned}$$

since, for each  $j \in \{1, \dots, d\}$ ,

$$\begin{aligned}
& \mathbb{E} \left( \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} \left| \phi_k(Y_{s-,i} + (\mathbb{1}_{\{u \leq X_{s-,j}\}} - \mathbb{1}_{\{u \leq X'_{s-,j}\}})z_i) - \phi_k(Y_{s-,i}) \right| \right. \\
& \quad \left. \times \mathbb{1}_{\{\|z\| \geq 1\}} ds \mu_j(dz) du \right) \\
&= \mathbb{E} \left( \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} \left| \phi_k(Y_{s-,i} + z_i) - \phi_k(Y_{s-,i}) \right| \mathbb{1}_{\{\|z\| \geq 1\}} \mathbb{1}_{\{X'_{s-,j} < u \leq X_{s-,j}\}} \right. \\
& \quad \left. \times \mathbb{1}_{\{Y_{s-,j} > 0\}} ds \mu_j(dz) du \right) \\
&+ \mathbb{E} \left( \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} \left| \phi_k(Y_{s-,i} - z_i) - \phi_k(Y_{s-,i}) \right| \mathbb{1}_{\{\|z\| \geq 1\}} \mathbb{1}_{\{X_{s-,j} < u \leq X'_{s-,j}\}} \right. \\
& \quad \left. \times \mathbb{1}_{\{Y_{s-,j} < 0\}} ds \mu_j(dz) du \right) \\
&\leq \mathbb{E} \left( \int_0^{t \wedge \tau_m} \int_{U_d} z_i \mathbb{1}_{\{\|z\| \geq 1\}} |Y_{s-,j}| ds \mu_j(dz) \right) \leq 2mt \int_{U_d} z_i \mathbb{1}_{\{\|z\| \geq 1\}} \mu_j(dz) < \infty,
\end{aligned}$$

where we used that, by properties (ii) and (iii) of the function  $\phi_k$ , we have  $\phi'_k(u) \in [0, 1]$  for all  $u \in \mathbb{R}$ , and hence, by mean value theorem,

$$-z \leq \phi_k(y-z) - \phi_k(y) \leq 0 \leq \phi_k(y+z) - \phi_k(y) \leq z, \quad y \in \mathbb{R}, \quad z \in \mathbb{R}_+, \quad k \in \mathbb{N}. \quad (4.2)$$

One can check that the process  $(I_{i,m,k,2}(t) + I_{i,m,k,4}(t) + I_{i,m,k,6,1}(t))_{t \in \mathbb{R}_+}$  is a martingale. Indeed, by properties (ii) and (iii) of the function  $\phi_k$  and the definition of  $\tau_m$ ,

$$\begin{aligned}
\mathbb{E} \left( \int_0^{t \wedge \tau_m} \left( \phi'_k(Y_{s,i}) \sqrt{2c_i} \left( \sqrt{X_{s,i}} - \sqrt{X'_{s,i}} \right) \right)^2 ds \right) &\leq 2c_i \mathbb{E} \left( \int_0^{t \wedge \tau_m} (X_{s,i} + X'_{s,i}) ds \right) \\
&\leq 4c_i mt < \infty,
\end{aligned}$$

hence, by [Ikeda and Watanabe \(1989, page 55\)](#),  $(I_{i,m,k,2}(t))_{t \in \mathbb{R}_+}$  is a martingale.

Next we show

$$\begin{aligned}
& \mathbb{E} \left( \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} |\phi_k(Y_{s-,i} + z_i) - \phi_k(Y_{s-,i})|^2 \right. \\
& \quad \left. \times \mathbb{1}_{\{\|z\| < 1\}} \mathbb{1}_{\{X'_{s-,j} < u \leq X_{s-,j}\}} \mathbb{1}_{\{Y_{s-,j} > 0\}} ds \mu_j(dz) du \right) < \infty,
\end{aligned}$$

and

$$\begin{aligned}
& \mathbb{E} \left( \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} |\phi_k(Y_{s-,i} - z_i) - \phi_k(Y_{s-,i})|^2 \right. \\
& \quad \left. \times \mathbb{1}_{\{\|z\| < 1\}} \mathbb{1}_{\{X_{s-,j} < u \leq X'_{s-,j}\}} \mathbb{1}_{\{Y_{s-,j} < 0\}} ds \mu_j(dz) du \right) < \infty
\end{aligned}$$

for all  $j \in \{1, \dots, d\}$ , which yield that the functions

$$\begin{aligned} \mathbb{R}_+ \times U_d \times U_1 \times \Omega \ni (s, \mathbf{z}, u, \omega) \mapsto & (\phi_k(Y_{s-,i}(\omega) + z_i) - \phi_k(Y_{s-,i}(\omega))) \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \\ & \times \mathbb{1}_{\{X'_{s-,j}(\omega) < u \leq X_{s-,j}(\omega)\}} \mathbb{1}_{\{Y_{s-,j}(\omega) > 0\}} \mathbb{1}_{\{s \leq \tau_m(\omega)\}} \end{aligned}$$

and

$$\begin{aligned} \mathbb{R}_+ \times U_d \times U_1 \times \Omega \ni (s, \mathbf{z}, u, \omega) \mapsto & (\phi_k(Y_{s-,i}(\omega) - z_i) - \phi_k(Y_{s-,i}(\omega))) \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \\ & \times \mathbb{1}_{\{X_{s-,j}(\omega) < u \leq X'_{s-,j}(\omega)\}} \mathbb{1}_{\{Y_{s-,j}(\omega) < 0\}} \mathbb{1}_{\{s \leq \tau_m(\omega)\}} \end{aligned}$$

belong to the class  $\mathbf{F}_p^2$ , and then  $(I_{i,m,k,4}(t))_{t \in \mathbb{R}_+}$  is a martingale, again by page 62 in [Ikeda and Watanabe \(1989\)](#). By (2.13) and (4.2),

$$\begin{aligned} & \mathbb{E} \left( \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} |\phi_k(Y_{s-,i} + z_i) - \phi_k(Y_{s-,i})|^2 \right. \\ & \quad \times \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \mathbb{1}_{\{X'_{s-,j} < u \leq X_{s-,j}\}} \mathbb{1}_{\{Y_{s-,j} > 0\}} ds \mu_j(d\mathbf{z}) du \Big) \\ & \leq \mathbb{E} \left( \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} z_i^2 \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \mathbb{1}_{\{X'_{s-,j} < u \leq X_{s-,j}\}} \mathbb{1}_{\{Y_{s-,j} > 0\}} ds \mu_j(d\mathbf{z}) du \right) \\ & = \mathbb{E} \left( \int_0^{t \wedge \tau_m} \int_{U_d} z_i^2 \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} Y_{s-,j} \mathbb{1}_{\{Y_{s-,j} > 0\}} ds \mu_j(d\mathbf{z}) \right) \\ & \leq 2mt \int_{U_d} z_i^2 \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \mu_j(d\mathbf{z}) < \infty. \end{aligned}$$

In the same way one can get the finiteness of the other expectation. Finally, we show

$$\begin{aligned} & \mathbb{E} \left( \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} |\phi_k(Y_{s-,i} + z_i) - \phi_k(Y_{s-,i})| \right. \\ & \quad \times \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mathbb{1}_{\{X'_{s-,j} < u \leq X_{s-,j}\}} \mathbb{1}_{\{Y_{s-,j} > 0\}} ds \mu_j(d\mathbf{z}) du \Big) < \infty, \end{aligned}$$

and

$$\begin{aligned} & \mathbb{E} \left( \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} |\phi_k(Y_{s-,i} - z_i) - \phi_k(Y_{s-,i})| \right. \\ & \quad \times \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mathbb{1}_{\{X_{s-,j} < u \leq X'_{s-,j}\}} \mathbb{1}_{\{Y_{s-,j} < 0\}} ds \mu_j(d\mathbf{z}) du \Big) < \infty \end{aligned}$$

for all  $j \in \{1, \dots, d\}$ , which yield that the functions

$$\begin{aligned} \mathbb{R}_+ \times U_d \times U_1 \times \Omega \ni (s, \mathbf{z}, u, \omega) \mapsto & (\phi_k(Y_{s-,i}(\omega) + z_i) - \phi_k(Y_{s-,i}(\omega))) \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \\ & \times \mathbb{1}_{\{X'_{s-,j}(\omega) < u \leq X_{s-,j}(\omega)\}} \mathbb{1}_{\{Y_{s-,j}(\omega) > 0\}} \mathbb{1}_{\{s \leq \tau_m(\omega)\}} \end{aligned}$$

and

$$\begin{aligned} \mathbb{R}_+ \times U_d \times U_1 \times \Omega \ni (s, \mathbf{z}, u, \omega) \mapsto & (\phi_k(Y_{s-,i}(\omega) - z_i) - \phi_k(Y_{s-,i}(\omega))) \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \\ & \times \mathbb{1}_{\{X_{s-,j}(\omega) < u \leq X'_{s-,j}(\omega)\}} \mathbb{1}_{\{Y_{s-,j}(\omega) < 0\}} \mathbb{1}_{\{s \leq \tau_m(\omega)\}} \end{aligned}$$

belong to the class  $F_p^1$ , and then  $(I_{i,m,k,6,1}(t))_{t \in \mathbb{R}_+}$  is a martingale, again by Ikeda and Watanabe (1989, page 62). By (2.2) and (4.2),

$$\begin{aligned}
& \mathbb{E} \left( \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} |\phi_k(Y_{s-,i} + z_i) - \phi_k(Y_{s-,i})| \right. \\
& \quad \times \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mathbb{1}_{\{X'_{s-,j} < u \leq X_{s-,j}\}} \mathbb{1}_{\{Y_{s-,j} > 0\}} \, ds \, \mu_j(d\mathbf{z}) \, du \Big) \\
& \leq \mathbb{E} \left( \int_0^{t \wedge \tau_m} \int_{U_d} \int_{U_1} z_i \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mathbb{1}_{\{X'_{s-,j} < u \leq X_{s-,j}\}} \mathbb{1}_{\{Y_{s-,j} > 0\}} \, ds \, \mu_j(d\mathbf{z}) \, du \right) \\
& = \mathbb{E} \left( \int_0^{t \wedge \tau_m} \int_{U_d} z_i \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} Y_{s-,j} \mathbb{1}_{\{Y_{s-,j} > 0\}} \, ds \, \mu_j(d\mathbf{z}) \right) \\
& \leq 2mt \int_{U_d} z_i \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_j(d\mathbf{z}) < \infty,
\end{aligned}$$

and the finiteness of the other expectation can be shown in the same way.

Using the assumption  $\beta \leq \beta'$ , the property that the matrix  $\mathbf{D}$  has non-negative off-diagonal entries and the properties (ii) and (iii), we obtain

$$\begin{aligned}
I_{i,m,k,1}(t) &= \int_0^{t \wedge \tau_m} \phi'_k(Y_{s,i}) \left( \beta_i - \beta'_i + \sum_{j=1}^d d_{i,j} Y_{s,j} \right) \, ds \\
&\leq \int_0^{t \wedge \tau_m} \phi'_k(Y_{s,i}) \left( d_{i,i} Y_{s,i} + \sum_{j \in \{1, \dots, d\} \setminus \{i\}} d_{i,j} Y_{s,j}^+ \right) \mathbb{1}_{\mathbb{R}_+}(Y_{s,i}) \, ds \\
&\leq \int_0^{t \wedge \tau_m} \left( |d_{i,i}| Y_{s,i}^+ + \sum_{j \in \{1, \dots, d\} \setminus \{i\}} d_{i,j} Y_{s,j}^+ \right) \, ds = \sum_{j=1}^d |d_{i,j}| \int_0^{t \wedge \tau_m} Y_{s,j}^+ \, ds.
\end{aligned}$$

By (iv),

$$I_{i,m,k,3}(t) \leq (t \wedge \tau_m) c_i \frac{2}{k} \leq \frac{2c_i t}{k}.$$

Now we estimate

$$\begin{aligned}
I_{i,m,k,5}(t) &= \sum_{j=1}^d \int_0^{t \wedge \tau_m} \int_{U_d} \left[ \phi_k(Y_{s-,i} + z_i) - \phi_k(Y_{s-,i}) - \phi'_k(Y_{s-,i}) z_i \right] \\
& \quad \times \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} Y_{s-,j} \mathbb{1}_{\{Y_{s-,j} > 0\}} \, ds \, \mu_j(d\mathbf{z}) \\
& \quad + \sum_{j=1}^d \int_0^{t \wedge \tau_m} \int_{U_d} \left[ \phi_k(Y_{s-,i} - z_i) - \phi_k(Y_{s-,i}) + \phi'_k(Y_{s-,i}) z_i \right] \\
& \quad \times \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} (-Y_{s-,j}) \mathbb{1}_{\{Y_{s-,j} < 0\}} \, ds \, \mu_j(d\mathbf{z}).
\end{aligned}$$



By (4.2) and (iii), we obtain

$$\begin{aligned} & \int_0^{t \wedge \tau_m} \int_{U_d} \left[ \phi_k(Y_{s-,i} - z_i) - \phi_k(Y_{s-,i}) + \phi'_k(Y_{s-,i})z_i \right] \\ & \quad \times \mathbb{1}_{\{\|z\| < 1\}}(-Y_{s-,j}) \mathbb{1}_{\{Y_{s-,j} < 0\}} \, ds \, \mu_j(dz) \leq 0 \end{aligned}$$

for all  $i, j \in \{1, \dots, d\}$ . By (2.12),  $\int_{U_d} z_i \mathbb{1}_{\{\|z\| < 1\}} \mu_j(dz) < \infty$  for all  $i, j \in \{1, \dots, d\}$  with  $i \neq j$ , hence using (iii), we obtain

$$\begin{aligned} I_{i,m,k,5}(t) & \leq \int_0^{t \wedge \tau_m} \int_{U_d} \left[ \phi_k(Y_{s-,i} + z_i) - \phi_k(Y_{s-,i}) - \phi'_k(Y_{s-,i})z_i \right] \\ & \quad \times \mathbb{1}_{\{\|z\| < 1\}} Y_{s-,i}^+ \, ds \, \mu_i(dz) \\ & \quad + \sum_{j \in \{1, \dots, d\} \setminus \{i\}} \int_0^{t \wedge \tau_m} \int_{U_d} \left[ \phi_k(Y_{s-,i} + z_i) - \phi_k(Y_{s-,i}) \right] \\ & \quad \times \mathbb{1}_{\{\|z\| < 1\}} Y_{s-,j}^+ \, ds \, \mu_j(dz). \end{aligned}$$

By (4.2), for  $i \neq j$ ,

$$\begin{aligned} & \int_0^{t \wedge \tau_m} \int_{U_d} \left[ \phi_k(Y_{s-,i} + z_i) - \phi_k(Y_{s-,i}) \right] \mathbb{1}_{\{\|z\| < 1\}} Y_{s-,j}^+ \, ds \, \mu_j(dz) \\ & \leq \int_0^{t \wedge \tau_m} Y_{s-,j}^+ \, ds \int_{U_d} z_i \mathbb{1}_{\{\|z\| < 1\}} \mu_j(dz). \end{aligned}$$

Applying (iv) with  $y = 0$ , we have  $x\phi_k''(x) \leq 2/k$  for all  $x \in \mathbb{R}_+$  and  $k \in \mathbb{N}$ . By Taylor's theorem, for all  $y \in \mathbb{R}_{++}$ ,  $z \in \mathbb{R}_+$  and  $k \in \mathbb{N}$ , there exists some  $\vartheta = \vartheta(y, z) \in [0, 1]$  such that

$$\phi_k(y+z) - \phi_k(y) - \phi'_k(y)z = \phi_k''(y+\vartheta z) \frac{z^2}{2} \leq \frac{2z^2}{2k(y+\vartheta z)} \leq \frac{z^2}{ky}.$$

Hence, using (2.13), we obtain

$$\begin{aligned} & \int_0^{t \wedge \tau_m} \int_{U_d} \left[ \phi_k(Y_{s-,i} + z_i) - \phi_k(Y_{s-,i}) - \phi'_k(Y_{s-,i})z_i \right] \mathbb{1}_{\{\|z\| < 1\}} Y_{s-,i}^+ \, ds \, \mu_i(dz) \\ & \leq \int_0^{t \wedge \tau_m} \int_{U_d} \frac{z_i^2}{kY_{s-,i}} \mathbb{1}_{\{Y_{s-,i} > 0\}} \mathbb{1}_{\{\|z\| < 1\}} Y_{s-,i}^+ \, ds \, \mu_i(dz) \\ & \leq \frac{t}{k} \int_{U_d} z_i^2 \mathbb{1}_{\{\|z\| < 1\}} \mu_i(dz). \end{aligned}$$

Using (4.1), one can easily check that

$$\begin{aligned} I_{i,m,k,6,2}(t) &= \sum_{j=1}^d \int_0^{t \wedge \tau_m} \int_{U_d} \left[ \phi_k(Y_{s-,i} + z_i) - \phi_k(Y_{s-,i}) \right] \\ &\quad \times \mathbb{1}_{\{\|z\| \geq 1\}} Y_{s-,j} \mathbb{1}_{\{Y_{s-,j} > 0\}} ds \mu_j(dz) \\ &\quad + \sum_{j=1}^d \int_0^{t \wedge \tau_m} \int_{U_d} \left[ \phi_k(Y_{s-,i} - z_i) - \phi_k(Y_{s-,i}) \right] \\ &\quad \times \mathbb{1}_{\{\|z\| \geq 1\}} (-Y_{s-,j}) \mathbb{1}_{\{Y_{s-,j} < 0\}} ds \mu_j(dz). \end{aligned}$$

By (4.2), we obtain

$$\int_0^{t \wedge \tau_m} \int_{U_d} \left[ \phi_k(Y_{s-,i} - z_i) - \phi_k(Y_{s-,i}) \right] \mathbb{1}_{\{\|z\| \geq 1\}} (-Y_{s-,j}) \mathbb{1}_{\{Y_{s-,j} < 0\}} ds \mu_j(dz) \leq 0$$

for all  $i, j \in \{1, \dots, d\}$ . By (2.2),  $\int_{U_d} z_i \mathbb{1}_{\{\|z\| \geq 1\}} \mu_j(dz) < \infty$  for all  $i, j \in \{1, \dots, d\}$ , thus applying (4.2), we obtain

$$\begin{aligned} I_{i,m,k,6,2}(t) &\leq \sum_{j=1}^d \int_0^{t \wedge \tau_m} \int_{U_d} \left[ \phi_k(Y_{s-,i} + z_i) - \phi_k(Y_{s-,i}) \right] \mathbb{1}_{\{\|z\| \geq 1\}} Y_{s-,j}^+ ds \mu_j(dz) \\ &\leq \sum_{j=1}^d \int_0^{t \wedge \tau_m} Y_{s-,j}^+ ds \int_{U_d} z_i \mathbb{1}_{\{\|z\| \geq 1\}} \mu_j(dz). \end{aligned}$$

Summarizing, we have

$$\begin{aligned} \phi_k(Y_{t \wedge \tau_m, i}) &\leq \phi_k(Y_{0, i}) + C_i \sum_{j=1}^d \int_0^{t \wedge \tau_m} Y_{s-,j}^+ ds + \frac{2c_i t}{k} + \frac{t}{k} \int_{U_d} z_i^2 \mathbb{1}_{\{\|z\| < 1\}} \mu_i(dz) \\ &\quad + I_{i,m,k,2}(t) + I_{i,m,k,4}(t) + I_{i,m,k,6,1}(t), \quad t \in \mathbb{R}_+, \end{aligned} \tag{4.3}$$

where

$$C_i := \max_{j \in \{1, \dots, d\}} |d_{i,j}| + \max_{j \in \{1, \dots, d\} \setminus \{i\}} \int_{U_d} z_i \mu_j(dz) + \int_{U_d} z_i \mathbb{1}_{\{\|z\| \geq 1\}} \mu_i(dz).$$

By (iii), we obtain  $\mathbb{P}(\phi_k(Y_{0,i}) \leq 0) = 1$ ,  $i \in \{1, \dots, d\}$ . By (i), the non-negativeness of  $\phi_k$  and monotone convergence theorem yield  $\mathbb{E}(\phi_k(Y_{t \wedge \tau_m, i})) \rightarrow \mathbb{E}(Y_{t \wedge \tau_m, i}^+)$  as  $k \rightarrow \infty$  for all  $t \in \mathbb{R}_+$ ,  $m \in \mathbb{N}$ , and  $i \in \{1, \dots, d\}$ . We have  $\int_0^{t \wedge \tau_m} Y_{s-,j}^+ ds \leq \int_0^t Y_{s \wedge \tau_m, j}^+ ds$ , hence taking the expectation of (4.3) and letting  $k \rightarrow \infty$ , we obtain

$$\mathbb{E} \left( \sum_{i=1}^d Y_{t \wedge \tau_m, i}^+ \right) \leq C \int_0^t \mathbb{E} \left( \sum_{i=1}^d Y_{s \wedge \tau_m, i}^+ \right) ds,$$

with  $C := \sum_{i=1}^d C_i$ . By Gronwall's inequality, we conclude

$$\mathbb{E} \left( \sum_{i=1}^d Y_{t \wedge \tau_m, i}^+ \right) = 0$$

for all  $t \in \mathbb{R}_+$  and  $m \in \mathbb{N}$ . Hence  $\mathbb{P}(\mathbf{X}_{t \wedge \tau_m, i} \leq \mathbf{X}'_{t \wedge \tau_m, i}) = 1$  for all  $t \in \mathbb{R}_+$ ,  $m \in \mathbb{N}$  and  $i \in \{1, \dots, d\}$ , and then  $\mathbb{P}(\mathbf{X}_{t \wedge \tau_m, i} \leq \mathbf{X}'_{t \wedge \tau_m, i}$  for all  $m \in \mathbb{N}) = 1$  for all  $t \in \mathbb{R}_+$  and  $i \in \{1, \dots, d\}$ . Since  $\mathbf{X}$  and  $\mathbf{X}'$  have càdlàg trajectories, these trajectories are bounded almost surely on  $[0, T]$  for all  $T \in \mathbb{R}_+$ , hence  $\tau_m \xrightarrow{\text{a.s.}} \infty$  as  $m \rightarrow \infty$ . This yields  $\mathbb{P}(\mathbf{X}_t \leq \mathbf{X}'_t) = 1$  for all  $t \in \mathbb{R}_+$ . Since the set of non-negative rational numbers  $\mathbb{Q}_+$  is countable, we obtain  $\mathbb{P}(\mathbf{X}_t \leq \mathbf{X}'_t$  for all  $t \in \mathbb{Q}_+) = 1$ . Using again that  $\mathbf{X}$  and  $\mathbf{X}'$  have càdlàg trajectories almost surely, we get  $\mathbb{P}(\mathbf{X}_t \leq \mathbf{X}'_t$  for all  $t \in \mathbb{R}_+) = 1$ .  $\square$

*Remark 4.3.* We note that Dawson and Li (2012, Theorem 2.3) provided a comparison theorem for SDEs with jumps in a much more general setting, but only for 1-dimensional processes.  $\square$

Consider the following objects:

- (E1) a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ ;
- (E2) a  $d$ -dimensional standard Brownian motion  $(\mathbf{W}_t)_{t \in \mathbb{R}_+}$ ;
- (E3) a stationary Poisson point process  $p$  on  $V$  with characteristic measure  $m$  given in (3.1);
- (E4) a random vector  $\xi$  with values in  $\mathbb{R}_+^d$ , independent of  $\mathbf{W}$  and  $p$ .

*Remark 4.4.* Note that if conditions (E1)–(E4) are satisfied, then  $\xi$ ,  $\mathbf{W}$  and  $p$  are automatically mutually independent according to Remark 3.3.  $\square$

Provided that the objects (E1)–(E4) are given, let  $(\mathcal{F}_t^{\xi, \mathbf{W}, p})_{t \in \mathbb{R}_+}$  be the augmented filtration generated by  $\xi$ ,  $\mathbf{W}$  and  $p$ , i.e., for each  $t \in \mathbb{R}_+$ ,  $\mathcal{F}_t^{\xi, \mathbf{W}, p}$  is the  $\sigma$ -field generated by  $\sigma(\xi; \mathbf{W}_s, s \in [0, t]; p(s), s \in (0, t] \cap D(p))$  and by the  $\mathbb{P}$ -null sets from  $\sigma(\xi; \mathbf{W}_s, s \in \mathbb{R}_+; p(s), s \in \mathbb{R}_{++} \cap D(p))$  (which is similar to the definition in Karatzas and Shreve (1991, page 285)). One can check that  $(\mathcal{F}_t^{\xi, \mathbf{W}, p})_{t \in \mathbb{R}_+}$  satisfies the usual hypotheses,  $(\mathbf{W}_t)_{t \in \mathbb{R}_+}$  is a standard  $(\mathcal{F}_t^{\xi, \mathbf{W}, p})_{t \in \mathbb{R}_+}$ -Brownian motion, and  $p$  is a stationary  $(\mathcal{F}_t^{\xi, \mathbf{W}, p})_{t \in \mathbb{R}_+}$ -Poisson point process on  $V$  with characteristic measure  $m$ , see, e.g., Barczy et al. (2015).

**Definition 4.5.** Suppose that the objects (E1)–(E4) are given. An  $\mathbb{R}_+^d$ -valued strong solution of the SDE (3.2) on  $(\Omega, \mathcal{F}, \mathbb{P})$  and with respect to the standard Brownian motion  $\mathbf{W}$ , the stationary Poisson point process  $p$  and initial value  $\xi$ , is an  $\mathbb{R}_+^d$ -valued  $(\mathcal{F}_t^{\xi, \mathbf{W}, p})_{t \in \mathbb{R}_+}$ -adapted càdlàg process  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  with  $\mathbb{P}(\mathbf{X}_0 = \xi) = 1$  satisfying (D4)(b)–(e).

Clearly,  $(\Omega, \mathcal{F}, (\mathcal{F}_t^{\xi, \mathbf{W}, p})_{t \in \mathbb{R}_+}, \mathbb{P}, \mathbf{W}, p, \mathbf{X})$  is an  $\mathbb{R}_+^d$ -valued weak solution, whenever  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  is an  $\mathbb{R}_+^d$ -valued strong solution.

**Theorem 4.6.** Let  $(d, \mathbf{c}, \beta, \mathbf{B}, \nu, \boldsymbol{\mu})$  be a set of admissible parameters in the sense of Definition 2.2 such that the moment condition (2.7) holds. Suppose that objects (E1)–(E4) are given. If  $\mathbb{E}(\|\xi\|) < \infty$ , then there is a pathwise unique  $\mathbb{R}_+^d$ -valued strong solution to the SDE (3.2) with initial value  $\xi$ , and the solution is a CBI process with parameters  $(d, \mathbf{c}, \beta, \mathbf{B}, \nu, \boldsymbol{\mu})$ .

*Proof:* The pathwise uniqueness among  $\mathbb{R}_+^d$ -valued weak solutions follows from Lemma 4.2. Then, by Theorem 5.5 in Barczy et al. (2015) (Yamada-Watanabe type result for SDEs with jumps) and Theorem 3.7, we conclude that the SDE (3.2) has a pathwise unique  $\mathbb{R}_+^d$ -valued strong solution.  $\square$

## 5. Special cases

In this section we specialize our results to dimension 1 and 2. Moreover, we consider a special case of the SDE (3.2) with  $\nu = 0$ ,  $\mu_i = 0$ ,  $i \in \{1, \dots, d\}$ , i.e., without integrals with respect to (compensated) Poisson random measures, and another special case with  $\mathbf{c} = \mathbf{0}$ , i.e., without integral with respect to a Wiener process.

First we rewrite the SDE (3.2) in a form which is more comparable with the results of Li (2011, Theorem 9.31) (one-dimensional case) and Ma (2013, Theorem 3.2) (two-dimensional case).

For each  $j \in \{0, 1, \dots, d\}$ , the thinning  $p_j$  of  $p$  onto  $\mathcal{R}_j$  is again a stationary  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -Poisson point process on  $\mathcal{R}_j$ , and its characteristic measure is the restriction  $m|_{\mathcal{R}_j}$  of  $m$  onto  $\mathcal{R}_j$  (this can be checked calculating its conditional Laplace transform, see Ikeda and Watanabe (1989, page 44)). Using these Poisson point processes, we obtain the useful decomposition

$$\begin{aligned} & \int_0^t \int_{V_0} f(\mathbf{X}_{s-}, \mathbf{r}) \tilde{N}(ds, d\mathbf{r}) + \int_0^t \int_{V_1} g(\mathbf{X}_{s-}, \mathbf{r}) N(ds, d\mathbf{r}) \\ &= \sum_{j=1}^d \int_0^t \int_{\mathcal{R}_{j,0}} z \mathbb{1}_{\{u \leq X_{s-,j}\}} \tilde{N}_j(ds, d\mathbf{r}) \\ & \quad + \sum_{j=1}^d \int_0^t \int_{\mathcal{R}_{j,1}} z \mathbb{1}_{\{u \leq X_{s-,j}\}} N_j(ds, d\mathbf{r}) + \int_0^t \int_{\mathcal{R}_0} \mathbf{r} M(ds, d\mathbf{r}), \end{aligned} \quad (5.1)$$

where, for each  $j \in \{1, \dots, d\}$ ,  $N_j(ds, d\mathbf{r})$  is the counting measure of  $p_j$  on  $\mathbb{R}_{++} \times \mathcal{R}_j$ ,  $\tilde{N}_j(ds, d\mathbf{r}) := N_j(ds, d\mathbf{r}) - ds(\mu_j(d\mathbf{z})du)$ , and  $M(ds, d\mathbf{r})$  is the counting measure of  $p_0$  on  $\mathbb{R}_{++} \times \mathcal{R}_0$ . Indeed,

$$\begin{aligned} \int_0^t \int_{\mathcal{R}'} F(s, \mathbf{r}) \tilde{N}(ds, d\mathbf{r}) &= \int_0^t \int_{\mathcal{R}'} F(s, \mathbf{r}) \tilde{N}'(ds, d\mathbf{r}), & F \in \mathbf{F}_p^{2,loc}, \\ \int_0^t \int_{\mathcal{R}'} G(s, \mathbf{r}) N(ds, d\mathbf{r}) &= \int_0^t \int_{\mathcal{R}'} G(s, \mathbf{r}) N'(ds, d\mathbf{r}), & G \in \mathbf{F}_p, \end{aligned}$$

are valid for the thinning  $p'$  of  $p$  onto any measurable subset  $\mathcal{R}' \subset \mathcal{R}$ , where  $N'(ds, d\mathbf{r})$  denotes the counting measure of the stationary  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -Poisson point process  $p'$ , and  $\tilde{N}'(ds, d\mathbf{r}) := N'(ds, d\mathbf{r}) - \mathbb{1}_{\{\mathbf{r} \in \mathcal{R}'\}} ds m(d\mathbf{r})$ .

Remark that for any  $\mathbb{R}_+^d$ -valued weak solution of the SDE (3.2), the Brownian motion  $\mathbf{W}$  and the stationary Poisson point processes  $p_j$ ,  $j \in \{0, 1, \dots, d\}$  are mutually independent according again to Theorem 6.3 in Chapter II of Ikeda and Watanabe (1989). Indeed, the intensity measures of  $p_j$ ,  $j \in \{0, 1, \dots, d\}$ , are deterministic, and condition (6.11) of this theorem is satisfied, because  $p_j$ ,  $j \in \{0, 1, \dots, d\}$ , live on disjoint subsets of  $\mathcal{R}$ .

For  $d = 1$ , applying (5.1), the SDE (3.2) takes the form

$$\begin{aligned} X_t &= X_0 + \int_0^t (\beta + dX_s) ds + \int_0^t \sqrt{2cX_s^+} dW_s + \int_0^t \int_{\mathcal{R}_{1,0}} z \mathbb{1}_{\{u \leq X_{s-}\}} \tilde{N}_1(ds, d\mathbf{r}) \\ & \quad + \int_0^t \int_{\mathcal{R}_{1,1}} z \mathbb{1}_{\{u \leq X_{s-}\}} N_1(ds, d\mathbf{r}) + \int_0^t \int_{\mathcal{R}_0} r M(ds, d\mathbf{r}) \end{aligned}$$

for  $t \in \mathbb{R}_+$ , where  $\beta \in \mathbb{R}_+$ ,  $d = \tilde{b} - \int_0^\infty z \mathbb{1}_{\{z \geq 1\}} \mu_1(dz)$ ,  $\tilde{b} = b + \int_0^\infty (z-1)^+ \mu_1(dz)$ ,  $b \in \mathbb{R}$ ,  $c \in \mathbb{R}_+$ ,  $\mathcal{R}_{1,0} = \{0\} \times \{z \in \mathbb{R}_{++} : z < 1\} \times \mathbb{R}_{++}$ ,  $\mathcal{R}_{1,1} = \{0\} \times \{z \in \mathbb{R}_{++} : z \geq 1\} \times \mathbb{R}_{++}$ ,  $\mathcal{R}_0 = \mathbb{R}_{++} \times \{(0, 0)\}$ . We have

$$\begin{aligned} I_0 &:= \int_0^t \int_{\mathcal{R}_{1,0}} z \mathbb{1}_{\{u \leq X_{s-}\}} \tilde{N}_1(ds, dr) = \int_0^t \int_0^\infty \int_0^\infty z \mathbb{1}_{\{z < 1\}} \mathbb{1}_{\{u \leq X_{s-}\}} \tilde{N}_1(ds, dz, du), \\ I_1 &:= \int_0^t \int_{\mathcal{R}_{1,1}} z \mathbb{1}_{\{u \leq X_{s-}\}} N_1(ds, dr) = \int_0^t \int_0^\infty \int_0^\infty z \mathbb{1}_{\{z \geq 1\}} \mathbb{1}_{\{u \leq X_{s-}\}} \bar{N}_1(ds, dz, du), \\ I_2 &:= \int_0^t \int_{\mathcal{R}_0} r M(ds, dr) = \int_0^t \int_0^\infty z \bar{M}(ds, dz), \end{aligned}$$

where  $\bar{N}_1$  and  $\bar{M}$  are Poisson random measures on  $\mathbb{R}_{++} \times \mathbb{R}_{++}^2$  and on  $\mathbb{R}_{++} \times \mathbb{R}_{++}$  with intensity measures  $ds \mu_1(dz) du$  and  $ds \nu(dz)$ , respectively, and  $\tilde{N}_1(ds, dz, du) := \bar{N}_1(ds, dz, du) - ds \mu_1(dz) du$ . Under the moment conditions (2.2),

$$I_0 + I_1 = \int_0^t \int_0^\infty \int_0^\infty z \mathbb{1}_{\{u \leq X_{s-}\}} \tilde{N}_1(ds, dz, du) + \int_0^t X_s ds \int_0^\infty z \mathbb{1}_{\{z \geq 1\}} \mu_1(dz).$$

Consequently, the SDE (3.2) can be rewritten in the form

$$\begin{aligned} X_t &= X_0 + \int_0^t (\beta + \tilde{b} X_s) ds + \int_0^t \sqrt{2c X_s^+} dW_s \\ &\quad + \int_0^t \int_0^\infty \int_0^\infty z \mathbb{1}_{\{u \leq X_{s-}\}} \tilde{N}_1(ds, dz, du) + \int_0^t \int_0^\infty z \bar{M}(ds, dz), \quad t \in \mathbb{R}_+, \end{aligned}$$

hence, taking into account the form (2.14) of the infinitesimal generator of the process  $(X_t)_{t \in \mathbb{R}_+}$ , we obtain equation (9.46) of Li (2011).

In a similar way, for  $d = 2$ , applying (5.1), the SDE (3.2) takes the form

$$\begin{aligned} \mathbf{X}_t &= \mathbf{X}_0 + \int_0^t (\boldsymbol{\beta} + \mathbf{D} \mathbf{X}_s) ds + \sum_{i=1}^2 \int_0^t \sqrt{2c_i X_{s,i}^+} \mathbf{e}_i \mathbf{e}_i^\top d\mathbf{W}_s + \int_0^t \int_{\mathcal{R}_0} \mathbf{r} M(ds, d\mathbf{r}) \\ &\quad + \sum_{j=1}^2 \int_0^t \int_{\mathcal{R}_{j,0}} z \mathbb{1}_{\{u \leq X_{s-,j}\}} \tilde{N}_j(ds, d\mathbf{r}) + \sum_{j=1}^2 \int_0^t \int_{\mathcal{R}_{j,1}} z \mathbb{1}_{\{u \leq X_{s-,j}\}} N_j(ds, d\mathbf{r}) \end{aligned}$$

for  $t \in \mathbb{R}_+$ , where  $\boldsymbol{\beta} \in \mathbb{R}_+^2$ ,  $\mathbf{D}$  is given in (2.8),  $(c_1, c_2)^\top \in \mathbb{R}_+^2$ ,

$$\begin{aligned} \mathcal{R}_0 &= U_2 \times \{(0, 0, 0)\} \times \{(0, 0, 0)\}, \\ \mathcal{R}_{1,0} &= \{(0, 0)\} \times \{\mathbf{z} \in U_2 : \|\mathbf{z}\| < 1\} \times \mathbb{R}_{++} \times \{(0, 0, 0)\}, \\ \mathcal{R}_{2,0} &= \{(0, 0)\} \times \{(0, 0, 0)\} \times \{\mathbf{z} \in U_2 : \|\mathbf{z}\| < 1\} \times \mathbb{R}_{++}, \\ \mathcal{R}_{1,1} &= \{(0, 0)\} \times \{\mathbf{z} \in U_2 : \|\mathbf{z}\| \geq 1\} \times \mathbb{R}_{++} \times \{(0, 0, 0)\}, \\ \mathcal{R}_{2,1} &= \{(0, 0)\} \times \{(0, 0, 0)\} \times \{\mathbf{z} \in U_2 : \|\mathbf{z}\| \geq 1\} \times \mathbb{R}_{++}. \end{aligned}$$

For each  $j \in \{1, 2\}$ , we have

$$\begin{aligned}
I_{j,0} &:= \int_0^t \int_{\mathcal{R}_{j,0}} \mathbf{z} \mathbb{1}_{\{u \leq X_{s-,j}\}} \tilde{N}_j(ds, d\mathbf{r}) \\
&= \int_0^t \int_{U_2} \int_0^\infty \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| < 1\}} \mathbb{1}_{\{u \leq X_{s-,j}\}} \tilde{N}_j(ds, d\mathbf{z}, du), \\
I_{j,1} &:= \int_0^t \int_{\mathcal{R}_{j,1}} \mathbf{z} \mathbb{1}_{\{u \leq X_{s-,j}\}} N_j(ds, d\mathbf{r}) \\
&= \int_0^t \int_{U_2} \int_0^\infty \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mathbb{1}_{\{u \leq X_{s-,j}\}} \bar{N}_j(ds, d\mathbf{z}, du), \\
I_2 &:= \int_0^t \int_{\mathcal{R}_0} \mathbf{r} M(ds, d\mathbf{r}) = \int_0^t \int_{U_2} \mathbf{z} \bar{M}(ds, d\mathbf{z}),
\end{aligned}$$

where  $\bar{N}_j$  and  $\bar{M}$  are Poisson random measures on  $\mathbb{R}_{++} \times U_2 \times \mathbb{R}_{++}$  and on  $\mathbb{R}_{++} \times U_2$  with intensity measures  $ds \mu_j(d\mathbf{z}) du$  and  $ds \nu(d\mathbf{z})$ , respectively, and  $\tilde{N}_j(ds, d\mathbf{z}, du) := \bar{N}_j(ds, d\mathbf{z}, du) - ds \mu_j(d\mathbf{z}) du$ . Under the moment conditions (2.2),

$$\begin{aligned}
I_{j,0} + I_{j,1} &= \int_0^t \int_{U_2} \int_0^\infty \mathbf{z} \mathbb{1}_{\{u \leq X_{s-,j}\}} \tilde{N}_j(ds, d\mathbf{z}, du) \\
&\quad + \int_0^t X_{s,j} ds \int_{U_2} \mathbf{z} \mathbb{1}_{\{\|\mathbf{z}\| \geq 1\}} \mu_j(d\mathbf{z}).
\end{aligned}$$

Consequently, the SDE (3.2) can be rewritten in the form

$$\begin{aligned}
\mathbf{X}_t &= \mathbf{X}_0 + \int_0^t (\boldsymbol{\beta} + \tilde{\mathbf{B}} \mathbf{X}_s) ds + \sum_{i=1}^2 \int_0^t \sqrt{2c_i X_{s,i}^+} dW_{s,i} \mathbf{e}_i \\
&\quad + \sum_{j=1}^2 \int_0^t \int_{U_2} \int_0^\infty \mathbf{z} \mathbb{1}_{\{u \leq X_{s-,j}\}} \tilde{N}_j(ds, d\mathbf{z}, du) + \int_0^t \int_{U_2} \mathbf{z} \bar{M}(ds, d\mathbf{z})
\end{aligned}$$

for  $t \in \mathbb{R}_+$ . Due to (2.12), we have

$$\begin{aligned}
X_{t,1} &= X_{0,1} + \int_0^t \left( \beta_1 + \tilde{b}_{1,1} X_{s,1} + \left( \tilde{b}_{1,2} - \int_{U_2} z_1 \mu_2(d\mathbf{z}) \right) X_{s,2} \right) ds \\
&\quad + \int_0^t \sqrt{2c_1 X_{s,1}^+} dW_{s,1} + \int_0^t \int_{U_2} \int_0^\infty z_1 \mathbb{1}_{\{u \leq X_{s-,1}\}} \tilde{N}_1(ds, d\mathbf{z}, du) \\
&\quad + \int_0^t \int_{U_2} \int_0^\infty z_1 \mathbb{1}_{\{u \leq X_{s-,2}\}} \bar{N}_2(ds, d\mathbf{z}, du) + \int_0^t \int_{U_2} z_1 \bar{M}(ds, d\mathbf{z}), \quad t \in \mathbb{R}_+,
\end{aligned}$$

and

$$\begin{aligned}
X_{t,2} &= X_{0,2} + \int_0^t \left( \beta_2 + \left( \tilde{b}_{2,1} - \int_{U_2} z_2 \mu_1(d\mathbf{z}) \right) X_{s,1} + \tilde{b}_{2,2} X_{s,2} \right) ds \\
&\quad + \int_0^t \sqrt{2c_2 X_{s,2}^+} dW_{s,2} + \int_0^t \int_{U_2} \int_0^\infty z_2 \mathbb{1}_{\{u \leq X_{s-,2}\}} \tilde{N}_2(ds, d\mathbf{z}, du) \\
&\quad + \int_0^t \int_{U_2} \int_0^\infty z_2 \mathbb{1}_{\{u \leq X_{s-,1}\}} \bar{N}_1(ds, d\mathbf{z}, du) + \int_0^t \int_{U_2} z_2 \bar{M}(ds, d\mathbf{z}), \quad t \in \mathbb{R}_+.
\end{aligned}$$

In the special case  $\nu = 0$ , we obtain equations (2.1) and (2.2) of [Ma \(2013\)](#). Indeed, due to (2.12), one can rewrite the infinitesimal generator (2.14) of the process  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  in the following form

$$\begin{aligned} (\mathcal{A}_{\mathbf{X}} f)(\mathbf{x}) &= \sum_{i=1}^2 c_i x_i f''_{i,i}(x) + \sum_{i=1}^2 x_i \int_{U_2} (f(\mathbf{x} + \mathbf{z}) - f(\mathbf{x}) - z_i f'_i(x)) \mu_i(d\mathbf{z}) \\ &\quad + \langle \beta + \tilde{\mathbf{B}}\mathbf{x}, \mathbf{f}'(\mathbf{x}) \rangle + \int_{U_2} (f(\mathbf{x} + \mathbf{z}) - f(\mathbf{x})) \nu(d\mathbf{z}) \\ &\quad - x_1 f'_2(x) \int_{U_2} z_2 \mu_1(d\mathbf{z}) - x_2 f'_1(x) \int_{U_2} z_1 \mu_2(d\mathbf{z}) \\ &= \sum_{i=1}^2 c_i x_i f''_{i,i}(x) + \sum_{i=1}^2 x_i \int_{U_2} (f(\mathbf{x} + \mathbf{z}) - f(\mathbf{x}) - z_i f'_i(x)) \mu_i(d\mathbf{z}) \\ &\quad + \langle \beta + \tilde{\mathbf{B}}\mathbf{x}, \mathbf{f}'(\mathbf{x}) \rangle + \int_{U_2} (f(\mathbf{x} + \mathbf{z}) - f(\mathbf{x})) \nu(d\mathbf{z}) \end{aligned}$$

for  $f \in C_c^2(\mathbb{R}_+^d, \mathbb{R})$  and  $\mathbf{x} \in \mathbb{R}_+^d$ , where

$$\tilde{\mathbf{B}} := \begin{bmatrix} \tilde{b}_{1,1} & \tilde{b}_{1,2} - \int_{U_2} z_1 \mu_2(d\mathbf{z}) \\ \tilde{b}_{2,1} - \int_{U_2} z_2 \mu_1(d\mathbf{z}) & \tilde{b}_{2,2} \end{bmatrix}.$$

This form of the infinitesimal generator  $\mathcal{A}_{\mathbf{X}}$  is readily comparable with the corresponding one in [Ma \(2013, equation \(1.5\)\)](#).

In what follows, we consider a special form of the SDE (3.2) without integrals with respect to (compensated) Poisson random measures. Namely, if  $\nu = 0$ ,  $\mu_i = 0$ ,  $i \in \{1, \dots, d\}$ , then the SDE (3.2) takes the form

$$\begin{aligned} \mathbf{X}_t &= \mathbf{X}_0 + \int_0^t b(\mathbf{X}_s) ds + \int_0^t \sigma(\mathbf{X}_s) d\mathbf{W}_s \\ &= \mathbf{X}_0 + \int_0^t (\beta + \mathbf{B}\mathbf{X}_s) ds + \sum_{i=1}^d \int_0^t \sqrt{2c_i X_{s,i}} \mathbf{e}_i \mathbf{e}_i^\top d\mathbf{W}_s, \quad t \in \mathbb{R}_+, \end{aligned}$$

and consequently,

$$X_{t,i} = \int_0^t \left( \beta_i + \sum_{j=1}^d b_{i,j} X_{s,j} \right) dt + \int_0^t \sqrt{2c_i X_{s,i}} dW_{s,i}, \quad t \in \mathbb{R}_+, \quad i \in \{1, \dots, d\}.$$

If  $\mathbf{B}$  is diagonal, then the process  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  is known to be a multi-factor Cox-Ingersoll-Ross process, see, e.g., [Jagannathan et al. \(2003\)](#).

Finally, Theorem 4.6 is valid also if the SDE (3.2) does not contain integral with respect to a Wiener process, i.e., if  $\mathbf{c} = \mathbf{0}$ . We note that in the proof of Theorem 3.7 we applied Theorem 7.1' in Chapter II of [Ikeda and Watanabe \(1989\)](#), which is valid in case  $\mathbf{c} = \mathbf{0}$  as well.

## Appendix

### Appendix A. Extension of a probability space

We recall the definition of extensions of probability spaces, see, e.g., [Ikeda and Watanabe \(1989, Chapter II, Definition 7.1\)](#).

**Definition A.1.** We say that a filtered probability space  $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{F}}_t)_{t \in \mathbb{R}_+}, \tilde{\mathbb{P}})$  is an extension of a filtered probability space  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P})$ , if there exists an  $\tilde{\mathcal{F}}/\mathcal{F}$ -measurable mapping  $\pi : \tilde{\Omega} \rightarrow \Omega$  such that  $\pi^{-1}(\mathcal{F}_t) \subset \tilde{\mathcal{F}}_t$  for all  $t \in \mathbb{R}_+$ ,  $\mathbb{P}(A) = \tilde{\mathbb{P}}(\pi^{-1}(A))$  for all  $A \in \mathcal{F}$ , and  $\tilde{\mathbb{E}}(\tilde{X} | \tilde{\mathcal{F}}_t)(\tilde{\omega}) = \mathbb{E}(X | \mathcal{F}_t)(\pi(\tilde{\omega}))$   $\tilde{\mathbb{P}}$ -almost surely for each essentially bounded  $(\mathcal{F}/\mathcal{B}(\mathbb{R}^d))$ -measurable random variable  $X : \Omega \rightarrow \mathbb{R}^d$ , where we set  $\tilde{X}(\tilde{\omega}) := X(\pi(\tilde{\omega}))$ ,  $\tilde{\omega} \in \tilde{\Omega}$ .

*Remark A.2.* With the notations of Definition A.1, if  $(\mathbf{X}_t)_{t \in \mathbb{R}_+}$  is an  $\mathbb{R}^d$ -valued  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -adapted stochastic process, then  $(\tilde{\mathbf{X}}_t)_{t \in \mathbb{R}_+}$  is  $(\tilde{\mathcal{F}}_t)_{t \in \mathbb{R}_+}$ -adapted. Indeed, for each  $t \in \mathbb{R}_+$  and  $B \in \mathcal{B}(\mathbb{R}^d)$ , we have

$$\begin{aligned} \tilde{\mathbf{X}}_t^{-1}(B) &= \{\tilde{\omega} \in \tilde{\Omega} : \tilde{\mathbf{X}}_t(\tilde{\omega}) \in B\} = \{\tilde{\omega} \in \tilde{\Omega} : \mathbf{X}_t(\pi(\tilde{\omega})) \in B\} \\ &= \pi^{-1}(\mathbf{X}_t^{-1}(B)) \in \tilde{\mathcal{F}}_t, \end{aligned}$$

since  $\mathbf{X}_t^{-1}(B) \in \mathcal{F}_t$ .  $\square$

**Lemma A.3.** Let  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P})$  be a filtered probability space, and let  $(\mathbf{W}_t)_{t \in \mathbb{R}_+}$  be a  $d$ -dimensional  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -Brownian motion. Let  $(\tilde{\Omega}, \tilde{\mathcal{F}}, (\tilde{\mathcal{F}}_t)_{t \in \mathbb{R}_+}, \tilde{\mathbb{P}})$  be an extension of  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P})$  with the mapping  $\pi : \tilde{\Omega} \rightarrow \Omega$ . Let  $\tilde{\mathbf{W}}_t(\tilde{\omega}) := \mathbf{W}_t(\pi(\tilde{\omega}))$  for all  $\tilde{\omega} \in \tilde{\Omega}$  and  $t \in \mathbb{R}_+$ . Then  $(\tilde{\mathbf{W}}_t)_{t \in \mathbb{R}_+}$  is a  $d$ -dimensional  $(\tilde{\mathcal{F}}_t)_{t \in \mathbb{R}_+}$ -Brownian motion.

*Proof:* According to Ikeda and Watanabe (1989, Chapter I, Definition 7.2), we have to check that the process  $(\tilde{\mathbf{W}}_t)_{t \in \mathbb{R}_+}$  has continuous trajectories, it is  $(\tilde{\mathcal{F}}_t)_{t \in \mathbb{R}_+}$ -adapted, and satisfies

$$\tilde{\mathbb{E}}(\exp\{i\langle \mathbf{u}, \tilde{\mathbf{W}}_t - \tilde{\mathbf{W}}_s \rangle\} | \tilde{\mathcal{F}}_s) = e^{-(t-s)\|\mathbf{u}\|^2/2} \quad \tilde{\mathbb{P}}\text{-almost surely}$$

for every  $\mathbf{u} \in \mathbb{R}^d$  and  $s, t \in \mathbb{R}_+$  with  $s < t$ . Clearly,  $\mathbb{R}_+ \ni t \mapsto \tilde{\mathbf{W}}_t(\tilde{\omega}) = \mathbf{W}_t(\pi(\tilde{\omega}))$  is continuous for all  $\tilde{\omega} \in \tilde{\Omega}$ . By Remark A.2,  $(\tilde{\mathbf{W}}_t)_{t \in \mathbb{R}_+}$  is  $(\tilde{\mathcal{F}}_t)_{t \in \mathbb{R}_+}$ -adapted. Finally, for every  $\mathbf{u} \in \mathbb{R}^d$  and  $s, t \in \mathbb{R}_+$  with  $s < t$ ,

$$\begin{aligned} \tilde{\mathbb{E}}(\exp\{i\langle \mathbf{u}, \tilde{\mathbf{W}}_t - \tilde{\mathbf{W}}_s \rangle\} | \tilde{\mathcal{F}}_s)(\tilde{\omega}) &= \mathbb{E}(\exp\{i\langle \mathbf{u}, \mathbf{W}_t - \mathbf{W}_s \rangle\} | \mathcal{F}_s)(\pi(\tilde{\omega})) \\ &= e^{-(t-s)\|\mathbf{u}\|^2/2} \end{aligned}$$

$\tilde{\mathbb{P}}$ -almost surely, since we have  $\xi(\omega) = c$   $\mathbb{P}$ -almost surely with  $\xi := \mathbb{E}(\exp\{i\langle \mathbf{u}, \mathbf{W}_t - \mathbf{W}_s \rangle\} | \mathcal{F}_s)$  and  $c := e^{-(t-s)\|\mathbf{u}\|^2/2}$ , which implies  $\xi(\pi(\tilde{\omega})) = c$   $\tilde{\mathbb{P}}$ -almost surely, because  $\tilde{\mathbb{P}}(\{\tilde{\omega} \in \tilde{\Omega} : \xi(\pi(\tilde{\omega})) = c\}) = \tilde{\mathbb{P}}(\pi^{-1}(\xi^{-1}(\{c\}))) = \mathbb{P}(\xi^{-1}(\{c\})) = 1$ .  $\square$

**Lemma A.4.** Let  $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P})$  be a filtered probability space, let  $(\mathbf{W}_t)_{t \in \mathbb{R}_+}$  be a  $d$ -dimensional  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -Brownian motion, and let  $p$  be a stationary  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -Poisson point process on  $V = \mathbb{R}_+^d \times (\mathbb{R}_+^d \times \mathbb{R}_+)^d$  with characteristic measure  $m$ , where  $m$  is given in (3.1). Let

$$\mathcal{G}_t := \bigcap_{\varepsilon > 0} \sigma(\mathcal{F}_{t+\varepsilon} \cup \mathcal{N}), \quad t \in \mathbb{R}_+,$$

where  $\mathcal{N}$  denotes the collection of null sets under the probability measure  $\mathbb{P}$ . Then  $(\mathbf{W}_t)_{t \in \mathbb{R}_+}$  is a  $d$ -dimensional  $(\mathcal{G}_t)_{t \in \mathbb{R}_+}$ -Brownian motion, and  $p$  is a stationary  $(\mathcal{G}_t)_{t \in \mathbb{R}_+}$ -Poisson point process on  $V$  with characteristic measure  $m$ .

*Proof:* The proof is essentially the same as the proof of Lemma A.5 in Barczy et al. (2015).  $\square$



## Acknowledgements

We would like to thank the referee for his/her comments that helped us to improve the presentation of the Introduction.

## References

- M. Barczy, Z. Li and G. Pap. Yamada-Watanabe results for stochastic differential equations with jumps. *Int. J. Stoch. Anal.* **2015** (Article ID 460472), 23 pages (2015). [MR3298537](#).
- D. A. Dawson and Z. Li. Skew convolution semigroups and affine Markov processes. *Ann. Probab.* **34** (3), 1103–1142 (2006). [MR2243880](#).
- D. A. Dawson and Z. Li. Stochastic equations, flows and measure-valued processes. *Ann. Probab.* **40** (2), 813–857 (2012). [MR2952093](#).
- D. Duffie, D. Filipović and W. Schachermayer. Affine processes and applications in finance. *Ann. Appl. Probab.* **13** (3), 984–1053 (2003). [MR1994043](#).
- M. C. Fittipaldi and J. Fontbona. On sde associated with continuous-state branching processes conditioned to never be extinct. *Electron. Commun. Probab.* **17** (49), 1–13 (2012). [MR2988395](#).
- Z. Fu and Z. Li. Stochastic equations of non-negative processes with jumps. *Stochastic Process. Appl.* **120** (3), 306–330 (2010). [MR2584896](#).
- N. Ikeda and S. Watanabe. *Stochastic Differential Equations and Diffusion Processes*. North-Holland/Kodansha, Amsterdam/Tokyo (1989). 2nd ed. [MR1011252](#).
- R. Jagannathan, A. Kaplin and S. Sun. An evaluation of multi-factor CIR models using LIBOR, swap rates, and cap and swaption prices. *J. Econometrics* **116** (1–2), 113–146 (2003). [MR2011151](#).
- M. Jiřina. Stochastic branching processes with continuous state space. *Czechoslovak Math. J.* **8** (2), 292–313 (1958). [MR0101554](#).
- I. Karatzas and S. E. Shreve. *Brownian Motion and Stochastic Calculus*. Springer, Berlin (1991). 2nd ed. [MR1121940](#).
- K. Kawazu and S. Watanabe. Branching processes with immigration and related limit theorems. *Theory Probab. Appl.* **16** (1), 36–54 (1971). [MR0290475](#).
- Z. Li. A limit theorem for discrete Galton-Watson branching processes with immigration. *J. Appl. Probab.* **43** (1), 289–295 (2006). [MR2225068](#).
- Z. Li. *Measure-Valued Branching Markov Processes*. Springer-Verlag, Heidelberg (2011). [MR2760602](#).
- R. Ma. Stochastic equations for two-type continuous-state branching processes with immigration. *Acta Math. Sin. (Engl. Ser.)* **29** (2), 287–294 (2013). [MR3016529](#).
- D. W. Stroock. *Probability Theory. An Analytic View*. Cambridge University Press, Cambridge (2011). 2nd ed. [MR2760872](#).
- S. Watanabe. On two dimensional Markov processes with branching property. *Trans. Amer. Math. Soc.* **136**, 447–466 (1969). [MR0234531](#).