



ORAL PRESENTATION

Open Access

# Stratification of patients with autoinflammatory phenotypes by interferon (IFN) score suggests a new group of IFN mediated autoinflammatory diseases with overlapping clinical phenotypes

A Almeida de Jesus<sup>1\*</sup>, Z Deng<sup>2</sup>, S Brooks<sup>2</sup>, H Kim<sup>1</sup>, G Montealegre<sup>1</sup>, D Chapelle<sup>1</sup>, Y Liu<sup>2</sup>, B Marrero<sup>1</sup>, L Malle<sup>1</sup>, M O'Brien<sup>1</sup>, W Goodspeed<sup>1</sup>, Y Huang<sup>1</sup>, P Hashkes<sup>3</sup>, G Nasrullayeva<sup>4</sup>, MT Terreri<sup>5</sup>, C Silva<sup>6</sup>, B Arabshahi<sup>7</sup>, K O'Neill<sup>8</sup>, M Punaro<sup>9</sup>, L Moorthy<sup>10</sup>, A Reinhardt<sup>11</sup>, V Lilleby<sup>12</sup>, J Niemela<sup>13</sup>, S Rosenzweig<sup>13</sup>, T Fleisher<sup>13</sup>, R Goldbach-Mansky<sup>1,13</sup>

From 8th International Congress of Familial Mediterranean Fever and Systemic Autoinflammatory Diseases Dresden, Germany. 30 September - 3 October 2015

## Background

We have identified mutations in proteasome components as the cause of CANDLE syndrome and in *TMEM173/STING* as the cause for a severe vasculopathy and lung disease, SAVI. CANDLE and SAVI patients do not respond to IL-1 inhibition and consistently demonstrate marked up-regulation of IFN-inducible genes. Our data suggest innate immune dysregulation caused by chronic Type I IFN signaling in both conditions.

## Objective

We hypothesize that the presence of IFN signature may identify patients with autoinflammatory disease (AID) who have genetic mutations in other IFN regulating genes.

## Methods

To identify patients with IFN signatures, RNA sequencing (RNA-seq) from whole blood RNA was performed using HiSeq 2000 Illumina<sup>®</sup> platform. Heatmaps with 64 IFN response genes were assessed. Whole exome sequencing (WES) was performed from whole blood DNA.

## Results

We identified 19 patients with marked upregulation of IFN inducible genes. WES was performed in 14 patients

and parents (trios) and in 5 individual patients. Of the probands, 9/19 were female, 8/19 were Caucasian, 3 Asian, 2 Hispanic, 2 Norwegian and 4 had other ethnicities. All patients presented with immunodysregulatory phenotypes with clinical similarities to the previously described interferonopathies, including skin vasculitis/vasculopathy (9/19), panniculitis (12/19), myositis (5/19) and basal ganglion calcifications (5/19), but had no genetic diagnosis prior to NIH evaluation. The bioinformatics variant annotation, analysis and filtering workflow successfully identified mutations in IFN-regulating genes in 7 of the 19 probands. In one patient, we found a disease causing *de novo* and somatic mutation in *TREX1*. This patient also presented with an in-frame deletion in *DHX9* inherited from her mother and a missense mutation in *MAVS* inherited from her father. In one patient, we identified a *de novo* mutation in *DHX9* and this patient is also a compound heterozygous for mutations in *IFIH1/MDA5*. In a third patient, we found a missense mutation in *TREX1* inherited from the mother and a heterozygous variant in *MB21D1* (gene encoding cGAS) inherited from the father. A fourth patient with a clinical phenotype of CANDLE had two novel compound heterozygous mutations in *PSMG2*. Additionally, a male patient with lupus-like clinical and laboratory findings was found to have an X-linked mutation in *TREX2* gene. All mutations described were confirmed by Sanger sequencing.

<sup>1</sup>National Institutes of Health, Translational Autoinflammatory Diseases Section, Bethesda, USA

Full list of author information is available at the end of the article

## Conclusion

RNA-seq can be a tool for the identification of patients with an IFN signature and guide the search for disease causing variants in IFN-regulating genes by WES. However, disease causality of these mutations needs to be assessed in functional assays. Moreover the identification of patients with a type I interferon signature and a set of clinical features that are not seen in IL-1- mediated-AIDs allow stratification of a subset of AIDs that are typically “poor IL-1 responsive”. Whether the IFN signature identifies a subset of patients that respond to the blockade of Type I IFN signaling needs to be further validated.

## Authors' details

<sup>1</sup>National Institutes of Health, Translational Autoinflammatory Diseases Section, Bethesda, USA. <sup>2</sup>National Institutes of Health, NIAMS, Bethesda, USA. <sup>3</sup>Hebrew University School of Medicine, Jerusalem, Israel. <sup>4</sup>Azerbaijan Medical University, Department of Immunology, Baku, Azerbaijan. <sup>5</sup>Federal University of Sao Paulo, Sao Paulo, Brazil. <sup>6</sup>University of Sao Paulo, Sao Paulo, Brazil. <sup>7</sup>Inova Fairfax Hospital, Fairfax, USA. <sup>8</sup>Riley Children's Hospital, Indianapolis, USA. <sup>9</sup>Children's Medical Center Dallas, Dallas, USA. <sup>10</sup>Robert Wood Johnson University Hospital, New Brunswick, USA. <sup>11</sup>University of Nebraska Medical Center, Omaha, USA. <sup>12</sup>Rikshospitalet University Hospital, Oslo, Norway. <sup>13</sup>National Institutes of Health, Department of Laboratory Medicine, Bethesda, USA.

Published: 28 September 2015

doi:10.1186/1546-0096-13-S1-O35

**Cite this article as:** de Jesus *et al.*: Stratification of patients with autoinflammatory phenotypes by interferon (IFN) score suggests a new group of IFN mediated autoinflammatory diseases with overlapping clinical phenotypes. *Pediatric Rheumatology* 2015 **13**(Suppl 1):O35.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

