

Strong law of large numbers for supercritical superprocesses under second moment condition

Zhen-Qing CHEN¹, Yan-Xia REN², Renming SONG³, Rui ZHANG⁴

¹ Department of Mathematics, University of Washington, Seattle, WA 98195, USA

² LMAM School of Mathematical Sciences & Center for Statistical Science, Peking University, Beijing 100871, China

³ Department of Mathematics, University of Illinois, Urbana, IL 61801, USA

⁴ LMAM School of Mathematical Sciences, Peking University, Beijing 100871, China

© Higher Education Press and Springer-Verlag Berlin Heidelberg 2015

Abstract Consider a supercritical superprocess $X = \{X_t, t \geq 0\}$ on a locally compact separable metric space (E, m) . Suppose that the spatial motion of X is a Hunt process satisfying certain conditions and that the branching mechanism is of the form

$$\psi(x, \lambda) = -a(x)\lambda + b(x)\lambda^2 + \int_{(0, +\infty)} (e^{-\lambda y} - 1 + \lambda y)n(x, dy), \quad x \in E, \lambda > 0,$$

where $a \in \mathcal{B}_b(E)$, $b \in \mathcal{B}_b^+(E)$, and n is a kernel from E to $(0, +\infty)$ satisfying $\sup_{x \in E} \int_0^{+\infty} y^2 n(x, dy) < +\infty$. Put $T_t f(x) = \mathbb{P}_{\delta_x} \langle f, X_t \rangle$. Suppose that the semigroup $\{T_t; t \geq 0\}$ is compact. Let λ_0 be the eigenvalue of the (possibly non-symmetric) generator L of $\{T_t\}$ that has the largest real part among all the eigenvalues of L , which is known to be real-valued. Let ϕ_0 and $\hat{\phi}_0$ be the eigenfunctions of L and \hat{L} (the dual of L) associated with λ_0 , respectively. Assume $\lambda_0 > 0$. Under some conditions on the spatial motion and the ϕ_0 -transform of the semigroup $\{T_t\}$, we prove that for a large class of suitable functions f ,

$$\lim_{t \rightarrow +\infty} e^{-\lambda_0 t} \langle f, X_t \rangle = W_\infty \int_E \hat{\phi}_0(y) f(y) m(dy), \quad \mathbb{P}_\mu\text{-a.s.},$$

for any finite initial measure μ on E with compact support, where W_∞ is the martingale limit defined by $W_\infty := \lim_{t \rightarrow +\infty} e^{-\lambda_0 t} \langle \phi_0, X_t \rangle$. Moreover, the exceptional set in the above limit does not depend on the initial measure μ and the function f .

Received February 5, 2015; accepted March 18, 2015

Corresponding author: Yan-Xia REN, E-mail: yxren@math.pku.edu.cn

Keywords Superprocess, scaling limit theorem, Hunt process, spectral gap, h -transform, martingale measure

MSC 60J80, 60F15, 60J25

1 Introduction

Recently, there have been quite a few papers on law of large numbers for superdiffusions. In [11,13,14], some weak laws of large numbers (convergence in law or in probability) were established. The strong law of large numbers for superprocesses was first studied in [7] followed by [10,20,22,31]. The continuity of the sample paths of the spatial motions played an important role in all the papers mentioned above except [7,20]. It is more difficult to establish strong law of large numbers for superprocesses with discontinuous spatial motions. For a good survey on recent developments in laws of large numbers for branching Markov processes and superprocesses, see [10]. In the papers mentioned above, either the spatial motion is assumed to be a diffusion, or the spatial motion is assumed to be a symmetric Hunt process. In [7], where the spatial motion is a symmetric Hunt process, a condition on the smallness at ‘infinity’ of the linear term in the branching mechanism of the superprocess has to be assumed. The purpose of this paper is to give a different setup under which the strong law of large numbers for superprocesses holds. The setup of this paper complements the previous setups. In particular, the spatial motion may be discontinuous and non-symmetric. We will give some examples that satisfy the conditions of this paper.

Papers [7,10,22] dealt with strong law of large numbers for superprocesses with spatially dependent branching mechanism. The main ideas of the arguments of [7,10,22] are similar and consist of two steps. The first step is to prove an almost sure limit result along discrete times, and the second step is to prove that the result is true for continuous times. An essential difficulty comes from the second step. [22] gave a method for the transition from lattice times to continuous times based on resolvent operator and approximation of the indicator function of an open subset of E by resolvent functions. The reason that this approximation works for superdiffusions is that the sample paths of the spatial motion are continuous. [10] also used this idea to show that indicator functions can be approximated by resolvent functions. For general superprocesses with spatial motions which might be discontinuous, [7] is the first paper to establish a strong law of large numbers under a second moment condition. The paper [7] managed to overcome the difficulty of transition from discrete times to continuous times with a highly non-trivial application of the martingale formulation of superprocesses. However, the assumptions of [7] are restrictive in two aspects: the spatial motion is assumed to be symmetric and the linear term of the branching mechanism is assumed to satisfy a Kato class condition at ‘infinity’.

Papers [20,31] studied strong law of large numbers for super-Brownian

motions and super- α -stable processes with spatially independent branching mechanism, respectively. The key ingredients in the argument of [20,31] are Fourier analysis and stochastic analysis, and the conditions in [20,31] are quite different from those of [7,22]. The mean semigroup of the superprocess is assumed to have a spectral gap in [7,22], while the mean semigroups of the superprocesses of [20,31] have continuous spectra. In this paper, we assume that the spatial motion has a dual with respect to a certain measure and that the branching mechanism satisfies a second moment condition. Under the conditions of this paper, the mean semigroup of the superprocess automatically has a spectral gap.

1.1 Spatial process

Our assumptions on the underlying spatial process are similar to those in [25]. In this subsection, we lay out the assumptions on the spatial process.

Suppose that (E, m) is a locally compact separable metric space and m is a σ -finite Borel measure on E with full support. Let $E_\partial = E \cup \{\partial\}$ be the one-point compactification of E . Every function f on E is automatically extended to E_∂ by setting $f(\partial) = 0$. We will assume that $\xi = \{\xi_t, \Pi_x\}$ is a Hunt process on E and $\zeta := \inf\{t > 0: \xi_t = \partial\}$ is the lifetime of ξ . The transition semigroup of ξ will be denoted by $\{P_t, t \geq 0\}$. We will always assume that there exists a family of strictly positive continuous functions $\{p(t, x, y), t > 0\}$ on $E \times E$ such that

$$P_t f(x) = \int_E p(t, x, y) f(y) m(dy).$$

Define

$$a_t(x) := \int_E p^2(t, x, y) m(dy), \quad \widehat{a}_t(x) := \int_E p^2(t, y, x) m(dy). \quad (1.1)$$

In this paper, we make the following assumption.

Assumption 1.1 (a) For all $t > 0$ and $x \in E$, $\int_E p(t, y, x) m(dy) \leq 1$.

(b) For any $t > 0$, a_t and \widehat{a}_t are continuous $L^1(E; m)$ -integrable functions.

(c) There exists $t_0 > 0$ such that $a_{t_0}, \widehat{a}_{t_0} \in L^2(E; m)$.

By the Chapman-Kolmogorov equation and the Cauchy-Schwarz inequality,

$$p(t+s, x, y) = \int_E p(t, x, z) p(s, z, y) m(dz) \leq (a_t(x))^{1/2} (\widehat{a}_s(y))^{1/2}. \quad (1.2)$$

Therefore,

$$a_{t+s}(x) \leq a_t(x) \int_E \widehat{a}_s(y) m(dy), \quad \widehat{a}_{t+s}(x) \leq \widehat{a}_t(x) \int_E a_s(y) m(dy).$$

Thus, under condition (b), condition (c) above is equivalent to

(c') there exists $t_0 > 0$ such that for all $t \geq t_0$, $a_t, \widehat{a}_t \in L^2(E; m)$.

Under Assumption 1.1 (a), for every $t > 0$, both P_t and the operator \widehat{P}_t defined by

$$\widehat{P}_t f(x) = \int_E p(t, y, x) f(y) m(dy)$$

are contraction operators in $L^p(E; m)$ for every $p \in [1, +\infty]$, and they are dual to each other. Assumption 1.1 (b) implies that each P_t is a Hilbert-Schmidt operator in $L^2(E; m)$ and thus is compact. Hence, P_t has discrete spectrum.

1.2 Superprocesses

In this subsection, we introduce the superprocesses. Let $\mathcal{B}_b(E)$ (resp. $\mathcal{B}_b^+(E)$) be the family of bounded (resp. nonnegative bounded) Borel functions on E . Denote by $\langle \cdot, \cdot \rangle_m$ the inner product in $L^2(E; m)$.

The superprocess $X = \{X_t, t \geq 0\}$ is determined by three objects: a spatial motion $\xi = \{\xi_t, \Pi_x\}$ on E satisfying the assumptions of the previous subsection, a branching rate function $\beta(x)$ on E which is a nonnegative bounded Borel function, and a branching mechanism ψ of the form

$$\psi(x, \lambda) = -a(x)\lambda + b(x)\lambda^2 + \int_{(0, +\infty)} (e^{-\lambda y} - 1 + \lambda y) n(x, dy), \quad x \in E, \lambda > 0, \quad (1.3)$$

where $a \in \mathcal{B}_b(E)$, $b \in \mathcal{B}_b^+(E)$, and n is a kernel from E to $(0, +\infty)$ satisfying

$$\sup_{x \in E} \int_0^{+\infty} y^2 n(x, dy) < +\infty. \quad (1.4)$$

Let $\mathcal{M}_F(E)$ be the space of finite measures on E , equipped with the weak convergence topology. As usual,

$$\langle f, \mu \rangle := \int f(x) \mu(dx), \quad \|\mu\| := \langle 1, \mu \rangle.$$

By [21, Theorem 5.12], there is a Borel right process $X = \{\Omega, \mathcal{G}, \mathcal{G}_t, X_t, \mathbb{P}_\mu\}$ taking values in $\mathcal{M}_F(E)$, called superprocess, such that for every $f \in \mathcal{B}_b^+(E)$ and $\mu \in \mathcal{M}_F(E)$,

$$-\log \mathbb{P}_\mu(e^{-\langle f, X_t \rangle}) = \langle u_f(\cdot, t), \mu \rangle, \quad (1.5)$$

where $u_f(x, t)$ is the unique positive solution to the equation

$$u_f(x, t) + \Pi_x \int_0^t \psi(\xi_s, u_f(\xi_s, t-s)) \beta(\xi_s) ds = \Pi_x f(\xi_t), \quad (1.6)$$

where $\psi(\partial, \lambda) = 0$, $\lambda > 0$. Here, $(\mathcal{G}, \mathcal{G}_t)_{t \geq 0}$ are augmented, $(\mathcal{G}_t, t \geq 0)$ is right continuous and X satisfies the Markov property with respect to $(\mathcal{G}_t, t \geq 0)$. Moreover, such a superprocess X has a Hunt realization in $\mathcal{M}_F(E)$, see [21, Theorem 5.12]. In this paper, the superprocess we deal with always takes such a Hunt realization.

Define

$$\alpha(x) := \beta(x)a(x), \quad A(x) := \beta(x) \left(2b(x) + \int_0^{+\infty} y^2 n(x, dy) \right). \quad (1.7)$$

By our assumptions, $\alpha(x) \in \mathcal{B}_b(E)$ and $A(x) \in \mathcal{B}_b^+(E)$. Thus, there exists $K > 0$ such that

$$\sup_{x \in E} (|\alpha(x)| + A(x)) \leq K. \quad (1.8)$$

For any $f \in \mathcal{B}_b(E)$ and $(t, x) \in (0, +\infty) \times E$, define

$$T_t f(x) := \Pi_x [e^{\int_0^t \alpha(\xi_s) ds} f(\xi_t)]. \quad (1.9)$$

It is well known that

$$T_t f(x) = \mathbb{P}_{\delta_x} \langle f, X_t \rangle, \quad \forall x \in E.$$

It is known that (see, e.g., [25] and [27, Lemma 2.1]) $\{T_t, t \geq 0\}$ is a strongly continuous semigroup on $L^2(E; m)$ and there exists a function $q(t, x, y)$ on $(0, +\infty) \times E \times E$ which is continuous in (x, y) for each $t > 0$ such that

$$e^{-Kt} p(t, x, y) \leq q(t, x, y) \leq e^{Kt} p(t, x, y), \quad (t, x, y) \in (0, +\infty) \times E \times E, \quad (1.10)$$

and that for any bounded Borel function f on E and $(t, x) \in (0, +\infty) \times E$,

$$T_t f(x) = \int_E q(t, x, y) f(y) m(dy).$$

It follows immediately that, for any $f \in L^p(E, m)$ with $p \geq 1$,

$$\|T_t f\|_p \leq e^{Kt} \|P_t f\|_p \leq e^{Kt} \|f\|_p. \quad (1.11)$$

Define

$$b_t(x) := \int_E q^2(t, x, y) m(dy), \quad \widehat{b}_t(x) := \int_E q^2(t, y, x) m(dy). \quad (1.12)$$

Then b_t and \widehat{b}_t enjoy the following properties.

(i) For any $t > 0$, we have $b_t, \widehat{b}_t \in L^1(E; m)$. Moreover, $b_t(x)$ and $\widehat{b}_t(x)$ are continuous in $x \in E$.

(ii) There exists $t_0 > 0$ such that for all $t \geq t_0$, $b_t, \widehat{b}_t \in L^2(E; m)$.

Let $\{\widehat{T}_t, t > 0\}$ be the adjoint semigroup of $\{T_t, t \geq 0\}$ on $L^2(E, m)$ defined by

$$\widehat{T}_t g(x) = \int_E q(t, y, x) g(y) m(dy).$$

It is easy to see \widehat{T}_t is the dual operator of T_t in $L^2(E; m)$. It follows that $\{\widehat{T}_t, t > 0\}$ is also strongly continuous in $L^2(E, m)$. Since $q(t, \cdot, y)$ and a_t are

continuous, by (1.2) and (1.10), using the dominated convergence theorem, we get that for any $t > 0$ and $f \in L^2(E; m)$, $T_t f$ and $\widehat{T}_t f$ are continuous.

It follows from (i) above that, for any $t > 0$, T_t and \widehat{T}_t are compact operators in $L^2(E; m)$. Let L and \widehat{L} be the infinitesimal generators of the semigroups $\{T_t\}$ and $\{\widehat{T}_t\}$ in $L^2(E; m)$, respectively. Let $\sigma(L)$ and $\sigma(\widehat{L})$ be the spectra of L and \widehat{L} , respectively. It follows from [23, Theorem 2.2.4, Corollary 2.3.7] that both $\sigma(L)$ and $\sigma(\widehat{L})$ consist of eigenvalues, and that $\sigma(L)$ and $\sigma(\widehat{L})$ have the same number, say N , of eigenvalues. Let

$$\mathbb{I} = \begin{cases} \{0, 1, 2, \dots, N-1\}, & N < +\infty, \\ \{0, 1, 2, \dots\}, & \text{otherwise.} \end{cases}$$

For a complex number a , we use $\operatorname{Re}(a)$ to denote its real part. Define

$$\lambda_0 := \sup \operatorname{Re}(\sigma(L)) = \sup \operatorname{Re}(\sigma(\widehat{L})).$$

By Jentzsch's theorem ([29, Theorem V.6.6]), λ_0 is an eigenvalue of multiplicity 1 for both L and \widehat{L} ; see [8, Theorem 7.1] and its proof for details. Assume that ϕ_0 and $\widehat{\phi}_0$ are eigenfunctions of L and \widehat{L} , respectively, associated with λ_0 . Functions ϕ_0 and $\widehat{\phi}_0$ can be chosen to be continuous strictly positive and satisfy

$$\|\phi_0\|_2 = 1, \quad \langle \phi_0, \widehat{\phi}_0 \rangle_m = 1.$$

We list the eigenvalues of $\{\lambda_k, k \in \mathbb{I}\}$ of L in an order so that

$$\lambda_0 > \operatorname{Re}(\lambda_1) \geq \operatorname{Re}(\lambda_2) \geq \dots$$

Then $\{\bar{\lambda}_k, k \in \mathbb{I}\}$ are the eigenvalues of \widehat{L} . For convenience, we define, for any positive integer not in \mathbb{I} ,

$$\lambda_k = \bar{\lambda}_k = -\infty.$$

For $t > 0$,

$$T_t \phi_0(x) = e^{\lambda_0 t} \phi_0(x),$$

and thus,

$$\phi_0(x) \leq e^{-\lambda_0 t} b_t^{1/2}(x). \quad (1.13)$$

Similarly, we have

$$\widehat{T}_t \widehat{\phi}_0(x) = e^{\lambda_0 t} \widehat{\phi}_0(x), \quad \widehat{\phi}_0(x) \leq e^{-\lambda_0 t} \|\widehat{\phi}_0\|_2 \widehat{b}_t^{1/2}(x).$$

Therefore, by Assumption 1.1 (c),

$$\phi_0 \in L^2(E; m) \cap L^4(E; m).$$

In this paper, we always assume that the superprocess X is supercritical, that is, $\lambda_0 > 0$. Define

$$W_t := e^{-\lambda_0 t} \langle \phi_0, X_t \rangle.$$

By the Markov property of X , $\{W_t, t \geq 0\}$ is a nonnegative martingale with respect to $\{\mathcal{G}_t, t \geq 0\}$, and thus,

$$W_\infty := \lim_{t \rightarrow +\infty} W_t$$

exists. Under our assumptions, W_t is a L^2 -bounded martingale, and thus, W_∞ is non-degenerate, that is,

$$\mathbb{P}_\mu(W_\infty > 0) > 0.$$

1.3 Main results

In this subsection, we state our main results. In the remainder of this paper, whenever we talk about an initial configuration $\mu \in \mathcal{M}_F(E)$, we always implicitly assume that it has compact support.

For $q > \max\{K, \lambda_0\}$ and $f \in L^p(E; m)$ with $p \geq 1$, define

$$U_q f(x) := \begin{cases} \int_0^{+\infty} e^{-qs} T_s f(x) ds, & \int_0^{+\infty} e^{-qs} T_s |f|(x) ds < +\infty, \\ 0, & \text{otherwise.} \end{cases}$$

Note that for $p \geq 1$, by (1.11),

$$\begin{aligned} \left(\int_E \left(\int_0^{+\infty} e^{-qs} T_s |f|(x) ds \right)^p m(dx) \right)^{1/p} &\leq \int_0^{+\infty} e^{-qs} \|T_s(|f|)\|_p ds \\ &\leq \int_0^{+\infty} e^{-qs} e^{Ks} ds \|f\|_p \\ &< +\infty, \end{aligned} \quad (1.14)$$

which implies

$$\int_0^{+\infty} e^{-qs} T_s |f|(x) ds \in L^p(E; m),$$

and thus,

$$\int_0^{+\infty} e^{-qs} T_s |f|(x) ds < +\infty, \quad m\text{-a.e.}$$

Consequently, $U_q f \in L^p(E; m)$. In Lemma 2.2 below, we will show that if $f \in L^2(E, m) \cap L^4(E, m)$, then $\langle U_q f, X_t \rangle$ is well defined.

Theorem 1.2 *Assume that Assumption 1.1 holds. If $g = U_q f$ for some $f \in L^2(E, m) \cap L^4(E, m)$ and $q > \max\{K, \lambda_0\}$, then for any $\mu \in \mathcal{M}_F(E)$, as $t \rightarrow +\infty$,*

$$e^{-\lambda_0 t} \langle g, X_t \rangle \rightarrow \langle g, \widehat{\phi}_0 \rangle_m W_\infty, \quad \mathbb{P}_\mu\text{-a.s.} \quad (1.15)$$

For any $f \geq 0$, define

$$T_t^{\phi_0} f(x) = \frac{e^{-\lambda_0 t}}{\phi_0(x)} \Pi_x \left[\exp \left(\int_0^t \alpha(\xi_s) ds \right) (f \phi_0)(\xi_t) \right]. \quad (1.16)$$

Let $C_0(E; \mathbb{R})$ denote the family of real-valued continuous functions f on E with the property that $\lim_{x \rightarrow \partial} f(x) = 0$.

We will also make the following assumption in this paper.

Assumption 1.3 The semigroup $\{T_t^{\phi_0}, t \geq 0\}$ has the following properties: for any $f \in C_0(E; \mathbb{R})$,

$$\lim_{t \rightarrow 0} \|T_t^{\phi_0} f - f\|_{\infty} = 0. \quad (1.17)$$

The following theorem is the main result of this paper.

Theorem 1.4 Under Assumptions 1.1 and 1.3, there exists $\Omega_0 \subset \Omega$ of probability one (that is, $\mathbb{P}_{\mu}(\Omega_0) = 1$ for every $\mu \in \mathcal{M}_F(E)$) such that, for every $\omega \in \Omega_0$ and for every bounded Borel function f on E satisfying

- (a) $|f| \leq c\phi_0$ for some $c > 0$,
- (b) the set of discontinuous points of f has zero m -measure,

we have

$$\lim_{t \rightarrow +\infty} e^{-\lambda_0 t} \langle f, X_t \rangle(\omega) = W_{\infty}(\omega) \int_E \widehat{\phi}_0(y) f(y) m(dy). \quad (1.18)$$

Assumption 1.3 will be used to extend the test functions from resolvent functions $g = U_q f$ with $f \in L^2(E, m) \cap L^4(E; m)$ to functions of the form $g = f\phi_0$ with $f \in C_0(E; \mathbb{R})$. We will give some examples in Section 4 to show that Assumptions 1.1 and 1.3 are satisfied by many interesting superprocesses including super Ornstein-Uhlenbeck processes (both inward and outward) and superprocesses with discontinuous spatial motions.

Remark 1.5 (1) Compared with [7], our spatial motion can be nonsymmetric and we do not assume that $\alpha(x) = \beta(x)a(x)$ is in the Kato class $K_{\infty}(\xi)$. The latter would require α be small at $+\infty$ in some sense (see [7] for the definition of $K_{\infty}(\xi)$). In [7], a compact embedding condition (see [7, 2.4]) is also assumed to ensure that the generator of the semigroup $\{T_t, t \geq 0\}$ has a spectral gap. In this paper, we assume instead Assumption 1.1, which implies that the generator of $\{T_t, t \geq 0\}$ has discrete spectrum.

(2) Compared with [22] where the spatial motion is a diffusion, our spatial motion may be discontinuous. The setup of [22] and the setup of the present paper are also different in the following ways. In [22], the semigroup of the spatial motion is assumed to be intrinsic ultracontractive. This condition is pretty strong and it excludes some interesting examples including the OU processes. In this paper, we assume Assumption 1.1 instead, which is weaker than the intrinsic ultracontractive property and is enough to ensure that, for resolvent functions g , the limit $\lim_{t \rightarrow +\infty} e^{-\lambda_0 t} \langle g, X_t \rangle$ exists almost surely. In [22], the branching mechanism is assumed to satisfy an $L \log L$ condition, while in this paper, we assume that the branching mechanism satisfies a second moment condition.

2 Preliminaries

2.1 Moment estimates

By [25, Lemma 2.2] with $k = 1$, for any $t_1 > 0$ and $a < -\operatorname{Re}(\lambda_1)$, there exists a constant $c = c(a, t_1) > 0$ such that for all $(t, x, y) \in (2t_1, +\infty) \times E \times E$,

$$|q(t, x, y) - e^{\lambda_0 t} \phi_0(x) \widehat{\phi}_0(y)| \leq c e^{-at} b_{t_1}^{1/2}(x) \widehat{b}_{t_1}^{1/2}(y). \quad (2.1)$$

Multiplying both sides by $e^{-\lambda_0 t}$, we get that for all $(t, x, y) \in (2t_1, +\infty) \times E \times E$,

$$|e^{-\lambda_0 t} q(t, x, y) - \phi_0(x) \widehat{\phi}_0(y)| \leq c e^{-(a+\lambda_0)t} b_{t_1}^{1/2}(x) \widehat{b}_{t_1}^{1/2}(y).$$

Note that $a < -\operatorname{Re}(\lambda_1)$ is equivalent to

$$a + \lambda_0 < \lambda_0 - \operatorname{Re}(\lambda_1).$$

Thus, for any $\tilde{a} \in (0, \lambda_0 - \operatorname{Re}(\lambda_1))$ and $t_1 > 0$, there exists $c_1 = c_1(\tilde{a}, t_1) > 0$ such that for all $(t, x, y) \in (2t_1, +\infty) \times E \times E$,

$$|e^{-\lambda_0 t} q(t, x, y) - \phi_0(x) \widehat{\phi}_0(y)| \leq c_1 e^{-\tilde{a}t} b_{t_1}^{1/2}(x) \widehat{b}_{t_1}^{1/2}(y). \quad (2.2)$$

Thus, for $f \in L^2(E; m)$, and for all $(t, x) \in (2t_1, +\infty) \times E$, we have

$$|e^{-\lambda_0 t} T_t f(x) - \phi_0(x) \langle f, \widehat{\phi}_0 \rangle_m| \leq c_1 \|\widehat{b}_{t_1}^{1/2}\|_2 \|f\|_2 e^{-\tilde{a}t} b_{t_1}^{1/2}(x),$$

which implies that there exists $c_2 = c_2(\tilde{a}, t_1) > 0$ such that for all $(t, x) \in (2t_1, +\infty) \times E$,

$$|e^{-\lambda_0 t} T_t f(x) - \phi_0(x) \langle f, \widehat{\phi}_0 \rangle_m| \leq c_2 \|f\|_2 e^{-\tilde{a}t} b_{t_1}^{1/2}(x). \quad (2.3)$$

Hence, by (1.13) and the fact that $\tilde{a} > 0$, we have

$$\begin{aligned} e^{-\lambda_0 t} |T_t f(x)| &\leq \phi_0(x) |\langle f, \widehat{\phi}_0 \rangle_m| + c_2 \|f\|_2 e^{-\tilde{a}t} b_{t_1}^{1/2}(x) \\ &\leq (e^{-\lambda_0 t_1} \|\widehat{\phi}_0\|_2 + c_2) \|f\|_2 b_{t_1}^{1/2}(x). \end{aligned}$$

Thus, there exists $c_3 = c_3(\tilde{a}, t_1) > 0$ such that for all $(t, x) \in (2t_1, +\infty) \times E$,

$$|T_t f(x)| \leq c_3 \|f\|_2 e^{\lambda_0 t} b_{t_1}^{1/2}(x). \quad (2.4)$$

We now recall the second moment formula for the superprocess $\{X_t, t \geq 0\}$ (see, for example, [24]): for $f \in L^2(E; m) \cap L^4(E; m)$ and $\mu \in \mathcal{M}_F(E)$, for any $t > 0$, we have

$$\operatorname{Var}_\mu \langle f, X_t \rangle = \langle \operatorname{Var}_\delta \langle f, X_t \rangle, \mu \rangle = \int_E \int_0^t T_s [A(T_{t-s} f)^2](x) ds \mu(dx), \quad (2.5)$$

where $\mathbb{V}\text{ar}_\mu$ stands for the variance under \mathbb{P}_μ and $A(x)$ is the function defined in (1.7). Moreover, for $f \in L^2(E; m) \cap L^4(E; m)$,

$$\mathbb{V}\text{ar}_{\delta_x} \langle f, X_t \rangle \leq e^{Kt} T_t(f^2)(x) \in L^2(E; m). \quad (2.6)$$

In the following lemma, we give a useful estimate on the second moment of X . If we choose the constant $\tilde{a} \in (0, \lambda_0 - \text{Re}(\lambda_1))$ small enough, we can get the next lemma by [25, Lemma 2.5]. Here, we give a direct proof.

Lemma 2.1 *Suppose that Assumption 1.1 holds. For any $\tilde{a} \in (0, (\lambda_0 - \text{Re}(\lambda_1)) \wedge (\lambda_0/2))$ and $f \in L^2(E; m) \cap L^4(E; m)$ with $\langle f, \hat{\phi}_0 \rangle_m = 0$, there exists $c_4 = c_4(t_0, \tilde{a}, f) > 0$ such that*

$$\sup_{t > 10t_0} e^{2(-\lambda_0 + \tilde{a})t} \mathbb{V}\text{ar}_{\delta_x} \langle f, X_t \rangle \leq c_4 b_{t_0}^{1/2}(x). \quad (2.7)$$

Proof In the following proof, we use $c = c(t_0, \tilde{a}, f)$ to denote a constant whose value may change from one appearance to another. Recall that

$$\mathbb{V}\text{ar}_{\delta_x} \langle f, X_t \rangle = \left(\int_0^{2t_0} + \int_{2t_0}^{t-2t_0} + \int_{t-2t_0}^t \right) T_s[A(T_{t-s}f)^2](x) ds.$$

In the following, we will deal with the above three parts separately.

(i) For $t > 10t_0$ and $s < 2t_0$, using (2.3) with $t_1 = 4t_0$ and noticing that $\langle f, \hat{\phi}_0 \rangle_m = 0$, we get

$$|T_{t-s}f(x)| \leq ce^{(\lambda_0 - \tilde{a})(t-s)} b_{4t_0}^{1/2}(x).$$

Thus,

$$\int_0^{2t_0} T_s[A(T_{t-s}f)^2](x) ds \leq ce^{2(\lambda_0 - \tilde{a})t} \int_0^{2t_0} T_s[b_{4t_0}](x) ds.$$

If we can prove that

$$\int_0^{2t_0} T_s[b_{4t_0}](x) ds \leq cb_{t_0}^{1/2}(x), \quad (2.8)$$

then we will get

$$\int_0^{2t_0} T_s[A(T_{t-s}f)^2](x) ds \leq ce^{2(\lambda_0 - \tilde{a})t} b_{t_0}^{1/2}(x). \quad (2.9)$$

Now, we prove (2.8). By Fubini's theorem and Hölder's inequality, we get

$$\begin{aligned} a_{t+s}(x) &= \int_E p(t+s, x, y) \int_E p(t, x, z) p(s, z, y) m(dz) m(dy) \\ &= \int_E p(t, x, z) \int_E p(t+s, x, y) p(s, z, y) m(dy) m(dz) \\ &\leq a_{t+s}^{1/2}(x) \int_E p(t, x, z) a_s^{1/2}(z) m(dz), \end{aligned}$$

which implies

$$a_{t+s}(x) \leq \left(\int_E p(t, x, z) a_s^{1/2}(z) m(dz) \right)^2 \leq \int_E p(t, x, z) a_s(z) m(dz). \quad (2.10)$$

By (2.10), we get

$$b_{4t_0}(x) \leq e^{8Kt_0} a_{4t_0}(x) \leq e^{10Kt_0} T_{2t_0}(a_{2t_0})(x).$$

Thus, by Assumption 1.1 (c') and (2.4), we have

$$\begin{aligned} \int_0^{2t_0} T_s(b_{4t_0})(x) ds &\leq e^{10Kt_0} \int_0^{2t_0} T_{s+2t_0}(a_{2t_0})(x) ds \\ &\leq c \int_0^{2t_0} e^{\lambda_0(s+2t_0)} ds b_{t_0}^{1/2}(x) \\ &\leq c b_{t_0}^{1/2}(x). \end{aligned} \quad (2.11)$$

Therefore, (2.8) holds.

(ii) For $t > 10t_0$ and $s \in (2t_0, t - 2t_0)$, by (2.3), (2.4), and Assumption 1.1 (c'),

$$T_s[A(T_{t-s}f)^2](x) \leq ce^{2(\lambda_0 - \tilde{a})(t-s)} T_s(b_{t_0})(x) \leq ce^{2(\lambda_0 - \tilde{a})(t-s)} e^{\lambda_0 s} b_{t_0}^{1/2}(x).$$

Thus, using the fact $\lambda_0 - 2\tilde{a} > 0$,

$$\begin{aligned} \int_{2t_0}^{t-2t_0} T_s[A(T_{t-s}f)^2](x) ds &\leq ce^{2(\lambda_0 - \tilde{a})t} \int_{2t_0}^{t-2t_0} e^{-(\lambda_0 - 2\tilde{a})s} ds b_{t_0}^{1/2}(x) \\ &\leq ce^{2(\lambda_0 - \tilde{a})t} b_{t_0}^{1/2}(x). \end{aligned} \quad (2.12)$$

(iii) For $t > 10t_0$ and $s > t - 2t_0$, since

$$|T_{t-s}f(x)|^2 \leq e^{K(t-s)} T_{t-s}(f^2)(x),$$

we have

$$\begin{aligned} T_s[A(T_{t-s}f)^2](x) &\leq K e^{K(t-s)} T_t(f^2)(x) \\ &\leq K e^{2t_0K} c_3 e^{\lambda_0 t} b_{t_0}^{1/2}(x) \\ &\leq K e^{2t_0K} c_3 e^{2(\lambda_0 - \tilde{a})t} b_{t_0}^{1/2}(x), \end{aligned}$$

where, in the last equality, we use the fact $\lambda_0 - 2\tilde{a} > 0$. Thus,

$$\int_{t-2t_0}^t T_s[A(T_{t-s}f)^2](x) ds \leq ce^{2(\lambda_0 - \tilde{a})t} b_{t_0}^{1/2}(x). \quad (2.13)$$

Combining (2.9), (2.12), and (2.13), we get (2.7). \square

2.2 Martingale measure for superprocesses

In this subsection, we recall the associated martingale measure for the superprocess X . For more details, see, for instance, [21, Chapter 7]. The martingale measure for superprocesses is a very useful tool in the proof of our main theorems.

For our superprocess X , there exists a worthy (\mathcal{G}_t) -martingale measure

$$\{M_t(B) = M(t, B); t \geq 0, B \in \mathcal{B}(E)\}$$

with covariation measure

$$\nu(ds, dx, dy) := ds \int_E A(z) \delta_z(dx) \delta_z(dy) X_s(dz)$$

such that for $t \geq 0$ and $f \in L^2(E; m) \cap L^4(E; m)$, we have, \mathbb{P}_μ -a.s.,

$$\langle f, X_t \rangle = \langle T_t f, \mu \rangle + \int_0^t \int_E T_{t-s} f(z) M(ds, dz). \quad (2.14)$$

For any $u > 0$ and $0 \leq t \leq u$, we define

$$M_t^{(u)} := \int_0^t \int_E T_{u-s} f(x) M(ds, dx).$$

Then, for any $\mu \in \mathcal{M}_F(E)$, $\{M_t^{(u)}, 0 \leq t \leq u\}$ is a cadlag square-integrable martingale under \mathbb{P}_μ with

$$\langle M^u \rangle_t = \int_0^t \langle A(T_{u-s} f)^2, X_s \rangle ds. \quad (2.15)$$

Here, cadlag means ‘right continuous having left limits’. Note that

$$\mathbb{P}_\mu(M_u^{(u)})^2 = \mathbb{P}_\mu \langle M^u \rangle_u = \mathbb{V}ar_\mu \langle f, X_u \rangle. \quad (2.16)$$

In the remainder of this paper, we will always assume that $q > \max\{K, \lambda_0\}$.

Lemma 2.2 *Assume that Assumption 1.1 holds. If $f \in L^2(E; m) \cap L^4(E; m)$, then for any $\mu \in \mathcal{M}_F(E)$,*

$$\mathbb{P}_\mu(\langle U_q f | f \rangle, X_t) < +\infty \text{ for } t \geq 0) = \mathbb{P}_\mu(\langle U_q f, X_t \rangle \text{ is finite for } t \geq 0) = 1.$$

Moreover, \mathbb{P}_μ -a.s., $\langle U_q f, X_t \rangle$ is cadlag on $[0, +\infty)$, and for all $t > 0$,

$$\langle U_q f, X_t \rangle = \langle T_t(U_q f), \mu \rangle + e^{qt} \int_t^{+\infty} e^{-qu} M_t^{(u)} du. \quad (2.17)$$

Proof When the spatial motion ξ is symmetric, this lemma has been established in [26, Lemmas 2.4, 2.5]. The proof for the non-symmetric case is almost the same. For reader’s convenience, we include a proof here. We can

check that the argument in the proof of [26, Lemma 2.4] works without the assumption that ξ is m -symmetric, so $\langle U_q f, X_t \rangle$ is right continuous on $[0, +\infty)$, \mathbb{P}_μ -a.s.

For $f \in L^2(E; m) \cap L^4(E; m)$,

$$U_q f \in L^2(E; m) \cap L^4(E; m).$$

By (2.14), for $t > 0$ and $\mu \in \mathcal{M}_F(E)$, we have, \mathbb{P}_μ -a.s.,

$$\begin{aligned} \langle U_q f, X_t \rangle &= \langle T_t(U_q f), \mu \rangle + \int_0^t \int_E T_{t-s}(U_q f)(z) M(ds, dz) \\ &= \langle T_t(U_q f), \mu \rangle + \int_0^t \int_E \int_0^{+\infty} e^{-qu} T_{u+t-s} f(z) du M(ds, dz) \\ &= \langle T_t(U_q f), \mu \rangle + e^{qt} \int_0^t \int_E \int_t^{+\infty} e^{-qu} T_{u-s} f(z) du M(ds, dz) \\ &= \langle T_t(U_q f), \mu \rangle + e^{qt} \int_t^{+\infty} e^{-qu} du \int_0^t \int_E T_{u-s} f(z) M(ds, dz) \\ &=: J_1^f(t) + e^{qt} J_2^f(t), \end{aligned} \quad (2.18)$$

where the fourth equality follows from the stochastic Fubini's theorem for martingale measures (see, for instance, [21, Theorem 7.24]). Thus, for $t > 0$ and $\mu \in \mathcal{M}_F(E)$,

$$\mathbb{P}_\mu(\langle U_q f, X_t \rangle = J_1^f(t) + e^{qt} J_2^f(t)) = 1. \quad (2.19)$$

Then, in light of (2.19), to prove (2.17), it suffices to prove that $J_1^f(t)$ and $J_2^f(t)$ are all cadlag in $(0, +\infty)$, \mathbb{P}_μ -a.s. For $J_1^f(t)$, by Fubini's theorem, for $t > 0$,

$$J_1^f(t) = e^{qt} \int_t^{+\infty} e^{-qs} \langle T_s f, \mu \rangle ds.$$

Thus, it is easy to see that $J_1^f(t)$ is continuous in $t \in (0, +\infty)$. Now, we consider $J_2^f(t)$. We claim that, for any $t_1 > 0$,

$$\mathbb{P}_\mu(J_2^f(t) \text{ is cadlag in } [t_1, +\infty)) = 1. \quad (2.20)$$

By the definition of J_2^f , for $t \geq t_1$,

$$J_2^f(t) = \int_{t_1}^{+\infty} e^{-qu} M_t^{(u)} \mathbf{1}_{t < u} du.$$

Since $t \mapsto M_t^{(u)} \mathbf{1}_{t < u}$ is right continuous, by the dominated convergence theorem, to prove (2.20), it suffices to show that

$$\mathbb{P}_\mu \left(\int_{t_1}^{+\infty} e^{-qu} \sup_{t \geq t_1} (|M_t^{(u)}| \mathbf{1}_{t < u}) du < +\infty \right) = 1. \quad (2.21)$$

By the L^p -maximum inequality and (2.16), we have

$$\begin{aligned}
& \mathbb{P}_\mu \left(\int_{t_1}^{+\infty} e^{-qu} \sup_{t \geq t_1} (|M_t^{(u)}| 1_{t < u}) du \right) \\
& \leq 2 \int_{t_1}^{+\infty} e^{-qu} \sqrt{\mathbb{P}_\mu |M_u^{(u)}|^2} du \\
& = 2 \int_{t_1}^{+\infty} e^{-qu} \sqrt{\int_E \text{Var}_{\delta_x} \langle f, X_u \rangle \mu(dx)} du. \tag{2.22}
\end{aligned}$$

By (2.6) and (2.4), for $u > t_1$, we have

$$\begin{aligned}
\int_E \text{Var}_{\delta_x} \langle f, X_u \rangle \mu(dx) & \leq e^{Ku} \int_E T_u(f^2)(x) \mu(dx) \\
& \leq ce^{Ku} e^{\lambda_0 u} \int_E b_{t_1/2}^{1/2}(x) \mu(dx),
\end{aligned}$$

where $c = c(t_1, \tilde{a}, f)$ is a positive constant and $b_t(x)$ is the function defined in (1.12). Since $x \mapsto b_{t_1/2}(x)$ is continuous and μ has compact support, we have

$$\int_E b_{t_1/2}^{1/2}(x) \mu(dx) < +\infty.$$

Thus, by (2.22), we have

$$\begin{aligned}
& \mathbb{P}_\mu \left(\int_{t_1}^{+\infty} e^{-qu} \sup_{t \geq t_1} (|M_t^{(u)}| 1_{t < u}) du \right) \\
& \leq 2\sqrt{c} \int_{t_1}^{+\infty} e^{-qu} e^{(K+\lambda_0)u/2} du \sqrt{\int_E b_{t_1/2}^{1/2}(x) \mu(dx)} \\
& < +\infty.
\end{aligned}$$

Now, (2.21) follows immediately. Since $t_1 > 0$ is arbitrary, we have

$$\mathbb{P}_\mu(J_2^f(t) \text{ is cadlag in } (0, +\infty)) = 1.$$

The proof is now complete. \square

3 Strong law of large numbers

In this section, we give the proofs of Theorems 1.2 and 1.4. We start with a lemma.

Lemma 3.1 *Suppose that Assumption 1.1 holds and $f \in L^2(E; m) \cap L^4(E; m)$ with $\langle f, \hat{\phi}_0 \rangle_m = 0$. Then for any $\mu \in \mathcal{M}_F(E)$ and $\tilde{a} \in (0, (\lambda_0 - \text{Re}(\lambda_1)) \wedge (\lambda_0/2))$,*

$$\sup_{n > 10t_0} e^{(-\lambda_0 + \tilde{a})n} \mathbb{P}_\mu \left(\sup_{n \leq t \leq n+1} |\langle U_q f, X_t \rangle| \right) < +\infty. \tag{3.1}$$

Proof In this proof, we always assume that $n > 10t_0$ and c is a positive constant whose value does not depend on n and may change from one appearance to another. Define

$$J_1^f(t) := \langle T_t U_q f, \mu \rangle, \quad J_2^f(t) := \int_t^{+\infty} e^{-qu} M_t^{(u)} du.$$

By (2.17), for any $t > 0$,

$$\mathbb{P}_\mu \left(\sup_{n \leq t \leq n+1} |\langle U_q f, X_t \rangle| \leq \sup_{n \leq t \leq n+1} |J_1^f(t)| + e^{q(n+1)} \mathbb{P}_\mu \left(\sup_{n \leq t \leq n+1} |J_2^f(t)| \right) \right).$$

First, we consider $J_1^f(t)$. Since $\langle U_q f, \hat{\phi}_0 \rangle_m = 0$, by (2.3), we have

$$|T_t U_q f|(x) \leq ce^{(\lambda_0 - \tilde{a})t} b_{t_0}^{1/2}(x).$$

Thus, for $n > 10t_0$,

$$\begin{aligned} \sup_{n \leq t \leq n+1} |J_1^f(t)| &\leq \sup_{n \leq t \leq n+1} \langle |T_t U_q f|, \mu \rangle \\ &\leq c \sup_{n \leq t \leq n+1} e^{(\lambda_0 - \tilde{a})t} \langle b_{t_0}^{1/2}, \mu \rangle \\ &\leq ce^{(\lambda_0 - \tilde{a})n}. \end{aligned} \tag{3.2}$$

Next, we deal with $J_2^f(t)$. For $t \in [n, n+1]$,

$$J_2^f(t) = \int_t^{+\infty} e^{-qu} M_t^{(u)} du = \int_n^{+\infty} e^{-qu} M_t^{(u)} 1_{t < u} du.$$

Thus, for $n > 10t_0$,

$$\begin{aligned} \mathbb{P}_\mu \left(\sup_{n \leq t \leq n+1} |J_2^f(t)| \right) &\leq \int_n^{+\infty} e^{-qu} \mathbb{P}_\mu \left(\sup_{n \leq t \leq n+1} (|M_t^{(u)}| 1_{t < u}) \right) du \\ &\leq 2 \int_n^{+\infty} e^{-qu} \sqrt{\mathbb{P}_\mu(M_u^{(u)})^2} du \\ &\leq 2\sqrt{c \langle b_{t_0}^{1/2}, \mu \rangle} \int_n^{+\infty} e^{-qu} e^{(\lambda_0 - \tilde{a})u} du \\ &\leq c(q - \lambda_0 + \tilde{a})^{-1} e^{-(q - \lambda_0 + \tilde{a})n}, \end{aligned}$$

where the third equality follows from (2.15), (2.16), and (2.7). It follows that for $n > 10t_0$,

$$e^{q(n+1)} \mathbb{P}_\mu \left(\sup_{n \leq t \leq n+1} J_2^f(t) \right) \leq ce^{(\lambda_0 - \tilde{a})n}. \tag{3.3}$$

Combining (3.2) and (3.3), this yields (3.1). The proof is now complete. \square

Proof of Theorem 1.2 Put

$$\tilde{f} = f - \langle f, \widehat{\phi}_0 \rangle_m \phi_0.$$

Note that

$$U_q \phi_0(x) = \int_0^{+\infty} e^{-qt} T_t \phi_0(x) dt = \int_0^{+\infty} e^{-qt} e^{\lambda_0 t} dt \phi_0(x) = (q - \lambda_0)^{-1} \phi_0(x)$$

and

$$\begin{aligned} \langle U_q f, \widehat{\phi}_0 \rangle_m &= \int_0^{+\infty} e^{-qt} \langle T_t f, \widehat{\phi}_0 \rangle_m dt \\ &= \int_0^{+\infty} e^{-qt} e^{\lambda_0 t} dt \langle f, \widehat{\phi}_0 \rangle_m \\ &= (q - \lambda_0)^{-1} \langle f, \widehat{\phi}_0 \rangle_m. \end{aligned} \quad (3.4)$$

Thus,

$$U_q f(x) = \langle f, \widehat{\phi}_0 \rangle_m U_q \phi_0(x) + U_q(\tilde{f})(x) = \langle U_q f, \widehat{\phi}_0 \rangle_m \phi_0(x) + U_q(\tilde{f})(x).$$

Hence, to prove (1.15), we only need to show that

$$e^{-\lambda_0 t} \langle U_q(\tilde{f}), X_t \rangle \rightarrow 0, \quad \mathbb{P}_\mu\text{-a.s.} \quad (3.5)$$

Let

$$M_n := \sup_{n \leq t \leq n+1} e^{-\lambda_0 t} |\langle U_q(\tilde{f}), X_t \rangle|.$$

By (3.1), there is a constant $c > 0$ so that

$$\mathbb{P}_\mu M_n \leq c e^{-\tilde{a}n}, \quad \forall n > 10t_0.$$

We conclude by the Borel-Cantelli lemma that $M_n \rightarrow 0$, as $n \rightarrow +\infty$, $\mathbb{P}_\mu\text{-a.s.}$, from which, (3.5) follows immediately. The proof is now complete. \square

For any $f \geq 0$ and $q > \max\{K, \lambda_0\}$, define

$$U_q^{\phi_0} f(x) = \int_0^{+\infty} e^{-qt} T_t^{\phi_0} f(x) dt, \quad x \in E,$$

where $T_t^{\phi_0}$ is defined in (1.16). It is easy to see that

$$\phi_0(x) U_q^{\phi_0} f(x) = U_{q+\lambda_0}(\phi_0 f).$$

Proposition 3.2 *Suppose that Assumptions 1.1 and 1.3 hold. For any $0 \leq f \in C_0(E; \mathbb{R})$ and $\mu \in \mathcal{M}_F(E)$,*

$$\lim_{t \rightarrow +\infty} e^{-\lambda_0 t} \langle \phi_0 f, X_t \rangle = \langle f \phi_0, \widehat{\phi}_0 \rangle_m W_\infty, \quad \mathbb{P}_\mu\text{-a.s.} \quad (3.6)$$

Proof By Theorem 1.2,

$$\begin{aligned} \lim_{t \rightarrow +\infty} e^{-\lambda_0 t} \langle \phi_0 U_q^{\phi_0} f, X_t \rangle &= \lim_{t \rightarrow +\infty} e^{-\lambda_0 t} \langle U_{q+\lambda_0}(\phi_0 f), X_t \rangle \\ &= \langle U_{q+\lambda_0}(\phi_0 f), \widehat{\phi}_0 \rangle_m W_\infty, \quad \mathbb{P}_\mu\text{-a.s.} \end{aligned}$$

According to (3.4),

$$\langle U_{q+\lambda_0}(\phi_0 f), \widehat{\phi}_0 \rangle_m = \frac{1}{q} \langle \phi_0 f, \widehat{\phi}_0 \rangle_m.$$

Therefore, for any $q > \max\{K, \lambda_0\}$,

$$\lim_{t \rightarrow +\infty} e^{-\lambda_0 t} \langle \phi_0 q U_q^{\phi_0} f, X_t \rangle = \langle f \phi_0, \widehat{\phi}_0 \rangle_m W_\infty, \quad \mathbb{P}_\mu\text{-a.s.} \quad (3.7)$$

Choose a sequence $q_k > \max\{K, \lambda_0\}$ so that $\lim_{k \rightarrow +\infty} q_k = +\infty$. Put

$$\begin{aligned} \Omega^* &:= \bigcap_{k \geq 1} \left\{ \lim_{t \rightarrow +\infty} e^{-\lambda_0 t} \langle \phi_0 q_k U_{q_k}^{\phi_0} f, X_t(\omega) \rangle = \langle f \phi_0, \widehat{\phi}_0 \rangle_m W_\infty(\omega) \right\} \\ &\quad \bigcap \left\{ \lim_{t \rightarrow +\infty} W_t(\omega) = W_\infty(\omega) \right\}. \end{aligned}$$

Then $\mathbb{P}_\mu(\Omega^*) = 1$. Note that, for any $\omega \in \Omega^*$,

$$\begin{aligned} &|e^{-\lambda_0 t} \langle \phi_0 q_k U_{q_k}^{\phi_0} f, X_t(\omega) \rangle - e^{-\lambda_0 t} \langle \phi_0 f, X_t(\omega) \rangle| \\ &\leq e^{-\lambda_0 t} \langle \phi_0 |q_k U_{q_k}^{\phi_0} f - f|, X_t(\omega) \rangle \\ &\leq \|q_k U_{q_k}^{\phi_0} f - f\|_\infty e^{-\lambda_0 t} \langle \phi_0, X_t(\omega) \rangle, \end{aligned}$$

where $\|\cdot\|_\infty$ is the L^∞ norm. Letting $t \rightarrow +\infty$, we obtain that

$$\limsup_{t \rightarrow +\infty} |e^{-\lambda_0 t} \langle \phi_0 q_k U_{q_k}^{\phi_0} f, X_t(\omega) \rangle - e^{-\lambda_0 t} \langle \phi_0 f, X_t(\omega) \rangle| \leq \|q_k U_{q_k}^{\phi_0} f - f\|_\infty W_\infty(\omega). \quad (3.8)$$

By Assumption 1.3,

$$\lim_{k \rightarrow +\infty} \|q_k U_{q_k}^{\phi_0} f - f\|_\infty = 0.$$

Thus, (3.8) implies that, for $\omega \in \Omega^*$,

$$\lim_{k \rightarrow +\infty} \limsup_{t \rightarrow +\infty} |e^{-\lambda_0 t} \langle \phi_0 q_k U_{q_k}^{\phi_0} f, X_t(\omega) \rangle - e^{-\lambda_0 t} \langle \phi_0 f, X_t(\omega) \rangle| = 0. \quad (3.9)$$

Now, combining (3.7) and (3.9), we get (3.6). \square

Proof of Theorem 1.4 Note that E_∂ is a compact separable metric space. According to [30, Exercise 9.1.16 (iii)], $C_b(E_\partial; \mathbb{R})$, the space of bounded continuous \mathbb{R} -valued functions f on E , is separable. Therefore, $C_0(E; \mathbb{R})$ is also a separable space. Let $\{f_n, n \geq 1\}$ be a countable dense subset of $C_0(E; \mathbb{R})$. Define Ω_0 to be the intersection of the following two events:

$$\left\{ \omega \in \Omega: \lim_{t \rightarrow +\infty} W_t(\omega) = W_\infty(\omega) \right\},$$

$$\bigcap_{k \geq 1} \left\{ \omega \in \Omega : \lim_{t \rightarrow +\infty} e^{-\lambda_0 t} \langle f_k \phi_0, X_t \rangle(\omega) = W_\infty(\omega) \int_E f_k(y) \phi_0(y) \widehat{\phi}_0(y) m(dy) \right\}.$$

By Proposition 3.2, $\mathbb{P}_\mu(\Omega_0) = 1$ for any $\mu \in \mathcal{M}_F(E)$.

We first consider (1.18) on $\{W_\infty > 0\}$. For each $\omega \in \Omega_0 \cap \{W_\infty > 0\}$ and $t \geq 0$, we define two probability measures ν_t and ν on E , respectively, by

$$\nu_t(F)(\omega) = \frac{e^{-\lambda_0 t} \langle 1_F \phi_0, X_t \rangle(\omega)}{W_t(\omega)},$$

$$\nu(F) = \int_F \phi_0(y) \widehat{\phi}_0(y) m(dy), \quad F \in \mathcal{B}(E).$$

Note that the measure ν_t is well defined for every $t \geq 0$, and ν_t and ν are probability measures. By the definition of Ω_0 , we know that ν_t converges weakly to ν as $t \rightarrow +\infty$. Since ϕ_0 is strictly positive and continuous on E , if f is a function on E such that $|f| \leq c\phi_0$ for some $c > 0$ and that the discontinuity set of f has zero m -measure (equivalently, zero ν -measure), then $g := f/\phi_0$ is a bounded function with the same set of discontinuity. We thus have

$$\lim_{t \rightarrow +\infty} \int_E g(x) \nu_t(dx) = \int_E g(x) \nu(dx),$$

which is equivalent to

$$\lim_{t \rightarrow +\infty} e^{-\lambda_0 t} \langle f, X_t \rangle(\omega) = W_\infty(\omega) \int_E \widehat{\phi}_0(y) f(y) m(dy), \quad \omega \in \Omega_0 \cap \{W_\infty(\phi) > 0\}.$$

If $|f| \leq c\phi_0$ for some positive constant $c > 0$, (1.18) holds automatically on $\{W_\infty = 0\}$. This completes the proof of the theorem. \square

4 Examples

In this section, we give some examples. The main purpose is to illustrate the diverse situations where the main result of this paper can be applied. We will not try to give the most general examples possible.

Example 4.1 (Super inward Ornstein-Uhlenbeck processes) Let $d \geq 1$, $E = \mathbb{R}^d$. Suppose that the spatial motion $\xi = \{\xi_t, \Pi_x\}$ is an Ornstein-Uhlenbeck (OU) process on \mathbb{R}^d with infinitesimal generator

$$\mathcal{L} = \frac{1}{2} \sigma^2 \Delta - cx \cdot \nabla \quad \text{on } \mathbb{R}^d,$$

where $\sigma, c > 0$. Without loss of generality, we assume $\sigma = 1$. Let

$$\varphi(x) := \left(\frac{c}{\pi}\right)^{d/2} e^{-c|x|^2}, \quad m(dx) = \varphi(x) dx.$$

Then ξ is symmetric with respect to the probability measure $m(dx)$. Suppose that the branching rate function $\beta(x) = \beta$ is a positive constant, and the branching mechanism ψ is given by

$$\psi(x, \lambda) = -\lambda + b(x)\lambda^2 + \int_{(0, +\infty)} (e^{-\lambda y} - 1 + \lambda y)n(x, dy), \quad x \in \mathbb{R}^d, \lambda > 0, \quad (4.1)$$

where $b \in \mathcal{B}_b^+(\mathbb{R}^d)$ and n is a kernel from \mathbb{R}^d to $(0, +\infty)$ satisfying

$$\sup_{x \in \mathbb{R}^d} \int_0^{+\infty} y^2 n(x, dy) < +\infty.$$

Then for the corresponding superprocess,

$$T_t f(x) = e^{\beta t} \Pi_x [f(\xi_t)] = e^{\beta t} P_t f(x).$$

It is easy to see that $\lambda_0 = \beta$, $\phi_0 = \widehat{\phi}_0 = 1$ and then $T_t^{\phi_0} = P_t$.

It is well known that, for any $x \in \mathbb{R}^d$, under Π_x , ξ_t is of Gaussian distribution with mean $x e^{-ct}$ and variance σ_t^2 , where

$$\sigma_t^2 := \frac{1 - e^{-2ct}}{2c}.$$

The transition density of ξ_t with respect to the probability measure $m(dx)$ on \mathbb{R}^d is given by

$$p(t, x, y) := \left(\frac{1}{2c\sigma_t^2} \right)^{d/2} \exp \left(c|y|^2 - \frac{|y - x e^{-ct}|^2}{2\sigma_t^2} \right).$$

Note that

$$p(t, x, x) = \frac{(2\pi\sigma_t^2)^{-d/2} \exp(-c \frac{1-e^{-ct}}{1+e^{-ct}} |x|^2)}{\varphi(x)}.$$

Thus, we have

$$a(t) = p(2t, x, x) \in L^1(\mathbb{R}^d; m), \quad \forall t > 0,$$

and there is some $t_0 > 0$ so that $a(t) \in L^2(\mathbb{R}^d; m)$ for $t \geq t_0$. Hence, Assumption 1.1 holds for ξ .

For any $f \in C_0(\mathbb{R}^d; \mathbb{R})$, we have

$$P_t f(x) = \int_{\mathbb{R}^d} p(t, x, y) f(y) m(dy) = \int_{\mathbb{R}^d} (2\pi)^{-d/2} \exp \left(-\frac{|y|^2}{2} \right) f(\sigma_t y + x e^{-ct}) dy.$$

Using the dominated convergence theorem, one can easily check that $P_t f \in C_0(\mathbb{R}^d; \mathbb{R})$. Suppose that f is a continuous function with compact support. Let

$M_0 > 0$ so that $f(x) = 0$ for $|x| \geq M_0$. For any $M > 0$,

$$\begin{aligned} |P_t f(x) - f(x)| &= \left| \int_{\mathbb{R}^d} (2\pi)^{-d/2} \exp\left(-\frac{|y|^2}{2}\right) [f(\sigma_t y + x e^{-ct}) - f(x)] dy \right| \\ &\leq \int_{\mathbb{R}^d} (2\pi)^{-d/2} \exp\left(-\frac{|y|^2}{2}\right) |f(\sigma_t y + x e^{-ct}) - f(x)| dy \\ &\leq \int_{|y| \leq M} (2\pi)^{-d/2} \exp\left(-\frac{|y|^2}{2}\right) |f(\sigma_t y + x e^{-ct}) - f(x)| dy \\ &\quad + 2\|f\|_\infty \int_{|y| \geq M} (2\pi)^{-d/2} \exp\left(-\frac{|y|^2}{2}\right) dy \\ &=: I + II. \end{aligned}$$

For any $\varepsilon > 0$, we choose $M > 0$ such that $II \leq \varepsilon/2$. For part I , we claim that, for any $\varepsilon > 0$, there exists δ , for $t \leq \delta$,

$$\sup_{|y| \leq M} \sup_{x \in \mathbb{R}^d} |f(\sigma_t y + x e^{-ct}) - f(x)| \leq \frac{\varepsilon}{2}.$$

Therefore, $I < \varepsilon/2$, and then $\|P_t f - f\|_\infty \rightarrow 0$ as $t \rightarrow 0$.

Now, we prove the claim. Note that

$$|f(\sigma_t y + x e^{-ct}) - f(x)| \leq |f(\sigma_t y + x e^{-ct}) - f(x e^{-ct})| + |f(x e^{-ct}) - f(x)|.$$

Since f is uniformly continuous on \mathbb{R}^d , there is a constant $\delta_0 > 0$ such that $|f(y) - f(x)| \leq \varepsilon/4$ for any x, y satisfying $|x - y| \leq \delta_0$. Since $\sigma_t \rightarrow 0$ as $t \rightarrow 0$, there exists $\delta_1 > 0$ such that, for $t < \delta_1$, $|\sigma_t| \leq \delta_0/M$, and then

$$\sup_{|y| \leq M} \sup_{x \in \mathbb{R}^d} |f(\sigma_t y + x e^{-ct}) - f(x e^{-ct})| \leq \frac{\varepsilon}{4}.$$

Choose δ_2 , such that for $t \leq \delta_2$, $e^{ct} - 1 \leq \delta_0/M_0$. Then, for $t \leq \delta_2$,

$$|f(x e^{-ct}) - f(x)| \leq |f(x e^{-ct}) - f(x)| 1_{|x| \leq M_0 e^{ct}} \leq \frac{\varepsilon}{4},$$

where, in the second inequality, we use the fact that

$$|x e^{-ct} - x| = |x|(1 - e^{-ct}) \leq M_0(e^{ct} - 1) \leq \delta_0.$$

Then, choosing $\delta = \delta_1 \wedge \delta_2$, we prove the claim.

For general $f \in C_0(\mathbb{R}^d; \mathbb{R})$, there exist continuous functions f_n with compact support such that $\|f_n - f\|_\infty \rightarrow 0$ as $n \rightarrow +\infty$. Then

$$\begin{aligned} \|P_t f - f\|_\infty &\leq \|P_t f - P_t f_n\|_\infty + \|P_t f_n - f_n\|_\infty + \|f_n - f\|_\infty \\ &\leq \|P_t f_n - f_n\|_\infty + 2\|f_n - f\|_\infty. \end{aligned}$$

Letting $t \rightarrow 0$ and then $n \rightarrow +\infty$, we get that $\|P_t f - f\|_\infty \rightarrow 0$ as $t \rightarrow 0$. Since $T_t^{\phi_0} = P_t$, Assumption 1.3 is satisfied. Therefore, for the superprocess in this example, all our assumptions are satisfied.

Example 4.1 covers [10, Examples 4.1, 4.6]. For variable $\alpha(x) = \beta(x)a(x)$, see Example 4.9.

Example 4.2 (Super outward Ornstein-Uhlenbeck processes) Let $d \geq 1$, $E = \mathbb{R}^d$. Suppose that the spatial motion $\xi = \{\xi_t, \Pi_x\}$ is an OU process on \mathbb{R}^d with infinitesimal generator

$$\mathcal{L} = \frac{1}{2} \sigma^2 \Delta + cx \cdot \nabla \quad \text{on } \mathbb{R}^d,$$

where $\sigma, c > 0$. Without loss of generality, we assume $\sigma = 1$. Under Π_x , ξ_t is of Gaussian distribution with mean xe^{ct} and variance $(e^{2ct} - 1)/(2c)$.

Let

$$\tilde{\varphi}(x) := \left(\frac{c}{\pi}\right)^{-d/2} e^{c|x|^2}, \quad m(dx) = \tilde{\varphi}(x)dx.$$

Then ξ is symmetric with respect to the σ -finite measure $m(dx)$. As in Example 4.1, we suppose that the branching rate function $\beta(x) = \beta$ is a positive constant, and the branching mechanism ψ is given by (4.1). Then for the corresponding superprocess,

$$T_t f(x) = e^{\beta t} \Pi_x [f(\xi_t)] = e^{\beta t} P_t f(x).$$

The generator of $\{T_t : t \geq 0\}$ is $\mathcal{L} + \beta$.

The transition density of ξ with respect to the measure m is

$$p(t, x, y) = \left(\frac{1}{e^{2ct} - 1}\right)^{d/2} \exp\left(-\frac{c}{1 - e^{-2ct}} (|y|^2 + |x|^2 - 2x \cdot ye^{-ct})\right).$$

Thus,

$$a_t(x) = p(2t, x, x) = \left(\frac{1}{e^{4ct} - 1}\right)^{d/2} \exp\left(-\frac{2c|x|^2}{1 + e^{-2ct}}\right).$$

It is obvious that $a_t \in L^1(\mathbb{R}^d; m) \cap L^2(\mathbb{R}^d; m)$. Thus, Assumption 1.1 is satisfied. Suppose $\beta(x) = \beta \in (cd, +\infty)$.

The operator $\mathcal{L} + cd$ is the formal adjoint of the inward OU process with infinitesimal generator $\frac{1}{2} \sigma^2 \Delta - cx \cdot \nabla$ on \mathbb{R}^d . Since $\varphi(x)$ defined in Example 4.1 is the invariant density of $\frac{1}{2} \sigma^2 \Delta - cx \cdot \nabla$ on \mathbb{R}^d ,

$$(\mathcal{L} + cd)\varphi = 0.$$

Thus, we have

$$(\mathcal{L} + \beta)\varphi = (\beta - cd)\varphi.$$

Since $\varphi \in L^2(\mathbb{R}^d, m)$ and φ is strictly positive everywhere, we know that

$$\phi_0 = \hat{\phi}_0 = \varphi, \quad \lambda_0 = \beta - cd.$$

Thus,

$$T_t^{\phi_0} f(x) = \frac{e^{cdt} P_t(f\varphi)(x)}{\varphi(x)} = \tilde{P}_t f(x),$$

where \tilde{P}_t is the semigroup of the inward OU-process with infinitesimal generator

$$\frac{1}{2} \Delta - cx \cdot \nabla \quad \text{on } \mathbb{R}^d.$$

From the discussion in Example 4.1, we see that Assumption 1.3 is satisfied. Thus, when $\beta(x) = \beta \in (cd, +\infty)$, the superprocess of this example satisfies all our assumptions.

Example 4.2 covers [10, Example 4.2].

Example 4.3 Suppose that $\eta = \{\eta_t, \Pi_x\}$ is an m -symmetric Hunt process on E and that η has a transition density $\tilde{p}(t, x, y)$ with respect to m . Suppose also that \tilde{p} is strictly positive, continuous, and satisfies Assumption 1.1. Let $\{\tilde{P}_t, t \geq 0\}$ be the transition semigroup of η on $L^2(E; m)$. Since, for each $t > 0$, \tilde{P}_t is compact, the infinitesimal generator $\tilde{\mathcal{L}}$ of $\{\tilde{P}_t, t \geq 0\}$ has discrete spectrum:

$$0 \geq \tilde{\lambda}_0 > \tilde{\lambda}_1 \geq \dots.$$

Denote the corresponding normalized eigenfunctions by $\{\tilde{\phi}_k; k \geq 0\}$, with

$$\|\tilde{\phi}_k\|_{L^2(E; m)} = 1, \quad \forall k \geq 0.$$

We can choose $\tilde{\phi}_0$ so that it is strictly positive and continuous. By the spectral representation, we can express $\tilde{p}(t, x, y)$ by $\sum_{k=0}^{+\infty} e^{-\tilde{\lambda}_k t} \tilde{\phi}_k(x) \tilde{\phi}_k(y)$. It follows that $\tilde{p}(t, x, x)$ is decreasing in $t > 0$; see [9, Section 2]. Define

$$\tilde{P}_t^{\tilde{\phi}_0} f := e^{-\tilde{\lambda}_0 t} \frac{\tilde{P}_t(f\tilde{\phi}_0)(x)}{\tilde{\phi}_0(x)}.$$

Assume that $\tilde{P}_t^{\tilde{\phi}_0}$ satisfies Assumption 1.3.

Let S_t be a subordinator, independent of η , with drift $b > 0$. Then $S_t \geq bt$. Let ϕ be the Laplace exponent of S , that is,

$$\mathbb{E}(e^{-\theta S_t}) = e^{-t\phi(\theta)}, \quad \theta > 0.$$

Suppose that $\alpha(x) = \alpha$ is a constant function and satisfies $\alpha > \phi(-\tilde{\lambda}_0)$. We put $\xi_t := \eta_{S_t}$. Let P_t be the semigroup of ξ , and let $p(t, x, y)$ be the transition density of ξ with respect to m . Then

$$p(t, x, y) = \mathbb{E}\tilde{p}(S_t, x, y).$$

Since $t \rightarrow \tilde{p}(t, x, x)$ is a decreasing function, we have

$$p(2t, x, x) = \mathbb{E}\tilde{p}(S_{2t}, x, x) \leq \tilde{p}(2bt, x, x),$$

which implies that η satisfies Assumption 1.1. Note that $T_t = e^{\alpha t} P_t$, and

$$P_t \tilde{\phi}_0(x) = \mathbb{E}(\tilde{P}_{S_t} \tilde{\phi}_0(x)) = \mathbb{E} e^{\tilde{\lambda}_0 S_t} \tilde{\phi}_0(x) = e^{-t\phi(-\tilde{\lambda}_0)} \tilde{\phi}_0(x).$$

Thus,

$$\lambda_0 = \alpha - \phi(-\tilde{\lambda}_0) > 0, \quad \phi_0 = \tilde{\phi}_0.$$

Then

$$T_t^{\phi_0} f(x) = e^{t\phi(-\tilde{\lambda}_0)} \frac{P_t(f\phi_0)(x)}{\phi_0(x)} = e^{t\phi(-\tilde{\lambda}_0)} \mathbb{E} \left[\frac{\tilde{P}_{S_t}(f\phi_0)(x)}{\phi_0(x)} \right].$$

Thus, we have

$$\begin{aligned} |T_t^{\phi_0} f(x) - f(x)| &\leq e^{t\phi(-\tilde{\lambda}_0)} \mathbb{E} \left| \frac{\tilde{P}_{S_t}(f\phi_0)(x)}{\phi_0(x)} - e^{\tilde{\lambda}_0 S_t} f(x) \right| \\ &\leq e^{bt\tilde{\lambda}_0} e^{t\phi(-\tilde{\lambda}_0)} \mathbb{E} [\|\tilde{P}_{S_t}^{\phi_0} f - f\|_\infty]. \end{aligned}$$

Since

$$\|\tilde{P}_{S_t}^{\phi_0} f - f\|_\infty \rightarrow 0, \quad t \rightarrow 0,$$

and

$$\|\tilde{P}_{S_t}^{\phi_0} f - f\|_\infty \leq 2\|f\|_\infty,$$

using the dominated convergence theorem, we get

$$\lim_{t \rightarrow 0} \|T_t^{\phi_0} f - f\|_\infty = 0.$$

Thus, the superprocess of this example satisfies all our assumptions.

In particular, this example is applicable when η is the outward Ornstein-Uhlenbeck process or inward Ornstein-Uhlenbeck process dealt with in Examples 4.1 and 4.2.

The next two examples give the cases when α is not a constant function.

Example 4.4 (Pure jump subordinate Brownian motion) Suppose that $S = \{S_t, t \geq 0\}$ is a drift-free subordinator. The Laplace exponent ϕ of S can be written in the form

$$\phi(\lambda) = \int_0^{+\infty} (1 - e^{-\lambda t}) u(dt), \quad (4.2)$$

where u is a measure on $(0, +\infty)$ satisfying

$$\int_0^{+\infty} (1 \wedge t) u(dt) < +\infty.$$

The measure u is the Lévy measure of the subordinator (or of ϕ). In this example, we will assume that ϕ is a complete Bernstein function, that is, the measure u has a completely monotone density, which we also denote by u .

Let $W = \{W_t, t \geq 0\}$ be a Brownian motion in \mathbb{R}^d independent of the subordinator S . The subordinate Brownian motion $Y = \{Y_t, t \geq 0\}$ is defined by $Y_t := W_{S_t}$, which is a rotationally symmetric Lévy process with Lévy exponent $\phi(|\xi|^2)$ (see [28, Theorem 30.1]). It is known that the Lévy measure of the process Y has a density given by $x \rightarrow j(|x|)$, where

$$j(r) := \int_0^{+\infty} (4\pi t)^{-d/2} e^{-r^2/(4t)} u(t) dt, \quad r > 0. \quad (4.3)$$

Note that the function $r \mapsto j(r)$ is continuous and decreasing on $(0, +\infty)$.

Suppose that ϕ satisfies the following growth condition at infinity.

(A) There exist constants $\delta_1, \delta_2 \in (0, 1)$, $a_1 \in (0, 1)$, $a_2 \in (1, +\infty)$, and $R_0 > 0$ such that

$$a_1 \lambda^{\delta_1} \phi(r) \leq \phi(\lambda r) \leq a_2 \lambda^{\delta_2} \phi(r), \quad \lambda \geq 1, r \geq R_0.$$

See [3] for examples of a large class of symmetric Lévy processes satisfying condition (A).

Suppose that D is a bounded $C^{1,1}$ open set with characteristics (R_0, Λ) , and let ξ be the subprocess of Y killed upon leaving D . It is known that ξ is a Feller process with strong Feller property in D . Moreover, by [3, Corollary 1.6], ξ has a jointly continuous transition density function $p_D(t, x, y)$ with respect to the Lebesgue measure on D so that for every $T > 0$, there exist

$$c_1 = c_1(R_0, \Lambda, T, d, \phi) \geq 1, \quad c_2 = c_2(R_0, \Lambda, T, d, \phi) > 0,$$

such that for $0 < t \leq T$, $x, y \in D$,

$$\begin{aligned} & c_1^{-1} \left(1 \wedge \frac{\Phi(\delta_D(x))}{t}\right)^{1/2} \left(1 \wedge \frac{\Phi(\delta_D(y))}{t}\right)^{1/2} \left(\Phi^{-1}(t)^{-d} \wedge t j(|x - y|)\right) \\ & \leq p_D(t, x, y) \\ & \leq c_2 \left(1 \wedge \frac{\Phi(\delta_D(x))}{t}\right)^{1/2} \left(1 \wedge \frac{\Phi(\delta_D(y))}{t}\right)^{1/2} \left(\Phi^{-1}(t)^{-d} \wedge t j\left(\frac{c_2|x - y|}{4}\right)\right). \end{aligned} \quad (4.4)$$

Here, $\Phi(r) := 1/\phi(r^{-2})$, j is the function defined in (4.3), and $\delta_D(x)$ is the Euclidean distance between x and ∂D . Since $p_D(t, x, y)$ is symmetric,

$$a_t(x) = p_D(2t, x, x) \leq c_2 \Phi^{-1}(2t)^{-d}.$$

Thus, Assumption 1.1 is satisfied.

Suppose that the branching rate function β and the branching mechanism satisfy the assumptions of Section 1.2, and that the corresponding superprocess X is supercritical. The corresponding semigroup $\{T_t: t \geq 0\}$ has a continuous density $q(t, x, y)$ satisfying the same two-sided estimates (4.4) with possibly different $c_1 \geq 1$ and c_2 . Since

$$\phi_0(x) = e^{\lambda_0 t} T_t \phi_0(x),$$

by (4.4), there exists $c_3 > 1$ such that

$$c_3^{-1}\Phi^{1/2}(\delta_D(x)) < \phi_0(x) < c_3\Phi^{1/2}(\delta_D(x)).$$

We now show that Assumption 1.3 holds. Suppose $f \in C_0(D)$. For any given $\varepsilon > 0$, there exists $\delta > 0$ so that $|f(x) - f(y)| < \varepsilon$ whenever $|x - y| < \delta$. Hence, by the display above and (4.4), for small $t > 0$,

$$\begin{aligned} & \sup_{x \in D} |T_t^{\phi_0} f(x) - f(x)| \\ &= \sup_{x \in D} \frac{e^{-\lambda_0 t} |\Pi_x[e^{\int_0^t \alpha(\xi_s) ds} \phi_0(\xi_t)(f(\xi_t) - f(\xi_0))]|}{\phi_0(x)} \\ &\leq \varepsilon + \sup_{x \in D} \frac{e^{-\lambda_0 t} |\Pi_x[e^{\int_0^t \alpha(\xi_s) ds} \phi_0(\xi_t)|f(\xi_t) - f(\xi_0)|; |\xi_t - \xi_0| \geq \delta]|}{\phi_0(x)} \\ &\leq \varepsilon + \sup_{x \in D} c \frac{e^{(-\lambda_0 + \|\alpha\|_\infty)t} \|\phi_0\|_\infty \|f\|_\infty \Pi_x(|\xi_t - \xi_0| \geq \delta)}{\phi_0(x)} \\ &\leq \varepsilon + \sup_{x \in D} c \frac{\Phi^{1/2}(\delta_D(x)) t^{-1/2} \int_{y \in D: |y-x| > \delta} t j(c_2|y-x|/4) dy}{\Phi^{1/2}(\delta_D(x))} \\ &\leq \varepsilon + c\sqrt{t} \int_{|z| \geq c_2\delta/4} (1 \wedge |z|^2) j(|z|) dz. \end{aligned} \tag{4.5}$$

It follows that

$$\lim_{t \rightarrow 0} \|T_t^{\phi_0} f - f\|_\infty = 0$$

and Assumption 1.3 is satisfied.

Example 4.5 (Subordinate Brownian motion with Gaussian component) Suppose that $S = \{S_t, t \geq 0\}$ is a subordinator with drift $b > 0$. The Laplace exponent ϕ of S can be written in the form

$$\phi(\lambda) = b\lambda + \int_0^{+\infty} (1 - e^{-\lambda t}) u(dt), \tag{4.6}$$

where u is a measure on $(0, +\infty)$ satisfying

$$\int_0^{+\infty} (1 \wedge t) u(dt) < +\infty.$$

Without loss of generality, we assume that $b = 1$. In this example, we will assume that ϕ is a complete Bernstein function and that the Lévy density $u(t)$ of S satisfies the following growth condition near zero: for any $M > 0$, there exists $c = c(M) > 1$ such that

$$u(r) \leq cu(2r), \quad r \in (0, M). \tag{4.7}$$

Let $W = \{W_t, t \geq 0\}$ be a Brownian motion in \mathbb{R}^d independent of the subordinator S . The subordinate Brownian motion $Y = \{Y_t, t \geq 0\}$ is defined by $Y_t := W_{S_t}$, which is a rotationally symmetric Lévy process with Lévy exponent $\phi(|\xi|^2)$. It is known that the Lévy measure of the process Y has a density $j(|x|)$ given by (4.3).

For any open set $D \subset \mathbb{R}^d$ and positive constants c_1 and c_2 , we define

$$h_{D,c_1,c_2}(t,x,y) := \left(1 \wedge \frac{\delta_D(x)}{\sqrt{t}}\right) \left(1 \wedge \frac{\delta_D(y)}{\sqrt{t}}\right) (t^{-d/2} e^{-c_1|x-y|^2/t} + t^{-d/2} \wedge tj(c_2|x-y|)). \quad (4.8)$$

Suppose that D is a bounded $C^{1,1}$ open set with characteristics (R_0, Λ) , and let ξ be the subprocess of Y killed upon leaving D . It is known that ξ is a Hunt process symmetric with respect to the Lebesgue measure on D and that ξ has a strictly positive continuous transition density $p_D(t,x,y)$ with respect to the Lebesgue measure on D . We assume the following upper bound condition on the transition density function $\tilde{p}(t,|x|)$ of Y : for any $T > 0$, there exist $C_j \geq 1$, $j = 1, 2, 3$, such that for all $(t,r) \in (0,T] \times [0, \text{diam}(D)]$,

$$\tilde{p}(t,r) \leq C_1 \left(t^{-d/2} e^{-r^2/C_2 t} + t^{-d/2} \wedge tj\left(\frac{r}{C_3}\right) \right). \quad (4.9)$$

It was established in [6] that the above estimate holds for a large class of symmetric diffusion processes with jumps with $D = \mathbb{R}^d$. Using Meyer's method of removing and adding jumps, it can be shown that (4.9) is true for a larger class of symmetric Markov processes, including subordinate Brownian motions with Gaussian components under some additional condition. See the paragraph containing (1.12) in [5] for more information.

The following is proved in [5, Theorem 1].

(i) For every $T > 0$, there exist

$$c_1 = c_1(R_0, \Lambda_0, \lambda_0, T, \psi, d) > 0, \quad c_2 = c_2(R_0, \Lambda_0, \lambda_0, d) > 0,$$

such that for all $(t,x,y) \in (0,T] \times D \times D$,

$$p_D(t,x,y) \geq c_1 h_{D,c_2,1}(t,x,y). \quad (4.10)$$

(ii) If D satisfies (4.9), then for every $T > 0$, there exists

$$c_3 = c_3(R_0, \Lambda_0, T, d, \psi, C_1, C_2, C_3, d) > 1$$

such that for all $(t,x,y) \in (0,T] \times D \times D$,

$$p_D(t,x,y) \leq c_3 h_{D,C_4,C_5}(t,x,y), \quad (4.11)$$

where

$$C_4 = (16C_2)^{-1}, \quad C_5 = (8 \vee 4C_3)^{-1}.$$

Let $E = D$, and let m be the Lebesgue measure on D . Since $p_D(t, x, y)$ is symmetric, we have

$$a_t(x) = p_D(2t, x, x) \leq ct^{-d/2}.$$

Thus, Assumption 1.1 is satisfied.

Suppose that the branching rate function β and the branching mechanism satisfy the assumptions of Section 1.2, and that the corresponding superprocess X is supercritical. Using the above two-sided heat kernel estimate for ξ , we can establish in a similar way as in Example 4.4 that Assumption 1.3 also holds.

Remark 4.6 In fact, in Examples 4.4 and 4.5, ξ does not need to be a subordinate Brownian motion killed upon leaving D . All we need are the heat kernel estimates like (4.4) or (4.10)-(4.11). For example, suppose that Y^D is the subprocess of some subordinate Brownian motion Y killed upon leave D that has property (4.4) or (4.10)-(4.11). Let ξ be a Markov process obtained from Y^D though a Feynman-Kac transform with bounded potential function. Then ξ enjoys property (4.4) or (4.10)-(4.11). For other examples of processes that satisfy two-sided bounds similar to (4.4), including censored stable processes in $C^{1,1}$ open sets and their local and non-local Feynman-Kac transforms, see [4]. Our main results are applicable to these processes as well.

In all the examples above, the spatial motion ξ is symmetric. Now, we give two examples where the spatial motion ξ is not symmetric.

Example 4.7 Suppose $d \geq 3$ and that $\nu = (\nu^1, \nu^2, \dots, \nu^d)$, where each ν^j is a signed measure on \mathbb{R}^d such that

$$\limsup_{r \rightarrow 0} \sup_{x \in \mathbb{R}^d} \int_{B(x,r)} \frac{|\nu^j|(dy)}{|x-y|^{d-1}} = 0.$$

Let $\xi^{(1)} = \{\xi_t^{(1)}, t \geq 0\}$ be a Brownian motion with drift ν in \mathbb{R}^d , see [1]. Suppose that D is a bounded domain in \mathbb{R}^d . Let $M > 0$ so that $B(0, M/2) \supset D$. Put $B = B(0, M)$. Let G_B be the Green function of $\xi^{(1)}$ in B and define

$$H(x) := \int_B G_B(y, x) dy.$$

Then H is a strictly positive continuous function on B . Let ξ be the process obtained by killing $\xi^{(1)}$ upon exiting D . ξ is a Hunt process and it has a strictly positive continuous transition density $\tilde{p}(t, x, y)$ with respect to the Lebesgue measure on D . Let $E = D$, and let m be the measure defined by

$$m(dx) = H(x) dx.$$

It follows from [16,17] that ξ has a dual process with respect to m . The transition density of ξ with respect to m is given by

$$p(t, x, y) = \frac{\tilde{p}(t, x, y)}{H(y)}.$$

Suppose further that D is $C^{1,1}$. Then it follows from [15, Theorem 4.6] that there exist $c_1 > 1$ and $c_2 > c_3 > 0$ such that for all $(t, x, y) \in (0, 1] \times D \times D$,

$$\begin{aligned} & c_1^{-1} t^{-d/2} \left(1 \wedge \frac{\delta_D(x)}{\sqrt{t}}\right) \left(1 \wedge \frac{\delta_D(y)}{\sqrt{t}}\right) \exp\left(-\frac{c_2|x-y|^2}{t}\right) \\ & \leq \tilde{p}(t, x, y) \\ & \leq c_1 t^{-d/2} \left(1 \wedge \frac{\delta_D(x)}{\sqrt{t}}\right) \left(1 \wedge \frac{\delta_D(y)}{\sqrt{t}}\right) \exp\left(-\frac{c_3|x-y|^2}{t}\right). \end{aligned}$$

It follows from the display above and the semigroup property that, for any $t > 0$, $\tilde{p}(t, x, y)$ is bounded. By [17, (2.6)], there exists $c_4 > 1$ such that

$$c_4^{-1} \delta_B(x) < H(x) < c_4 \delta_B(x).$$

So for $x \in D$, $c \leq H(x) \leq C$, where $c, C > 0$. Thus, $p(t, x, y)$ is also bounded in D and m is a finite measure. Thus, Assumption 1.1 is satisfied.

Suppose that the branching rate function β and the branching mechanism satisfy the assumptions of Section 1.2, and that the corresponding superprocess X is supercritical. Using the above two-sided heat kernel estimate for ξ , we can establish in a similar way as in Example 4.4 that Assumption 1.3 also holds.

Example 4.8 Suppose $d \geq 2$, $\alpha \in (1, 2)$, and that $\nu = (\nu^1, \nu^2, \dots, \nu^d)$, where each ν^j is a signed measure on \mathbb{R}^d such that

$$\limsup_{r \rightarrow 0} \sup_{x \in \mathbb{R}^d} \int_{B(x, r)} \frac{|\nu^j|(dy)}{|x-y|^{d-\alpha+1}} = 0.$$

Let $\xi^{(2)} = \{\xi_t^{(2)}, t \geq 0\}$ be an α -stable process with drift ν in \mathbb{R}^d , see [18]. Suppose that D is a bounded open set in \mathbb{R}^d and suppose that $M > 0$ is such that $D \subset B(0, M/2)$. Put $B = B(0, M)$. Let G_B be the Green function of $\xi^{(2)}$ in B and define

$$H(x) := \int_B G_B(y, x) dy.$$

Then H is a strictly positive continuous function on B . Let ξ be the process obtained by killing $\xi^{(2)}$ upon exiting D . ξ is a Hunt process and it has a strictly positive continuous transition density $\tilde{p}(t, x, y)$ with respect to the Lebesgue measure on D . Let $E = D$, and let m be the measure defined by

$$m(dx) = H(x) dx.$$

It follows from [2, Section 5] and [18] that ξ has a dual process with respect to m . The transition density of ξ with respect to m is given by

$$p(t, x, y) = \frac{\tilde{p}(t, x, y)}{H(y)}.$$

By [2, Corollary 1.4] and [19], we can check that

$$c_0^{-1} \delta_B^{\alpha/2}(x) < H(x) < c_0 \delta_B^{\alpha/2}(x),$$

where $c_0 > 1$. Thus, for $x \in D$, $c \leq H(x) \leq C$ for some $c, C > 0$.

Suppose further that D is $C^{1,1}$. Then it follows from [2, Theorem 1.3] and [19] that there exists $c_1 > 1$ such that for all $(t, x, y) \in (0, 1] \times D \times D$,

$$\begin{aligned} & c_1^{-1} \left(1 \wedge \frac{\delta_D^{\alpha/2}(x)}{\sqrt{t}}\right) \left(1 \wedge \frac{\delta_D^{\alpha/2}(y)}{\sqrt{t}}\right) \left(t^{-d/\alpha} \wedge \frac{t}{|x-y|^{d+\alpha}}\right) \\ & \leq \tilde{p}(t, x, y) \\ & \leq c_1 \left(1 \wedge \frac{\delta_D^{\alpha/2}(x)}{\sqrt{t}}\right) \left(1 \wedge \frac{\delta_D^{\alpha/2}(y)}{\sqrt{t}}\right) \left(t^{-d/\alpha} \wedge \frac{t}{|x-y|^{d+\alpha}}\right). \end{aligned}$$

It follows from the display above and the semigroup property that, for any $t > 0$, $\tilde{p}(t, x, y)$ is bounded. Since H is bounded between two positive constants, Assumption 1.1 is satisfied.

Suppose that the branching rate function β and the branching mechanism satisfy the assumptions of Section 1.2, and that the corresponding superprocess X is supercritical. Using the above two-sided heat kernel estimate for ξ , we can establish in a similar way as in Example 4.4 that Assumption 1.3 also holds.

In the following example, our main result does not apply directly. However, we could apply our main result after a transform.

Example 4.9 Suppose that the spatial motion $\xi = \{\xi_t, \Pi_x\}$ is an OU-process on \mathbb{R}^d with infinitesimal generator

$$\mathcal{L} = \frac{1}{2} \sigma^2 \Delta - cx \cdot \nabla \quad \text{on } \mathbb{R}^d,$$

where $\sigma, c > 0$. Without loss of generality, we assume $\sigma = 1$. Let

$$\varphi(x) := \left(\frac{c}{\pi}\right)^{d/2} e^{-c|x|^2}, \quad m(dx) = \varphi(x)dx.$$

Then (ξ, Π_x) is symmetric with respect to the probability measure $m(dx)$.

Let

$$a(x) = c_1|x|^2 + c_2, \quad c_1, c_2 > 0,$$

and let P_t^a be the Feynman-Kac semigroup:

$$P_t^a f(x) := \Pi_x[e^{\int_0^t a(\xi_s) ds} f(\xi_t)].$$

Suppose $c > \sqrt{2c_1}$ and write

$$v = \frac{1}{2} (c - \sqrt{c^2 - 2c_1}).$$

Let

$$\lambda_c := \inf\{\lambda \in \mathbb{R} : \text{there exists } u > 0 \text{ such that } (\mathcal{L} + a - \lambda)u = 0 \text{ in } \mathbb{R}^d\}$$

be the generalized principal eigenvalue. Let h denote the corresponding ground state, i.e., $h > 0$ such that

$$(\mathcal{L} + a - \lambda_c)h = 0.$$

As was indicated in [12],

$$\lambda_c = c_2 + dv > 0, \quad h(x) = \left(\frac{c-2v}{c}\right)^{d/2} \exp\{v|x|^2\}.$$

Note that

$$h = e^{-\lambda_c t} P_t^a h$$

on \mathbb{R}^d . Let Π_x^h be defined as in (1.16) with ϕ_0 replaced by h . The transformed process (ξ, Π_x^h) is also an OU-process with infinitesimal generator

$$\frac{1}{2} \Delta - (c-2v)x \cdot \nabla \quad \text{on } \mathbb{R}^d.$$

Let

$$\psi(x, z) = -a(x)z + b(x)z^2,$$

where $b \in C^\eta(\mathbb{R}^d)$, $b(x) > 0$ for all $x \in \mathbb{R}^d$. A superprocess X with spacial motion ξ , branching rate $\beta(x) = 1$, and branching mechanism ψ , can be defined by

$$X = \frac{1}{h} X^h,$$

where X^h is the superprocess with spacial motion (ξ, Π_x^h) , branching rate $\beta(x) = 1$, and branching mechanism

$$\psi^h(x, z) = -\lambda_c z + h(x)b(x)z^2.$$

Assume that hb is bounded in \mathbb{R}^d . Then, for X^h , we have

$$m^h(dx) = \left(\frac{c-2v}{\pi}\right)^{d/2} e^{-(c-2v)|x|^2} dx, \quad \lambda_0^h = \lambda_c, \quad \phi_0^h = 1.$$

From the discussion in Example 4.1, we see that Assumptions 1.1 and 1.3 are satisfied for the superprocess X^h . Then, there exists $\Omega_0 \subset \Omega$ of probability one (that is, $\mathbb{P}_\mu(\Omega_0) = 1$ for every $\mu \in \mathcal{M}_F(\mathbb{R}^d)$) such that, for every $\omega \in \Omega_0$ and for every bounded Borel measurable function $f \geq 0$ on \mathbb{R}^d with $f/h \leq c$ for some $c > 0$ and that the set of discontinuous points of f has zero m -measure, we have

$$\begin{aligned} \lim_{t \rightarrow +\infty} e^{-\lambda_0^h t} \left\langle \frac{f}{h}, X_t^h \right\rangle(\omega) &= W_\infty(\omega) \int_{\mathbb{R}^d} \left(\frac{f}{h}\right)(y) m^h(dy) \\ &= W_\infty(\omega) \left(\frac{c}{\pi}\right)^{d/2} \int_{\mathbb{R}^d} f(y) e^{(v-c)|y|^2} dy, \end{aligned} \quad (4.12)$$

where $W_\infty(\omega)$ is the limit of the martingale

$$W_t := e^{-\lambda_c t} \langle 1, X_t^h \rangle = e^{-\lambda_c t} \langle h, X_t \rangle$$

as $t \rightarrow +\infty$. We rewrite (4.12) to get the limit result on X :

$$\begin{aligned} \lim_{t \rightarrow +\infty} e^{-\lambda_c t} \langle f, X_t \rangle(\omega) &= W_\infty(\omega) \int_{\mathbb{R}^d} \left(\frac{c}{\pi}\right)^{d/2} e^{(v-c)|y|^2} f(y) dy \\ &= W_\infty(\omega) \int_{\mathbb{R}^d} \tilde{\phi}_0(y) f(y) dy, \end{aligned} \quad (4.13)$$

where

$$\tilde{\phi}_0 = \left(\frac{c}{\pi}\right)^{d/2} e^{(v-c)|y|^2}.$$

Since h is bounded from below, in the weak topology,

$$e^{-\lambda_c t} X_t \rightarrow W_\infty(\omega) \tilde{\phi}_0(x) dx, \quad \mathbb{P}_\mu\text{-a.s.},$$

for any $\mu \in \mathcal{M}_F(\mathbb{R}^d)$. This example covers [10, Example 4.7].

Acknowledgements The research of Zhen-Qing Chen is partially supported by NSF grant DMS-1206276 and the National Natural Science Foundation of China (Grant No. 11128101). The research of Yan-Xia Ren was supported by the National Natural Science Foundation of China (Grant Nos. 11271030, 11128101). The research of Renming Song was supported in part by a grant from the Simons Foundation (208236), and also in part by a project funded by the Priority Academic Program Development (PAPD) of Jiangsu Higher Education Institutions during his stay in Jiangsu Normal University.

References

1. Bass R F, Chen Z -Q. Brownian motion with singular drift. *Ann Probab*, 2013, 31: 791–817
2. Chen Z -Q, Kim P, Song R. Dirichlet heat kernel estimates for fractional Laplacian with gradient perturbation. *Ann Probab*, 2012, 40: 2483–2538
3. Chen Z -Q, Kim P, Song R. Dirichlet heat kernel estimates for rotationally symmetric Lévy processes. *Proc Lond Math Soc*, 2014, 109: 90–120
4. Chen Z -Q, Kim P, Song R. Stability of Dirichlet heat kernel estimates for non-local operators under Feynman-Kac perturbations. *Trans Amer Math Soc*, 2015, 367: 5237–5270
5. Chen Z -Q, Kim P, Song R. Dirichlet heat kernel estimates for subordinate Brownian motions with Gaussian components. *J Reine Angew Math*, Ahead of print, DOI:10.1515/crelle-2013-0090
6. Chen Z -Q, Kumagai T. A priori Hölder estimate, parabolic Harnack principle and heat kernel estimates for diffusions with jumps. *Rev Mat Iberoam*, 2010, 26: 551–589
7. Chen Z -Q, Ren Y -X, Wang H. An almost sure scaling limit theorem for Dawson-Watanabe superprocesses. *J Funct Anal*, 2008, 254: 1988–2019
8. Chen Z -Q, Yang T. Dirichlet heat kernel estimates for fractional Laplacian under non-local perturbation. Preprint, 2015, arXiv: 1503.05302

9. Davies E B, Simon B. Ultracontractivity and the kernel for Schrödinger operators and Dirichlet Laplacians. *J Funct Anal*, 1984, 59: 335–395
10. Eckhoff M, Kyprianou A E, Winkel M. Spine, skeletons and the strong law of large numbers for superdiffusion. *Ann Probab (To appear)*, arXiv: 1309.6196
11. Engländer J. Law of large numbers for superdiffusions: The non-ergodic case. *Ann Inst Henri Poincaré Probab Stat*, 2009, 45: 1–6
12. Engländer J, Harris S C, Kyprianou A E. Strong law of large numbers for branching diffusions. *Ann Inst Henri Poincaré Probab Stat*, 2010, 46(1): 279–298
13. Engländer J, Turaev D. A scaling limit theorem for a class of superdiffusions. *Ann Probab*, 2002, 30: 683–722
14. Engländer J, Winter A. Law of large numbers for a class of superdiffusions. *Ann Inst Henri Poincaré Probab Stat*, 2006, 42: 171–185
15. Kim P, Song R. Two-sided estimates on the density of Brownian motion with singular drift. *Illinois J Math*, 2006, 50: 635–688
16. Kim P, Song R. On dual processes of non-symmetric diffusions with measure-valued drifts. *Stochastic Process Appl*, 2008, 118: 790–817
17. Kim P, Song R. Intrinsic ultracontractivity of non-symmetric diffusions with measure-valued drifts and potentials. *Ann Probab*, 2008, 36: 1904–1945
18. Kim P, Song R. Stable process with singular drift. *Stochastic Process Appl*, 2014, 124: 2479–2516
19. Kim P, Song R. Dirichlet heat kernel estimates for stable processes with singular drift in unbounded $C^{1,1}$ open sets. *Potential Anal*, 2014, 41: 555–581
20. Kouritzin M A, Ren Y -X. A strong law of large numbers for super-stable Processes. *Stochastic Process Appl*, 2014, 121: 505–521
21. Li Z. *Measure-valued Branching Markov Processes*. Heidelberg: Springer, 2011
22. Liu R -L, Ren Y -X, Song R. Strong law of large numbers for a class of superdiffusions. *Acta Appl Math*, 2013, 123: 73–97
23. Pazy A. *Semigroups of Linear Operators and Applications to Partial Differential Equations*. New York: Springer, 1983
24. Ren Y -X, Song R, Zhang R. Central limit theorems for supercritical superprocesses. *Stochastic Process Appl*, 2015, 125: 428–457
25. Ren Y -X, Song R, Zhang R. Central limit theorems for supercritical branching nonsymmetric Markov processes. *Ann Probab (to appear)*, arXiv: 1404.0116
26. Ren Y -X, Song R, Zhang R. Functional central limit theorems for supercritical superprocesses. Preprint, 2014, arXiv: 1410.1598
27. Ren Y -X, Song R, Zhang R. Limit theorems for some critical superprocesses. Preprint, 2014, arXiv: 1403.1342
28. Sato K -I. *Lévy Processes and Infinitely Divisible Distribution*. Cambridge: Cambridge University Press, 1999
29. Schaefer H H. *Banach Lattices and Positive Operators*. New York: Springer, 1974
30. Stroock D W. *Probability Theory. An Analytic View*. 2nd ed. Cambridge: Cambridge University Press, 2011
31. Wang L. An almost sure limit theorem for super-Brownian motion. *J Theoret Probab*, 2010, 23: 401–416