

## **Structure, mechanism and ensemble formation of the Alkylhydroperoxide Reductase subunits**

### **AhpC and AhpF from *Escherichia coli***

Phat Vinh Dip<sup>1,#</sup>, Neelagandan Kamariah<sup>2,#</sup>, Malathy Sony Subramanian Manimekalai<sup>1</sup>, Wilson Nartey<sup>1</sup>, Asha Manikkoth Balakrishna<sup>1</sup>, Frank Eisenhaber<sup>2,3,4</sup>, Birgit Eisenhaber<sup>2</sup> and Gerhard Grüber<sup>1,2,\*</sup>

<sup>1</sup> Nanyang Technological University, School of Biological Sciences, 60 Nanyang Drive, Singapore 637551, Republic of Singapore

<sup>2</sup> Bioinformatics Institute, Agency for Science, Technology and Research (A\*STAR), 30 Biopolis Street, #07-01 Matrix, Singapore 138671, Republic of Singapore

<sup>3</sup> School of Computer Engineering, Nanyang Technological University (NTU), 50 Nanyang Drive, Singapore 637553, Republic of Singapore

<sup>4</sup> Department of Biological Sciences, National University of Singapore, 8 Medical Drive, Singapore 117597, Republic of Singapore

# Authors contributed equally to this work

\* To whom correspondence should be addressed: Prof. Dr. Gerhard Grüber, Tel.: + 65 - 6316 2989;

Fax: + 65 - 6791 3856, E-mail: [ggrueber@ntu.edu.sg](mailto:ggrueber@ntu.edu.sg)

## Supplemental Figure Legends

**Figure S1:** (A) The homo-dimeric form of *EcAhpF*, revealing a maximum length of 170 Å. The second symmetry related *EcAhpF* molecule is depicted with lighter shaded color. The 90° view shows the distance (125 Å) between the NTD redox centers (C129/132 and C129'/132') of the molecule. The dimer interface covers mainly the FAD-FAD'-domain with the solvent accessible area of 2,520 Å<sup>2</sup>, which is around 10% of the total monomeric solvent accessible area of 25,103 Å<sup>2</sup>. The interface is stabilized by hydrophobic- and hydrophilic interactions, like hydrogen bonds and salt bridges. (B) The N-terminal domain redox-active dithiol (C129/132) is reduced in the 2 Å resolution structure, while oxidized in the 2.65 Å one. Stereo view of the omit map ( $\sigma$ -weighted Fo-Fc (mFo-DFc) electron density map) contoured at 3  $\sigma$  for the region around the N-terminal domain redox center (C129/132) shows that NTD is oxidized in the 2.65 Å structure (i) and reduced in the 2 Å resolution structure (ii). (C) The *E. coli* AhpF crystal structure at 2 Å and 2.65 Å resolution. The four segments of both *EcAhpF* structures, the N-terminal domain (orange), linker segment (green), the FAD-domain (yellow) and the NADH-domain (brown) were aligned (the structure at 2.65 Å is depicted in lighter shades). The NTD adopts different conformation and the redox-active dithiol (C129/132) (purple and red) is reduced in the 2 Å resolution structure, while oxidized in the 2.65 Å structure. (D) Cadmium ion is located in the interface between neighboring *EcAhpF* molecules. Two highly conserved and catalytically important histidine side chains (H130 and H85) are protruded on the same side to make a positively charged groove, wherein one cadmium ion from the crystallizing solution is bound. The cadmium ion forms six coordinated bonds, two to E110' and each to H85, His114', H130 and one water molecule, respectively. The symbol (') highlight the symmetry related residue.

**Figure S2:** In order to identify the NADH binding pocket, the *EcAhpF* structures were compared with the *E. coli* TrxR (PDB ID: 1TDF), which has a bound NADP<sup>+</sup>. The NADP<sup>+</sup>-binding motif

(<sub>152</sub>GGGNTA<sub>157</sub>) is conserved with the sequence of Gly-X-Gly-X-X-Ala in all NADP-binding domains, in which the alanine is replaced with glycine in *EcAhpF* structure (<sub>362</sub>GGGNSG<sub>367</sub>). Another important substitution is the replacement of the polar R176 to non-polar F386 in the *EcAhpF* structure. R176 in *EcTrxR* interacts with the phosphate group of NADP<sup>+</sup> and this interaction is hindered by the substitution to F386 in the *EcAhpF* structure, which might explain its higher preference for NADH over NADPH. A significant structural difference was observed in the nicotinamide interacting loop region in the FAD-domain. In *EcAhpF*, the loop is displaced away from the nicotinamide ring by about 3.5 Å relative to *EcTrxR*, wherein the nicotinamide interacting residues, H290 and R293, are replaced by V492 and K495 in *EcAhpF*, respectively. In the 2.65 Å *EcAhpF* structure, two sulphate molecules are found. The first sulphate is hydrogen bonded to NADP binding motif residues N365 and S366 and the second one is close to the FAD molecule and is hydrogen bonded with W326, Q448 and G450. In the 2.0 Å *EcAhpF* structure, the NADH binding pocket is filled with one sulfate, two glycerol and one PEG molecules. The sulfate occupies similar position as that in the 2.65 Å *EcAhpF* structure and has interactions with N365 and S366. The first glycerol has interaction with K391 and the other one has interaction with F386 and E385. The PEG molecule interacts with V447, I449, G450, L451, S366 and R327. These interacting residues may likely represent the NADH binding pocket residues of AhpF. It is also seen from both the structures that the sulfate molecule bound in the NADH binding pocket may have an additional role by enhancing the NADH binding and hence in the catalytic activity of AhpF.

**Figure S3:** Solution X-ray scattering studies of *EcAhpF*. (A) Small angle X-ray scattering pattern (○) and its corresponding fitting curves (—; green: experimental, red: calculated) at 8 mg/ml concentration. The curves of *E. coli* AhpF are displayed in logarithmic unit for clarity. (B) Distance distribution functions of *EcAhpF* at 8 mg/ml concentration. (C) *EcAhpF* solution shape as calculated by GASBOR program. (D) Theoretical scattering curves for the monomer (magenta) and dimer (orange) of open conformation and monomer (blue) and dimer (cyan) of closed conformation for *EcAhpF* calculated

using the CRY SOL program fits with experimental scattering pattern ( $\circ$ ) with a  $\chi$ -value of 4.4 and 2.1 and 5.1 and 2.2, respectively.

**Figure S4:** (A) Electron density for the redox active C<sub>p</sub>47'/C<sub>R</sub>166 disulfide bond between molecule A (green) and molecule B (cyan). The omit map ( $\sigma$ -weighted Fo-Fc (mFo-DFc) electron density map) is contoured at 2  $\sigma$  (green lines) around the disulphide bond. The symbol ' is used for residues of the symmetry subunit in the dimer formation. (B) The redox active cysteines (C47/C166' and C47'/C166) of *StAhpC* are depicted in magenta and are not involved in an intermolecular disulfide bond. The C-terminal amino acids of *StAhpC* are ordered and form intermolecular hydrogen bonds.

**Figure S5:** The FAD- and NADH-domain are responsible for the head-to-tail dimer formation in the *EcAhpF*. The second symmetry related *EcAhpF* molecule is depicted with lighter shaded color. The dimer interface is stabilized by the hydrophobic, hydrogen and salt bridge interactions. There is no significant difference observed in the dimer interface between the *EcAhpF*- and *StAhpF* structures, except for an additional salt bridge between K234 and D373 and a bifurcated hydrogen bond between the side chain nitrogen atom of R233 with main chain oxygen atoms of H347 and C348, that could have occurred in *EcAhpF*. Such bridge is missing in *StAhpF*, due to the rotational motion of the NADH-domain. The symbol \* indicates the residues, belonging to symmetry molecule.

**Figure S6:** The high purity of *E. coli* AhpF-CTD (A) and NTD<sub>1-196</sub> (B) are shown in the SDS-PAGE. Small angle X-ray scattering pattern ( $\circ$ ) of *EcAhpF*-CTD (F) and *EcNTD*<sub>1-196</sub> (C), and the corresponding experimental fitting curves (green; the red curve represents the calculated one). (D, G) Distance distribution function of *EcAhpF*-CTD (G) and NTD<sub>1-196</sub> (D). The solution shape of *EcAhpF*-CTD and *EcNTD*<sub>1-196</sub> are revealed in column (H) and (E), which have been calculated by GASBOR program.

(A)





Figure S1A

**(B(i))**



**(B(ii))**




Figure S1B





Figure S1C-D






Figure S2



(A)



(B)



(C)




Figure S3A-C





Figure S3D

(A)



(B)




Figure S4A-B




Figure S5




Figure S6A-B







**(C)****(D)****(E)****(F)****(G)****(H)**

Figure S6C-H