

Published in final edited form as:

Nature. 2004 June 10; 429(6992): 671–675. doi:10.1038/nature02588.

Structure of a complex between a voltage-gated calcium channel β -subunit and an α -subunit domain

Filip Van Petegem, Kimberly A. Clark, Franck C. Chatelain, and Daniel L. Minor Jr
 Cardiovascular Research Institute, Departments of Biochemistry and Biophysics, and Cellular and Molecular Pharmacology, University of California San Francisco, 513 Parnassus Avenue, Box 0130, San Francisco, California 94143, USA

Abstract

Voltage-gated calcium channels (Ca_Vs) govern muscle contraction, hormone and neurotransmitter release, neuronal migration, activation of calcium-dependent signalling cascades, and synaptic input integration¹. An essential Ca_V intracellular protein, the β -subunit ($\text{Ca}_V\beta$)^{1,2}, binds a conserved domain (the α -interaction domain, AID) between transmembrane domains I and II of the pore-forming α_1 subunit³ and profoundly affects multiple channel properties such as voltage-dependent activation², inactivation rates², G-protein modulation⁴, drug sensitivity⁵ and cell surface expression^{6,7}. Here, we report the high-resolution crystal structures of the $\text{Ca}_V\beta_{2a}$ conserved core, alone and in complex with the AID. Previous work suggested that a conserved region, the β -interaction domain (BID), formed the AID-binding site^{3,8}; however, this region is largely buried in the $\text{Ca}_V\beta$ core and is unavailable for protein–protein interactions. The structure of the AID– $\text{Ca}_V\beta_{2a}$ complex shows instead that $\text{Ca}_V\beta_{2a}$ engages the AID through an extensive, conserved hydrophobic cleft (named the α -binding pocket, ABP). The ABP–AID interaction positions one end of the $\text{Ca}_V\beta$ near the intracellular end of a pore-lining segment, called IS6, that has a critical role in Ca_V inactivation^{9,10}. Together, these data suggest that $\text{Ca}_V\beta\text{s}$ influence Ca_V gating by direct modulation of IS6 movement within the channel pore.

The 1.97 Å resolution structure of the $\text{Ca}_V\beta_{2a}$ core shows that $\text{Ca}_V\beta\text{s}$ comprise two well-conserved domains (Fig. 1a). The first, an SH3 fold, contains five antiparallel β -strands ($\beta 1$ – $\beta 5$), a 3_{10} helix ($\eta 1$), and two α -helices ($\alpha 1$ and $\alpha 2$) that lie amino-terminal to $\beta 1$ and carboxy-terminal to $\beta 4$, respectively. The strand that completes the SH3 fold, $\beta 5$ (residues 217–224), is separated in the primary structure from the core of the SH3 domain by approximately 70 residues (variable domain 2, V2, a site of splice variation and amino acid insertions and deletions²) that are absent from the structure (Fig. 1b). The second conserved domain consists of a five-stranded parallel β -sheet ($\beta 6$ – $\beta 10$), surrounded by six α -helices ($\alpha 3$ – $\alpha 8$) and two 3_{10} helices ($\eta 2$ and $\eta 3$), and is related to the core of nucleotide kinase enzymes.

$\text{Ca}_V\beta\text{s}$ share structural features with membrane-associated guanylate kinases (MAGUKs), a protein scaffold family that organizes signalling components near membranes¹¹. MAGUKs contain one or more PDZ domains N-terminal to an SH3 domain, a bridging region known as a HOOK domain and a nucleotide kinase domain^{11,12}. PDZ domains are approximately

© 2004 Nature Publishing Group

Correspondence and requests for materials should be addressed to D.L.M. (minor@itsa.ucsf.edu).

Supplementary Information accompanies the paper on www.nature.com/nature.

Competing interests statement The authors declare that they have no competing financial interests.

Coordinates and structure factors have been deposited in the Protein Data Bank under accession codes 1T0H and 1T0J.

100 residues long. The $\text{Ca}_v\beta_{2a}$ structure indicates that $\text{Ca}_v\beta$ s lack N-terminal PDZ domains. There are too few residues N-terminal to the SH3 domain (even in $\text{Ca}_v\beta_{1b}$, the $\text{Ca}_v\beta$ with the longest (55 amino acids) N-terminal variable region 1, V1) to fold as a PDZ domain. The absence of a PDZ domain distinguishes $\text{Ca}_v\beta$ s from canonical MAGUK proteins.

Comparison of $\text{Ca}_v\beta_{2a}$ with a representative MAGUK, PSD-95 (refs 12, 13), reveals other differences. Superposition of the nucleotide kinase domains shows that the relative orientations of the SH3 and nucleotide kinase domains differ by approximately 90° , an arrangement that makes $\text{Ca}_v\beta_{2a}$ a more elongated structure (Fig. 2a). The nucleotide kinase domain of MAGUKs is homologous to guanylate kinases and retains guanosine monophosphate (GMP) binding, but key residues for enzymatic function are missing¹². The four-stranded β -sheet nucleotide kinase subdomain that binds GMP in MAGUKs is absent in $\text{Ca}_v\beta_{2a}$ (Fig. 2a). Furthermore, two $\text{Ca}_v\beta_{2a}$ loops (between $\beta 7$ – $\eta 2$ and $\beta 8$ – $\beta 9$) occlude part of the binding site for the GMP guanosine ring. Thus, the $\text{Ca}_v\beta$ nucleotide kinase domain seems to have lost the ability to bind nucleotides.

The structures of $\text{Ca}_v\beta_{2a}$ and PSD-95 SH3 domains are similar (Fig. 2b). Neither is compatible with canonical modes of proline-rich ligand binding. Both lack the aromatic residues necessary for ligand engagement¹³, and the surface that would bind polyproline ligands is blocked by the $\alpha 2$ helix^{12,13}. In PSD-95, residues C-terminal to the nucleotide kinase domain contribute an extra SH3 β -strand that is absent from canonical SH3 domains¹³ and absent in $\text{Ca}_v\beta_{2a}$. The HOOK domain, present in MAGUKs and $\text{Ca}_v\beta_{2a}$, bridges SH3 β -strands $\beta 4$ and $\beta 5$ and comprises $\alpha 2$ and variable domain 2 of $\text{Ca}_v\beta_{2a}$. HOOK domains are important regulatory regions for interactions of MAGUKs with other proteins^{11,12,14} and may serve a similar function for $\text{Ca}_v\beta$ protein–protein interactions.

$\text{Ca}_v\beta$ s exert their effects on Ca_v function by binding the pore-forming α_1 subunit at a conserved, 18-residue sequence located between membrane domains I and II (the AID)^{3,15} (Fig. 3a). Interpretation of mutagenesis and biochemical studies suggested that $\text{Ca}_v\beta$ –AID interactions occur through a 41-residue segment ($\text{Ca}_v\beta_{2a}$ residues 212–252) termed the β -interaction domain (BID)^{8,16}. The $\text{Ca}_v\beta_{2a}$ structure shows that the central region of the BID, which includes the residues previously thought to be important for BID–AID interactions⁸, is entirely buried and cannot participate directly in protein–protein interactions (Fig. 3b). Two putative BID phosphorylation sites^{8,16} are also buried in this region. Given the extent of burial, mutations used to determine the relative importance of residues involved in BID–AID interactions (for example, proline to arginine)⁸ are likely to have abolished AID binding by disrupting the folded structure of the nucleotide kinase domain rather than by perturbing direct AID contacts. Although it would appear that the $\text{Ca}_v\beta_{2a}$ structure conflicts with previous data, most of the supporting evidence for the BID–AID interaction relied on indirect functional experiments and direct BID–AID binding was never demonstrated^{8,16}.

If the BID does not interact directly with the AID, how do $\text{Ca}_v\beta_{2a}$ and the AID interact? To answer this, we solved the high-resolution (2.00 Å) structure of a complex between the conserved $\text{Ca}_v\beta_{2a}$ core and an 18-residue peptide containing the AID from the L-type channel $\text{Ca}_v1.2$. The electron density reveals the first 16 residues of the AID and the location of the binding pocket on $\text{Ca}_v\beta_{2a}$ (Fig. 3c). Overall, the $\text{Ca}_v\beta_{2a}$ structure is very similar to the unbound form (root-mean-square deviation (RMSD)_{C α} = 0.397 Å) and bears only a few conformational changes in side chains near the AID-binding pocket ($\text{Ca}_v\beta_{2a}$ residues M244, N390, E393 and R351). The AID forms an α -helix that is anchored to the binding pocket through a set of conserved residues (AID residues L434, G436, Y437, W440 and I441) that are important for $\text{Ca}_v\beta$ binding and for α -subunit modulation by $\text{Ca}_v\beta$ s^{8,15,17}. These residues bind a deep groove that we call the α -binding pocket (ABP), formed by helices $\alpha 3$,

$\alpha 5$ and $\alpha 8$ of the $\text{Ca}_v\beta_{2a}$ nucleotide kinase domain at a site distal to the SH3 domain (Figs 1a, 3b and 4a).

The complex buries approximately 730 \AA^2 of ABP surface, of which about 350 \AA^2 are hydrophobic. AID side chains D433, W440 and Q443 make direct hydrogen bonds to the ABP (Supplementary Fig. 1). The depth and extent of burial of the aromatic AID positions Y437 and W440 is particularly striking (Fig. 4b, c). The AID Y437 hydroxyl group is central to a buried hydrogen bond network comprising three water molecules, AID residue D433 and five $\text{Ca}_v\beta_{2a}$ residues (Supplementary Fig. 1). Mutation of this tyrosine to phenylalanine greatly diminishes AID– $\text{Ca}_v\beta$ binding¹⁵. The extensive AID–ABP interactions are consistent with biochemical experiments demonstrating strong AID– $\text{Ca}_v\beta$ interaction (dissociation constant $\sim 6\text{--}20 \text{ nM}$)¹⁸. The $\text{Ca}_v\beta$ side chains that contact the AID are highly conserved among $\text{Ca}_v\beta$ isoforms (see Figs 1b, 3a and 4a; see also Supplementary Fig. 1). Thus, both binding partners engage each other through conserved residues to create the AID–ABP interaction.

Interactions between the $\text{Ca}_v\alpha_1$ and $\text{Ca}_v\beta$ subunits markedly influence the cell surface expression of functional channels^{6,7}. Control of Ca_v trafficking by regulating $\text{Ca}_v\alpha_1$ – $\text{Ca}_v\beta$ interactions is emerging as an important means of modulating cellular excitability⁷. Ca_v channel subtypes are major clinical targets for drugs that treat cardiovascular disease, migraine and pain¹⁹. Development of compounds that could interfere with the AID–ABP binding interactions might provide new ways to modulate Ca_v function in pathological states.

The $\text{Ca}_v\beta$ –AID structure provides a starting point for understanding how $\text{Ca}_v\beta$ modulates numerous channel properties. G-protein $\beta\gamma$ subunits ($\text{G}\beta\gamma$) inhibit Ca_v function^{1,4}; however, the sites of $\text{G}\beta\gamma$ – Ca_v interactions and precise inhibitory mechanisms remain highly controversial⁴. Biochemical experiments show that $\text{G}\beta\gamma$ binds the Ca_v2 AID and that mutations at AID positions Q1, Q2, R5, L7, G9 and Y10 (corresponding to $\text{Ca}_v1.2$ AID residues 428, 429, 432, 434, 436 and 437) abolish $\text{G}\beta\gamma$ –AID binding²⁰. In contrast, other studies suggest that the *in vitro* $\text{G}\beta\gamma$ –AID interaction is functionally irrelevant and that the relevant $\text{G}\beta\gamma$ -binding determinants lie elsewhere in the channel cytoplasmic domains^{21,22}. The structure of the $\text{Ca}_v\beta_{2a}$ –AID complex shows that three of the putative $\text{G}\beta\gamma$ –AID interacting positions (L7, G9 and Y10), which are invariant in Ca_v1 and Ca_v2 channels (Fig. 3a), are deeply buried by interactions with $\text{Ca}_v\beta$ (Fig. 4a, b; see also Supplementary Fig. 2 and Table 2). The extent of burial of these residues, which are critical for maintaining $\text{Ca}_v\beta$ –AID association^{3,15}, suggests that $\text{G}\beta\gamma$ and $\text{Ca}_v\beta$ cannot bind to the AID simultaneously. Taken together with the observation that the $\text{G}\beta\gamma$ –AID affinity is at least 10–20-fold weaker than the $\text{Ca}_v\beta$ –AID²⁰ affinity, the structure also indicates that it is unlikely that $\text{G}\beta\gamma$ could effectively compete with $\text{Ca}_v\beta$ for AID binding without drastic structural rearrangement of the $\text{Ca}_v\beta$ –AID complex. Thus, our data lend support to the view that the major $\text{G}\beta\gamma$ interaction sites lie in other Ca_v cytoplasmic domains^{21–23}.

How might $\text{Ca}_v\beta$ s affect channel gating? Although the detailed mechanisms of Ca_v inactivation processes remain unknown, functional experiments show that the IS6 pore-lining segment has a key role¹⁰. Motions of pore-lining transmembrane segments are a common theme emerging in ion channel gating^{24,25}. Twenty-two residues separate the AID helix N terminus from the cytoplasmic end of IS6. We do not know the structure, but sequence evaluation suggests that these residues have a high helix propensity and could readily form a continuous helix between the AID helix and the presumed transmembrane helix of IS6 (Fig. 5a). AID residue 5 (E432, here) slows inactivation when negatively charged (as in Ca_v1 channels) and speeds inactivation when positively charged (as in Ca_v2 channels)^{26,27}. This residue is exposed on the surface of the AID helix (Fig. 4a) where it

would be available to interact with other parts of the channel and could influence the rates of movement of the $\text{Ca}_V\beta$ -AID complex and therefore IS6. The profound inactivation rate slowing caused by $\text{Ca}_V\beta_{2a}$ requires anchoring of the N terminus to the membrane by palmitoylation²⁸. Orientation of the AID helix towards IS6 places the N-terminal membrane anchor on the periphery of the α_1 subunit and suggests that $\text{Ca}_V\beta_{2a}$ slows channel inactivation by restricting the movement of the IS6 transmembrane domain. $\text{Ca}_V\beta$ s have a deep groove between the SH3 and nucleotide kinase domains that is on the same face as the V2 domain. These features may be used to engage other cytoplasmic parts of the channel^{29,30} and allow $\text{Ca}_V\beta$ to couple motions in other channel domains directly to IS6 through AID attachment.

The structures presented here represent the first high-resolution view of any part of the voltage-gated calcium channel, and provide an important step towards understanding the detailed molecular mechanism models for how Ca_V s function. This work strongly suggests that $\text{Ca}_V\beta$ affects Ca_V gating properties by directly influencing conformational changes that are likely to occur in the channel pore¹⁰.

Methods

The crystal structures of recombinant $\text{Ca}_V\beta_{2a}$ and $\text{Ca}_V\beta_{2a}$ -AID complexes were solved to resolutions of 1.97 Å and 2.00 Å, respectively. The final R/R_{free} values are 18.55%/21.32% for $\text{Ca}_V\beta_{2a}$ and 19.97%/24.15% for the $\text{Ca}_V\beta_{2a}$ -AID complex. Figures were prepared with PyMOL, MOLSCRIPT and RASTER3D. The experimental details for protein expression, purification, crystallization, structure solution, statistics of data collection, phasing and refinement are available as Supplementary Information.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank J. M. Berger, K. Brejc, D. Fass, D. Julius, E. A. Lumpkin and B. A. Schulman for comments on the manuscript; J. Holton at beamline 8.3.1 at the Advanced Light Source for help with data collection; R. W. Tsien and D. T. Yue for the calcium channel clones; and members of the Minor laboratory for support at all stages of this work. This work was supported by awards to D.L.M. from the McKnight Foundation for Neuroscience, the March of Dimes Basil O'Connor Scholar program, the Alfred P. Sloan Foundation, and the Rita Allen Foundation. D.L.M. is a McKnight Foundation Scholar, an Alfred P. Sloan Research Fellow and a Rita Allen Foundation Scholar.

References

1. Catterall WA. Structure and regulation of voltage-gated Ca^{2+} channels. *Annu. Rev. Cell Dev. Biol.* 2000; 16:521–555. [PubMed: 11031246]
2. Dolphin AC. β -subunits of voltage-gated calcium channels. *J. Bioenerg. Biomembr.* 2003; 35:599–620. [PubMed: 15000522]
3. Pragnell M, et al. Calcium channel β -subunit binds to a conserved motif in the I–II cytoplasmic linker of the α_1 -subunit. *Nature.* 1994; 368:67–70. [PubMed: 7509046]
4. Dolphin AC. G protein modulation of voltage-gated calcium channels. *Pharmacol. Rev.* 2003; 55:607–627. [PubMed: 14657419]
5. Hering S. β -Subunits: fine tuning of Ca^{2+} channel block. *Trends Pharmacol. Sci.* 2002; 23:509–513. [PubMed: 12413805]
6. Bichet D, et al. The I–II loop of the Ca^{2+} channel α_1 subunit contains an endoplasmic reticulum retention signal antagonized by the β subunit. *Neuron.* 2000; 25:177–190. [PubMed: 10707982]
7. Beguin P, et al. Regulation of Ca^{2+} channel expression at the cell surface by the small G-protein kir/*Gem.* *Nature.* 2001; 411:701–706. [PubMed: 11395774]

8. De Waard M, Scott VE, Pragnell M, Campbell KP. Identification of critical amino acids involved in $\alpha 1$ - β interaction in voltage-dependent Ca^{2+} channels. *FEBS Lett.* 1996; 380:272–276. [PubMed: 8601439]
9. Stotz SC, Hamid J, Spaetgens RL, Jarvis SE, Zamponi GW. Fast inactivation of voltage-dependent calcium channels. A hinged-lid mechanism? *J. Biol. Chem.* 2000; 275:24575–24582. [PubMed: 10823819]
10. Zhang JF, Ellinor PT, Aldrich RW, Tsien RW. Molecular determinants of voltage-dependent inactivation in calcium channels. *Nature.* 1994; 372:97–100. [PubMed: 7969428]
11. Dimitratos SD, Woods DF, Stathakis DG, Bryant PJ. Signaling pathways are focused at specialized regions of the plasma membrane by scaffolding proteins of the MAGUK family. *Bioessays.* 1999; 21:912–921. [PubMed: 10517864]
12. Tavares GA, Panepucci EH, Brunger AT. Structural characterization of the intramolecular interaction between the SH3 and guanylate kinase domains of PSD-95. *Mol. Cell.* 2001; 8:1313–1325. [PubMed: 11779506]
13. McGee AW, et al. Structure of the SH3-guanylate kinase module from PSD-95 suggests a mechanism for regulated assembly of MAGUK scaffolding proteins. *Mol. Cell.* 2001; 8:1291–1301. [PubMed: 11779504]
14. Paarmann I, Spangenberg O, Lavie A, Konrad M. Formation of complexes between Ca^{2+} .calmodulin and the synapse-associated protein SAP97 requires the SH3 domain-guanylate kinase domain-connecting HOOK region. *J. Biol. Chem.* 2002; 277:40832–40838. [PubMed: 12189141]
15. Witcher DR, De Waard M, Liu H, Pragnell M, Campbell KP. Association of native Ca^{2+} channel β subunits with the $\alpha 1$ subunit interaction domain. *J. Biol. Chem.* 1995; 270:18088–18093. [PubMed: 7629119]
16. De Waard M, Pragnell M, Campbell KP. Ca^{2+} channel regulation by a conserved β subunit domain. *Neuron.* 1994; 13:495–503. [PubMed: 8060623]
17. Berrou L, Klein H, Bernatchez G, Parent L. A specific tryptophan in the I–II linker is a key determinant of β -subunit binding and modulation in $\text{Ca}(\text{V})_{2.3}$ calcium channels. *Biophys. J.* 2002; 83:1429–1442. [PubMed: 12202369]
18. Opatowsky Y, Chomsky-Hecht O, Kang MG, Campbell KP, Hirsch JA. The voltage-dependent calcium channel β subunit contains two stable interacting domains. *J. Biol. Chem.* 2003; 278:52323–52332. [PubMed: 14559910]
19. Kochegarov AA. Pharmacological modulators of voltage-gated calcium channels and their therapeutical application. *Cell Calcium.* 2003; 33:145–162. [PubMed: 12600802]
20. DeWaard M, et al. Direct binding of G-protein $\beta\gamma$ complex to voltage-dependent calcium channels. *Nature.* 1997; 385:446–450. [PubMed: 9009193]
21. Zhang JF, Ellinor PT, Aldrich RW, Tsien RW. Multiple structural elements in voltage-dependent Ca^{2+} channels support their inhibition by G proteins. *Neuron.* 1996; 17:991–1003. [PubMed: 8938130]
22. Qin N, Platano D, Olcese R, Stefani E, Birnbaumer L. Direct interaction of $\text{G}\beta\gamma$ with a C-terminal $\text{G}\beta\gamma$ -binding domain of the Ca^{2+} channel $\alpha 1$ subunit is responsible for channel inhibition by G protein-coupled receptors. *Proc. Natl Acad. Sci. USA.* 1997; 94:8866–8871. [PubMed: 9238069]
23. Ivanina T, Blumenstein Y, Shistik E, Barzilai R, Dascal N. Modulation of L-type Ca^{2+} channels by $\text{G}\beta\gamma$ and calmodulin via interactions with N and C termini of $\alpha 1\text{C}$. *J. Biol. Chem.* 2000; 275:39846–39854. [PubMed: 10995757]
24. Miyazawa A, Fujiyoshi Y, Unwin N. Structure and gating mechanism of the acetylcholine receptor pore. *Nature.* 2003; 424:949–955. [PubMed: 12827192]
25. Jiang Y, et al. The open pore conformation of potassium channels. *Nature.* 2002; 417:523–526. [PubMed: 12037560]
26. Herlitz S, Hockerman GH, Scheuer T, Catterall WA. Molecular determinants of inactivation and G protein modulation in the intracellular loop connecting domains I and II of the calcium channel $\alpha 1\text{A}$ subunit. *Proc. Natl Acad. Sci. USA.* 1997; 94:1512–1516. [PubMed: 9037084]
27. Berrou L, Bernatchez G, Parent L. Molecular determinants of inactivation within the I–II linker of $\alpha 1\text{E}$ ($\text{CaV}_{2.3}$) calcium channels. *Biophys. J.* 2001; 80:215–228. [PubMed: 11159396]

28. Restituito S, et al. The $\beta 2a$ subunit is a molecular groom for the Ca^{2+} channel inactivation gate. *J. Neurosci.* 2000; 20:9046–9052. [PubMed: 11124981]
29. Walker D, et al. A new β subtype-specific interaction in $\alpha 1A$ subunit controls P/Q-type Ca^{2+} channel activation. *J. Biol. Chem.* 1999; 274:12383–12390. [PubMed: 10212211]
30. Walker D, Bichet D, Campbell KP, De Waard M. A $\beta 4$ isoform-specific interaction site in the carboxyl-terminal region of the voltage-dependent Ca^{2+} channel $\alpha 1A$ subunit. *J. Biol. Chem.* 1998; 273:2361–2367. [PubMed: 9442082]




Figure 1.

Structure of the $\text{Ca}_v\beta_{2a}$ - $\text{Ca}_v1.2$ AID complex. **a**, Ribbon diagram of the complex. Dashed lines indicate regions absent from the structures. SH3 and nucleotide kinase (NK) domains are shown in green and blue, respectively. The AID is shown in red. $\text{Ca}_v\beta_{2a}$ α -helices are labelled. Variable regions V1, V2 and V3 are indicated. The $\text{Ca}_v\beta_{2a}$ unbound structure is similar to that shown here for the complex. The arrow indicates where the AID connects to transmembrane segment IS6. **b**, Sequence alignment of representatives of each $\text{Ca}_v\beta$ isoform. The top sequence shows residues 40–425 of rat $\text{Ca}_v\beta_{2a}$. Numbers on the right denote each line's terminal residue. Shading denotes residues identical among isoforms. The two $\text{Ca}_v\beta_{2a}$ domains used for crystallization are indicated in green and blue, respectively. Secondary structure elements are indicated: α , α -helix; η , 3_{10} helix; β , β -strand. Dashed lines indicate residues present in the crystallized constructs but absent in the electron density. Location of the V2 and part of the V3 regions are shown. Asterisks identify residues that contribute side-chain contacts to the AID-binding pocket; diamonds mark side chains with direct hydrogen bonds to the AID.




Figure 2.

Structural comparisons between PSD-95 (gold) and $\text{CaV}\beta_{2a}$ (blue). **a**, Superposition of $\text{CaV}\beta_{2a}$ and PSD-95 nucleotide kinase domains ($\text{RMSD}_{\text{C}\alpha} = 3.9 \text{ \AA}$). The dashed circle indicates the guanosine-monophosphate (GMP)-binding domain present in PSD-95 but absent in $\text{CaV}\beta_{2a}$. The guanosine monophosphate molecule bound to PSD-95 is displayed in space-filling representation. Nucleotide kinase (NK) and SH3 domains are indicated. The relative change in SH3 domain orientation is indicated. **b**, Superposition of PSD-95 and $\text{CaV}\beta_{2a}$ SH3 domains ($\text{RMSD}_{\text{C}\alpha} = 1.6 \text{ \AA}$). Position of the polyproline ligand from a superposition with the Sem5 SH3 domain (Protein Data Bank code 2SEM) ($\text{RMSD}_{\text{C}\alpha} = 1.8$

Å) is shown in space-filling representation. The Sem5 SH3 is not shown. The DALI server generated the superpositions (<http://www.ebi.ac.uk/dali/>).




Figure 3. Features of the AID–Cav β_{2a} interaction and location of the previously described BID. **a**, Sequence alignment of AID domains (Cav1.2 residues 428–445) and neighbouring residues. The positions of the last transmembrane segment of transmembrane domain I (IS6) and the first transmembrane segment of transmembrane domain II (IIS1) are shown. Secondary structure of the AID from the co-crystal structure is indicated (red). Dashed lines indicate residues absent from the electron density. Asterisks identify side-chain contacts with Cav β_{2a} closer than 4 Å. **b**, Position of the previously described BID (residues 212–252; yellow)^{3,8}. Residues previously proposed to mediate AID–BID interactions (P224, P228, P234, Y239) are indicated and have relative accessibilities of 1.4%, 0%, 0% and 32.4%. Putative PKC

sites S225, S235 and S345 are also shown (magenta) and have relative accessibilities of 8.8%, 0% and 35.4%, respectively. S345 accessibility reduces to 12% in the complex. Accessibility values are relative to a tripeptide, Gly-X-Gly. **c**, The left panel shows $F_o - F_c$ electron density, contoured at 2σ , for the AID-Ca $\nu\beta_{2a}$ complex before building the AID. The right panel shows final $2F_o - F_c$ density, contoured at 1σ , for the AID from the refined AID-Ca $\nu\beta_{2a}$ structure (right). In both panels the final AID model is shown.




Figure 4. AID–ABP interactions. **a**, Surface representation of the $Ca_v\beta_{2a}$ ABP, bound to the AID. The AID (gold) is shown in stick representation. Y437 and W440 are white. $Ca_v\beta_{2a}$ residues that contribute hydrophobic (blue) and hydrogen bond (red) side-chain contacts to the AID are labelled. Select residues of the AID are labelled to orient the reader. **b**, **c**, Slices through the AID–ABP interaction at AID positions Y437 and W440 (gold). Labels indicate the AID residues.




Figure 5.

Cartoon of proposed model for how $\text{Ca}_V\beta$ affects $\text{Ca}_V\alpha_1$ gating. **a**, Orientation of $\text{Ca}_V\beta_2a$ with respect to the I-II loop (red), pore-forming subunit, and connection to IS6. In $\text{Ca}_V\beta_2a$, variable region 1 (V1) is tethered to the membrane. The I-II loop between the AID N terminus and IS6 is depicted as a helix. $\text{Ca}_V\beta_2a$ SH3 and nucleotide kinase domains are coloured green and blue, respectively. The arrow indicates that $\text{Ca}_V\beta$ couples to IS6 movements (rotations, translations or both). **b**, View from the opposite side of **a**. The groove between SH3 and nucleotide kinase domains (demarcated by the arrow) and two flexible $\text{Ca}_V\beta_2a$ regions, the 275-284 loop and variable region 2 (V2), are on the same $\text{Ca}_V\beta$ face,

opposite the ABP. These regions may interact with other pore-forming subunit cytoplasmic domains.