

ORIGINAL INVESTIGATION

Erich Roessler · Laureane Mittaz · Yangzhu Du  
Hamish S. Scott · Jenny Chang · Colette Rossier  
Michel Guipponi · Seiichi P.T. Matsuda  
Maximilian Muenke · Stylianos E. Antonarakis

## Structure of the human *Lanosterol Synthase* gene and its analysis as a candidate for holoprosencephaly (*HPE1*)

Received: 1 June 1999 / Accepted: 16 August 1999 / Published online: 26 October 1999

**Abstract** Holoprosencephaly (HPE) is the most common birth defect of the brain in humans. It involves various degrees of incomplete separation of the cerebrum into distinct left and right halves, and it is frequently accompanied by craniofacial anomalies. The *HPE1* locus in human chromosome 21q22.3 is one of a dozen putative genetic

loci implicated in causing HPE. Here, we report the complete gene structure of the human *lanosterol synthase* (*LS*) gene, which is located in this interval, and present its mutational analysis in HPE patients. We considered *LS* an excellent candidate HPE gene because of the requirement for cholesterol modification of the Sonic Hedgehog protein for the correct patterning activity of this HPE-associated protein. Despite extensive pedigree analysis of numerous polymorphisms, as well as complementation studies in yeast on one of the missense mutations, we find no evidence that the *LS* gene is in fact *HPE1*, implicating another gene located in this chromosomal region in HPE pathogenesis.

E. Roessler and L. Mittaz contributed equally to this work.

E. Roessler · J. Chang · M. Muenke (✉)  
Medical Genetics Branch,  
National Human Genome Research Institute,  
National Institutes of Health, 10 Center Drive,  
MSC 1852, Bld. 10 10C101, Bethesda MD, 20892–1852, USA  
e-mail: mmuenke@nhgri.nih.gov,  
Tel.: +1-301-4028167, Fax: +1-301-4967157

E. Roessler · Y. Du · M. Muenke  
Department of Pediatrics,  
University of Pennsylvania School of Medicine,  
and The Children's Hospital of Philadelphia,  
Philadelphia, PA, USA

E. Roessler · Y. Du · M. Muenke  
Department of Genetics,  
University of Pennsylvania School of Medicine,  
and The Children's Hospital of Philadelphia,  
Philadelphia, PA, USA

M. Muenke  
Department of Neurology,  
University of Pennsylvania School of Medicine,  
and The Children's Hospital of Philadelphia,  
Philadelphia, PA, USA

L. Mittaz · H. S. Scott · C. Rossier · M. Guipponi  
S. E. Antonarakis  
Division of Medical Genetics,  
University of Geneva Medical School, Geneva, Switzerland  
e-mail: Stylianos.Antonarakis@medecine.unige.ch,  
Tel.: +41-22-702-5708, Fax: +41-22-702-5706

S. P. T. Matsuda  
Department of Chemistry, Rice University, Houston, TX, USA

S. P. T. Matsuda  
Department of Biochemistry, Rice University, Houston, TX, USA

S. P. T. Matsuda  
Department of Cell Biology, Rice University, Houston, TX, USA

### Introduction

Holoprosencephaly (HPE) is a clinically heterogeneous brain malformation, which can be caused by genetic and/or environmental factors (Ming et al. 1998a; Roessler and Muenke 1998). In cyclopia, the most severe form, there is a single eye, a superiorly placed nasal structure called a proboscis, and the lack of separation of the cerebral hemispheres into discrete left and right halves (Cohen 1989). The clinical HPE spectrum is variable even in autosomal dominant families segregating mutations in a single HPE gene such as *Sonic Hedgehog* (Ming and Muenke 1998). Craniofacial findings of microcephaly, hypotelorism, or a single central incisor can be clinical clues to underlying brain anomalies comprising HPE (DeMyer et al. 1964). Recently, several genes have been identified with a role in HPE pathogenesis, including *Sonic Hedgehog* (*HPE3*) (Belloni et al. 1996; Roessler et al. 1996, 1997a, 1997b), the transcription factor *ZIC2* (Brown et al. 1998a, 1998b), *SIX3* (*HPE2*) (Wallis et al. 1999), and several others currently under investigation (Gripp et al. 1998; Ming et al. 1998b). The *HPE1* locus on human chromosome 21q22.3 is one of several named HPE loci (Frézal and Schinzel 1991) and is defined by a set of three overlapping cytogenetic deletions (Muenke et al. 1995).

The lanosterol synthase (*LS*) enzyme catalyzes the cyclization step in the biosynthesis of cholesterol. Among

the essential functions of cholesterol in preserving the integrity of cell membranes and, as the substrate for the synthesis of sterols and hormones, there is a novel role recently described for cholesterol as an adduct to signaling molecules such as Sonic hedgehog (Porter et al. 1996a, 1996b). Undermodification of the Sonic Hedgehog protein is suspected to be a contributing factor in the pathogenesis of Smith-Lemli-Opitz (SLO) syndrome, which in turn is associated with an increased incidence of HPE (Kelley et al. 1996). SLO has been shown to result from the biochemical deficiency of cholesterol during fetal development (Cunniff et al. 1997) and is associated with mutations in the enzyme that catalyzes the penultimate step in cholesterol biosynthesis, 7- $\delta$  sterol reductase (Fitzky et al. 1998; Moebius et al. 1998; Wassif et al. 1998; Waterham et al. 1998). We speculated that mutations in the human *LS* gene might lead to similar patterning deficiencies of the Sonic Hedgehog protein and contribute to brain malformations within the HPE spectrum. Now, we report on the gene structure of the human *LS* gene and a strategy to amplify each coding exon for mutational analysis. Here, we show the mutational analysis of the human *LS* gene (Young et al. 1996), which maps within the 5 cM minimal critical region for *HPE1* (Muenke et al. 1995).

## Materials and methods

### Determination of the *LS* gene structure

Having previously identified a cosmid 7G8 (Young et al. 1996) that contained the 5' end of the *LS* gene, we used the primer pairs LSH2830F (5' ggcagtagtgatgcacccgt 3') and HSL3000R (5' gacttgggatgttccatgacagac 3'), derived from the 3' UTR sequence, as a probe to screen a chromosome-21-specific cosmid library (LL2INCO2-Q, Soeda et al. 1995) for the 3' end of the gene. A single cosmid, 72H3, was chosen for further analysis, since it completed the contig for the entire *LS* gene. Putative exon-specific primers based on the cDNA sequence and the initial 5 trapped exons (Young et al. 1996) were then used to sequence the cosmids and to determine the first set of intron/exon boundaries. Additional primers were then synthesized to determine the sequence of both DNA strands adjacent to the intron/exon boundaries as each new set was empirically determined. Note that the initiator methionine is in exon 2. Genbank numbers are as follows: accession no. AJ239021 (exons 2, 3, and 4), AJ239022 (exons 5 and 6), AJ23903 (exon 7), AJ239024 (exons 8, 9, and 10), AJ239025 (exon 11), AJ239026 (exon 12), AJ239027 (exons 13 and 14), AJ239028 (exons 15, 16, 17, and 18), AJ239029 (exons 19 and 20), AJ239030 (exon 21), AJ239031 (exons 22 and 23). Sequencing was performed on an ABI PRISM 377 automated sequencer.

### Patients and DNA preparation

Patient samples were obtained after being given informed consent according to the guidelines of the Children's Hospital of Philadelphia Institutional Review Board, and DNA samples were prepared from blood or lymphoblastoid cells lines using routine methods. Three HPE patients had various deletions, including the distal long arm of chromosome 21. Two of these were previously published: 46,XX,del(21)(q22.3->qter) and 46,XY,-21,+der(21)t(10;21)(p11.2;q22.3)mat (Muenke et al. 1995). A third patient with semilobar HPE had the following karyotype: 46,XY,r(21).

A total of 30 HPE patients from 20 unrelated families were chosen for single-strand conformational polymorphism (SSCP)

analysis. These HPE families were chosen for their apparent autosomal recessive pedigrees with clinically normal parents and multiple affected children. These families included the typical range of HPE phenotypes. When SSCP band shifts were detected, these samples were sequenced to determine the potential mutations. Where possible, the sequence variations were examined in the parents and siblings to determine the pattern of transmission and also examined in a panel of normal controls to determine any correlation with the disease state or uniqueness of the sequence variations. All families were too small for effective linkage analysis. As has been shown, some of these families could also be consistent with germline mosaicism for a dominant HPE gene, such as *SHH* (Roessler et al. 1996, 1997a) or *SIX3* (Wallis et al. 1999).

### Reverse-transcriptase polymerase chain reaction analysis

Poly-A selected RNA was isolated from lymphoblastoid cell lines (FastTrack, InVitrogen), synthesized into complementary DNA using random primers (cDNA cycle kit, InVitrogen), and amplified with the three primer sets as depicted in Fig. 1. The first primer pair is LSF2 (5' cccaagcttaggggctctgaacgggatgaca 3') and LSR2 (5' cccaagcttcgatgctgatgctcttggatgaat 3'), which overlaps with the second primer pair LSF3 (5' ggggaattcccagcaccgattcacca 3') and LSR3 (5' cccaagcttaggagcagcacagcctcaa 3'), which overlaps with the last amplicon LSF4 (5' ggggaattcccagataaccctccgactacc 3') and LSR4 (5' cccaagctttctcaccagcccgacaagc 3'). Amplicons were directionally subcloned into Bluescript KS<sup>-</sup> (Stratagene) using the *Hind*III or *Eco*RI sites nested within them and pools of at least 20 clones were subjected to automated sequencing. Candidate sequence changes were confirmed by individual exon amplification using patient genomic DNA and re-sequencing. The reverse-transcription polymerase chain reaction (RT-PCR) products were sufficient to screen exons 5 to 23; the first three coding exons (2, 3 and 4) were analyzed by direct amplification and sequencing of the patient genomic DNA.


### Mutational analysis

Amplification of genomic DNA was performed in a 15- $\mu$ l reaction volume, using 60–100 ng DNA template, 200  $\mu$ M each of dATP, dGTP, and dTTP, 125  $\mu$ M dCTP, 3.5  $\mu$ Ci  $\alpha$ <sup>32</sup>P-dCTP (800 Ci/mmol [10 mCi/ml]), 30 pmol each primer, 1.5  $\mu$ l 10 $\times$  PCR buffer (Gibco), 1.25  $\mu$ l 10 $\times$  PCR Enhancer (Gibco), 1.5 mM MgSO<sub>4</sub> (Gibco), and 1 U *Amplitaq* polymerase (Perkin Elmer). All of the PCR reactions were performed in a PTC-100 thermal cycler (MJ Research, Inc.).

For SSCP analysis, following amplification, 5  $\mu$ l of each sample was diluted with 10  $\mu$ l stop solution [9.5 ml deionized formamide (Fluka), 20 mM ethylene diamine tetraacetic acid (EDTA), 0.5% (w/v) bromophenol blue, and 0.5% (w/v) xylene cyanol], denatured for 10 min at 94°C, then immediately chilled on ice. The samples were run either with or without 5% glycerol at room temperature on 0.5 $\times$  MDE gel solution (FMC Bioproducts) sequencing-grade polyacrylamide gels. Sequencing of the amplicons demonstrating SSCP band shifts was performed by the Protein and DNA Core Facility of the Children's Hospital of Philadelphia on an ABI Prism 377 analyzer.

### Functional analysis of *LS* activity

The yeast *LS* cDNA (Corey et al. 1994) was modified by site-directed mutagenesis (Ausubel et al. 1999) to include the corresponding K625Q mutation and was expressed in a yeast mutant for functional complementation analysis. The yeast *LS* mutant SMY8 was transformed using this construct, and the resultant strain's ability to grow on media without supplemental ergosterol was tested as described previously (Corey et al. 1996).



**Fig. 1** In the *top panel*, a diagram of the *lanosterol synthase* (*LS*) cDNA is shown aligned with a set of three amplicons used in reverse-transcription polymerase chain reaction (RT-PCR) experiments to subclone the *EcoRI* (R1) and *HindIII* (HIII) fragments a, b, c, and d for sequencing. In the *bottom panel*, the same *LS* cDNA diagram and its relationship to the genomic structure of individual exons is shown. Note that the coding exons extend from exon 2 to 23

## Results

### Structure of the human *LS* gene

A contig of two cosmids (7G8 and 72H3) was analyzed to identify 23 individual exons spanning 40–50 kb of genomic DNA for the *LS* gene as diagramed in Fig. 1. The genomic structure shown in Fig. 1, predicts a total of 22 coding exons and is in agreement with the cDNA sequence (Baker et al. 1995). Note that the 5' untranslated region represented as exon 1 has not been completely characterized. Also note that the sequence variants seen in the hemizygous probands (see below) were detected as expressed transcripts as well as from genomic DNA, indicating that this is a functional gene. The relevant sequences of the intron/exon boundaries of the *LS* gene are shown in Table 1.

### RT-PCR analysis of HPE patients with deletions of 21q22.3

Poly-A RNA was isolated from an affected member of each of the three different 21q22.3 hemizygous deletions (these unrelated families are denoted here as families A, B, and C), which served to define the *HPE1* locus. In all three cases, the deletion includes the human *LS* gene based on the comparison of the somatic cell hybrid data defining *HPE1* (Muenke et al. 1995) and the genetic position of the YACs and cosmids containing the *LS* gene (Young et al. 1996). We reasoned that these would be the most likely HPE individuals to have a loss of function mutation in the remaining *LS* allele. As shown in Table 2 (number 6), we found a sequence variation 863C>G in family A, predicting no change in the coding sequence (P288P) and this change was also noted in four additional HPE families (G, H, M, P) as a common polymorphism that failed to segregate with the disease. This sequence variant was also readily detectable in either the heterozygous or homozygous state in the normal control population. In family A, we also noted a 1949T>C sequence variation in exon 20, predicting no change in the coding sequence (H650H). This variation was seen as a homozy-

**Table 1** Sequences of the lanosterol synthase (LS) intron/exon boundaries

Exon	Splice donor	Size bp (Intron)	Splice acceptor	Size bp (Exon)
1	5'UTR	> 300 (1)		> 133 (1)
2	ACGGAGGGCAC gtagt	130 (2)	ccggtggcctcag GTGTCTGCGG	29 (2)
3	GGCTGGACACC gtaagt	730 (3)	cgatctcttgcag AAGAATTACT	166 (3)
4	CCTCCTGCCAG gtagga	> 480 (4)	ctcctgccacacag GCCTCCTGAT	139 (4)
5	GGCTGGGGCCT gtagt	650 (5)	ctctgctcctgtag GCACATTGAG	109 (5)
6	CACAAGAAAG gtacgg	> 970 (6)	ctctccccgctcag GTGGTGCTGT	121 (6)
7	CAGAGATGTG gtatgt	> 900 (7)	tgctgttgccccag GCTGTTTCTT	97 (7)
8	GCCTCCGCCAG gtagga	600 (8)	actgccccaccag GAGCTCTATG	136 (8)
9	CGTGGTATATG gtcac	390 (9)	cctgtcccctgtag CGCTCCTCAA	109 (9)
10	GCATCGGCCCG gtcagt	> 850 (10)	tgtctctcctgtag ATCTCGAAAA	119 (10)
11	GACTATCTCTG gtagt	> 540 (11)	tttctctcacag GATGGGCCTT	98 (11)
12	TGAAAATGCAG gtaagg	> 500 (12)	tcctttgtgtgtag GGCACCAACG	28 (12)
13	CTCTGCTTGTG gttcgt	1060 (13)	gtctctctctccag GCGGGCGGGC	57 (13)
14	GGCTCTCACAG gtagg	> 920 (14)	tcgatgctgcccag GTCCCAGATA	72 (14)
15	AGATGCGCAAAG gtagt	670 (15)	cttgtgctcctgtag GGTGGCTTCT	51 (15)
16	CTGTGGCTGTG gtaagg	660 (16)	ctctgtgatttcag CTGCTGAACA	150 (16)
17	GGAGGTCTTCC gtagt	730 (17)	cctccccttcccag GGGACATCAT	97 (17)
18	GCGGAGATCCG gtaagg	760 (18)	atctcgtgtccag GGAGACCCTC	106 (18)
19	TCCTGGGAAGG gtagt	450 (19)	ctgtttccctccag CTCCTGGGGA	66 (19)
20	TACCGAGATGG gtagt	> 430 (20)	tctggtctttgtag GACTGCCTGT	81 (20)
21	ATGGCCGTTCG gtgggg	> 450 (21)	ttctctccaccag GCATCCTGAC	170 (21)
22	ACTGGCCGCAG gtagt	640 (22)	ctgtgttcttttag GAAAACATTG	79 (22)
23	TGA codon at position 130–132			> 960 (23)

**Table 2** Sequence variations in the human *lanosterol synthase* (LS) gene. For the normal controls, the first number is the number of individuals who were heterozygous for the normal and the sequence variation, followed by the number of individuals homozygous for the sequence variation, followed by the number of indi-

viduals studied by means of single-strand conformational polymorphism (SSCP). All band-shift patterns detected in the normal controls were verified by sequencing. *N.D.* not detected under these gel conditions. Notation follows the recommendations of Antonarakis and the Nomenclature Workshop Group (1998)

Sequence variation	Intron	Exon	Codon	Position	Family	Normal controls hetero/homo/total
1. GCC to GCT		4	A121A	362C>T	D, E, F	12/2/56
2. CGA to CAA		5	R175Q	523G>A	D, E, F	10/5/54
3. cccc(c/g)ctcag	5		Acceptor	IVS5–5C>G	K, M	16/12/54
4. cccc(c/t)accag	7		Acceptor	IVS7–6C>T	E <sup>b</sup> , F, G, H, M, P	13/13/53
5. CCT to CCC		8	P282P	845T>C	G	N.D.
6. CCC to CCG		8	P288P	863C>G	A <sup>a</sup> , G, H, M <sup>b</sup> , P	14/17/53
7. ccctgtc(c/t)cctgtag	8		Acceptor	IVS8–8C>T	M	N.D.
8. CAC to CGC		9	H310R	928A>G	D, E, F	11/1/50
9. ttgc(a/g)cct	10		None	IVS10–29A>G	D, E, F	6/0/50
10. acgt(a/g)agc	12		None	IVS12+32A>G	L, O	18/16/52
11. agatgca(c/t)ggg	18		None	IVS18+36C>T	D <sup>b</sup> , G <sup>b</sup> , I	28/12/52
12. atcca(c/g)aag	18		None	IVS18+45C>G	E <sup>b</sup> , F <sup>b</sup> , H <sup>b</sup> , I <sup>b</sup>	22/20/52
13. CTG to CGG		20	W576R	1728T>G	K, R <sup>b</sup>	N.D.
14. CGG to CAG		20	R622Q	1864G>A	C <sup>a</sup>	2/0/103
15. TTG to GTG		20	L641V	1923T>G	B <sup>a</sup> , E <sup>b</sup> , F <sup>b</sup> , K <sup>b</sup> , P	19/20/53
16. CAT to CAC		20	H650H	1949T>C	A <sup>a</sup> , F <sup>b</sup> , K <sup>b</sup>	N.D.
17. CCG to CTG		21	P688L	2062C>T	O	0/0/57

<sup>a</sup>Hemizygous

<sup>b</sup>Homozygous variant: individuals were detected in the holoprosencephaly (HPE) families who were homozygous for the rare allele without clear correlation with HPE

gous variant in clinically normal individuals in families F and K, although, it was not readily detected in the analysis of the normal group of patients either because the band shift was minor under these gel conditions or that it was

indistinguishable from the other band shifts in exon 20. There were no other sequence changes noted in either the cDNA fragments or the genomic amplicons for exons 2, 3 or 4 in family A.

**Table 3.** Sequences of the primers used in this study

Primer	5' to 3' Sequence	Exon	Amplicon size (bp)	Annealing temperature (°C)
HLSLF1	gccagcattagagcactgcagcag	1 to 2	212	62
LSH200R	gcgggctcggctctgtagg	1 to 2		
065F	ccctacaagaccgagccccgc	2	195	63
066R	ctcaccctcagggtcccag	2		
111F	ggctgtatgtgaagagggttct	3	259	58
112R	gatccaagggtgatccagggtg	3		
067F	aggaaggggctgtcttagcag	4	210	61
068R	catgtgcagtacacaggggcagg	4		
114F	ggtttctacctgtcctctgctcc	5	178	58
115R	gcccagcacatgctgcattcc	5		
069F	gctgacaggaagcagatgacgtgg	6	174	63
070R	cgaccaacaaccaatcaacagc	6		
071F	gcaggctgagcttgggtctcc	7	214	61
072R	ccttactttgtccctgatgagg	7		
073F	gaggccaccctcctggttgagc	8	235	63
074R	accctgacctgacctgggctgc	8		
075F	agtgtcatggtgtctgcatccac	9	219	61
076R	cactttcggacctcaacccccagg	9		
160F	ggtctggagcggagccttgacagg	10	191	60
161R	ctcgggtccagtctctgtgagagg	10		
162F	cctgtttccatagagtgtggc	11	230	59
163R	gcagccttacctgcatttcat	11		
061F	gcatctgctgctgtaacacc	12	199	56
062R	gaggagtggcaagtgtgtgg	12		
077F	cctctctgctgctgtctctcc	13	200	56
078R	tagggcagggtggaggtgaggtgg	13		
204F	tgtcctctgcttccacattgc	14	120	58
205R	cgggacagggatgggctggctcc	14		
079F	ctgtgggaggttctgagaactgg	15	259	61
080R	ggatggacgggctgctgggacc	15		
121F	gtgtgtgctcagcacatgggc	16	172	58
122R	cgcgcgagtgtgctggccgacc	16		
081F	tgcttgtgtggcttgcctgag	17	216	56
082R	acaactgaatggctgagaccctcc	17		
145F	ttcactggaccaatctcgtgtcc	18	162	59
146R	acgcagtgtgtgagcagaaacc	18		
123F	ggtggaagggtgaggctctccc	19	167	58
124R	acagagcattgggttgagaacc	19		
125F	tgtgatggtgtcctctctctgg	20	239	60
126R	accaggctcaggacggtcc	20		
147F	gggtgtgttcagccttctctcc	21	160	60
148R	cacgctggaggtcagtgtctgg	21		
127F	aggctcaccacaagccagccaagc	22	268	60
128R	acagcgggaccaccagcaggtaggc	22		

Similarly, for family B, we identified only one missense mutation in exon 20, predicting a conservative L641V change, that was also seen in four additional pedigrees (E, F, K, P); this 1923T>G mutation segregated as a common polymorphism and not with the affected status in these families (data not shown). This mutation is also commonly seen in the control population.

In family C, we identified a potentially significant missense mutation (Table 2, number 14) in exon 20. This 1864G>A sequence variation was not observed initially in over 100 normal chromosomes and predicts the substitu-

tion of a Q for an R residue at a position conserved as the positively charged residues R or K in the family of *LS* genes. When we tested the ability of this mutation to complement the ability of a LS mutant yeast strain to grow on media lacking ergosterol, the mutant form of the yeast protein allowed the transformed strain to grow normally (data not shown). This demonstrates that the mutant enzyme retains catalytic function. Subsequent analysis of 53 additional normal control individuals revealed that this is a rare polymorphism, since it was eventually detected in two normal control individuals.

## SSCP analysis

We examined the entire coding region of the *LS* gene by means of SSCP in a total of 30 additional patients from 20 unrelated families with apparent autosomal recessive inheritance of HPE and present the results in Table 2. The primers used for the amplification are described in Table 3. We identified a total of 17 different sequence variations, which we number sequentially for clarity. The majority of these sequence variations were also detected in the normal control population (1, 2, 3, 4, 6, 8, 9, 10, 11, 12, 14, and 15). For the remainder, either the analysis of the pedigree transmission and/or the benign nature of the sequence change predicts no correlation with the HPE phenotype.

Seven of these sequence variations (3, 4, 7, 9–12) occur in non-coding intronic sequences. Only one of these intronic variants was not readily detected in the control population. This intronic variation, number 7, predicts the substitution of a T for C in the pyrimidine tract of the consensus splice acceptor. This change is likely to be functionally equivalent.

Of the ten remaining sequence variations, four are in the wobble position, and the codons are predicted not to alter the primary structure of the protein (1, 5, 6, 16). Of the six remaining missense mutations, one of these (number 14) has been excluded as a likely cause for HPE by functional studies in yeast; note also that we eventually detected this variant when we screened 53 additional normal controls. The sequence variant 13, the W576R substitution, is observed in an asymptomatic parent (family R) in the homozygous state. The identical mutation fails to segregate with HPE in a second family, K, where it is detected in a clinically normal mother and not either of her children with HPE. Therefore, this missense mutation is unlikely to exhibit a significant effect. The mutation 17 predicts a P688L change, which is observed in a normal parent and not his child with HPE. Although we did not detect this variant in the control population, the lack of a correlation with HPE makes this missense mutation unlikely to be responsible for HPE in this family.

The remaining three missense mutations are part of an interesting haplotype consisting of at least seven different sequence variations (1, 2, 4, 8, 9, 12, and 15), which are transmitted as a unit from parent to child in three different HPE families. All of these sequence variants are detectable as common variants in the normal control population. In one of these families, D, the haplotype is observed only in the phenotypically normal mother and not her son with HPE. This mother has mutations 1, 2, 8, and 9 by sequencing and the remaining changes seen in the other two families (E and F) were not detected using SSCP. Furthermore, in family E, the haplotype is present in only one of two affected sibs, suggesting that these changes are irrelevant to the HPE phenotype, despite the fact that this haplotype accounts for the final three missense mutations.

---

**Discussion**

The *Sonic Hedgehog* gene was the first of several HPE genes to be identified, and there is growing evidence for many additional genes (Roessler and Muenke 1998). We considered the *LS* gene as an excellent candidate since it was located within the *HPE1* minimal critical region in 21q22.3, and cholesterol is an essential adduct to the carboxy-terminus of the signaling domain of SHH-N. This modification serves to limit the diffusion of the patterning activity away from the site of synthesis. Furthermore, teratogenic agents, which inhibit the synthesis of cholesterol, are an effective means of producing HPE in experimental animals such as the rat (Roux et al. 1980; Lanoue et al. 1997). We further suspected that defects in *LS* would be at least as severe as those associated with SLO syndrome, which is caused by deficiency of the final step in cholesterol synthesis. This study did not provide any evidence for the *LS* gene in HPE pathogenesis, at least in live-born infants, which were the only samples available for analysis. Since we are aware that an incidence of HPE of 1:250 conceptuses (Matsunaga and Shiota 1977) differs from the 1:16,000 incidence at birth (Roach et al. 1975) by 50-fold, we cannot exclude a role for cholesterol-synthesis defects that might not be detectable in our sample of patients. Furthermore, in contrast to the SLO defect, complete loss of LS activity would severely limit sterol intermediates, suggesting that complete lack of LS activity might be lethal to the developing embryo. Although our premise was that LS defects would be recessively inherited, even modeling the mutations that we have identified as dominant is incompatible with the pedigree transmission and does not alter our conclusion that *LS* is not the *HPE1* gene.

Analysis of the functional domains of the Sonic Hedgehog protein (Hall et al. 1995, 1997) has led us to speculate that defective cholesterol modification of the protein due to intrinsic abnormalities could be one of several mechanisms causing HPE. In addition to structural mutations in SHH-N, we are attempting to identify a category of mutations in HPE patients that could affect cholesterol transferase activity. Experiments are in progress to substantiate this hypothesis. Therefore, despite the negative results in this report, there continues to be the possibility that examination of the cholesterol synthesis pathway may yet provide additional clues to HPE pathogenesis.

**Acknowledgements** We wish to thank the patients and their families for their participation in these studies. This work was supported by grants from the Swiss National Science Research Foundation 31.40500.94 and the European Union/OPES/PL 950302 (S.E.A), NIH grants AI41598 (S.P.T.M.), HD28732 and HD29862 (M.M), and by the Division of Intramural Research, National Human Genome Research Institute, NIH.

## References

- Antonarakis SE and the Nomenclature Working Group (1998) Recommendations for a nomenclature system for human gene mutations. *Hum Mutat* 11:1–5
- Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, Struhl K (1999) Current protocols in molecular biology. Wiley-Interscience, New York
- Baker CH, Matsuda SPT, Liu DR, Corey EJ (1995) Molecular cloning of the human gene encoding lanosterol synthase from a liver cDNA library. *Biochem Biophys Res Commun* 213:154–160
- Belloni E, Muenke M, Roessler E, Traverso G, Siegel-Bartelt J, Frumkin A, Mitchell HF, et al. (1996) Identification of *Sonic Hedgehog* as a candidate gene responsible for holoprosencephaly. *Nat Genet* 14:353–356
- Brown SA, Brown LY, Yu CC, Warburton D, Muenke M (1998a) *ZIC2*, a human homologue of *odd-paired*, is in the 13q32 critical deletion region and mutations are associated with holoprosencephaly (abstract). *Am J Hum Genet* 63[suppl]:A3
- Brown SA, Warburton D, Brown LY, Yu C, Roeder ER, Stengel-Rutkowski S, Hennekam RCM, et al. (1998b) Holoprosencephaly due to mutations in *ZIC2*, a homologue of *Drosophila odd-paired*. *Nat Genet* 20:180–183
- Cohen MM, Jr. (1989) Perspectives on holoprosencephaly: part I. Epidemiology, genetics, and syndromology. *Teratology* 40:211–235
- Corey EJ, Matsuda SPT, Bartel B (1994) Molecular cloning, characterization, and overexpression of *ERG7*, the *Saccharomyces cerevisiae* gene encoding lanosterol synthase. *Proc Natl Acad Sci U S A* 91:2211–2215
- Corey EJ, Matsuda SPT, Baker CH, Ting AY, Cheng H (1996) Molecular cloning of a *Shizosaccharomyces pombe* cDNA encoding lanosterol synthase and investigation of conserved tryptophan residues. *Biochem Biophys Res Commun* 219:327–331
- Cunniff C, Krantz IE, Moser A, Natowitz MR, Kelley R (1997) The clinical and biochemical spectrum of patients with Smith-Lemli-Opitz syndrome and abnormal cholesterol metabolism. *Am J Med Genet* 68:263–269
- DeMyer W, Zeman W, Palmer CG (1964) The face predicts the brain: diagnostic significance of median facial anomalies for holoprosencephaly (arhinencephaly). *Pediatrics* 34:256–263
- Fitzky BU, Witsch-Baumgartner M, Erdel M, Lee JN, Paik YK, Glossmann H, Utermann G, et al. (1998) Mutations in the delta-7-sterol reductase gene in patients with the Smith-Lemli-Opitz syndrome. *Proc Natl Acad Sci U S A* 95:8181–8186
- Frézal J, Schinzel A (1991) Report on the committee on clinical disorders, chromosome aberration, and uniparental disomy. *Cytogenet Cell Genet* 58:986–1052
- Gripp KW, Edwards MC, Mowat D, Meinecke P, Richieri-Costa A, Zackai EH, Elledge S, et al. (1998) Mutations in the transcription factor *TGIF* in holoprosencephaly (abstract). *Am J Hum Genet* 63[suppl]:A32
- Hall TMT, Porter JA, Beachy PA, Leahy DJ (1995) A potential catalytic site revealed by the 1.7-Å crystal structure of the amino-terminal signaling domain of Sonic hedgehog. *Nature* 378:212–216
- Hall TMT, Porter JA, Young KE, Koonin EV, Beachy PA, Leahy DJ (1997) Crystal structure of a hedgehog autoprocessing domain: homology between hedgehog and self-splicing proteins. *Cell* 91:85–97
- Kelley RI, Roessler E, Hennekam RC, Feldam GL, Kosaki K, Jones MC, Palumbos JC, et al. (1996) Holoprosencephaly in RSH/Smith-Lemli-Opitz syndrome: does abnormal cholesterol metabolism affect the function of *Sonic hedgehog*? *Am J Med Genet* 66:478–484
- Lanoue J, Denhart DB, Hinsdale ME, Maeda N, Tint GS, Sulik KK (1997) Limb, genital, CNS, and facial malformations result from gene/environment-induced cholesterol deficiency; further evidence for a link to Sonic hedgehog. *Am J Med Genet* 73:24–37
- Matsunaga E, Shiota K (1977) Holoprosencephaly in human embryos. Epidemiologic study of 150 cases. *Teratology* 16:261–272
- Ming JE, Muenke M (1998) Holoprosencephaly: from Homer to Hedgehog. *Clin Genet* 53:155–163
- Ming JE, Roessler E, Muenke M (1998a) Human developmental disorders and the Sonic hedgehog pathway. *Mol Med Today* 8:343–349
- Ming JE, Kaupas ME, Roessler E, Brunner HG, Nance WE, Stratton RF, Sujansky E, Bale SJ, Muenke M (1998b) Mutations of *PATCHED* in holoprosencephaly (abstract). *Am J Hum Genet* 63[suppl]:A27
- Moebius FF, Fitzky BU, Lee JN, Park YK, Glossmann H (1998) Molecular cloning and expression of the human delta7-sterol reductase. *Proc Natl Acad Sci U S A* 95:1899–1902
- Muenke M, Bone LJ, Mitchell HF, et al. (1995) Physical mapping of the holoprosencephaly critical region in 21q22.3, exclusion of *SIM2* as a candidate gene for holoprosencephaly, and mapping of *SIM2* to a region of chromosome 21 important for Down's syndrome. *Am J Hum Genet* 57:1074–1079
- Porter JA, Ekker SC, Park W-J, von Kessler DP, Young KE, Chen C-H, Ma Y, et al. (1996a) Hedgehog patterning activity: role of lipophilic modification mediated by the carboxy-terminal autoprocessing domain. *Cell* 86:21–34
- Porter JA, Young KE, Beachy PA (1996b) Cholesterol modification of hedgehog signaling proteins in animal development. *Science* 274:255–259
- Roach E, DeMeyer W, Conneally PM, Palmer C, Merritt AD (1975) Holoprosencephaly: birth data, genetic and demographic analysis of 30 families. *Birth Defects Original Ser* 11:294–313
- Roessler E, Muenke M (1998) Holoprosencephaly: a paradigm for the complex genetics of brain development. *J Inher Metab Dis* 21:481–497
- Roessler E, Belloni E, Gaudenz K, Jay P, Berta P, Scherer SW, Tsui L-C, et al. (1996) Mutations in the human *Sonic Hedgehog* gene cause holoprosencephaly. *Nat Genet* 14:357–360
- Roessler E, Belloni E, Gaudenz K, Vargas F, Scherer SW, Tsui L-C, Muenke M (1997a) Mutations in the C-terminal domain of Sonic Hedgehog cause holoprosencephaly. *Hum Mol Genet* 6:1847–1853
- Roessler E, Ward DE, Gaudenz K, Belloni E, Scherer SW, Donnai D, Siegel-Bartelt J, et al. (1997b) Cytogenetic rearrangements involving the loss of the Sonic Hedgehog gene at 7q36 cause holoprosencephaly. *Hum Genet* 100:172–181
- Roux C, Dupuis R, Horrath C, Talbot JN (1980) Teratogenic effect of an inhibitor of cholesterol synthesis (AY9944) in rats. Correlation with maternal cholesterolemia. *J Nutr* 110:2310–2312
- Soeda E, Osoegawa K, Atsuchi Y, Yamagata T, Simokawa T, Kishida H, Soeda E, Okano S, Chumakov J (1995) Cosmid assembly and anchoring to human chromosome 21. *Genomics* 25:73–84
- Wallis DE, Roessler E, Hehr U, Nanni L, Gillessen-Kaesbach G, Richieri-Costa A, Zackai E, Rommens J, Muenke M (1999) Missense mutations in the homeodomain of the human *SIX3* gene cause holoprosencephaly. *Nat Genet* 22:196–198
- Wassif CA, Masten C, Kachiliele-Linjewile S, Lin D, Linck LM, Connor WE, Steiner RD, Porter FD (1998) Mutations in the human sterol delta7-reductase gene at 11q12–13 cause Smith-Lemli-Opitz syndrome. *Am J Hum Genet* 63:55–62
- Waterham HR, Wijburg FA, Hennekam RC, Vrokon F, Pott-The BT, Dorland L, Duran M, Jira PE, Smeltink JA, Wevers RA, Wanders RJ (1998) Smith-Lemli-Opitz syndrome is caused by mutations in the 7-dehydrocholesterol reductase gene. *Am J Hum Genet* 63:329–338
- Young M, Chen H, Lalioti MD, Antonarakis SE (1996) The human lanosterol synthase gene maps to chromosome 21q22.3. *Hum Genet* 97:620–624