

## Study of resonances in exclusive $B$ decays to $\bar{D}^{(*)}D^{(*)}K$

B. Aubert,<sup>1</sup> M. Bona,<sup>1</sup> D. Boutigny,<sup>1</sup> Y. Karyotakis,<sup>1</sup> J. P. Lees,<sup>1</sup> V. Poireau,<sup>1</sup> X. Prudent,<sup>1</sup> V. Tisserand,<sup>1</sup> A. Zghiche,<sup>1</sup> J. Garra Tico,<sup>2</sup> E. Grauges,<sup>2</sup> L. Lopez,<sup>3</sup> A. Palano,<sup>3</sup> M. Pappagallo,<sup>3</sup> G. Eigen,<sup>4</sup> B. Stugu,<sup>4</sup> L. Sun,<sup>4</sup> G. S. Abrams,<sup>5</sup> M. Battaglia,<sup>5</sup> D. N. Brown,<sup>5</sup> J. Button-Shafer,<sup>5</sup> R. N. Cahn,<sup>5</sup> Y. Groyzman,<sup>5</sup> R. G. Jacobsen,<sup>5</sup> J. A. Kadyk,<sup>5</sup> L. T. Kerth,<sup>5</sup> Yu. G. Kolomoisky,<sup>5</sup> G. Kukartsev,<sup>5</sup> D. Lopes Pegna,<sup>5</sup> G. Lynch,<sup>5</sup> L. M. Mir,<sup>5</sup> T. J. Orimoto,<sup>5</sup> I. L. Osipenkov,<sup>5</sup> M. T. Ronan,<sup>5,\*</sup> K. Tackmann,<sup>5</sup> T. Tanabe,<sup>5</sup> W. A. Wenzel,<sup>5</sup> P. del Amo Sanchez,<sup>6</sup> C. M. Hawkes,<sup>6</sup> A. T. Watson,<sup>6</sup> H. Koch,<sup>7</sup> T. Schroeder,<sup>7</sup> D. Walker,<sup>8</sup> D. J. Asgeirsson,<sup>9</sup> T. Cuhadar-Donszelmann,<sup>9</sup> B. G. Fulsom,<sup>9</sup> C. Hearty,<sup>9</sup> T. S. Mattison,<sup>9</sup> J. A. McKenna,<sup>9</sup> A. Khan,<sup>10</sup> M. Saleem,<sup>10</sup> L. Teodorescu,<sup>10</sup> V. E. Blinov,<sup>11</sup> A. D. Bukin,<sup>11</sup> V. P. Druzhinin,<sup>11</sup> V. B. Golubev,<sup>11</sup> A. P. Onuchin,<sup>11</sup> S. I. Serednyakov,<sup>11</sup> Yu. I. Skovpen,<sup>11</sup> E. P. Solodov,<sup>11</sup> K. Yu. Todyshev,<sup>11</sup> M. Bondioli,<sup>12</sup> S. Curry,<sup>12</sup> I. Eschrich,<sup>12</sup> D. Kirkby,<sup>12</sup> A. J. Lankford,<sup>12</sup> P. Lund,<sup>12</sup> M. Mandelkern,<sup>12</sup> E. C. Martin,<sup>12</sup> D. P. Stoker,<sup>12</sup> S. Abachi,<sup>13</sup> C. Buchanan,<sup>13</sup> S. D. Foulkes,<sup>14</sup> J. W. Gary,<sup>14</sup> F. Liu,<sup>14</sup> O. Long,<sup>14</sup> B. C. Shen,<sup>14</sup> G. M. Vitug,<sup>14</sup> L. Zhang,<sup>14</sup> H. P. Paar,<sup>15</sup> S. Rahatlou,<sup>15</sup> V. Sharma,<sup>15</sup> J. W. Berryhill,<sup>16</sup> C. Campagnari,<sup>16</sup> A. Cunha,<sup>16</sup> B. Dahmes,<sup>16</sup> T. M. Hong,<sup>16</sup> D. Kovalskyi,<sup>16</sup> J. D. Richman,<sup>16</sup> T. W. Beck,<sup>17</sup> A. M. Eisner,<sup>17</sup> C. J. Flacco,<sup>17</sup> C. A. Heusch,<sup>17</sup> J. Kroseberg,<sup>17</sup> W. S. Lockman,<sup>17</sup> T. Schalk,<sup>17</sup> B. A. Schumm,<sup>17</sup> A. Seiden,<sup>17</sup> M. G. Wilson,<sup>17</sup> L. O. Winstrom,<sup>17</sup> E. Chen,<sup>18</sup> C. H. Cheng,<sup>18</sup> F. Fang,<sup>18</sup> D. G. Hitlin,<sup>18</sup> I. Narsky,<sup>18</sup> T. Piatenko,<sup>18</sup> F. C. Porter,<sup>18</sup> R. Andreassen,<sup>19</sup> G. Mancinelli,<sup>19</sup> B. T. Meadows,<sup>19</sup> K. Mishra,<sup>19</sup> M. D. Sokoloff,<sup>19</sup> F. Blanc,<sup>20</sup> P. C. Bloom,<sup>20</sup> S. Chen,<sup>20</sup> W. T. Ford,<sup>20</sup> J. F. Hirschauer,<sup>20</sup> A. Kreisel,<sup>20</sup> M. Nagel,<sup>20</sup> U. Nauenberg,<sup>20</sup> A. Olivas,<sup>20</sup> J. G. Smith,<sup>20</sup> K. A. Ulmer,<sup>20</sup> S. R. Wagner,<sup>20</sup> J. Zhang,<sup>20</sup> A. M. Gabareen,<sup>21</sup> A. Soffer,<sup>21,†</sup> W. H. Toki,<sup>21</sup> R. J. Wilson,<sup>21</sup> F. Winklmeier,<sup>21</sup> D. D. Altenburg,<sup>22</sup> E. Feltresi,<sup>22</sup> A. Hauke,<sup>22</sup> H. Jasper,<sup>22</sup> J. Merkel,<sup>22</sup> A. Petzold,<sup>22</sup> B. Spaan,<sup>22</sup> K. Wacker,<sup>22</sup> V. Klose,<sup>23</sup> M. J. Kobel,<sup>23</sup> H. M. Lacker,<sup>23</sup> W. F. Mader,<sup>23</sup> R. Nogowski,<sup>23</sup> J. Schubert,<sup>23</sup> K. R. Schubert,<sup>23</sup> R. Schwierz,<sup>23</sup> J. E. Sundermann,<sup>23</sup> A. Volk,<sup>23</sup> D. Bernard,<sup>24</sup> G. R. Bonneaud,<sup>24</sup> E. Latour,<sup>24</sup> V. Lombardo,<sup>24</sup> Ch. Thiebaut,<sup>24</sup> M. Verderi,<sup>24</sup> P. J. Clark,<sup>25</sup> W. Gradl,<sup>25</sup> F. Muheim,<sup>25</sup> S. Playfer,<sup>25</sup> A. I. Robertson,<sup>25</sup> J. E. Watson,<sup>25</sup> Y. Xie,<sup>25</sup> M. Andreotti,<sup>26</sup> D. Bettoni,<sup>26</sup> C. Bozzi,<sup>26</sup> R. Calabrese,<sup>26</sup> A. Cecchi,<sup>26</sup> G. Cibinetto,<sup>26</sup> P. Franchini,<sup>26</sup> E. Luppi,<sup>26</sup> M. Negrini,<sup>26</sup> A. Petrella,<sup>26</sup> L. Piemontese,<sup>26</sup> E. Prencipe,<sup>26</sup> V. Santoro,<sup>26</sup> F. Anulli,<sup>27</sup> R. Baldini-Ferrolì,<sup>27</sup> A. Calcaterra,<sup>27</sup> R. de Sangro,<sup>27</sup> G. Finocchiaro,<sup>27</sup> S. Pacetti,<sup>27</sup> P. Patteri,<sup>27</sup> I. M. Peruzzi,<sup>27,‡</sup> M. Piccolo,<sup>27</sup> M. Rama,<sup>27</sup> A. Zallo,<sup>27</sup> A. Buzzo,<sup>28</sup> R. Contri,<sup>28</sup> M. Lo Vetere,<sup>28</sup> M. M. Macri,<sup>28</sup> M. R. Monge,<sup>28</sup> S. Passaggio,<sup>28</sup> C. Patrignani,<sup>28</sup> E. Robutti,<sup>28</sup> A. Santroni,<sup>28</sup> S. Tosi,<sup>28</sup> K. S. Chaisanguanthum,<sup>29</sup> M. Morii,<sup>29</sup> J. Wu,<sup>29</sup> R. S. Dubitzky,<sup>30</sup> J. Marks,<sup>30</sup> S. Schenk,<sup>30</sup> U. Uwer,<sup>30</sup> D. J. Bard,<sup>31</sup> P. D. Dauncey,<sup>31</sup> R. L. Flack,<sup>31</sup> J. A. Nash,<sup>31</sup> W. Panduro Vazquez,<sup>31</sup> M. Tibbetts,<sup>31</sup> P. K. Behera,<sup>32</sup> X. Chai,<sup>32</sup> M. J. Charles,<sup>32</sup> U. Mallik,<sup>32</sup> J. Cochran,<sup>33</sup> H. B. Crawley,<sup>33</sup> L. Dong,<sup>33</sup> V. Eyges,<sup>33</sup> W. T. Meyer,<sup>33</sup> S. Prell,<sup>33</sup> E. I. Rosenberg,<sup>33</sup> A. E. Rubin,<sup>33</sup> Y. Y. Gao,<sup>34</sup> A. V. Gritsan,<sup>34</sup> Z. J. Guo,<sup>34</sup> C. K. Lae,<sup>34</sup> A. G. Denig,<sup>35</sup> M. Fritsch,<sup>35</sup> G. Schott,<sup>35</sup> N. Arnaud,<sup>36</sup> J. Béquilleux,<sup>36</sup> A. D'Orazio,<sup>36</sup> M. Davier,<sup>36</sup> G. Grosdidier,<sup>36</sup> A. Höcker,<sup>36</sup> V. Lepeltier,<sup>36</sup> F. Le Diberder,<sup>36</sup> A. M. Lutz,<sup>36</sup> S. Pruvot,<sup>36</sup> S. Rodier,<sup>36</sup> P. Roudeau,<sup>36</sup> M. H. Schune,<sup>36</sup> J. Serrano,<sup>36</sup> V. Sordini,<sup>36</sup> A. Stocchi,<sup>36</sup> W. F. Wang,<sup>36</sup> G. Wormser,<sup>36</sup> D. J. Lange,<sup>37</sup> D. M. Wright,<sup>37</sup> I. Bingham,<sup>38</sup> C. A. Chavez,<sup>38</sup> J. R. Fry,<sup>38</sup> E. Gabathuler,<sup>38</sup> R. Gamet,<sup>38</sup> D. E. Hutchcroft,<sup>38</sup> D. J. Payne,<sup>38</sup> K. C. Schofield,<sup>38</sup> C. Touramanis,<sup>38</sup> A. J. Bevan,<sup>39</sup> K. A. George,<sup>39</sup> F. Di Lodovico,<sup>39</sup> R. Sacco,<sup>39</sup> G. Cowan,<sup>40</sup> H. U. Flaecher,<sup>40</sup> D. A. Hopkins,<sup>40</sup> S. Paramesvaran,<sup>40</sup> F. Salvatore,<sup>40</sup> A. C. Wren,<sup>40</sup> D. N. Brown,<sup>41</sup> C. L. Davis,<sup>41</sup> J. Allison,<sup>42</sup> D. Bailey,<sup>42</sup> N. R. Barlow,<sup>42</sup> R. J. Barlow,<sup>42</sup> Y. M. Chia,<sup>42</sup> C. L. Edgar,<sup>42</sup> G. D. Lafferty,<sup>42</sup> T. J. West,<sup>42</sup> J. I. Yi,<sup>42</sup> J. Anderson,<sup>43</sup> C. Chen,<sup>43</sup> A. Jawahery,<sup>43</sup> D. A. Roberts,<sup>43</sup> G. Simi,<sup>43</sup> J. M. Tuggle,<sup>43</sup> G. Blaylock,<sup>44</sup> C. Dallapiccola,<sup>44</sup> S. S. Hertzbach,<sup>44</sup> X. Li,<sup>44</sup> T. B. Moore,<sup>44</sup> E. Salvati,<sup>44</sup> S. Saremi,<sup>44</sup> R. Cowan,<sup>45</sup> D. Dujmic,<sup>45</sup> P. H. Fisher,<sup>45</sup> K. Koeneke,<sup>45</sup> G. Sciolla,<sup>45</sup> M. Spitznagel,<sup>45</sup> F. Taylor,<sup>45</sup> R. K. Yamamoto,<sup>45</sup> M. Zhao,<sup>45</sup> Y. Zheng,<sup>45</sup> S. E. Mclachlin,<sup>46,\*</sup> P. M. Patel,<sup>46</sup> S. H. Robertson,<sup>46</sup> A. Lazzaro,<sup>47</sup> F. Palombo,<sup>47</sup> J. M. Bauer,<sup>48</sup> L. Cremaldi,<sup>48</sup> V. Eschenburg,<sup>48</sup> R. Godang,<sup>48</sup> R. Kroeger,<sup>48</sup> D. A. Sanders,<sup>48</sup> D. J. Summers,<sup>48</sup> H. W. Zhao,<sup>48</sup> S. Brunet,<sup>49</sup> D. Côté,<sup>49</sup> M. Simard,<sup>49</sup> P. Taras,<sup>49</sup> F. B. Viaud,<sup>49</sup> H. Nicholson,<sup>50</sup> G. De Nardo,<sup>51</sup> F. Fabozzi,<sup>51,§</sup> L. Lista,<sup>51</sup> D. Monorchio,<sup>51</sup> C. Sciacca,<sup>51</sup> M. A. Baak,<sup>52</sup> G. Raven,<sup>52</sup> H. L. Snoek,<sup>52</sup> C. P. Jessop,<sup>53</sup> K. J. Knoepfel,<sup>53</sup> J. M. LoSecco,<sup>53</sup> G. Benelli,<sup>54</sup> L. A. Corwin,<sup>54</sup> K. Honscheid,<sup>54</sup> H. Kagan,<sup>54</sup> R. Kass,<sup>54</sup> J. P. Morris,<sup>54</sup> A. M. Rahimi,<sup>54</sup> J. J. Regensburger,<sup>54</sup> S. J. Sekula,<sup>54</sup> Q. K. Wong,<sup>54</sup> N. L. Blount,<sup>55</sup> J. Brau,<sup>55</sup> R. Frey,<sup>55</sup> O. Igonkina,<sup>55</sup> J. A. Kolb,<sup>55</sup> M. Lu,<sup>55</sup> R. Rahmat,<sup>55</sup> N. B. Sinev,<sup>55</sup> D. Strom,<sup>55</sup> J. Strube,<sup>55</sup> E. Torrence,<sup>55</sup> N. Gagliardi,<sup>56</sup> A. Gaz,<sup>56</sup> M. Margoni,<sup>56</sup> M. Morandin,<sup>56</sup> A. Pompili,<sup>56</sup> M. Posocco,<sup>56</sup> M. Rotondo,<sup>56</sup> F. Simonetto,<sup>56</sup> R. Stroili,<sup>56</sup> C. Voci,<sup>56</sup> E. Ben-Haim,<sup>57</sup> H. Briand,<sup>57</sup> G. Calderini,<sup>57</sup> J. Chauveau,<sup>57</sup> P. David,<sup>57</sup> L. Del Buono,<sup>57</sup> Ch. de la Vaissière,<sup>57</sup> O. Hamon,<sup>57</sup> Ph. Leruste,<sup>57</sup> J. Malcès,<sup>57</sup> J. Ocariz,<sup>57</sup> A. Perez,<sup>57</sup> J. Prendki,<sup>57</sup> L. Gladney,<sup>58</sup> M. Biasini,<sup>59</sup> R. Covarelli,<sup>59</sup> E. Manoni,<sup>59</sup> C. Angelini,<sup>60</sup> G. Batignani,<sup>60</sup> S. Bettarini,<sup>60</sup> M. Carpinelli,<sup>60</sup> R. Cenci,<sup>60</sup> A. Cervelli,<sup>60</sup> F. Forti,<sup>60</sup> M. A. Giorgi,<sup>60</sup> A. Lusiani,<sup>60</sup> G. Marchiori,<sup>60</sup>

M. A. Mazur,<sup>60</sup> M. Morganti,<sup>60</sup> N. Neri,<sup>60</sup> E. Paoloni,<sup>60</sup> G. Rizzo,<sup>60</sup> J. J. Walsh,<sup>60</sup> J. Biesiada,<sup>61</sup> P. Elmer,<sup>61</sup> Y. P. Lau,<sup>61</sup> C. Lu,<sup>61</sup> J. Olsen,<sup>61</sup> A. J. S. Smith,<sup>61</sup> A. V. Telnov,<sup>61</sup> E. Baracchini,<sup>62</sup> F. Bellini,<sup>62</sup> G. Cavoto,<sup>62</sup> D. del Re,<sup>62</sup> E. Di Marco,<sup>62</sup> R. Faccini,<sup>62</sup> F. Ferrarotto,<sup>62</sup> F. Ferroni,<sup>62</sup> M. Gaspero,<sup>62</sup> P. D. Jackson,<sup>62</sup> L. Li Gioi,<sup>62</sup> M. A. Mazzoni,<sup>62</sup> S. Morganti,<sup>62</sup> G. Piredda,<sup>62</sup> F. Polci,<sup>62</sup> F. Renga,<sup>62</sup> C. Voena,<sup>62</sup> M. Ebert,<sup>63</sup> T. Hartmann,<sup>63</sup> H. Schröder,<sup>63</sup> R. Waldi,<sup>63</sup> T. Adye,<sup>64</sup> G. Castelli,<sup>64</sup> B. Franek,<sup>64</sup> E. O. Olaiya,<sup>64</sup> W. Roethel,<sup>64</sup> F. F. Wilson,<sup>64</sup> S. Emery,<sup>65</sup> M. Escalier,<sup>65</sup> A. Gaidot,<sup>65</sup> S. F. Ganzhur,<sup>65</sup> G. Hamel de Monchenault,<sup>65</sup> W. Kozanecki,<sup>65</sup> G. Vasseur,<sup>65</sup> Ch. Yèche,<sup>65</sup> M. Zito,<sup>65</sup> X. R. Chen,<sup>66</sup> H. Liu,<sup>66</sup> W. Park,<sup>66</sup> M. V. Purohit,<sup>66</sup> R. M. White,<sup>66</sup> J. R. Wilson,<sup>66</sup> M. T. Allen,<sup>67</sup> D. Aston,<sup>67</sup> R. Bartoldus,<sup>67</sup> P. Bechtel,<sup>67</sup> R. Claus,<sup>67</sup> J. P. Coleman,<sup>67</sup> M. R. Convery,<sup>67</sup> J. C. Dingfelder,<sup>67</sup> J. Dorfan,<sup>67</sup> G. P. Dubois-Felsmann,<sup>67</sup> W. Dunwoodie,<sup>67</sup> R. C. Field,<sup>67</sup> T. Glanzman,<sup>67</sup> S. J. Gowdy,<sup>67</sup> M. T. Graham,<sup>67</sup> P. Grenier,<sup>67</sup> C. Hast,<sup>67</sup> W. R. Innes,<sup>67</sup> J. Kaminski,<sup>67</sup> M. H. Kelsey,<sup>67</sup> H. Kim,<sup>67</sup> P. Kim,<sup>67</sup> M. L. Kocian,<sup>67</sup> D. W. G. S. Leith,<sup>67</sup> S. Li,<sup>67</sup> S. Luitz,<sup>67</sup> V. Luth,<sup>67</sup> H. L. Lynch,<sup>67</sup> D. B. MacFarlane,<sup>67</sup> H. Marsiske,<sup>67</sup> R. Messner,<sup>67</sup> D. R. Muller,<sup>67</sup> C. P. O'Grady,<sup>67</sup> I. Ofte,<sup>67</sup> A. Perazzo,<sup>67</sup> M. Perl,<sup>67</sup> T. Pulliam,<sup>67</sup> B. N. Ratcliff,<sup>67</sup> A. Roodman,<sup>67</sup> A. A. Salnikov,<sup>67</sup> R. H. Schindler,<sup>67</sup> J. Schwiening,<sup>67</sup> A. Snyder,<sup>67</sup> D. Su,<sup>67</sup> M. K. Sullivan,<sup>67</sup> K. Suzuki,<sup>67</sup> S. K. Swain,<sup>67</sup> J. M. Thompson,<sup>67</sup> J. Va'vra,<sup>67</sup> A. P. Wagner,<sup>67</sup> M. Weaver,<sup>67</sup> W. J. Wisniewski,<sup>67</sup> M. Wittgen,<sup>67</sup> D. H. Wright,<sup>67</sup> A. K. Yarritu,<sup>67</sup> K. Yi,<sup>67</sup> C. C. Young,<sup>67</sup> V. Ziegler,<sup>67</sup> P. R. Burchat,<sup>68</sup> A. J. Edwards,<sup>68</sup> S. A. Majewski,<sup>68</sup> T. S. Miyashita,<sup>68</sup> B. A. Petersen,<sup>68</sup> L. Wilden,<sup>68</sup> S. Ahmed,<sup>69</sup> M. S. Alam,<sup>69</sup> R. Bula,<sup>69</sup> J. A. Ernst,<sup>69</sup> V. Jain,<sup>69</sup> B. Pan,<sup>69</sup> M. A. Saeed,<sup>69</sup> F. R. Wappler,<sup>69</sup> S. B. Zain,<sup>69</sup> M. Krishnamurthy,<sup>70</sup> S. M. Spanier,<sup>70</sup> R. Eckmann,<sup>71</sup> J. L. Ritchie,<sup>71</sup> A. M. Ruland,<sup>71</sup> C. J. Schilling,<sup>71</sup> R. F. Schwitters,<sup>71</sup> J. M. Izen,<sup>72</sup> X. C. Lou,<sup>72</sup> S. Ye,<sup>72</sup> F. Bianchi,<sup>73</sup> F. Gallo,<sup>73</sup> D. Gamba,<sup>73</sup> M. Pelliccioni,<sup>73</sup> M. Bomben,<sup>74</sup> L. Bosisio,<sup>74</sup> C. Cartaro,<sup>74</sup> F. Cossutti,<sup>74</sup> G. Della Ricca,<sup>74</sup> L. Lanceri,<sup>74</sup> L. Vitale,<sup>74</sup> V. Azzolini,<sup>75</sup> N. Lopez-March,<sup>75</sup> F. Martinez-Vidal,<sup>75</sup> D. A. Milanes,<sup>75</sup> A. Oyanguren,<sup>75</sup> J. Albert,<sup>76</sup> Sw. Banerjee,<sup>76</sup> B. Bhuyan,<sup>76</sup> K. Hamano,<sup>76</sup> R. Kowalewski,<sup>76</sup> I. M. Nugent,<sup>76</sup> J. M. Roney,<sup>76</sup> R. J. Sobie,<sup>76</sup> P. F. Harrison,<sup>77</sup> J. Ilic,<sup>77</sup> T. E. Latham,<sup>77</sup> G. B. Mohanty,<sup>77</sup> H. R. Band,<sup>78</sup> X. Chen,<sup>78</sup> S. Dasu,<sup>78</sup> K. T. Flood,<sup>78</sup> J. J. Hollar,<sup>78</sup> P. E. Kutter,<sup>78</sup> Y. Pan,<sup>78</sup> M. Pierini,<sup>78</sup> R. Prepost,<sup>78</sup> S. L. Wu,<sup>78</sup> and H. Neal<sup>79</sup>

(BABAR Collaboration)

<sup>1</sup>Laboratoire de Physique des Particules, IN2P3/CNRS et Université de Savoie, F-74941 Annecy-Le-Vieux, France

<sup>2</sup>Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

<sup>3</sup>Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

<sup>4</sup>University of Bergen, Institute of Physics, N-5007 Bergen, Norway

<sup>5</sup>Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

<sup>6</sup>University of Birmingham, Birmingham, B15 2TT, United Kingdom

<sup>7</sup>Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

<sup>8</sup>University of Bristol, Bristol BS8 1TL, United Kingdom

<sup>9</sup>University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

<sup>10</sup>Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

<sup>11</sup>Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

<sup>12</sup>University of California at Irvine, Irvine, California 92697, USA

<sup>13</sup>University of California at Los Angeles, Los Angeles, California 90024, USA

<sup>14</sup>University of California at Riverside, Riverside, California 92521, USA

<sup>15</sup>University of California at San Diego, La Jolla, California 92093, USA

<sup>16</sup>University of California at Santa Barbara, Santa Barbara, California 93106, USA

<sup>17</sup>University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

<sup>18</sup>California Institute of Technology, Pasadena, California 91125, USA

<sup>19</sup>University of Cincinnati, Cincinnati, Ohio 45221, USA

<sup>20</sup>University of Colorado, Boulder, Colorado 80309, USA

<sup>21</sup>Colorado State University, Fort Collins, Colorado 80523, USA

<sup>22</sup>Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany

<sup>23</sup>Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

<sup>24</sup>Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France

<sup>25</sup>University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

<sup>26</sup>Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy

<sup>27</sup>Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy

<sup>28</sup>Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy

<sup>29</sup>Harvard University, Cambridge, Massachusetts 02138, USA

<sup>30</sup>Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany

<sup>31</sup>Imperial College London, London, SW7 2AZ, United Kingdom

- <sup>32</sup>University of Iowa, Iowa City, Iowa 52242, USA  
<sup>33</sup>Iowa State University, Ames, Iowa 50011-3160, USA  
<sup>34</sup>Johns Hopkins University, Baltimore, Maryland 21218, USA  
<sup>35</sup>Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany  
<sup>36</sup>Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B. P. 34, F-91898 ORSAY Cedex, France  
<sup>37</sup>Lawrence Livermore National Laboratory, Livermore, California 94550, USA  
<sup>38</sup>University of Liverpool, Liverpool L69 7ZE, United Kingdom  
<sup>39</sup>Queen Mary, University of London, E1 4NS, United Kingdom  
<sup>40</sup>University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom  
<sup>41</sup>University of Louisville, Louisville, Kentucky 40292, USA  
<sup>42</sup>University of Manchester, Manchester M13 9PL, United Kingdom  
<sup>43</sup>University of Maryland, College Park, Maryland 20742, USA  
<sup>44</sup>University of Massachusetts, Amherst, Massachusetts 01003, USA  
<sup>45</sup>Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA  
<sup>46</sup>McGill University, Montréal, Québec, Canada H3A 2T8  
<sup>47</sup>Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy  
<sup>48</sup>University of Mississippi, University, Mississippi 38677, USA  
<sup>49</sup>Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7  
<sup>50</sup>Mount Holyoke College, South Hadley, Massachusetts 01075, USA  
<sup>51</sup>Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy  
<sup>52</sup>NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands  
<sup>53</sup>University of Notre Dame, Notre Dame, Indiana 46556, USA  
<sup>54</sup>Ohio State University, Columbus, Ohio 43210, USA  
<sup>55</sup>University of Oregon, Eugene, Oregon 97403, USA  
<sup>56</sup>Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy  
<sup>57</sup>Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris6, Université Denis Diderot-Paris7, F-75252 Paris, France  
<sup>58</sup>University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA  
<sup>59</sup>Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy  
<sup>60</sup>Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy  
<sup>61</sup>Princeton University, Princeton, New Jersey 08544, USA  
<sup>62</sup>Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy  
<sup>63</sup>Universität Rostock, D-18051 Rostock, Germany  
<sup>64</sup>Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom  
<sup>65</sup>DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France  
<sup>66</sup>University of South Carolina, Columbia, South Carolina 29208, USA  
<sup>67</sup>Stanford Linear Accelerator Center, Stanford, California 94309, USA  
<sup>68</sup>Stanford University, Stanford, California 94305-4060, USA  
<sup>69</sup>State University of New York, Albany, New York 12222, USA  
<sup>70</sup>University of Tennessee, Knoxville, Tennessee 37996, USA  
<sup>71</sup>University of Texas at Austin, Austin, Texas 78712, USA  
<sup>72</sup>University of Texas at Dallas, Richardson, Texas 75083, USA  
<sup>73</sup>Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy  
<sup>74</sup>Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy  
<sup>75</sup>IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain  
<sup>76</sup>University of Victoria, Victoria, British Columbia, Canada V8W 3P6  
<sup>77</sup>Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom  
<sup>78</sup>University of Wisconsin, Madison, Wisconsin 53706, USA  
<sup>79</sup>Yale University, New Haven, Connecticut 06511, USA

(Received 11 August 2007; published 7 January 2008)

We present a study of resonances in exclusive decays of  $B$  mesons to  $\bar{D}^{(*)}D^{(*)}K$ . We report the observation of the decays  $B \rightarrow \bar{D}^{(*)}D_{s1}^{+}(2536)$  where the  $D_{s1}^{+}(2536)$  is reconstructed in the  $D^{*0}K^{+}$  and  $D^{*+}K_{S}^{0}$  decay channels. We report also the observation of the decays  $B \rightarrow \psi(3770)K$  where the  $\psi(3770)$

\*Deceased

†Now at Tel Aviv University, Tel Aviv, 69978, Israel

‡Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

§Also with Università della Basilicata, Potenza, Italy

|| Also with Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

decays to  $\bar{D}^0 D^0$  and  $D^- D^+$ . In addition, we present the observation of an enhancement for the  $\bar{D}^{*0} D^0$  invariant mass in the decays  $B \rightarrow \bar{D}^{*0} D^0 K$ , at a mass of  $(3875.1_{-0.5}^{+0.7} \pm 0.5)$  MeV/ $c^2$  with a width of  $(3.0_{-1.4}^{+1.9} \pm 0.9)$  MeV (the first errors are statistical and the second are systematic). Branching fractions and spin studies are shown for the three resonances. The results are based on 347 fb $^{-1}$  of data collected with the *BABAR* detector at the PEP-II *B* factory.

DOI: [10.1103/PhysRevD.77.011102](https://doi.org/10.1103/PhysRevD.77.011102)

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

In this article, we study the production of  $D_{s1}^+(2536)$ ,  $\psi(3770)$  and  $X(3872)$  resonances in decays of charged and neutral *B* mesons to  $\bar{D}^{(*)} D^{(*)} K$ . Here,  $D^{(*)}$  is either a  $D^0$ ,  $D^{*0}$ ,  $D^+$  or  $D^{*+}$ ,  $\bar{D}^{(*)}$  is the charge conjugate of  $D^{(*)}$  and *K* is either a  $K^+$  or a  $K_S^0$ . Both  $\bar{D}^{(*)}$  and  $D^{(*)}$  are fully reconstructed. Charge conjugate reactions are assumed throughout this article.

The  $D_{s1}^+(2536)$  resonance is the narrow *P*-wave  $D_s^+$  meson with  $J^P = 1^+$  assignment strongly favored. It can be produced in  $B \rightarrow \bar{D}^{(*)} D_{s1}^+(2536)$  decays and should decay dominantly to  $D^{*0} K^+$  and  $D^{*+} K^0$  [1]. Evidence for  $D_{s1}^+(2536)$  production in *B* decays was found by *BABAR* in the sum of all  $\bar{D}^{(*)} D^{*0} K^+$  final states [2] and more recently in the decay  $B^0 \rightarrow D^{*-} D^{*+} K_S^0$  [3]. For most of the  $B \rightarrow \bar{D}^{(*)} D_{s1}^+(2536)$  modes, only limits have been placed on the branching fractions [2]. We report herein branching fraction measurements of  $B \rightarrow \bar{D}^{(*)} D_{s1}^+(2536)$  decays, through a comprehensive study of both  $\bar{D}^{(*)} D^{*0} K^+$  and  $\bar{D}^{(*)} D^{*+} K_S^0$  final states.

The  $\psi(3770)$  meson is a charmonium state with  $J^P = 1^-$ , with a mass just above the open charm threshold. This meson is thought to be an admixture of the *D*-wave and *S*-wave of the angular momentum eigenstates of  $c\bar{c}$  system. Study of this state in *B* decays and branching fraction measurements could provide more information on the structure of the  $\psi(3770)$  wave function. This resonance decays dominantly to  $\bar{D}^0 D^0$  and  $D^- D^+$  [1], and was observed in *B* meson decays by the Belle experiment [4]. We present a study of the  $\bar{D}D$  mass distribution in  $\bar{D}DK$  events.

The  $X(3872)$  resonance was discovered by Belle in the invariant mass distribution of  $J/\psi\pi^+\pi^-$  produced in  $B \rightarrow J/\psi\pi^+\pi^-K$  decays, and was thereafter confirmed by *BABAR*, D0 and CDF [5]. This new meson has a mass of  $3871.2 \pm 0.5$  MeV/ $c^2$  and a natural width less than 2.3 MeV at 90% confidence level (C.L.). At present, the quantum numbers compatible with the observations are  $J^{PC} = 1^{++}$  or  $2^{-+}$  [6]. Recently, Belle showed an excess of events in the  $\bar{D}^0 D^0 \pi^0$  invariant mass in the channel  $B \rightarrow \bar{D}^0 D^0 \pi^0 K$ , with a mass of  $3875.2 \pm 0.7_{-1.6}^{+0.3} \pm 0.8$  MeV/ $c^2$  [7] (where the third error is due to the uncertainty in the neutral *D* mass). The  $X(3872)$  is probably not a charmonium state, given its measured mass and width, and several alternative interpretations have been proposed:  $\bar{D}^{*0} D^0$  molecule, tetraquark state, hybrid or gluonium states [8]. We present a search for  $X(3872)$  decays to  $\bar{D}^{*0} D^0$ .

The measurements reported here use 347 fb $^{-1}$  of data, corresponding to  $(383 \pm 4) \times 10^6$   $B\bar{B}$  pairs, collected at the  $Y(4S)$  resonance with the *BABAR* detector at the PEP-II *B* factory. The *BABAR* detector is described in detail elsewhere [9]. We use a Monte Carlo (MC) simulation based on GEANT4 [10] to study the relevant backgrounds and estimate the selection efficiencies.

The  $B^0$  and  $B^+$  mesons are reconstructed in a sample of hadronic events in the 22 possible  $\bar{D}^{(*)} D^{(*)} K$  modes. The selection criteria are optimized for each final state by maximizing the significance  $S/\sqrt{S+B}$ , where *S* and *B* refer to the expected number of signal and background events, based on MC simulation. The  $K_S^0$  candidates are reconstructed from two oppositely charged tracks consistent with coming from a common vertex and having an invariant mass within  $\pm 9.5$  MeV/ $c^2$  of the nominal  $K_S^0$  mass. For some channels, depending on the background level, we apply a requirement on the displacement of the  $K_S^0$  vertex in the plane transverse to the beam axis of at least 0.2 cm. The  $\pi^0$  candidates are reconstructed from pairs of photons with energies  $E_\gamma > 30$  MeV in the laboratory frame that have an invariant mass of  $115 < m_{\gamma\gamma} < 150$  MeV/ $c^2$ . We reconstruct *D* mesons in the modes  $D^0 \rightarrow K^- \pi^+$ ,  $K^- \pi^+ \pi^0$ ,  $K^- \pi^+ \pi^- \pi^+$ , and  $D^+ \rightarrow K^- \pi^+ \pi^+$ . The *K* and  $\pi$  tracks are required to originate from a common vertex. Charged kaon identification, based on the Cherenkov angle and the  $dE/dx$  measurements, is used for  $K^+$  from *B* decays and from most *D* decays. The invariant masses of the *D* candidates are required to be within  $\pm 2\sigma$  of the measured *D* mass. The measured *D* mass resolution,  $\sigma_D$ , is 13 MeV/ $c^2$  for  $D^0 \rightarrow K^- \pi^+ \pi^0$  and varies from 5.5 to 7 MeV/ $c^2$  for the other modes. The  $D^*$  candidates are reconstructed in the decay modes  $D^{*+} \rightarrow D^0 \pi^+$ ,  $D^{*+} \rightarrow D^+ \pi^0$ ,  $D^{*0} \rightarrow D^0 \pi^0$ , and  $D^{*0} \rightarrow D^0 \gamma$ . The  $\pi^0$  and the  $\pi^+$  must have momentum below 450 MeV/ $c$  in the  $Y(4S)$  rest frame, while the  $\gamma$  energy in the laboratory frame must be greater than 100 MeV. The mass difference between the  $D^*$  and *D* candidates is required to be within 3 MeV/ $c^2$  of the nominal value [1] for  $D^{*+}$  decays (4 MeV/ $c^2$  and 10 MeV/ $c^2$  for  $D^{*0} \rightarrow D^0 \pi^0$  and  $D^{*0} \rightarrow D^0 \gamma$ , respectively). The mode  $D^{*+} \rightarrow D^+ \pi^0$  is used only in the reconstruction of decays  $B^0 \rightarrow D^{*-} D^{*+} K_S^0$  and  $B^+ \rightarrow D^{*-} D^{*+} K^+$ .

*B* candidates are reconstructed by combining a  $\bar{D}^{(*)}$ , a  $D^{(*)}$  and a *K* candidate. For most of the modes involving two  $D^0$  mesons, at least one of them is required to decay to  $K^- \pi^+$ . During the optimization process, we remove *D*

decay modes if the significance for a particular  $B$  decay mode improves. In practice, only  $\bar{D}^{*0}D^{*+}K_S^0$  decays benefit from this optimization, for which we remove the combination with the first  $D^0$  meson of the decay chain going to  $K^- \pi^+$  and the second going to  $K^- \pi^+ \pi^- \pi^+$  (and vice-versa). A mass-constrained kinematic fit is applied to the intermediate particles ( $D^{*0}$ ,  $D^{*+}$ ,  $D^0$ ,  $D^+$ ,  $K_S^0$ ,  $\pi^0$ ) to improve their momentum resolution. For the  $D^0$  and the  $D^* - D^0$  mass differences, we use the recent CLEO measurement of the  $D^0$  mass [11]. To suppress the  $e^+e^- \rightarrow q\bar{q}$  ( $q = u, d, s, \text{ and } c$ ) continuum background, we perform a selection based on the ratio of the second to zeroth Fox-Wolfram moments of the event [12] and on the cosine of the angle between the thrust axis of the candidate decay and the thrust axis of the rest of the event. Signal events have  $m_{ES} = \sqrt{s/4 - p_B^{*2}}$ , where  $p_B^*$  is the center-of-mass momentum of the  $B$  candidate, compatible with the known  $B$  meson mass, and a difference between the candidate energy and the beam energy in the center-of-mass,  $\Delta E$ , compatible with 0. On average we have about 1.8 signal  $B$  candidates per event. If more than one candidate is selected in an event, we retain the one with the smallest  $|\Delta E|$ . In the final selection, we require  $|\Delta E|$  to be less than  $n\sigma_{\Delta E}$ , where the resolution  $\sigma_{\Delta E}$  varies between 7 and 14 MeV and  $n$  is determined for each mode by the optimization procedure ( $n$  ranges from 1 to 4). For each mode we define a  $B$  signal region  $m_{ES}^{\min} < m_{ES} < m_{ES}^{\max}$ , where  $m_{ES}^{\min}$  ranges from 5.268 to 5.277 GeV/ $c^2$ , and  $m_{ES}^{\max}$  from 5.284 to 5.290 GeV/ $c^2$ , and a control region,  $5.20 < m_{ES} < 5.26$  GeV/ $c^2$ . The signal purity obtained in the signal region ranges from 17% to 77%, depending on the mode.

We consider several sources of systematic errors. From the difference between data and Monte Carlo efficiencies we derive systematic errors of 0.5% per charged track, 2.2% per soft pion from  $D^*$  decays, 2.5% per  $K_S^0$ , 2% per  $K^+$ , 3% per  $\pi^0$  and 2% per single photon. As an example, the particle identification efficiency for  $K^+$  is measured using a  $D^{*+} \rightarrow D^0 \pi^+$  data control sample with  $D^0 \rightarrow K^- \pi^+$ . Other sources of systematic error are also taken into account: limited MC statistics (1%–3%), the estimate of the total number of  $B$  mesons in the data sample (1.1%), uncertainties on the  $D^{(*)}$  and  $K_S^0$  branching fractions (3%–8%) and uncertainties on the  $D^{(*)}$  and  $K$  masses (0.5%–6%). In addition, there are uncertainties in the fit procedure for the different resonances: when fixing a parameter, we repeat the fit varying the parameter by  $\pm 1\sigma$  of its error (where  $\sigma$  is the 68% C.L. standard deviation); these uncertainties include also variation of the background parametrization. Using the  $m_{ES}$  control region, we check that the combinatorial background events do not contain any significant  $D_{s1}^+(2536)$ ,  $\psi(3770)$  or  $X(3872)$  signals or additional peaking structures. Furthermore, for the mass and width measurements, we include effects from the energy loss in the tracking system, from the uncertainties

on the magnetic field and from the calibration and background of the calorimeter.

We search for  $D_{s1}^+(2536)$  decaying to  $D^{*+}K_S^0$  and  $D^{*0}K^+$  using  $B^0 \rightarrow \bar{D}^{(*)-}D^{*0}K^+$ ,  $\bar{D}^{(*)-}D^{*+}K_S^0$  and  $B^+ \rightarrow \bar{D}^{(*)0}D^{*0}K^+$ ,  $\bar{D}^{(*)0}D^{*+}K_S^0$  candidates. We show in Fig. 1(a) the  $D^{*+}K_S^0$  and  $D^{*0}K^+$  invariant mass distribution for the sum of the eight  $B$  modes. The overlaid curve is the result of a unbinned extended maximum likelihood fit. Since the  $D_{s1}^+(2536)$  meson is narrow ( $\Gamma_{D_{s1}^+(2536)} < 2.3$  MeV at 90% C.L. [1]), the invariant mass peak is described by the convolution of a nonrelativistic Breit-Wigner with a Gaussian function (called a Voigtian function) and the background by a threshold function  $a(m - m_0)^b \times e^{c(m - m_0)}$ , where  $m$  is the  $D^*K$  invariant mass,  $m_0$  is the sum of the  $D^*$  and  $K$  meson masses, and  $a$ ,  $b$ , and  $c$  are parameters of the fit. To take into account the large tails in the reconstruction of  $D_{s1}^+(2536) \rightarrow D^{*0}K^+$  decays, the probability density function (PDF) for the  $D^{*0}K^+$  modes is constructed from a sum of a Voigtian function and a Gaussian function, with a common mean. The  $D_{s1}^+(2536)$  mass and yield are floating parameters in the fit. The natural width of the  $D_{s1}^+(2536)$  is fixed to 1 MeV and varied from 0.1 to 2.0 MeV to estimate systematic errors. The other parameters (the second Gaussian function and the relative contribution of the Voigtian and the Gaussian function) are taken from the simulation. A significant signal is observed in each of the modes separately (see Table I). A fit to the eight  $B$  modes gives  $182 \pm 19$  events in the peak with a statistical significance of  $11.8\sigma$ . We compute an estimate of the statistical significance calculating  $\text{PROB}(2(L_0 - L_{\text{signal}}), N_{\text{dof}})$ , where  $L_0$  is the minimum of the likelihood without (with) the signal contribution,  $N_{\text{dof}}$  is the number of free parameters in the signal PDF and  $\text{PROB}$  is the upper tail probability of a chi-squared distribution, converting this probability into a number of standard deviations.

From the  $D_{s1}^+(2536)$  yields, we compute cross-feed-corrected branching fractions, using the signal efficiency and the relative contributions from cross-feed between the different  $B \rightarrow \bar{D}^{(*)}D_{s1}^+(2536)$  channels, as obtained from simulated events. The resulting branching fractions, the efficiencies, including the intermediate branching fractions, and the internal cross-feed contributions are given in Table I. In addition to the effects previously mentioned, systematics in the table include uncertainties from the cross-feed events (0%–3%), underestimate of the MC resolution (1%–10%) and uncertainty on the  $D_{s1}^+(2536)$  natural width (5%–18%). Using only modes containing  $D^{*+}K_S^0$  in order to minimize the systematic error, we also fit the  $D_{s1}^+(2536)$  mass,  $M(D_{s1}^+(2536)) = (2534.78 \pm 0.31 \pm 0.40)$  MeV/ $c^2$ . Our measurement is in good agreement with the world average [1].

The helicity angle distribution for the sum of the four  $B \rightarrow \bar{D}D_{s1}^+(2536)$  modes, determined by fitting the  $D^*K$

mass distribution for the  $D_{s1}^+(2536)$  yield in ten  $\cos\theta_{D_{s1}^+(2536)}$  bins, is shown in Fig. 1(b). Here  $\theta_{D_{s1}^+(2536)}$  is defined as the angle between the  $D^*$  direction and the  $B$  direction in the  $D_{s1}^+(2536)$  frame. We fit the helicity distribution to different spin-parity hypotheses (one free parameter for  $J^P = 1^-$  and  $2^+$ , and two free parameters for  $J^P = 1^+$  and  $2^-$ ). The fits to  $J^P = 1^+$  in pure  $S$  wave (flat distribution, not shown in the figure),  $J^P = 1^+$  with  $S$ - $D$  wave admixture, and  $J^P = 1^-$  are all in good agreement with data, with  $\chi^2/\text{n.d.f.}$  of 15.9/9, 9.3/8, and 9.6/9 respectively. Fits to  $J^P = 2^+$  and  $2^-$  are disfavored, with  $\chi^2/\text{n.d.f.}$  of 26.0/9 and 26.0/8 respectively.

We search for  $B^+(B^0) \rightarrow \psi(3770)K^+(K_S^0)$  with  $\psi(3770) \rightarrow D^-D^+$ ,  $\bar{D}^0D^0$ . In Fig. 1(c) we show the  $\bar{D}D$  invariant mass for the sum of  $B^+ \rightarrow \bar{D}^0D^0K^+$  and  $B^+ \rightarrow D^-D^+K^+$  candidates. No significant signal is observed in the  $\bar{D}DK_S^0$  modes. The curve shown is the result of a fit where the  $\psi(3770)$  peak is described by a  $P$ -wave Breit-Wigner with the mass as a free parameter in the fit and a natural width fixed to 23 MeV [1], while the background is described by the threshold function described previously. We observe  $57 \pm 11$  events, with a statistical significance of  $6.4\sigma$ , from which we obtain  $M(\psi(3770)) = (3775.5 \pm 2.4 \pm 0.5) \text{ MeV}/c^2$ , in good agreement with the recent high precision measurement of the BES collaboration [14]. We obtain the branching fractions, or limits, reported in Table I, by fitting separately the  $B^+ \rightarrow \bar{D}^0D^0K^+$ ,  $B^0 \rightarrow$

$\bar{D}^0D^0K_S^0$ ,  $B^+ \rightarrow D^-D^+K^+$  and  $B^0 \rightarrow D^-D^+K_S^0$  channels, with  $M(\psi(3770))$  fixed to the result of the combined fit.

For the two modes with significant signal, we study the  $\psi(3770)$  helicity angle,  $\theta_{\psi(3770)}$  [Fig. 1(d)], defined as the angle between the  $D^{0/+}$  direction and the  $B$  direction in the  $\psi(3770)$  frame. We confirm the spin 1 assignment of the  $\psi(3770)$  ( $\chi^2/\text{n.d.f.} = 2.9/9$ ). A spin 0 hypothesis gives  $\chi^2/\text{n.d.f.} = 22.0/9$ .

We search for decays  $B \rightarrow X(3872)K$ ,  $X(3872) \rightarrow \bar{D}^{*0}D^0$  in the  $B^+(B^0) \rightarrow \bar{D}^0D^{*0}K^+(K_S^0) + \bar{D}^{*0}D^0K^+(K_S^0)$  samples. We plot the  $\bar{D}^{*0}D^0$  invariant mass distribution for the sum of  $B^0$  and  $B^+$  candidates in Fig. 1(e). Because of the proximity of the threshold and to the fact that the natural width of the  $X(3872)$  is comparable to the  $\bar{D}^{*0}D^0$  mass resolution, there is no easy analytic parametrization of the reconstructed  $X(3872)$  mass spectrum. To measure the mass and width of the  $X(3872)$ , we generate and reconstruct high statistics MC samples of  $B \rightarrow X(3872)K$  events with various masses (from 3872 to 3877 MeV/ $c^2$ ) and widths (from 0 to 20 MeV), assuming a pure  $S$ -wave decay of a spin 1 resonance. We perform binned extended maximum likelihood fits to the measured  $\bar{D}^{*0}D^0$  invariant mass distributions using these different MC samples as signal PDFs combined with a threshold function for the background. We compare the agreement of each mass and width hypothesis to the data by computing the  $\chi^2$  of the fit for the sum of bins below 3.9 GeV/ $c^2$ . Figure 2 shows the




FIG. 1 (color online). Top: Invariant mass distributions of  $D^*K$  (a),  $\bar{D}D$  (c) and  $\bar{D}^{*0}D^0$  (e) in the data for events in the  $(m_{ES}, \Delta E)$   $B$  signal region. Points with statistical errors are data events, the solid line represents the fit to the data, the dashed line shows the contribution of the  $D_{s1}^+(2536)$  (a),  $\psi(3770)$  (c) and  $X(3872)$  (e) signals, and the dotted line shows the background contribution. Bottom: background-subtracted and efficiency-corrected helicity angle distributions for the  $D_{s1}^+(2536)$  (b) (containing only the  $B \rightarrow \bar{D}D_{s1}^+(2536)$  modes),  $\psi(3770)$  (d) and  $X(3872)$  (f) signals in the data (points with statistical errors) and fitted distributions for different spin hypotheses:  $J^P = 0^+$  (dashed lines),  $1^-$  (solid),  $1^+$  with  $S$ - $D$  wave admixture (dotted),  $2^-$  (long-dashed) and  $2^+$  (dotted-dashed).

TABLE I. Fitted event yields, number of events attributed to internal cross-feed events, efficiencies  $\epsilon$  (including intermediate branching fractions), final branching fractions,  $\mathcal{B}$ , and statistical significances for  $B \rightarrow D^{(*)}D_{s1}^+(2536)$  followed by  $D_{s1}^+(2536) \rightarrow D^{*0}K^+$  or  $D_{s1}^+(2536) \rightarrow D^{*+}K^0$  (first part of the table); for  $B \rightarrow \psi(3770)K$  followed by  $\psi(3770) \rightarrow \bar{D}^0D^0$  or  $\psi(3770) \rightarrow D^-D^+$  (second part of the table); for  $B \rightarrow X(3872)K$  followed by  $X(3872) \rightarrow \bar{D}^{*0}D^0$  (third part of the table). No branching fraction is given in the absence of signal observation. A 90% C.L. upper limit on the branching fraction is given when the statistical significance is less than or equal to  $3\sigma$  (these limits use the profile likelihood method [13] and include systematic errors). The first error on  $\mathcal{B}$  is statistical and the second is systematic.

$B$ mode	Yield	Cross-feed	$\epsilon(10^{-4})$	$\mathcal{B}(10^{-4})$	Limit ( $10^{-4}$ )	Significance
$B^0 \rightarrow D^-D_{s1}^+(2536) [D^{*0}K^+]$	$22.4 \pm 6.3$	$0.1 \pm 0.1$	3.4	$1.71 \pm 0.48 \pm 0.32$	—	$4.4\sigma$
$B^0 \rightarrow D^{*-}D_{s1}^+(2536) [D^{*0}K^+]$	$30.9 \pm 7.9$	$1.2 \pm 0.1$	2.3	$3.32 \pm 0.88 \pm 0.66$	—	$4.6\sigma$
$B^+ \rightarrow \bar{D}^0D_{s1}^+(2536) [D^{*0}K^+]$	$29.2 \pm 6.9$	$0.7 \pm 0.1$	3.5	$2.16 \pm 0.52 \pm 0.45$	—	$5.6\sigma$
$B^+ \rightarrow \bar{D}^{*0}D_{s1}^+(2536) [D^{*0}K^+]$	$66.7 \pm 13.0$	$6.3 \pm 0.1$	2.9	$5.46 \pm 1.17 \pm 1.04$	—	$6.1\sigma$
$B^0 \rightarrow D^-D_{s1}^+(2536) [D^{*+}K^0]$	$7.7 \pm 3.1$	$0.0 \pm 0.1$	0.8	$2.61 \pm 1.03 \pm 0.31$	—	$3.8\sigma$
$B^0 \rightarrow D^{*-}D_{s1}^+(2536) [D^{*+}K^0]$	$16.8 \pm 5.0$	$0.1 \pm 0.1$	0.9	$5.00 \pm 1.51 \pm 0.67$	—	$4.5\sigma$
$B^+ \rightarrow \bar{D}^0D_{s1}^+(2536) [D^{*+}K^0]$	$7.7 \pm 3.2$	$0.1 \pm 0.1$	0.9	$2.30 \pm 0.98 \pm 0.43$	—	$3.3\sigma$
$B^+ \rightarrow \bar{D}^{*0}D_{s1}^+(2536) [D^{*+}K^0]$	$4.8 \pm 2.7$	$0.6 \pm 0.1$	0.3	$3.92 \pm 2.46 \pm 0.83$	10.69	$2.3\sigma$
$B^+ \rightarrow \psi(3770)K^+ [\bar{D}^0D^0]$	$48.6 \pm 10.2$	—	9.0	$1.41 \pm 0.30 \pm 0.22$	—	$6.0\sigma$
$B^0 \rightarrow \psi(3770)K^0 [\bar{D}^0D^0]$	$0.0 \pm 1.5$	—	1.9	—	1.23	$0.0\sigma$
$B^+ \rightarrow \psi(3770)K^+ [D^-D^+]$	$8.9 \pm 3.4$	—	2.8	$0.84 \pm 0.32 \pm 0.21$	1.80	$3.0\sigma$
$B^0 \rightarrow \psi(3770)K^0 [D^-D^+]$	$2.2 \pm 1.6$	—	0.7	$0.81 \pm 0.59 \pm 0.09$	1.88	$1.6\sigma$
$B^0 \rightarrow X(3872)K^0 [\bar{D}^{*0}D^0]$	$5.8 \pm 2.7$	—	0.7	$2.22 \pm 1.05 \pm 0.42$	4.37	$1.3\sigma$
$B^+ \rightarrow X(3872)K^+ [\bar{D}^{*0}D^0]$	$27.4 \pm 5.9$	—	4.3	$1.67 \pm 0.36 \pm 0.47$	—	$4.6\sigma$

interpolated  $\chi^2 - \chi_{\min}^2$  contours versus the simulated masses and widths of the different signal samples, where  $\chi_{\min}^2$  is the  $\chi^2$  value for the best fit. This best fit gives  $33 \pm 7$  events in the  $X(3872)$  peak, with a statistical significance of  $4.9\sigma$ . Mass and width central values are obtained at the minimum of the  $\chi^2$  distribution, while the errors are given by the extreme points of the contour in the (mass, width) plane defined at  $\chi_{\min}^2 + 1$ . We obtain a mass of  $(3875.1_{-0.5}^{+0.7} \pm 0.5)$  MeV/ $c^2$  and a width of  $(3.0_{-1.4}^{+1.9} \pm 0.9)$  MeV, where the systematic errors include additional contributions from the choice of the bin width of the invariant mass distribution (0.14 MeV/ $c^2$  and 0.07 MeV, respectively, on the mass and on the width) and from the fact that in the MC we assume  $S$ -wave  $X(3872)$  decays to  $\bar{D}^{*0}D^0$  (0.20 MeV/ $c^2$  and 0.80 MeV, respectively, on the mass and on the width, determined using MC events with  $P$ -wave  $X(3872)$  decays). Independently of the mass value, the width measurement is  $1.8\sigma$  away from 0 MeV. The  $B^+$  and  $B^0$  branching fractions to  $X(3872)K$  (reported in Table I) are obtained by fitting the  $\bar{D}^{*0}D^0$  invariant mass spectra, separately for  $B^+$  and  $B^0$ , choosing the MC sample with  $M(X(3872)) = 3875$  MeV/ $c^2$  and  $\Gamma(X(3872)) = 3$  MeV, which is found to give the best fit to the data.

We study the helicity angle of the  $X(3872)$ ,  $\theta_{X(3872)}$ , for the sum of  $B^0$  and  $B^+$  modes [see Fig. 1(f)]. Here,  $\theta_{X(3872)}$  is defined as the angle between the  $D^0$  or  $D^{*0}$  direction and the  $B$  direction in the  $X(3872)$  frame. Comparing the curves obtained with different spin hypotheses with the data distribution, we obtain the following  $\chi^2/\text{n.d.f.}$ : 9.8/7 for  $J^P = 1^-$ , 3.9/7 for  $1^+$  assuming a pure  $S$  wave (flat distribution, not shown in the figure), 2.5/6 for  $1^+$  with

$S$ - $D$  wave admixture, 5.9/7 for  $2^+$  and 2.7/6 for  $2^-$ . On the basis of this data sample, we cannot distinguish the different spin assignments. However the  $\bar{D}^{*0}D^0$  decay would be suppressed by the angular momentum barrier for  $J = 2$ .

The ratio of  $X(3872)$  candidates reconstructed in the  $\bar{D}^0D^0\pi^0$  and  $\bar{D}^0D^0\gamma$  final states is  $1.37 \pm 0.56$  (statistical error only), while we expect 1.30 for a decay that proceeds exclusively via a  $D^{*0}$  meson. In addition, we measure parameters which can be used to differentiate various theoretical interpretations [8,15]:  $\Delta m$ , the mass difference




FIG. 2 (color online). Equidistant contours of  $\chi^2 - \chi_{\min}^2$  versus mass and width of the simulated  $X(3872)$  signals. The contours are separated by  $\chi^2$  values of 0.5. Solid contours represent values starting from 0.5, by step of 1, while dashed contours represent values starting from 1, by step of 1. The contour at  $\chi^2 - \chi_{\min}^2 = 1$  is indicated by a wider line. The cross shows the central values of the  $X(3872)$  mass and width. The vertical and horizontal dotted lines indicate the errors on the mass and on the width, respectively.

B. AUBERT *et al.*

PHYSICAL REVIEW D **77**, 011102(R) (2008)

between the state seen in  $B^0$  decays and  $B^+$  decays, as well as  $R_{0/+}$ , the ratio of branching fractions between  $B^0$  decays and  $B^+$  decays. Assuming that the signal seen in  $B^0$  decays is not a statistical fluctuation, we obtain  $\Delta m = (0.7 \pm 1.9 \pm 0.3) \text{ MeV}/c^2$  and  $R_{0/+} = 1.33 \pm 0.69 \pm 0.43$ .

In summary, we report the observation of the eight  $B \rightarrow \bar{D}^{(*)}D_{s1}^+(2536)$  decays, with a  $D_{s1}^+(2536)$  mass of  $(2534.78 \pm 0.31 \pm 0.40) \text{ MeV}/c^2$ . We observe the  $\psi(3770)$  resonance in  $B \rightarrow \bar{D}DK$  decays and measure its mass to be  $(3775.5 \pm 2.4 \pm 0.5) \text{ MeV}/c^2$ . We show that an enhancement of data is observed near the limit of phase space for the  $\bar{D}^{*0}D^0$  invariant mass, at a mass of  $(3875.1_{-0.5}^{+0.7} \pm 0.5) \text{ MeV}/c^2$ , with a width of  $(3.0_{-1.4}^{+1.9} \pm 0.9) \text{ MeV}$ . This enhancement could be interpreted as the  $X(3872)$ , although the observed mass is  $4.5\sigma$  away from

the mass measured in the  $J/\psi\pi^+\pi^-$  decay mode. Our mass value is in good agreement with the value measured by Belle in the  $\bar{D}^0D^0\pi^0$  final state.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), MEC (Spain), and STFC (United Kingdom). Individuals have received support from the Marie Curie EIF (European Union) and the A. P. Sloan Foundation.

- 
- [1] W.-M. Yao *et al.* (Particle Data Group), *J. Phys. G* **33**, 1 (2006).
  - [2] B. Aubert *et al.* (*BABAR* Collaboration), *Phys. Rev. D* **68**, 092001 (2003).
  - [3] B. Aubert *et al.* (*BABAR* Collaboration), *Phys. Rev. D* **74**, 091101 (2006).
  - [4] K. Abe *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **93**, 051803 (2004).
  - [5] S. K. Choi *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **91**, 262001 (2003); B. Aubert *et al.* (*BABAR* Collaboration), *Phys. Rev. D* **71**, 071103 (2005); D. Acosta *et al.* (CDF II Collaboration), *Phys. Rev. Lett.* **93**, 072001 (2004); V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **93**, 162002 (2004).
  - [6] A. Abulencia *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **96**, 102002 (2006).
  - [7] G. Gokhroo *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **97**, 162002 (2006).
  - [8] F. E. Close and P. R. Page, *Phys. Lett. B* **578**, 119 (2004); N. A. Tornqvist, *Phys. Lett. B* **590**, 209 (2004); L. Maiani *et al.*, *Phys. Rev. D* **71**, 014028 (2005); Bing An Li, *Phys. Lett. B* **605**, 306 (2005); K. K. Seth, *Phys. Lett. B* **612**, 1 (2005).
  - [9] B. Aubert *et al.* (*BABAR* Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **479**, 1 (2002).
  - [10] S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
  - [11] C. Cawlfeld *et al.* (CLEO Collaboration), *Phys. Rev. Lett.* **98**, 092002 (2007).
  - [12] G. C. Fox and S. Wolfram, *Nucl. Phys.* **B149**, 413 (1979).
  - [13] W. Rolke, A. Lopez, and J. Conrad, *Nucl. Instrum. Methods Phys. Res., Sect. A* **551**, 493 (2005).
  - [14] M. Ablikim *et al.* (BES Collaboration), *Phys. Rev. Lett.* **97**, 121801 (2006).
  - [15] E. Braaten and M. Kusunoki, *Phys. Rev. D* **71**, 074005 (2005); E. Swanson, *Phys. Rep.* **429**, 243 (2006).