

UC Irvine

UC Irvine Previously Published Works

Title

Subjective Socioeconomic Status Matters Less When Perceived Social Support Is High: A Study of Cortisol Responses to Stress

Permalink

<https://escholarship.org/uc/item/4vm2f078>

Authors

Hooker, Emily D.
Campos, Belinda
Zoccola, Peggy

Publication Date

2017

Peer reviewed

Subjective Socioeconomic Status Matters Less When Perceived Social Support is High:
A Study of Cortisol Responses to Stress

Emily D. Hooker¹

Belinda Campos¹

Peggy M. Zoccola²

Sally S. Dickerson³

University of California, Irvine¹

Ohio University²

Pace University³

Author Note

Emily D. Hooker, Psychology and Social Behavior, University of California, Irvine.
Belinda Campos, Chicano/Latino Studies, University of California, Irvine. Peggy M. Zoccola,
Psychology, Ohio University. Sally S. Dickerson, Psychology, Pace University.

Emily D. Hooker examines culture, social support, and concomitant psychological and physiological well-being. Sally S. Dickerson examines how social-evaluative threat and accompanying self-conscious emotional responses affect physiological outcomes. Belinda Campos focuses on how the sociocultural context shapes relationship experiences in ways that benefit health. Peggy M. Zoccola studies the cognitive and emotional factors that have the potential to extend psychological and physiological stress responses.

National Science Foundation (BCS-0720066) and University of California, Irvine, Academic Senate Council on Research, Computing, and Library Resources grants supported this

project for Dickerson. Hooker was supported by a National Science Foundation Graduate Research Fellowship (DGE-1321846) during this project. This work was completed, in part, while Dickerson was serving at the National Science Foundation. Any opinion, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation. This study was drawn from a dataset used in published reports. We thank the Campos and Pressman Laboratories and Marie Cross for feedback on earlier drafts of this manuscript.

Address correspondence to Emily D. Hooker, Psychology and Social Behavior, University of California, Irvine, CA 92697. E-mail: ehooker@uci.edu.

The final version of this manuscript was published in *Social Psychological and Personality Science*.

Direct link: <http://journals.sagepub.com/doi/abs/10.1177/1948550617732387>

Citation: Hooker, E. D., Campos, B., Zoccola, P. M. & Dickerson, S. S. (2017). Subjective socioeconomic status matters less when perceived social support is high: A study of cortisol responses to stress. *Social Psychological and Personality Science*. Advance online publication.

Abstract

Low objective and subjective socioeconomic status (SES) is associated with the experience of frequent stressors known to have physiological costs. We tested whether perceived social support, a key health-protective resource, buffered the association between lower subjective SES and cortisol responses to an acute stressor. Participants ($N=115$; 54.78% female; age $M=19.56$) reported subjective SES and perceived support, completed a social-evaluative stressor task, and provided saliva for cortisol assessment. There was a significant interaction of subjective SES with support predicting linear change in cortisol stress responses, $\gamma=0.08$, $z=2.34$, $p=.02$. When support was low, subjective SES was strongly related to cortisol, and those who reported lower subjective SES exhibited higher cortisol during recovery than those who reported higher subjective SES. When support was high, those who reported higher and lower subjective SES exhibited similar cortisol responses. These results highlight the important, protective role that supportive relationships can have when subjective SES is low.

Keywords: subjective socioeconomic status, perceived social support, cortisol, stress

Subjective socioeconomic status matters less when perceived social support is high: A study of cortisol responses to stress

Two decades of research have established that lower socioeconomic status (SES), whether measured *objectively* (e.g., income, educational history, or occupational prestige) or *subjectively* (i.e., perception of one's status; Adler, Epel, Castellazzo, & Ickovics, 2000), is robustly associated with higher mortality and morbidity (Adler, 2009; Adler et al., 1994). Pathways linking lower subjective and objective SES to poorer health include the experience of more frequent stressors, greater perceived stress, and/or heightened physiological responses to stressors (e.g., cortisol responses; Adler et al., 2000; Adler & Stewart, 2010; Senn, Walsh, & Carey, 2014). Emerging research also suggests that those with lower objective and subjective SES are more communally-focused (Kraus, Piff, Mendoza-Denton, Rheinschmidt, & Keltner, 2012) and are more likely to turn to others when faced with potential stressors (Piff, Stancato, Martinez, Kraus, & Keltner, 2012). Daily perceived social support, however, has been associated with *lower* cortisol responses to stressors (Eisenberger, Taylor, Gable, Hilmert, & Lieberman, 2007). Thus, we hypothesized that perceived social support would buffer cortisol stress responses for those who reported lower subjective SES.

Subjective SES and Health

While objective measures of SES may include self-reports of income, occupation, education background, or assets, subjective SES is typically measured by asking individuals to consider how they rank in terms of income, job prestige, and education *compared to others* in their country (Adler et al., 2000). Subjective SES reflects individuals' perceived SES relative to others and it is believed to "more accurately capture[s] subtle aspects of social status" (Operario, Adler, & Williams, 2004, p. 238). For example, a bachelor's degree from a prestigious university

may confer more social status than the same degree from a lesser-known university (Operario et al., 2004). Individuals may also incorporate their familial or spousal SES or future educational or occupational opportunities when determining their subjective SES (Singh-Manoux, Adler, & Marmot, 2003). Lower subjective SES in adulthood has been associated with poorer self-rated health, higher levels of depression, greater susceptibility to cold viruses, and higher mortality, as well as markers of angina and respiratory illness (Cohen, Alper, Adler, Treanor, & Turner, 2008; Goodman, Huang, Schafer-Kalkhoff, & Adler, 2007; Macleod, Davey Smith, Metcalfe, & Hart, 2005; Singh-Manoux et al., 2003). In *all* of these studies, subjective SES remained a significant predictor even when controlling for objective SES, and in some cases, was a better predictor (e.g., Cohen et al., 2008; Goodman et al., 2007). This research highlights the utility of incorporating both objective and subjective measures of SES into studies to clarify the relative importance of material resources versus social rank in understanding the link between social class context and health (Kraus et al., 2012).

Subjective SES and Stress

Lower subjective SES is believed to influence health, in part, via increased exposure to stressors (i.e., threatening events [Lazarus & Folkman, 1984]), greater perceived stress, heightened reactivity to stressors, and fewer resources to cope with stressors (Adler et al., 2000; Adler & Stewart, 2010; Senn et al., 2014). For example, if two individuals with equal qualifications are facing a job interview, if one perceives lower status, the stakes for the interview may feel higher, and therefore more threatening. Lower subjective SES is associated with heightened vigilance for potential threats (Kraus, Horberg, Goetz, & Keltner, 2011), which may have long-term physiological consequences. Recent fMRI research has demonstrated that lower subjective SES is associated with greater neural activity in areas related to mentalization

under social-evaluative conditions, which could indicate greater focus on the evaluation (Muscatell et al., 2016). Mentalization, in turn, predicted greater inflammatory responses, which may contribute to physiological wear and tear over time.

One type of threat strongly linked to physiological stress responses is social-evaluative threat. Social-evaluative threat, when one's status, self-esteem, or valued aspect of identity is threatened, is a potent elicitor of the stress hormone cortisol (Dickerson & Kemeny, 2004), which plays an important role in to day-to-day metabolic, immune, and autonomic regulation (Lovallo & Thomas, 2000). It has been proposed that individuals with either exaggerated or blunted cortisol responses to stressors could have future health risks (Phillips, Ginty, & Hughes, 2013), but longitudinal research has yet to precisely characterize the trajectories associated with health vulnerability. However, if lower subjective SES is associated with greater attunement to social-evaluative threats or stressors (Muscatell et al., 2016), it may also be associated with greater acute cortisol responses to these stressors.

A growing number of studies have examined the relationship between SES and cortisol responses to acute stressors, but the findings are mixed. A number of studies have found that objective SES was not associated with cortisol responses to acute stressors in a variety of populations (Carpenter, Shattuck, Tyrka, Geraciotti, & Price, 2011; Hackman, Betancourt, Brodsky, Hurt, & Farah, 2012; Lovallo, Farag, Sorocco, Cohoon, & Vincent, 2012; Steptoe, Kunz-Ebrecht, Wright, & Feldman, 2005), while others have found lower objective SES to be associated with smaller (Kristenson, Kucinskiene, Bergdahl, & Orth-Gomer, 2001) and larger (Fiocco, Joobar, & Lupien, 2007) increases in cortisol. In a study of college students, higher subjective status within individuals' dormitory floors was associated with larger cortisol responses to the stressor (Gruenewald, Kemeny, & Aziz, 2006). Lower subjective SES has also

been linked to less cortisol habituation to a social-evaluative laboratory stressor in middle-aged women (Adler et al., 2000), which is consistent with heightened threat vigilance associated with lower subjective SES. This mixed pattern of results suggests that the link between SES and cortisol reactivity to acute stressors may be moderated by a third variable. The support derived from one's social relationships may be one such variable.

Social Relationships as a Stress Buffer

Those who perceive that they have more social support tend to report lower life stress (Cohen & Hoberman, 1983) and exhibit a 35% *increased* likelihood of survival compared to those who perceive less support (Holt-Lunstad, Smith, & Layton, 2010). Neural evidence finds that perceived support is associated with reduced threat sensitivity and lower cortisol responses to acute social-evaluative stressors (Eisenberger et al., 2007).

Several theories of resilience propose that social support plays an important role in providing psychological and physiological buffering for those who report lower subjective or objective SES backgrounds. The "Shift-and-Persist" model theorizes that having a "stable positive role model" in childhood improves cognitive, emotional, and behavioral responses to the frequent and impactful stressors that those from lower subjective and objective SES backgrounds face (Chen & Miller, 2012). Consistent with this framework, several studies have found that childhood experiences of traumatic stressors (e.g., exposure to violence) are associated with dysregulated cortisol responses to stressors (i.e., greater increases cortisol in response to an acute stressor); however, this relationship may not emerge among children who have a nurturing family environment (although traumatic stressors, such as violence, may not be the same stressors children of low SES typically confront; e.g., Hibel, Granger, Blair, & Cox, 2011). Thus,

social relationships may be valuable psychosocial resources under stressful conditions for those reporting low subjective or objective SES.

When those who report lower objective or subjective SES or those who are made to feel lower subjective SES are confronted with potential stressors, they are more likely to turn toward their social network than those who have higher SES (Piff et al., 2012). This may occur because those who report lower subjective and objective SES tend to be more communally focused (Kraus et al., 2012), and they report greater meaning in life when spending time with others (Kushlev, Dunn, & Ashton-James, 2012). Although social relationships are a promising moderator in the relationship between SES and physiological responses to acute stressors, only one study, to our knowledge, has examined this. In this study of college-aged students, the presence of two supportive audience members during a social-evaluative stressor was associated with lower pro-inflammatory cytokine reactivity in lower subjective SES participants, which is believed to be a health protective pattern of reactivity (John-Henderson, Stellar, Mendoza-Denton, & Francis, 2015). Given that both cortisol and inflammatory responses are sensitive to the social-evaluative context (e.g., Dickerson, Gruenewald, & Kemeny, 2009; Dickerson & Kemeny, 2004), this suggests that a similar relationship may emerge with cortisol and warrants an examination of the relationship between subjective SES, perceived support, and acute cortisol stress responses.

The Present Study

We tested whether perceived social support may moderate the association of subjective SES with cortisol responses to an acute laboratory stressor. Drawing on theoretical and empirical evidence, our key hypothesis was that subjective SES and perceived support would interact to

predict more favorable cortisol trajectories for those who reported lower subjective SES and higher perceived support than lower subjective SES and lower perceived support.

We also sought to rule out alternative explanations for the hypothesized patterns. First, we examined the effect of objective SES to rule out the possibility that subjective SES represents the same construct rather than a measure that reflects a more nuanced understanding of one's social rank, which may be uniquely related to psychological processes.

Method

Participants

Eligible participants¹ ($N=115$) completed a larger study on physiological responses to laboratory and daily life stressors. They were recruited via the university student portal website and campus flyers. Eligible participants had to be 18 years or older, non-smokers, fluent in English, without serious mental or physical illnesses, and typically awake before 10:00 AM. Additionally, participants could not have been pregnant or taking hormonal contraceptives. These requirements ensured interpretability of cortisol assessments (Kirschbaum, Kudielka, Gaab, Schommer, & Hellhammer, 1999; Lovallo & Thomas, 2000; Rohleder & Kirschbaum, 2006). Participants were compensated \$20 for their participation. The final study sample was 54.78% female with a mean age of 19.56 ($SD=2.11$). Participants were of the following ethnic

¹One-hundred and twenty-four individuals consented to participate in the study, but nine participants' data were excluded from the analyses. One participant did not meet the medication eligibility criteria (as she was taking hormonal contraceptives), one declined to complete the speech task, one session was interrupted by a fire alarm, and one participant received a pilot version of the protocol. One participant was excluded due to baseline cortisol levels measuring more than four standard deviations above the mean and another because of missing cortisol values due to a laboratory error. Three participants were excluded because they did not complete the support measure.

background groups: 59.13% Asian, 13.91% European, 9.57% Middle Eastern, 9.57% Chicano/Latino, and 7.83% other.

Procedure

Participants came to the lab between 12:00-6:00 PM to account for the diurnal rhythm of cortisol. They gave informed consent and began a 40-minute baseline period during which they completed questionnaires. Afterward they participated in an adapted version of the Trier Social Stress Test (Kirschbaum, Pirke, & Hellhammer, 1993), which reliably elicits psychological and physiological stress responses. Participants had five minutes to prepare speeches as if they were interviewing for a job in their field. They then completed brief questionnaires and delivered their five-minute speech in front of two evaluators. Evaluators were trained to remain stoic during the task and refrain from giving verbal or nonverbal feedback to participants. The speech was followed by a 40-minute recovery period. Saliva samples were taken at the end of the baseline period, immediately after the speech, and 10, 25, and 40 minutes into the recovery period to capture peak reactivity and recovery.

Measures

Demographics. Age, ethnicity, and gender were assessed via self-report.

Body Mass Index (BMI). BMI was calculated based on self-reported height and weight.

Minutes since waking. Participants reported the time they woke up the morning of the laboratory session, and the difference between the time they woke up and the time of their session was calculated.

Subjective SES. Subjective SES was measured with the MacArthur Scale of Social Status (Adler et al., 2000). This scale shows a 10-rung ladder and asks participants to “think of this ladder as representing where people stand in the United States. At the top of the ladder are

the people who are the best off—those who have the most money, most education, and most respected jobs.” Participants ranked themselves by selecting a number from one to 10 that corresponded with each rung of the ladder.

Objective SES. Objective SES was measured by averaging mothers’ and fathers’ highest level of education based on the following possible options: elementary school (1), junior high (2), high school (3), some college/no degree (4), junior college (5), college (6), some graduate school/no degree (7), and graduate school (8). This measure was selected instead of a measure of occupation (also included in the surveys) because parental occupation can change over time, but parental education is relatively stable (Shavers, 2007).

Perceived Social Support. We used the 12-item positive portion of the Positive and Negative Social Exchanges Scale (Newsom, Nishishiba, Morgan, & Rook, 2003). Participants were asked to think about people in their lives and to indicate how often the items described their interactions with these people (0=never; 4=very often). Examples of items included: “In the past month, how often did the people you know...” “suggest ways that you could deal with problems you were having,” and “do social or recreational activities with you?” Items were averaged and higher scores indicate higher support ($\alpha=.92$; see supplemental materials for additional information about the selection of this measure).

Cortisol. Saliva was collected with a Salivette device (Sarstedt, Inc., Newton, NC) and stored at -20°C . Saliva samples were centrifuged and assayed using standard enzyme-linked immunoassay procedures (Diagnostic Systems Laboratories, Inc., Webster, TX) at the University General Clinical Research Center. Samples were assayed in duplicate and averaged. The sensitivity of the assay was $< 0.012 \mu\text{g/dL}$ and inter-assay and intra-assay coefficients of variance were less than eight percent. The natural log of cortisol was used to account for non-

normality. No other measures of endocrine activity were assessed.

Data Analytic Strategy and Initial Cortisol Analyses

Tools for determining sample size for adequately powered studies using multilevel modeling with continuous predictors are not well established. However, simulation studies suggest that coefficient and variance estimates are minimally biased when N is at least 50 at level two (Maas & Hox, 2005). The sample in this study is more than twice the recommended size ($N=115$).

Cortisol trajectories in response to the social-evaluative stressor were analyzed using multilevel modeling with the “xtmixed” command in STATA/IC version 13.1 (StataCorp, 2013). This method of analysis accounts for the nested nature of the data (i.e., cortisol assessments [Level 1] nested within each participant [Level 2]) and models both within-person change over time and between-person differences in cortisol responses over time (Blackwell, Mendes de Leon, & Miller, 2006; Hruschka, Kohrt, & Worthman, 2005; Snijders & Bosker, 2012). Thus, we were able to examine how individual level factors (i.e., perceived social support and subjective SES) influence change in cortisol over time. The interaction between the linear form of time, subjective SES, and perceived support (time*SES*support) represents the effect of the interaction between subjective SES and perceived support on the linear slope of cortisol throughout the study. The second interaction between the quadratic form of time, subjective SES, and perceived support (time*time*SES*support), represents the effect of the interaction between subjective SES and support on the quadratic slope of cortisol responses. We used maximum likelihood estimation, because it best tests for differences in fixed effects when random effects are held constant (Snijders & Bosker, 2012), and we specified an unstructured covariance matrix. A model with a random intercept for individuals and a random slope for the linear form of time

best fit the data.

Results

Sample Characteristics

The means, standard deviations, and correlations between the primary constructs and demographic and control variables are reported in Table 1. Only subjective SES was correlated with any other variables. It was positively correlated with both objective SES and social support.

Table 1

Correlations and Descriptive Statistics for SES, Support, and Covariates.

Variable	M(SD)	Subj. SES	Obj. SES	Support	Gender ^a	Age	Wake	BMI
Subj. SES	6.37(1.71)							
Obj. SES	5.26(1.45)	.25*						
Support	3.77(0.73)	.20*	.10					
Gender ^a	.45	.04	.02	-.04				
Age	19.56(2.11)	-.09	-.11	.07	-.03			
Wake	350.06(198.54)	.01	.08	.01	-.15	.002		
BMI	22.64(3.83)	.01	.07	.05	.17	.13	.01	

Note: Subj. SES = subjective SES. Obj. SES = objective SES. Support = perceived social support. Wake = minutes between awakening and session start time. BMI = body mass index. Point biserial correlation coefficients presented for correlations with gender.

^a0 = female, 1 = male.

* $p < .05$.

Cortisol Stress Responses

The model predicting cortisol responses to the stressor included gender as a covariate, the interaction between the linear form of time, subjective SES, and perceived support, and the interaction between the quadratic form of time, subjective SES, and perceived support as fixed effects. The interaction between subjective SES and perceived support was not significantly associated with the quadratic slope of cortisol trajectories, $p=.17^2$; therefore, the quadratic rate of change in cortisol responses was equal across levels of subjective SES and perceived support. However, the linear slope of cortisol was not equal across levels of subjective SES and perceived support. Consistent with hypotheses, there was a significant time*subjective SES*support interaction, $\gamma=0.08$, $z=2.32$, $p=.02$, 95% CI [0.01, 0.14].

To further probe the time*subjective SES*support interaction, we removed the four-way interaction between the quadratic form of time, subjective SES, and perceived support and re-ran the model³ (see Table 2), which eases the interpretation of model coefficients. In this model, the three-way interaction between time, subjective SES, and perceived support was significant, $\gamma=0.03$, $z=3.36$, $p=.001$, 95% CI [0.01, 0.05]. To understand this effect, we interpreted the main effect of subjective SES conditional on perceived support and time (Hoffman, 2015).⁴ Lower subjective SES was associated with higher overall cortisol responses to the stressor for an average level of perceived support at baseline, $\gamma=-0.01$, $p=.85$ (see footnote on page 13 regarding

² See Table 1 in the online supplemental materials for statistical information for non-significant findings.

³ We maintained the quadratic form of time because the model fit best when the change in cortisol over time was represented quadratically. The quadratic form of time simply does not *interact* with subjective SES and perceived support.

⁴ Some of effects described herein are not significant, because they reflect the effect of a given variable when all other variables are equal to 0 (i.e., the mean of each variable due to centering). Nevertheless, interpreting these, lower-order effects helps to elucidate the direction and pattern of the significant three-way interaction.

significance). However, as perceived support increased, subjective SES was *less strongly*, negatively associated with cortisol, $p=.98$. The significant time*subjective SES*support interaction indicated that the relationship between subjective SES and cortisol responses was stronger for those with lower perceived social support and became linearly stronger over the course of the study, $\gamma=0.03$, $z=3.36$, $p=.001$, 95% CI [0.01, 0.05]. In other words, subjective SES mattered more in predicting cortisol responses to the stressor when perceived support was low—especially late in recovery.

Table 2

Multilevel Between-Person Effects of the Interaction Between Subjective SES and Perceived Social Support on Cortisol Responses

Variable	Coefficients (RSE)	95% CI
Fixed Effects		
Intercept	-1.55(.08)***	[-1.70, -1.40]
Gender ^a	0.42(0.11)***	[0.21, 0.64]
Time	0.20(.05)***	[0.09, 0.30]
Time*Time	-0.05(0.01)***	[-0.07, -0.02]
Perceived Social Support	-0.03(0.08)	[-0.17, 0.12]
Time*Perceived Social Support	0.01(0.02)	[-0.03, 0.05]
Subjective SES	-0.01(0.04)	[-0.08, 0.07]
Time*Subjective SES	-0.01(0.01)	[-0.03, 0.01]
Subjective SES*Perceived Social Support	0.001(0.05)	[-0.09, 0.09]
Time*Subjective SES*Perceived Social Support	0.03(0.01)**	[0.01, 0.05]
Covariance Parameters		
	Estimate	
Random Effects		
Random Intercept ($\hat{\sigma}_0^2$)	0.23(0.05)	[0.15, 0.37]
Random Slope ($\hat{\sigma}_1^2$)	0.002(0.004)	[0.0001, 0.07]
Covariance ($\hat{\sigma}_{01}$)	0.02(0.01)	[-0.01, 0.04]
Residual Variance	0.20(0.03)	[0.15, 0.27]

Note: Based on 115 participants with 563 longitudinal records.

^a0 = female, 1 = male.

* $p < .05$. ** $p < .01$. *** $p < .001$

Figure 1 demonstrates the spreading throughout the study between those with lower (-1 *SD*) perceived support of varying subjective SES (the solid lines) and those with higher (+1 *SD*) perceived support and varying subjective SES (the dashed lines). To better understand the impact of this interaction, we tested the significance of the subjective SES and perceived support interaction at each timepoint. The interaction between subjective SES and perceived support was not significant at baseline, $p = .98$ (time zero), at time one, $p = .44$, or at time two, $p = .13$. The interaction between subjective SES and perceived support was, however, significant at time three, $z = 2.18$, $p = .03$, 95% CI [0.01, 0.18], and at time four. $z = 2.68$, $p = .01$, 95% CI [0.03, 0.22].

This indicates that at times three and four, subjective SES mattered more when perceived support was low than when perceived support was high.




Figure 1. Predicted cortisol responses when subjective SES and perceived support are set at one standard deviation above and below their means.

Next, we tested the effect of support when subjective SES was one standard deviation below the mean. When subjective SES was low, there was not a significant effect of perceived support at times three or four (i.e., the difference between triangles at time three and the difference between triangles at time four in Figure 1), $p\text{-values} \geq .11$. Next we tested the effect of SES when perceived support was one standard deviation above the mean. There was not a significant effect of subjective SES when perceived support was high at times three or four (i.e., the difference between the dashed lines at time three and the difference at time four in Figure 1),

p-values $\geq .52$. Finally, we tested the effect of subjective SES when support was one standard deviation below the mean. At both timepoints three and four, $z = -2.90$, $p = .004$, 95% CI [-0.18, -0.04] and $z = -3.62$, $p < .001$, 95% CI [-0.22, -0.06] respectively, there was a significant effect of subjective SES when support was low (i.e., the differences between the solid lines in Figure 1): when perceived support was low at timepoints three and four, the effect of subjective SES was significant. This indicates that for those who reported low support, subjective SES was a strong predictor of cortisol recovery. Figure 1 illustrates that those who reported lower subjective SES and lower support exhibited significantly greater cortisol during recovery than those who reported higher subjective SES and lower support.

Alternative Explanations

The next set of analyses tested whether other related variables explained the relationship between subjective SES, perceived support, and cortisol trajectories. In the model with time*time*subjective SES*support and objective SES, the three-way interaction between time, subjective SES, and perceived support remained the same, $z = -2.26$, $p = .02$, 95% CI [0.01, 0.14]. Furthermore, when subjective SES was replaced with objective SES, time*SES*support did not significantly predict cortisol trajectories, $p = .79$.

Discussion

Our results were consistent with the prediction that perceived support would be protective for those who reported lower subjective SES. When perceived support was high, subjective SES was not significantly associated with cortisol responses during recovery—higher perceived support buffered the association between lower subjective SES and cortisol responses. When perceived support was low, subjective SES was strongly associated with cortisol responses during recovery. Those who reported lower subjective SES and lower support exhibited

significantly greater cortisol during the last two recovery timepoints than those who report higher subjective SES and lower support.

These findings contribute to a small but emerging literature on the role of perceived support in buffering cortisol responses to stressors (e.g., Eisenberger et al., 2007; Wirtz et al., 2006). This study is also one of few examining the interactive effect that support and subjective SES can have on physiological responses to acute stressors (i.e., John-Henderson et al., 2015). Recent research has highlighted the important role of context in understanding and replicating psychological findings (van Bavel, Mende-Siedlecki, Brady, & Reinero, 2016), and our work underscores the importance of considering the social/relational context when evaluating the correlates of subjective SES. Social relationship norms that emphasize communality for those who report lower subjective SES may set a foundation that helps lower status individuals benefit from positive social relationships when facing stressors (Kraus et al., 2012; Piff et al., 2012). Our findings illustrate that social relationships can buffer against stressors, which may potentially offset the physiological costs of repeated stressor exposure for those who report lower subjective SES.

For those who reported higher support, subjective SES was not a robust predictor of cortisol responses. It is possible that those in our sample who reported lower subjective SES and higher support differed in meaningful ways from other low subjective SES samples. For example, our participants were in college, and it is possible that our sample may have been comprised of students who had “shifted-and-persisted” in the past (Chen & Miller, 2012). As a result, this group may be more likely to be impacted by current perceived support. It is beyond the scope of this study to tease apart the possibility that our sample was particularly resilient, but

future studies should examine community members (rather than college students) who report lower subjective SES.

Our finding that subjective SES was less strongly associated with cortisol responses when social support was high, especially in recovery, is in line with evidence suggesting that perceived support can be beneficial under conditions of stress (Eisenberger et al., 2007). The exact mechanism associated with the physiological stress-buffering effect of perceived support, however, is not known. One possibility is threat processing. Although those who report lower subjective SES are more threat vigilant (Kraus et al., 2011), neural research finds that perceived social support reduces threat sensitivity and threat responding and, subsequently, reduces cortisol responses to threat (Eisenberger et al., 2007). Without the potential threat-reducing effects of perceived support, those who reported lower subjective SES may have been at risk for perceiving more threat during the stressor and, therefore, mounting a larger increase in cortisol. Indeed, this was the cortisol pattern observed on the lower end of both subjective SES and support. Ancillary analyses (not reported) indicated that the interaction between subjective SES and perceived support was not associated with perceived threat/stress, which suggests that these factors were not mediators of cortisol responses. However, there is reason to believe that threat processing may operate implicitly at a neural level (Eisenberger et al., 2007). Future investigations should examine the neural underpinnings of threat to explore its potential mediational role.

Why was higher subjective SES and lower support linked to lower cortisol responses during recovery? Lower cortisol responses to acute stressors are sometimes characterized as health-protective responses and other times as representative of physiological dysregulation. The long-term consequences of varying cortisol responses to acute stressors on health are not well studied, and without knowing the long-term influence that each pattern has on health, it is

difficult to speculate about the potential mechanisms of each pattern of response on health. Furthermore, the mechanisms linking support with cortisol responses may vary by levels of subjective SES. Habituation, or exhibiting lower cortisol responses to an acute stressor experienced previously, may reflect “healthy adaptation to novel stress[ors]” (Adler et al., 2000, p. 588). In the future, findings from studies presenting the same stressor multiple times could clarify the potential health-impact of higher and lower cortisol responses to stressors (Adler et al., 2000).

In this study, objective SES did not interact with perceived support to predict cortisol responses, nor were the key findings explained by the contributions of objective SES. We interpret this as evidence that subjective SES may better capture subtle variations in perceived social standing than objective measures (Singh-Manoux et al., 2003). This may be particularly relevant to college samples because college is a transition period for students from childhood SES toward adult SES. Future work should continue to explore the influence of subjective SES and perceived support on acute stress responses in diverse community samples to better understand these processes. Studying younger adults, as we have, may provide meaningful information about the trajectory of the association between SES and responses to stressors across the lifespan and in those potentially shifting subjective SES.

A strength of the study is that the sample was ethnically diverse. However, this also complicated the interpretation of the relationship between SES, support, and ethnic background. Asian background samples are less likely to seek support (Taylor et al., 2004), but our findings are in line with research that suggests that perceived support buffers stressor responses in this group (Taylor, Welch, Kim, & Sherman, 2007). Future studies should replicate our findings in larger sub-samples of all ethnic background groups. Moreover, a larger study would allow for an

examination of the potential association of gender with the subjective SES-support interaction. Because women tend to be more communal (Cross & Madson, 1997), perceptions of lower support may be associated with greater cortisol responses in higher SES women and higher still in lower SES women.

Our work suggests that perceived support is important for understanding the relationship between subjective SES and cortisol responses to stressors. In the large body of literature on socioeconomic status, lower subjective and objective status contexts are often characterized as resource-insufficient in all domains. However, this study suggests that social relationships may be a powerful resource for those reporting lower subjective SES. Overall, the findings of this study illustrate that the impact of supportive relationships may vary by subjective social class. A better understanding of the long-term consequences of perceived support would allow it to be responsibly integrated into interventions aimed at improving the well-being individuals with lower social status.

References

- Adler, N. E. (2009). Health disparities through a psychological lens. *The American Psychologist*, *64*, 663–673.
- Adler, N. E., Boyce, T., Chesney, M. A., Cohen, S., Folkman, S., Kahn, R. L., & Syme, S. L. (1994). Socioeconomic status and health: The challenge of the gradient. *American Psychologist*, *49*, 15–24.
- Adler, N. E., Epel, E. S., Castellazzo, G., & Ickovics, J. R. (2000). Relationship of subjective and objective social status with psychological and physiological functioning: Preliminary data in healthy white women. *Health Psychology*, *19*, 586–592.
- Adler, N. E., & Stewart, J. (2010). Preface to the biology of disadvantage: Socioeconomic status and health. *Annals of the New York Academy of Sciences*, *1186*, 1–4.
- Carpenter, L. L., Shattuck, T. T., Tyrka, A. R., Geraciotti, T. D., & Price, L. H. (2011). Effect of childhood physical abuse on cortisol stress response. *Psychopharmacology*, *214*, 367–375.
- Chen, E., & Miller, G. E. (2012). “Shift-and-Persist” strategies: Why low socioeconomic status isn’t always bad for health. *Perspectives on Psychological Science*, *7*, 135–158.
- Cohen, S., Alper, C. M., Adler, N. E., Treanor, J. J., & Turner, R. B. (2008). Objective and subjective socioeconomic status and susceptibility to the common cold. *Health Psychology*, *27*, 268–274.
- Cohen, S., & Hoberman, H. M. (1983). Positive events and social support as buffers of life change stress. *Journal of Applied Social Psychology*, *13*, 99–125.
- Cross, S. E., & Madson, L. (1997). Models of the self: Self-construals and gender. *Psychological Bulletin*, *122*, 5–37.
- Dickerson, S. S., Gruenewald, T. L., & Kemeny, M. E. (2009). Psychobiological responses to

- social self threat: Functional or detrimental? *Self and Identity*, 8, 270–285.
- Dickerson, S. S., & Kemeny, M. E. (2004). Acute stressors and cortisol responses: A theoretical integration and synthesis of laboratory research. *Psychological Bulletin*, 130, 355–391.
- Eisenberger, N. I., Taylor, S. E., Gable, S. L., Hilmert, C. J., & Lieberman, M. D. (2007). Neural pathways link social support to attenuated neuroendocrine stress responses. *NeuroImage*, 35, 1601–1612.
- Fiocco, A. J., Jooper, R., & Lupien, S. J. (2007). Education modulates cortisol reactivity to the Trier Social Stress Test in middle-aged adults. *Psychoneuroendocrinology*, 32, 1158–63.
- Goodman, E., Huang, B., Schafer-Kalkhoff, T., & Adler, N. E. (2007). Perceived socioeconomic status: a new type of identity that influences adolescents' self-rated health. *The Journal of Adolescent Health*, 41, 479–87.
- Gruenewald, T. L., Kemeny, M. E., & Aziz, N. (2006). Subjective social status moderates cortisol responses to social threat. *Brain, Behavior, and Immunity*, 20, 410–419.
- Hackman, D. A., Betancourt, L. M., Brodsky, N. L., Hurt, H., & Farah, M. J. (2012). Neighborhood disadvantage and adolescent stress reactivity. *Frontiers in Human Neuroscience*, 6, 1–11.
- Hibel, L. C., Granger, D. a, Blair, C., & Cox, M. J. (2011). Maternal sensitivity buffers the adrenocortical implications of intimate partner violence exposure during early childhood. *Development and Psychopathology*, 23, 689–701.
- Hoffman, L. (2015). Between-person analysis and interpretation of interactions. In *Longitudinal Analysis* (pp. 30–78). New York, NY: Routledge.
- Holt-Lunstad, J., Smith, T. B., & Layton, J. B. (2010). Social relationships and mortality risk: A meta-analytic review. *PLoS Medicine*, 7, e1000316.

John-Henderson, N. A., Stellar, J. E., Mendoza-Denton, R., & Francis, D. D. (2015).

Socioeconomic status and social support: Social support reduces inflammatory reactivity for individuals whose early-life socioeconomic status was low. *Psychological Science*.

Kirschbaum, C., Kudielka, B. M., Gaab, J., Schommer, N. C., & Hellhammer, D. H. (1999).

Impact of gender, menstrual cycle phase, and oral contraceptives on the activity of the hypothalamus-pituitary-adrenal axis. *Psychosomatic Medicine*, *61*, 154–162.

Kirschbaum, C., Pirke, K.-M., & Hellhammer, D. H. (1993). The “Trier Social Stress Test”--A tool for investigating psychobiological stress responses in a laboratory setting.

Neuropsychobiology, *28*, 76–81.

Kraus, M. W., Horberg, E. J., Goetz, J. L., & Keltner, D. (2011). Social class rank, threat

vigilance, and hostile reactivity. *Personality and Social Psychology Bulletin*, *37*, 1376–88. h

Kraus, M. W., Piff, P. K., Mendoza-Denton, R., Rheinschmidt, M. L., & Keltner, D. (2012).

Social class, solipsism, and contextualism: How the rich are different from the poor.

Psychological Review, *119*, 546–72.

Kristenson, M., Kucinskiene, Z., Bergdahl, B., & Orth-Gomer, K. (2001). Risk factors for

coronary heart disease in different socioeconomic groups of Lithuania and Sweden--The LiVicordia study. *Scandinavian Journal of Public Health*, *29*, 140–150.

Kushlev, K., Dunn, E. W., & Ashton-James, C. E. (2012). Does affluence impoverish the

experience of parenting? *Journal of Experimental Social Psychology*, *48*, 1381–1384.

Lazarus, R. S., & Folkman, S. (1984). *Stress, appraisal, and coping*. New York, NY: Springer Publishing Company.

Lovallo, W. R., Farag, N. H., Sorocco, K. H., Cohoon, A. J., & Vincent, A. S. (2012). Lifetime

adversity leads to blunted stress axis reactivity: Studies from the Oklahoma Family Health

- Patterns Project. *Biological Psychiatry*, *71*, 344–349.
- Lovallo, W. R., & Thomas, T. L. (2000). Stress hormones in psychophysiological research: Emotional, behavioral, and cognitive implications. In J. T. Cacioppo, L. G. Tassinary, & G. G. Bernston (Eds.), *Handbook of psychophysiology* (2nd ed., pp. 342–367). Cambridge, UK: Cambridge University Press.
- Maas, C. J., & Hox, J. J. (2005). Sufficient sample sizes for multilevel modeling. *Journal of Research Methods for the Behavioral and Social Sciences*, *1*, 86–92.
- Macleod, J., Davey Smith, G., Metcalfe, C., & Hart, C. (2005). Is subjective social status a more important determinant of health than objective social status? Evidence from a prospective observational study of Scottish men. *Social Science & Medicine* (1982), *61*, 1916–29.
- Muscattell, K. A., Dedovic, K., Slavich, G. M., Jarcho, M. R., Breen, E. C., Bower, J. E., ... Eisenberger, N. I. (2016). Neural mechanisms linking social status and inflammatory responses to social stress. *Social Cognitive and Affective Neuroscience*, *11*, 915–922.
- Newsom, J. T., Nishishiba, M., Morgan, D. L., & Rook, K. S. (2003). The relative importance of three domains of positive and negative social exchanges: a longitudinal model with comparable measures. *Psychology and Aging*, *18*, 746–54.
- Operario, D., Adler, N. E., & Williams, D. R. (2004). Subjective social status: Reliability and predictive utility for global health. *Psychology & Health*, *19*, 237–246.
- Phillips, A. C., Ginty, A. T., & Hughes, B. M. (2013). The other side of the coin: Blunted cardiovascular and cortisol reactivity are associated with negative health outcomes. *International Journal of Psychophysiology*, *90*, 1–7.
- Piff, P. K., Stancato, D. M., Martinez, A. G., Kraus, M. W., & Keltner, D. (2012). Class, chaos, and the construction of community. *Journal of Personality and Social Psychology*, *103*,

949–62.

- Rohleder, N., & Kirschbaum, C. (2006). The hypothalamic-pituitary-adrenal (HPA) axis in habitual smokers. *International Journal of Psychophysiology*, *59*, 236–243.
- Senn, T. E., Walsh, J. L., & Carey, M. P. (2014). The mediating roles of perceived stress and health behaviors in the relation between objective, subjective, and neighborhood socioeconomic status and perceived health. *Annals of Behavioral Medicine*, *48*, 215–224.
- Shavers, V. L. (2007). Measurement of socioeconomic status in health disparities research. *Journal of the National Medical Association*, *99*, 1013–23.
- Singh-Manoux, A., Adler, N. E., & Marmot, M. G. (2003). Subjective social status: Its determinants and its association with measures of ill-health in the Whitehall II study. *Social Science & Medicine*, *56*, 1321–1333.
- StataCorp. (2013). *Stata Statistical Software: Release 13*. College Station, TX: StataCorp LP.
- Stephoe, A., Kunz-Ebrecht, S. R., Wright, C., & Feldman, P. J. (2005). Socioeconomic position and cardiovascular and neuroendocrine responses following cognitive challenge in old age. *Biological Psychology*, *69*, 149–66.
- Taylor, S. E., Sherman, D. K., Kim, H. S., Jarcho, J., Takagi, K., & Dunagan, M. S. (2004). Culture and social support: who seeks it and why? *Journal of Personality and Social Psychology*, *87*, 354–362.
- Taylor, S. E., Welch, W. T., Kim, H. S., & Sherman, D. K. (2007). Cultural differences in the impact of social support on psychological and biological stress responses. *Psychological Science*, *18*, 831–837.
- van Bavel, J. J., Mende-Siedlecki, P., Brady, W. J., & Reinero, D. A. (2016). Contextual sensitivity in scientific reproducibility Submission PDF Contextual sensitivity in scientific

reproducibility. *Proceedings of the National Academy of Sciences*, *113*, 6454–6459.

Wirtz, P. H., von Känel, R., Mohiyeddini, C., Emini, L., Ruedisueli, K., Groessbauer, S., & Ehlert, U. (2006). Low social support and poor emotional regulation are associated with increased stress hormone reactivity to mental stress in systemic hypertension. *The Journal of Clinical Endocrinology & Metabolism*, *91*, 3857–65.