

SUBNORMAL TUPLES QUASI-SIMILAR TO THE SZEGÖ TUPLE

Ameer Athavale

In what follows, if \mathcal{H} is a Hilbert space then $\mathcal{B}(\mathcal{H})$ denotes the set of bounded linear operators in \mathcal{H} . All the Hilbert spaces occurring below are separable and all the measures are compactly supported positive regular Borel measures on \mathbf{C}^m . Recall that if S_1, \dots, S_m are m commuting elements in $\mathcal{B}(\mathcal{H})$, then the operator tuple $S = (S_1, \dots, S_m)$ is called *subnormal* on \mathcal{H} if there exist a Hilbert space $\mathcal{H}' \supset \mathcal{H}$ and m commuting normal elements N_1, \dots, N_m in $\mathcal{B}(\mathcal{H}')$ such that $N_j \mathcal{H} \subset \mathcal{H}$ and $N_j|_{\mathcal{H}} = S_j$ for $1 \leq j \leq m$. If $H^2(\mathbf{B}^{2m})$ denotes the Hardy space of the open unit ball \mathbf{B}^{2m} in \mathbf{C}^m [i.e., $H^2(\mathbf{B}^{2m})$ is the completion of polynomials in $L^2(\sigma)$, σ being the surface area measure on the unit sphere \mathbf{S}^{2m-1}], and $M_{z_j}^{(\sigma)}$ denotes multiplication by z_j on $H^2(\mathbf{B}^{2m})$; then the multiplication tuple $M_z^{(\sigma)} = (M_{z_1}^{(\sigma)}, \dots, M_{z_m}^{(\sigma)})$, hereafter referred to as the Szegö tuple, is an example of a subnormal tuple. Moreover, $M_z^{(\sigma)}$ is cyclic. Recall that an operator tuple $S = (S_1, \dots, S_m)$ on \mathcal{H} is called *cyclic* if there exists a vector u in \mathcal{H} (called a *cyclic vector* for S) such that the smallest subspace of \mathcal{H} containing u and invariant under S_1, \dots, S_m is all of \mathcal{H} . The constant function 1 of course serves as a cyclic vector for $M_z^{(\sigma)}$. The following proposition is a well-known fact about cyclic subnormal tuples [3].

PROPOSITION 0. *Suppose $S = (S_1, \dots, S_m)$ is a subnormal tuple on \mathcal{H} with a cyclic vector of norm one. Then there exists a probability measure μ with compact support in \mathbf{C}^m and a unitary operator U from \mathcal{H} onto $H^2(\mu)$ [$H^2(\mu)$ is the completion of polynomials in $L^2(\mu)$] such that $Uu = 1$ and $S_j = U^* M_{z_j}^{(\mu)} U$, $1 \leq j \leq m$; where $M_{z_j}^{(\mu)}$ is multiplication by z_j on $H^2(\mu)$.*

DEFINITION. Let $S = (S_1, \dots, S_m)$ be a subnormal tuple on \mathcal{H} , and let $T = (T_1, \dots, T_m)$ be a subnormal tuple on \mathcal{K} . We say that S is *quasi-similar* to T if there exist bounded linear operators $A: \mathcal{H} \rightarrow \mathcal{K}$ and $B: \mathcal{K} \rightarrow \mathcal{H}$ such that $\text{Ker } A = \{0\}$, $\text{Ker } B = \{0\}$, $\overline{\text{Ran } A} = \mathcal{K}$, $\overline{\text{Ran } B} = \mathcal{H}$, and $AS = TA$, $SB = BT$; that is, $AS_j = T_j A$ and $SB_j = BT_j$ for $1 \leq j \leq m$.

A function-theoretic characterization of subnormal tuples quasi-similar to the multiplication tuple on the Hardy space of the unit polydisc was obtained in [3]. In this note we observe that a similar characterization holds for subnormal tuples quasi-similar to the Szegö tuple. Our characterization allows us in particular to recapture a result in [2] that the Bergman tuple is not quasi-similar to the Szegö tuple. [If $A^2(\mathbf{B}^{2m})$ denotes the completion of polynomials in $L^2(V)$, V being the volumetric measure on the closed unit ball \mathbf{B}^{2m} , then the multiplication tuple on $A^2(\mathbf{B}^{2m})$ is called the Bergman tuple.]

Received September 14, 1987.
Michigan Math. J. 35 (1988).

Crucial for our purposes is the following specialized version of an approximation theorem related to the solution of the inner function problem on the unit ball \mathbf{B}^{2m} (Theorem 3.5 in [4]).

THEOREM 0. *Suppose that*

- (a) $\chi: \overline{\mathbf{B}^{2m}} \rightarrow (0, \infty)$ is continuous, and
- (b) θ is a positive measure on \mathbf{S}^{2m-1} .

Then there exists a sequence $\{p_j\}$ of polynomials such that

- (1) $|p_j| < \chi$ on $\overline{\mathbf{B}^{2m}}$,
- (2) $\lim_{j \rightarrow \infty} p_j(z) = 0$ uniformly on every compact subset of \mathbf{B}^{2m} , and
- (3) $\lim_{j \rightarrow \infty} |p_j(\xi)| = \chi(\xi)$ a.e. $[\theta]$.

PROPOSITION 1. *Let $S = (S_1, \dots, S_m)$ and $T = (T_1, \dots, T_m)$ be subnormal tuples on $\mathfrak{H}\mathcal{C}$ and \mathfrak{K} respectively and let $A: \mathfrak{H}\mathcal{C} \rightarrow \mathfrak{K}$ and $B: \mathfrak{K} \rightarrow \mathfrak{H}\mathcal{C}$ be bounded linear operators such that $\overline{\text{Ran } A} = \mathfrak{K}$ and $\overline{\text{Ran } B} = \mathfrak{H}\mathcal{C}$, with $AS = TA$ and $SB = BT$. If T is cyclic then so is S , and A and B are injective; in particular, S is quasi-similar to T .*

Proof. The proof is a straightforward generalization of Lemma 2.4 in [1] and depends crucially on the fact that the commutant $\{T_1, \dots, T_m\}'$ of the polynomial algebra generated by T_1, \dots, T_m is equal to $\{M_\varphi: \varphi \in H^2(\mu) \cap L^\infty(\mu)\}$, where M_φ denotes multiplication by φ on $H^2(\mu)$ and where μ is the measure associated with T as in Proposition 0. (The last-mentioned fact is in turn a straightforward generalization to cyclic subnormal tuples of a result of Yoshino [5] for cyclic subnormal operators.) \square

Propositions 0 and 1 show that to discuss subnormal tuples S quasi-similar to the Szegő tuple, we need only consider $S = M_z^{(\mu)}$ for some compactly supported measure μ on \mathbf{C}^m . We choose to call a function f in $H^2(\mu)$ *cyclic* if f is a cyclic vector for $M_z^{(\mu)}$. Hereafter, ν will always stand for a fixed positive regular Borel measure on \mathbf{S}^{2m-1} .

PROPOSITION 2. *Let μ be a compactly supported measure on \mathbf{C}^m . Suppose there exists an operator $B: H^2(\nu) \rightarrow H^2(\mu)$ with dense range such that $M_z^{(\mu)}B = BM_z^{(\nu)}$. Then $\mu|_{\mathbf{S}^{2m-1}}$ is absolutely continuous with respect to ν and there is a cyclic function g in $H^2(\mu)$ such that $\int |p|^2 |g|^2 d\mu \leq \int |p|^2 d\nu$ for every m -variable polynomial p .*

Proof. We may assume B has norm one. If $g = B1$, then clearly g is cyclic in $H^2(\mu)$. Define, for any Borel set $E \subset \mathbf{S}^{2m-1}$, $\eta(E) = \int_E |g|^2 d\mu$. Then

$$\int |p|^2 d\eta \leq \int |p|^2 |g|^2 d\mu = \int |Bp|^2 d\mu \leq \int |p|^2 d\nu.$$

If χ is any positive continuous function on $\overline{\mathbf{B}^{2m}}$ then it can be extended to a positive continuous function on \mathbf{B}^{2m} (still denoted χ), and one can choose a sequence $\{p_j\}$ of polynomials corresponding to $\theta = \eta + \nu$ as in Theorem 0. In view of (1) and (3) of Theorem 0, it is clear that $\int \chi d\eta \leq \int \chi d\nu$. This shows that η is

absolutely continuous with respect to ν . Thus $\int_E |g|^2 d\mu = 0$ for every Borel set $E \subset \mathbf{S}^{2m-1}$ such that $\nu(E) = 0$. Since g is cyclic, however, g does not vanish on a set of positive μ -measure and it follows that $\mu(E) = 0$ for $\nu(E) = 0$. \square

PROPOSITION 3. *Let μ be a measure supported on $\overline{\mathbf{B}^{2m}}$. Suppose also there exists an operator $A: H^2(\mu) \rightarrow H^2(\nu)$ such that A has dense range and $AM_z^{(\mu)} = M_z^{(\nu)}A$. Then ν is absolutely continuous with respect to $\mu|_{\mathbf{S}^{2m-1}}$ and there exists a cyclic function f in $H^2(\nu)$ such that $\int |p|^2 |f|^2 d\nu \leq \int |p|^2 d(\mu|_{\mathbf{S}^{2m-1}})$ for every m -variable polynomial p .*

Proof. We may assume A has norm one. If $f = A1$, then clearly f is cyclic in $H^2(\nu)$. Let $\chi = 1$ in Theorem 0 and choose a sequence $\{p_j\}$ of polynomials corresponding to $\theta = \nu + (\mu|_{\mathbf{S}^{2m-1}})$. Then, for any m -variable polynomial p ,

$$\begin{aligned} \int |Ap|^2 d\nu &= \lim_{j \rightarrow \infty} \int |p_j|^2 |Ap|^2 d\nu = \lim_{j \rightarrow \infty} \int |Ap_j p|^2 d\nu \leq \lim_{j \rightarrow \infty} \int |p_j p|^2 d\mu \\ &= \lim_{j \rightarrow \infty} \int |p_j p|^2 d(\mu|_{\mathbf{S}^{2m-1}}) = \int |p|^2 d(\mu|_{\mathbf{S}^{2m-1}}). \end{aligned}$$

Thus $\int |p|^2 |f|^2 d\nu \leq \int |p|^2 d(\mu|_{\mathbf{S}^{2m-1}})$ for any m -variable polynomial p , and appealing to Theorem 0 again we conclude that $\int \chi |f|^2 d\nu \leq \int \chi d(\mu|_{\mathbf{S}^{2m-1}})$ for any positive continuous function χ on \mathbf{S}^{2m-1} . Since f is cyclic, however, f does not vanish on a set of positive ν -measure and it follows that ν is absolutely continuous with respect to $\mu|_{\mathbf{S}^{2m-1}}$. \square

THEOREM 1. *Let μ be a positive regular Borel measure with compact support in \mathbf{C}^m . Then $M_z^{(\mu)}$ is quasi-similar to $M_z^{(\nu)}$ if and only if*

(a) *there exists a cyclic function f in $H^2(\nu)$ such that*

$$\int |p|^2 |f|^2 d\nu \leq \int |p|^2 d(\mu|_{\mathbf{S}^{2m-1}})$$

for every m -variable polynomial p , and

(b) *there exists a cyclic function g in $H^2(\mu)$ such that*

$$\int |p|^2 |g|^2 d\mu \leq \int |p|^2 d\nu$$

for every m -variable polynomial p .

Proof. Suppose $M_z^{(\mu)}$ is quasi-similar to $M_z^{(\nu)}$. By Proposition 1 in [3], μ has its support in \mathbf{B}^{2m} . Condition (a) in Theorem 1 now follows from Proposition 3 and condition (b) from Proposition 2.

Conversely, suppose (a) and (b) are true. We define $A: H^2(\mu) \rightarrow H^2(\nu)$ and $B: H^2(\nu) \rightarrow H^2(\mu)$ by requiring $Ap = fp$ and $Bp = gp$ for every m -variable polynomial p . [The boundedness of A and B is of course a consequence of conditions in (a) and (b).] Since f and g are cyclic, A and B are seen to have dense range. Now apply Proposition 1. \square

REMARK 1. The characterization of subnormal tuples quasi-similar to the Szegő tuple is obtained by choosing ν in Theorem 1 to be the surface area measure σ on \mathbf{S}^{2m-1} .

REMARK 2. Since the restriction of the volumetric measure V on $\overline{\mathbf{B}^{2m}}$ to \mathbf{S}^{2m-1} is zero, σ is not absolutely continuous with respect to $V|_{\mathbf{S}^{2m-1}}$ and our observations above show that the Bergman tuple is not quasi-similar to the Szegő tuple.

REMARK 3. Condition (b) in Theorem 1 actually guarantees that μ has its support in $\overline{\mathbf{B}^{2m}}$. [Justification: Let α be any vector in \mathbf{C}^m with the Hermitian norm $\|\alpha\|$ less than or equal to one. If (b) holds and \cdot denotes the Hermitian inner product on \mathbf{C}^m , then for any positive integer n we have

$$\int |z \cdot \alpha|^{2n} |g|^2 d\mu \leq \int |z \cdot \alpha|^{2n} d\nu \leq \int \|z\|^{2n} \|\alpha\|^{2n} d\nu \leq \nu(\mathbf{S}^{2m-1}).$$

This shows that $|g|^2 d\mu$ has its support in $\bigcap_{\|\alpha\| \leq 1} \{z : |z \cdot \alpha| \leq 1\} = \overline{\mathbf{B}^{2m}}$. Because g is cyclic in $H^2(\mu)$, however, μ has its support in $\overline{\mathbf{B}^{2m}}$.]

ACKNOWLEDGMENTS. The author wishes to thank John B. Conway and Eric Bedford for some useful discussions.

REFERENCES

1. J. B. Conway, *On quasisimilarity for subnormal operators*, Illinois J. Math. 24 (1980), 689–702.
2. R. E. Curto and N. Salinas, *Generalized Bergman kernels and the Cowen–Douglas theory*, Amer. J. Math. 106 (1984), 447–488.
3. W. W. Hastings, *Commuting subnormal operators simultaneously quasisimilar to unilateral shifts*, Illinois J. Math. 22 (1978), 506–519.
4. W. Rudin, *New constructions of functions holomorphic in the unit ball of \mathbf{C}^n* , CBMS Regional Conference Series in Mathematics, Amer. Math. Soc., Providence, R.I., 1986.
5. T. Yoshino, *Subnormal operators with a cyclic vector*, Tohoku Math. J. 21 (1969), 47–55.

Department of Mathematical Sciences
 Indiana University
 Purdue University–Indianapolis
 Indianapolis, IN 46223