



University of Kentucky
UKnowledge

Physics and Astronomy Faculty Publications

Physics and Astronomy

1-25-2018

Subsystem Eigenstate Thermalization Hypothesis

Anatoly Dymarsky

University of Kentucky, a.dymarsky@uky.edu

Nima Lashkari

Massachusetts Institute of Technology

Hong Liu

Massachusetts Institute of Technology

Follow this and additional works at: https://uknowledge.uky.edu/physastron_facpub



Part of the [Quantum Physics Commons](#), and the [Statistical, Nonlinear, and Soft Matter Physics Commons](#)

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

Repository Citation

Dymarsky, Anatoly; Lashkari, Nima; and Liu, Hong, "Subsystem Eigenstate Thermalization Hypothesis" (2018). *Physics and Astronomy Faculty Publications*. 568.

https://uknowledge.uky.edu/physastron_facpub/568

This Article is brought to you for free and open access by the Physics and Astronomy at UKnowledge. It has been accepted for inclusion in Physics and Astronomy Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Subsystem Eigenstate Thermalization Hypothesis

Digital Object Identifier (DOI)

<https://doi.org/10.1103/PhysRevE.97.012140>

Notes/Citation Information

Published in *Physical Review E*, v. 97, issue 1, 012140, p. 1-7.

©2018 American Physical Society

The copyright holder has granted permission for posting the article here.


Subsystem eigenstate thermalization hypothesis

Anatoly Dymarsky,^{1,2} Nima Lashkari,³ and Hong Liu³

¹*Department of Physics and Astronomy, University of Kentucky, Lexington, Kentucky 40506, USA*

²*Skolkovo Institute of Science and Technology, Skolkovo Innovation Center, Moscow 143026, Russia*

³*Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

 (Received 27 March 2017; revised manuscript received 7 December 2017; published 25 January 2018)

Motivated by the qualitative picture of canonical typicality, we propose a refined formulation of the eigenstate thermalization hypothesis (ETH) for chaotic quantum systems. This formulation, which we refer to as subsystem ETH, is in terms of the reduced density matrix of subsystems. This strong form of ETH outlines the set of observables defined within the subsystem for which it guarantees eigenstate thermalization. We discuss the limits when the size of the subsystem is small or comparable to its complement. In the latter case we outline the way to calculate the leading volume-proportional contribution to the von Neumann and Renyi entanglement entropies. Finally, we provide numerical evidence for the proposal in the case of a one-dimensional Ising spin chain.

DOI: [10.1103/PhysRevE.97.012140](https://doi.org/10.1103/PhysRevE.97.012140)

I. INTRODUCTION AND MAIN RESULTS

During the last two decades there has been significant progress in understanding how quantum statistical physics emerges from the dynamics of an isolated quantum many-body system in a pure state. An important recent development was the realization that a typical pure state, when restricted to a small subsystem, is well approximated by the microcanonical ensemble [1–3]. More explicitly, for a system comprised of a sufficiently small subsystem A and its complement \bar{A} , for any random pure state Ψ from an energy shell $(E, E + \Delta E)$,

$$|\Psi\rangle = \sum_a c_a |E_a\rangle, \quad E_a \in (E, E + \Delta E), \quad (1)$$

the corresponding reduced density matrix $\rho_{\Psi}^A \equiv \text{Tr}_{\bar{A}} |\Psi\rangle\langle\Psi|$ is almost microcanonical. Taking the average $\langle \dots \rangle_{\Psi}$ over all states (1) with respect to the Haar measure one finds [3]

$$\left\| \left\| \rho_{\Psi}^A - \rho_{\text{micro}}^A \right\| \right\|_{\Psi} \leq \frac{1}{2} \frac{d_A}{\sqrt{d_{\Delta E}}}, \quad \|O\| = \frac{1}{2} \text{Tr} \sqrt{OO^\dagger}. \quad (2)$$

Here $\rho_{\text{micro}}^A = \text{Tr}_{\bar{A}} \rho_{\text{micro}}$ is the reduction of the microcanonical density matrix ρ_{micro} associated with the same energy shell $(E, E + \Delta E)$, $d_{\Delta E}$ is the number of energy levels inside it, and $d_A = \dim \mathcal{H}_A$ is the dimension of the Hilbert space of A .

Equation (2) implies that, when the system is sufficiently large, i.e., $\ln d_{\Delta E} \gg \ln d_A$, the subsystem of a typical pure state is well approximated by that of the microcanonical ensemble with an exponential precision. We refer to this mechanism as “canonical typicality” (CT). It is important to note that CT is a purely kinematic statement, and provides no insight into whether or how a nonequilibrium initial state thermalizes [4].

Heuristically, canonical typicality can be understood as a consequence of the entanglement between a sufficiently small subsystem and its complement [3]. While the full system evolves unitarily, a small subsystem can behave thermally as its complement plays the role of a large bath.

Another important development was the so-called eigenstate thermalization hypothesis (ETH) [5–7] which conjectures that a chaotic quantum system in a finitely excited energy eigenstate behaves thermally when probed by few-body operators. More explicitly, for a few-body operator \mathcal{O} , ETH postulates that [8,9]

$$\begin{aligned} \langle E_a | \mathcal{O} | E_b \rangle &= f_{\mathcal{O}}(E) \delta_{ab} + \Omega^{-1/2}(E) r_{ab}, \\ E &= (E_a + E_b)/2, \end{aligned} \quad (3)$$

where $|E_a\rangle$ denotes an energy eigenstate, $f_{\mathcal{O}}(E)$ is a smooth function of E , $\Omega(E) = e^{S(E)}$ is the density of states of the full system, and the fluctuations r_{ab} are of order 1, $r_{ab} \sim O(1)$. The big O here and in what follows refers to the limit when the size of the full system is taken to infinity.

Canonical typicality applies to all systems independent of the Hamiltonian, as opposed to ETH which only concerns chaotic systems, and does not apply to integrable or many-body localized systems. It is a stronger statement, as ETH implies the emergence of the microcanonical ensemble not only for random Ψ , but also for a wider class of states, including the linear combination of a few energy eigenstates.

The fact that ETH applies only to chaotic systems can be heuristically understood from the general picture of CT; only for chaotic systems energy eigenstates are “random enough” to be typical. This perspective thus motivates us to study the properties of the reduced density matrix of a subsystem in an energy eigenstate; see [10–12] for some earlier works.

Now consider a chaotic many-body system in an energy eigenstate $|E_a\rangle$ reduced to a spatial subsystem A which is smaller than its complement \bar{A} . We postulate the *subsystem ETH*:

(i) The reduced density matrix $\rho_a^A = \text{Tr}_{\bar{A}} |E_a\rangle\langle E_a|$ for region A in state $|E_a\rangle$ is exponentially close to some universal density matrix $\rho^A(E)$, which depends smoothly on E ,

$$\left\| \rho_a^A - \rho^A(E = E_a) \right\| \sim O[\Omega^{-1/2}(E_a)]. \quad (4)$$

(ii) The “off-diagonal” matrices $\rho_{ab}^A = \text{Tr}_{\bar{A}} |E_a\rangle\langle E_b|$ are exponentially small,

$$\|\rho_{ab}^A\| \sim O[\Omega^{-1/2}(E)], \quad E_a \neq E_b, \quad E = \frac{1}{2}(E_a + E_b). \quad (5)$$

The pre-exponential factors in (4) and (5) could depend on the size of subsystem A . Importantly, these factors should remain bounded for the fixed A . In the next section, we will give numerical support for the exponential suppression of (i) and (ii) using a spin system. Recently support for (4) and (5) was given in the context of CFTs in [13].

In the thermodynamic limit, i.e., with the system size taken to infinity, $V \rightarrow \infty$, while keeping the size of A and the energy density E/V finite and fixed, it can be readily seen from (i) and (ii) that

$$\|\rho_a^A - \rho_{\text{micro}}^A\| \sim O(\Delta E/E). \quad (6)$$

An implicit assumption here is that $\rho^A(E)$ is well defined in the thermodynamic limit, i.e., it is a function of E/V ¹ and the prefactor in (4) and (5) remains bounded in the limit $V \rightarrow \infty$. Note that while the suppressions in (4) and (5) are exponential in the system size, those in (6) are only power law suppressed.

Using $\|\rho\| = \max_{\mathcal{O}} \text{Tr}(\mathcal{O}\rho)/2$, where maximum is taken over all Hermitian operators of unit norm $\|\mathcal{O}\| = 1$, we conclude from (i) and (ii) that the matrix elements of ρ_a^A and $\rho^A(E = E_a)$ are exponentially close,

$$(\rho_a^A)_{ij} = (\rho^A)_{ij} + O(\Omega^{-1/2}), \quad (\rho_{ab}^A)_{ij} = O(\Omega^{-1/2}). \quad (7)$$

The formulation in (4) and (5) is stronger than the conventional form of ETH. In particular, for systems with an infinite-dimensional local Hilbert space (e.g., with harmonic oscillators at each lattice site) or continuum field theories it guarantees ETH for the particular class of observables, while for other observables ETH may not apply. In particular, subsystem ETH implies the exponential proximity between expectation values in an eigenstate $\langle E_a | \mathcal{O} | E_a \rangle$ and the universal value $f_{\mathcal{O}}(E_a) = \text{Tr}[\mathcal{O}\rho^A(E_a)]$ for any observable \mathcal{O} with the support in A .² This immediately follows from the Cauchy-Schwarz inequality,³

$$\begin{aligned} & \text{Tr} \{ [\rho_a^A - \rho^A(E_a)] \mathcal{O} \} \\ & \leq 2^{1/2} \sqrt{\text{Tr} \{ [\rho_a^A + \rho^A(E_a)] \mathcal{O}^2 \}} \|\rho_a^A - \rho^A(E_a)\|^{1/2}. \end{aligned} \quad (8)$$

Moreover, the subsystem ETH can be applied directly to nonlocal measures which are defined in terms of reduced density matrices, such as entanglement entropy, Renyi entropies, negativity, and so on. See, e.g., [12] for a recent

discussion. In particular, in case of finite-dimensional models it immediately leads to a natural interpretation of thermal entropy as the volume part of the entanglement entropy of a subsystem (see [15,16] for recent discussions). We should caution that when $\dim \mathcal{H}_A$ is infinite, arbitrarily close proximity of density matrices does not automatically imply equality for nonlocal observables. For example, in such cases, higher Renyi entropies for ρ_a^A may be different from those of the microcanonical or other thermal ensembles [13].

In the case of the spin model, for all matrix elements $(\rho^A)_{ij}$, we find strong evidence that r_{aa} of (3) are normally distributed. This is consistent with the heuristic picture of typical $|E_a\rangle$ and r_{ab} being a Gaussian random matrix.

It is tempting to ask whether one could further refine pre-exponential factors in (4) and (5), especially when the subsystem A is macroscopic. Motivated by the A -dependent prefactor in (2) and the average value of the “off-diagonal” matrices (23), it is natural to postulate that the prefactor in (4) and (5) should also be given by

$$\begin{aligned} \|\rho_a^A - \rho^A(E = E_a)\| & \sim O(e^{N_A - S(E)/2}), \\ \|\rho_{ab}^A\| & \sim O(e^{N_A - S(E)/2}), \end{aligned} \quad (9)$$

where N_A denotes the number of effective degrees of freedom in A . For a system of finite dimensional Hilbert space, such as a spin system, e^{N_A} simply corresponds to $d_A = \dim \mathcal{H}_A$, but for a system with an infinite dimensional Hilbert space at each lattice site or a continuum field theory we may view (9) as a definition of effective number of degrees of freedom. For a spin system we will give some numerical evidence for (9) in the next section.

In addition to (6) it is interesting to compare ρ_a^A with the reduced density matrices for other statistical ensembles. Of particular interest are the reduced state on the canonical ensemble for the whole system

$$\rho_C^A = \frac{\text{Tr}_{\bar{A}} e^{-\beta H}}{\text{Tr} e^{-\beta H}}, \quad (10)$$

and the local canonical ensemble for the subsystem A ,

$$\rho_G^A = \frac{e^{-\beta H_A}}{\text{Tr}_A e^{-\beta H_A}}. \quad (11)$$

Here, the Hamiltonian of the subsystem is the restriction of the Hamiltonian $H_A = \text{Tr}_{\bar{A}} H$. In (10), β is to be chosen so that the average energy of the total system is E_a . In (11), β can be interpreted as a local temperature of A (see also [17–19]). There is no canonical choice for β in this case. Below, we choose it to be the same as in (10). In the thermodynamic limit, $V \rightarrow \infty$ with the subsystem A and E/V kept fixed, the standard saddle point approximation argument provides equality between the canonical and the microcanonical ensembles leading to

$$\|\rho_{\text{micro}}^A - \rho_C^A\| = O(V^{-1}) \quad \Rightarrow \quad \|\rho_a^A - \rho_C^A\| = O(V^{-1}), \quad (12)$$

where we have also used (6). The reduced states ρ_C^A and ρ_G^A always remain different at the trace distance level, including thermodynamic limit [17]. Hence,

$$\rho_a^A \neq \rho_G^A, \quad V \rightarrow \infty. \quad (13)$$

¹At a technical level (6) requires a weaker condition of finite Lipschitz constant κ in the thermodynamic limit, $\|\rho^A(E_1) - \rho^A(E_2)\| \leq \kappa|E_1 - E_2|/V$. In (6) we also use that for finitely excited states $E \sim V$.

²In case of the continuous quantum field theory, when the full Hilbert space does not admit a tensor product structure of the Hilbert space of A and of its complement \bar{A} , operators \mathcal{O} should be defined in terms of the net of local operator algebras; see, e.g., [14].

³For any physically sensible observable \mathcal{O} the fluctuations of expectation value $\text{Tr}(\rho_a^A \mathcal{O})$, which are given by $\text{Tr}(\mathcal{O}^2 \rho_a^A)$, must be finite.

Finally, it is interesting to investigate whether (4) and (5) remain true in an alternative thermodynamic limit when the size of subsystem A scales proportionally with the full system. In this limit both the system volume V and the volume V_A for A go to infinity, but we keep the ratio fixed,

$$0 < p = \frac{V_A}{V} < \frac{1}{2}. \quad (14)$$

Note that for any fixed ratio $p < 1/2$ scaling (9) would imply the validity of ETH (4) and (5). In what follows we discuss a weaker version of this statement, which does not rely on (9). When A is scaled to infinity, we expect ρ_a^A to have a semiclassical description. We conjecture that in this limit ρ_a^A will be approaching $\rho^A(E_a)$ at the level of individual matrix elements,

$$(\rho_a^A)_{ij} = [\rho^A(E_a)]_{ij}. \quad (15)$$

Although individual matrix elements will scale as d_A^{-1} and go to zero, (15) is meaningful as it is satisfied with a precision controlled by $\Omega^{-1/2} \sim d^{-1/2} \ll d_A^{-1}$ for all $p < 1/2$. Furthermore, to the leading order in $1/V$, $\rho^A(E_a)$ will be diagonal in the eigenbasis $|\mathcal{E}_i\rangle$ of H_A , with the diagonal elements given by⁴

$$\langle \mathcal{E}_i | \rho^A(E_a) | \mathcal{E}_i \rangle = \langle \mathcal{E}_i | \rho_{\text{mic}}^A | \mathcal{E}_i \rangle = \frac{\Omega_{\bar{A}}(E_a - \mathcal{E}_i)}{\Omega(E_a)}, \quad (16)$$

where $\Omega_{\bar{A}}$ is the density of states of $H_{\bar{A}} = \text{Tr}_A H$. The expression (16) reflects the quasiclassical expectation that the probability to find the subsystem in a state with energy \mathcal{E}_i is proportional to the number of such states. Also for Hamiltonians with local interactions, $H = H_A + H_{\bar{A}}$ up to boundary terms, and in this limit we expect at the level of individual matrix elements

$$(\rho_C^A)_{ij} = (\rho_G^A)_{ij}. \quad (17)$$

As a self-consistency check, using the expression of $(\rho_a^A)_{ij}$ following from (15) and (16), one can calculate $(\rho_C^A)_{ij}$ using saddle point approximation to find that it is indeed equal to $(\rho_G^A)_{ij}$. Finally note that in the limit $V \rightarrow \infty$ with p fixed, $\rho_{\text{micro}}^A \neq \rho_C^A$ and thus we have at the level of individual matrix elements

$$\rho_a^A = \rho_{\text{micro}}^A \neq \rho_C^A = \rho_G^A. \quad (18)$$

The form of the density matrix (16) can be used to evaluate the leading contribution to the entanglement von Neumann and Renyi entropies in terms of the density of states Ω . This is discussed in Appendix Sec. 3. Curiously the leading volume-proportional behavior of the entanglement entropy of ρ_a^A and ρ_G^A is still the same, confirming previous observation of [12], while higher Renyi entropies are different.

In the second part of the paper we provide numerical supports for (4) and (5) as well as (3) and (16) in a one-dimensional spin chain model.

⁴The following form of $\rho_a^A(E_a)$ was previously observed and theoretical justified in [20] in the context of a particular model.

II. NUMERICAL RESULTS

Now we examine hypothesis (4) and (5) of the subsystem formulation of ETH by numerically simulating an Ising spin chain with a transverse and longitudinal magnetic field,

$$H = - \sum_{k=1}^{n-1} \sigma_z^k \otimes \sigma_z^{k+1} + g \sum_{k=1}^n \sigma_x^k + h \sum_{k=1}^n \sigma_z^k. \quad (19)$$

This system is known to be nonintegrable unless one of the coupling constants g or h is zero. We solve the system by exact diagonalization for $g = 1.05$ and various values of h ranging from $h = 0$ to $h = 1$. For this model, the range of the energy spectrum is roughly from $-n$ to n , where n is the total number of spins. The density of states is well approximated by a binomial function; see the Appendix. We will focus on the behavior of $|E_a\rangle$ for E_a near the central value $E_a \simeq 0$ of the spectrum, which corresponds to highly excited states.

We denote by m the number of leftmost consecutive spins which we take to be subsystem A . We introduce the difference between the reduced density matrices for two consecutive energy eigenstates $\Delta\rho_a = (\rho_{a+1}^A - \rho_a^A)/\sqrt{2}$, and define an average variance

$$\sigma_{m,n}^2 = \frac{1}{d_{\Delta E}} \sum_a \text{Tr}(\Delta\rho_a^2). \quad (20)$$

Here the sum is over all energy eigenstates inside the central band $|E_a| \leq \Delta E$, which is taken to be $\Delta E = 0.1n$ and $d_{\Delta E}$ is the total number of states within it. The exponential suppression of $\sigma_{m,n}$ with n is a necessary condition for (4), as follows from the second inequality below:

$$\text{Tr}(\Delta\rho^2) \leq 4\|\Delta\rho\|^2 \leq d_A \text{Tr}(\Delta\rho^2), \quad (21)$$

valid for any Hermitian $\Delta\rho$ supported on \mathcal{H}_A . Numerical results for $\ln(\sigma_{m,n})$ for different m as a function of total system size n are shown in Fig. 1(a). The numerical values are well approximated by a linear fit $\ln(\sigma_{m,n}) = -\alpha_m n + \beta_m$, with β_m increasing with m and the slope α_m for all m being numerically close (within 5% accuracy) to the theoretical value $\ln(2)/2$ suggested by (4).

To confirm that (4) for *each* individual E_a is exponentially small, we examine the maximal value of $\text{Tr}(\Delta\rho_a^2)$ for all E_a within the central band,

$$\mathcal{M}_{m,n} \equiv \max_a \text{Tr}(\Delta\rho_a^2). \quad (22)$$

The dependence of $\mathcal{M}_{m,n}$ for different m, n is shown in Fig. 1(b). We observe that indeed $\mathcal{M}_{m,n}$ is also exponentially suppressed in n .

Now let us examine (5). Similar to (20), we consider the mean variance, averaged over all states E_a . It can be calculated in full generality for any quantum system (see Appendix Sec. 2),

$$\frac{1}{d} \sum_b \text{Tr}[(\rho_{ab}^A)^\dagger \rho_{ab}^A] = \frac{d_A}{d}, \quad (23)$$

where d is the total dimension of the Hilbert space. In the case of spin chain $d_A/d = e^{-(n-m)\ln 2}$. This shows that the averaged $\|\rho_{ab}^A\|$ is always exponentially small, but there remains a possibility that a small number of $\text{Tr}(\rho_{ab}^2)$ for $a \neq b$ are actually




FIG. 1. (a) Values of $\ln(\sigma_{m,n})$ with superimposed linear fit functions $-\alpha_m n + \beta_m$ for $m = 1 \dots 8$, $n = 12 \dots 17$, and $\Delta E = 0.1n$, $g = 1.05$, $h = 0.1$. The slope of linear functions α_m for all m is within 5% close to the theoretical value $\ln(2)/2$. (b) The maximum value of $\text{Tr}(\Delta\rho^2)$ over all eigenstates inside the central band $|E_a| \leq \Delta E = 0.1n$.

not suppressed. This is the case for integrable systems. To eliminate this possibility, we further examine the following quantity:

$$M_m \equiv \max_{|E_a| < \Delta E} \max_b \text{Tr}[(\rho_{ab}^A)^\dagger \rho_{ab}^A], \quad (24)$$

where for a given E_a we first scan all $E_b \neq E_a$ to find the maximal value $L_A(a) \equiv \max_b \text{Tr}[(\rho_{ab}^A)^\dagger \rho_{ab}^A]$, and then find $M_A = \max_a L_A(a)$ by scanning all values of E_a within the window $|E_a| < \Delta E = 0.1n$. The restriction to $|E_a| < \Delta E$ is necessary as ETH is only expected to apply to the finitely excited states, not to the states from the edges of the spectrum. This is manifest from the plot of Fig. 2(a). Figure 2(b) indicates that M_A decreases exponentially with n . This provides strong numerical support for (5).

To study the fluctuations r_{aa} of individual matrix elements of ρ_E^A around the mean value we introduce eigenstates \mathcal{E}_a^A of the local Hamiltonian H_A and define

$$\Delta R_a^{ij} = \frac{1}{\sqrt{2}} (\mathcal{E}_i | \rho_{a+1}^A - \rho_a^A | \mathcal{E}_j). \quad (25)$$

In terms of the fluctuations $R_{ab} = \Omega^{-1/2} r_{ab}$ of (3), ΔR_a is simply the difference $(R_{(a+1)(a+1)} - R_{aa})/\sqrt{2}$. In Fig. 3(a), we show the distribution (histogram) $P(\Delta R)$ for E_a from the central band $|E_a| < \Delta E$ and one particular choice of i, j , and A consisting of $m = 1$ spin. The plot also contains a

superimposed normal distribution (continuous blue line) that is fitted to have the same variance (and the mean value, which is of order $d_{\Delta E}^{-1}$, i.e., exponentially small),

$$\sigma_n^{ij} = \frac{1}{d_{\Delta E}} \sum_a (\Delta R_a^{ij})^2. \quad (26)$$

Figure 3(a) shows that $P(\Delta R)$ is well approximated by the normal distribution. The situation for all other matrix elements for $m = 1, 2, 3$ is very similar.

Numerically, the standard deviation σ_n shows a robust independence of the width of the energy shell ΔE that includes a large number of states. We plot $\ln(\sigma_n)$ as a function of n in Fig. 3(b). We find that σ_n decreases exponentially with the system size n for all matrix elements of ρ_a^A for $m = 1, 2, 3$ and values of h which are not too close to the integrable point $h = 0$. The exponential suppression of ΔR_a follows from the exponential suppression of $|\Delta\rho|$. But (4) does not guarantee that different matrix elements of ρ_a would converge to those of $\rho(E_a)$ with the same rate. Numerics show that the convergence rate $\alpha = d \ln \sigma_n / dn$ is approximately the same, fluctuating around the numerical value $-\ln(2)/2$, for all matrix elements of $\rho_a^{m=1}$; see Fig. 3(b). The behavior for all matrix elements of $\rho_a^{m=2,3}$ is very similar. The numerical proximity of α to $-\ln(2)/2$ provides a strong numerical support for the form of the exponentially suppressed factor in (3), which was originally




FIG. 2. (a) Plot of $L_A(E)$ vs $\epsilon = E/n$ for $n = 15$ and $n = 17$. (b) $M_{m=1,2,3}$ all decrease exponentially with n . Here ΔE is chosen to be equal to $0.1n$ and $h = 0.1$.




FIG. 3. (a) Probability distribution $P(\Delta R)$ of the deviation $\Delta R = \Delta R_a^{11}$ corresponding to the matrix element $\langle \mathcal{E}_1 | \rho_a^{m=1} | \mathcal{E}_1 \rangle$ for $\Delta E = 0.1n$ and $h = 0.1$. It is superimposed with a Gaussian distribution fit. The vertical axis is the number of energy eigenstates within the energy shell $|E_a| < \Delta E$ with a particular value of ΔR . All matrix elements of $\rho_E^{m=1,2,3}$ show almost identical behavior. (b) Linear behavior of $\ln(\sigma_n)$ as a function of system size n for two matrix elements ΔR^{11} and ΔR^{12} for $m = 1$ and $h = 0.1$. Because of the approximate equality $\rho_C \approx \rho_G$ the typical magnitude of the diagonal terms of ρ_a is much larger than the off-diagonal ones. There is no qualitative difference between different matrix elements. Results for $m = 2, 3$ are similar.

introduced in [8]. Provided that $P(\Delta R)$ is well described by normal distribution, the probability of a given R_{aa} to be of order R or larger is given by $1 - \text{Erf}(R/\sqrt{2}\sigma_n) \sim e^{-2^n R^2/R_0^2}$, where R_0 is some constant. If the total number of eigenstates grows as 2^n , the probability of finding an energy eigenstate E_a which does not satisfy ETH and has large R_{aa} is given by $2^n e^{-2^n R^2/R_0^2}$. This probability quickly goes to zero with n , which explains the strong version of ETH recently discussed in [21].

Next, we investigate the prefactor in (4) to test the bound behavior outlined in (9). Namely, we are interested in the dependence of the exponential suppression factor on subsystem size m . To illustrate this behavior we plot $\ln \sigma_{m,n}$ for a fixed value of $n = 17$ and different m in Fig. 4(a). In terms of the spin chain, the bound (9) means the trace distance $\|\rho_a^A - \rho^A(E_a)\|$ should not grow faster than $O(e^{m \ln(2) - n \ln(2)/2})$. This follows from (21) if the second norm $\sqrt{\text{Tr}[\rho_a^A - \rho^A(E_a)]^2}$ is bounded by $O(e^{m \ln(2)/2 - n \ln(2)/2})$. The

actual slope of the linear fit of $\ln(\sigma_{m,n})$ as a function of m is ~ 0.219 . This is substantially smaller than $\ln(2)/2 \simeq 0.347$, providing numerical support for (9).

Finally, we consider the behavior of ρ_a^A when A becomes comparable to \bar{A} to probe the validity of (16) in the regime of fixed p . This is numerically more challenging. Nevertheless, our numerical results are still quite suggestive. We consider subspace A consisting of eight leftmost consecutive spins with $n = 17$ and $h = 0.1$. The numerical results comparing one diagonal matrix element $\langle \mathcal{E}_1 | \rho_a^A | \mathcal{E}_1 \rangle$ corresponding to the lowest energy level of H_A is given in Fig. 4(b). It shows that ρ_a^A follows (16) pretty well while it differs from $\rho_C^A \approx \rho_G^A$ significantly.

ACKNOWLEDGMENTS

This work was supported by funds provided by MIT-Skoltech Initiative. We would like to thank the University of




FIG. 4. (a) Dependence of $\ln \sigma_{m,17}$ on the subsystem size m with the superimposed linear fit $-4.455 + 0.219m$. (b) Comparison of matrix elements of $\rho_a^A, \rho_C^A, \rho_G^A$ and the quasiclassical result (16) which we refer to as ρ_Q^A . Blue dots are matrix elements $\langle \mathcal{E}_1 | \rho_a^{m=8} | \mathcal{E}_1 \rangle$ as a function of energy per site $\epsilon = E_a/n$ for $h = 0.1$ and $n = 17$. We see that $\langle \mathcal{E}_1 | \rho_a^{m=8} | \mathcal{E}_1 \rangle$ follows the semiclassical result $\langle \mathcal{E}_1 | \rho_Q^{m=8} | \mathcal{E}_1 \rangle$ as given by (16) well, while it differs significantly from $\langle \mathcal{E}_1 | \rho_C^{m=8} | \mathcal{E}_1 \rangle \approx \langle \mathcal{E}_1 | \rho_G^{m=8} | \mathcal{E}_1 \rangle$, which lie on top of each other. Quasiclassical result (16) is calculated using density of states Ω specified in Appendix Sec. 1. Other matrix elements show similar behavior.




FIG. 5. The density of states of the spin chain system for $g = 1.05$, $h = 0.1$, $n = 17$. The horizontal axis is energy per site $\epsilon = E/n$. The yellow bars which fill the plot are the histogram for the density of states calculated using direct diagonalization. The blue solid line is a theoretical fit by the binomial distribution function (A2) with $\kappa \approx 0.3489$; see Appendix Sec. 1.

Kentucky Center for Computational Sciences for computing time on the Lipscomb High Performance Computing Cluster.

APPENDIX

1. Density of states

A spin chain without nearest neighbor interactions exhibits a degenerate spectrum with the level spacing of order 1. In this case the density of states is given by the binomial distribution. Once the nearest neighbor interaction term is introduced, the spectrum becomes nondegenerate with the exponentially small level spacing. In this case the density of states can be described by a smooth function $\Omega(E)$, which would be reasonably approximated by the binomial distribution. For the spin chain in question,

$$H = - \sum_{i=1}^{n-1} \sigma_z^i \otimes \sigma_z^{i+1} + g \sum_{i=1}^n \sigma_x^i + h \sum_{i=1}^n \sigma_z^i, \quad (\text{A1})$$

we start with the binomial distribution,

$$\Omega_n(E) = \frac{\kappa n!}{(n/2 - \kappa E)!(n/2 + \kappa E)!}, \quad (\text{A2})$$

for some κ , and notice that it is properly normalized for any value of κ with an exponential precision, $\int dE \Omega_n(E) \simeq 2^n$. We fix the parameter κ using the value of the second moment,

$$\int dE E^2 \Omega_n(E) \simeq 2^{n-2} n \kappa^{-2} = \text{Tr } H^2. \quad (\text{A3})$$

The latter could be calculated exactly from (A1) yielding $\kappa = \frac{1}{2}(g^2 + h^2 + 1 - 1/n)^{-1/2}$. The resulting density of states provides a very accurate fit for the exact numerical result as depicted in Fig. 5. The expression for density of states (A2) is used to calculate $\langle \mathcal{E}_i | \rho^A | \mathcal{E}_i \rangle$ from (16) in Fig. 4(b).

2. Variance

Consider the variance

$$\Sigma_a^2 = \frac{1}{d} \sum_b \text{Tr}((\rho_{ab}^A)^\dagger \rho_{ab}^A) \quad (\text{A4})$$

for some fixed a and d being the dimension of the full Hilbert space. Since $|E_a\rangle$ is a complete basis,

$$\sum_b \langle E_b | \Psi_1 \rangle \langle \Psi_2 | E_b \rangle = \langle \Psi_2 | \Psi_1 \rangle. \quad (\text{A5})$$

Now let us introduce a basis in the Hilbert space $|i, \bar{j}\rangle = |i\rangle \otimes |\bar{j}\rangle$ associated with the decomposition $\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_{\bar{A}}$. Then

$$(\rho_{ab}^A)_{ij} = \langle i | \rho_{ab}^A | j \rangle = \sum_{\bar{k}} \langle i, \bar{k} | E_a \rangle \langle E_b | j, \bar{k} \rangle \quad (\text{A6})$$

and

$$\Sigma_a^2 = \frac{1}{d} \sum_b \sum_{i,j} \sum_{\bar{k}, \bar{\ell}} \langle i, \bar{k} | E_a \rangle \langle E_b | j, \bar{k} \rangle \langle j, \bar{\ell} | E_b \rangle \langle E_a | i, \bar{\ell} \rangle.$$

Now we use (A5) to get

$$\Sigma_a^2 = \frac{d_A}{d} \sum_i \sum_{\bar{j}} \langle E_a | i, \bar{j} \rangle \langle i, \bar{j} | E_a \rangle = \frac{d_A}{d}. \quad (\text{A7})$$

3. Semiclassical expression

We now discuss the properties of (16),

$$\langle \mathcal{E}_i | \rho_a^A | \mathcal{E}_i \rangle = \langle \mathcal{E}_i | \rho_{\text{micro}}^A | \mathcal{E}_i \rangle = \frac{\Omega_{\bar{A}}(E_a - \mathcal{E}_i)}{\Omega(E_a)}, \quad (\text{A8})$$

in the limit when

$$0 < p = \frac{V_A}{V} < \frac{1}{2} \quad (\text{A9})$$

is kept fixed and volume $V \rightarrow \infty$. In particular we show that at the leading order in $1/V$ the Von Neumann entropy associated with ρ_a^A , which is given by (A8), is the same as for ρ_G^A , despite the inequality

$$\rho_a^A = \rho_{\text{micro}}^A \neq \rho_C^A = \rho_G^A. \quad (\text{A10})$$

In the limit $V_A \rightarrow \infty$ we can treat the energy levels \mathcal{E}_i of A as a continuous variable \mathcal{E} , in terms of which

$$\Omega(E_a) = \int d\mathcal{E} \Omega_A(\mathcal{E}) \Omega_{\bar{A}}(E_a - \mathcal{E}), \quad (\text{A11})$$

where Ω_A is the density of states for A . Now introduce

$$\ln \Omega_A \equiv \mathcal{S}_A, \quad \ln \Omega_{\bar{A}} \equiv \mathcal{S}_{\bar{A}}, \quad \ln \Omega \equiv \mathcal{S} \quad (\text{A12})$$

with the conventional expectation that the density of states grows exponentially with the volume,

$$\mathcal{S}_A \propto V_A, \quad \mathcal{S}_{\bar{A}} \propto V_{\bar{A}}, \quad \mathcal{S} \propto V. \quad (\text{A13})$$

Since both \mathcal{S}_A and $\mathcal{S}_{\bar{A}}$ are proportional to V we can use the saddle point approximation in (A11) to obtain

$$\mathcal{S}(E) = \mathcal{S}_A(\bar{\mathcal{E}}_A) + \mathcal{S}_{\bar{A}}(\bar{\mathcal{E}}_{\bar{A}}), \quad (\text{A14})$$

where $\bar{\mathcal{E}}_A$ and $\bar{\mathcal{E}}_{\bar{A}}$ are determined by

$$\bar{\mathcal{E}}_A + \bar{\mathcal{E}}_{\bar{A}} = E_a, \quad \left. \frac{\partial \mathcal{S}_A}{\partial \mathcal{E}} \right|_{\bar{\mathcal{E}}_A} = \left. \frac{\partial \mathcal{S}_{\bar{A}}}{\partial \mathcal{E}} \right|_{\bar{\mathcal{E}}_{\bar{A}}}. \quad (\text{A15})$$

Using saddle point approximation for the canonical ensemble of the whole system we recover the conventional

relation between the inverse temperature β and the mean energy E ,

$$\beta = \frac{\partial \mathcal{S}(E_a)}{\partial E}. \quad (\text{A16})$$

Together with (A14) and (A15) this implies

$$\beta = \frac{\partial \mathcal{S}_A}{\partial \mathcal{E}} \Big|_{\bar{\mathcal{E}}_A} = \frac{\partial \mathcal{S}_{\bar{A}}}{\partial \mathcal{E}} \Big|_{\bar{\mathcal{E}}_A}. \quad (\text{A17})$$

Then it follows in a standard way that the entropy S_G^A associated with the diagonal density matrix ρ_G^A

$$\langle \mathcal{E} | \rho_G^A | \mathcal{E} \rangle \simeq e^{-\beta(\mathcal{E} - \bar{\mathcal{E}}_A) - S_A(\bar{\mathcal{E}}_A)} \quad (\text{A18})$$

is simply $S_G^A = S_A(\bar{\mathcal{E}}_A)$.

With the help of (A14) one can rewrite (16) and (A8) as follows:

$$\langle \mathcal{E} | \rho_a^A | \mathcal{E} \rangle \simeq e^{S_{\bar{A}}(E - \mathcal{E}) - S_{\bar{A}}(E - \bar{\mathcal{E}}_A) - S_A(\bar{\mathcal{E}}_A)}, \quad (\text{A19})$$

while off-diagonal matrix elements are negligible. Then the corresponding entropy S_a^A is given by

$$S_a^A = -\text{Tr}_A \rho_a^A \ln \rho_a^A = S_A(\bar{\mathcal{E}}_A) = S_G^A. \quad (\text{A20})$$

Applying saddle point approximation to powers of (A18) and (A19) one can readily calculate the leading volume-proportional contribution to Renyi entropies for ρ_a^A and ρ_G^A and see that they are different. This is consistent with the numerical results of [12].

-
- [1] J. Gemmer, A. Otte, and G. Mahler, Quantum Approach to a Derivation of the Second Law of Thermodynamics, *Phys. Rev. Lett.* **86**, 1927 (2001).
- [2] S. Goldstein, J. L. Lebowitz, R. Tumulka, and N. Zanghi, Canonical Typicality, *Phys. Rev. Lett.* **96**, 050403 (2006).
- [3] S. Popescu, A. J. Short, and A. Winter, Entanglement and the foundations of statistical mechanics, *Nat. Phys.* **2**, 754 (2006).
- [4] Nevertheless some general statements about an approach to the equilibrium can be made; see P. Reimann, Foundation of Statistical Mechanics under Experimentally Realistic Conditions, *Phys. Rev. Lett.* **101**, 190403 (2008); N. Linden, S. Popescu, A. J. Short, and A. Winter, Quantum mechanical evolution towards thermal equilibrium, *Phys. Rev. E* **79**, 061103 (2009); On the speed of fluctuations around thermodynamic equilibrium, *New J. Phys.* **12**, 055021 (2010); S. Goldstein, J. Lebowitz, C. Mastrodonato, R. Tumulka, and N. Zanghi, Approach to thermal equilibrium of macroscopic quantum systems, *Phys. Rev. E* **81**, 011109 (2010).
- [5] J. M. Deutsch, Quantum statistical mechanics in a closed system, *Phys. Rev. A* **43**, 2046 (1991).
- [6] M. Srednicki, Chaos and quantum thermalization, *Phys. Rev. E* **50**, 888 (1994).
- [7] M. Rigol, V. Dunjko, and M. Olshanii, Thermalization and its mechanism for generic isolated quantum systems, *Nature (London)* **452**, 854 (2008).
- [8] M. Srednicki, The approach to thermal equilibrium in quantized chaotic systems, *J. Phys. A: Math. Gen.* **32**, 1163 (1999).
- [9] L. D'Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics, *Adv. Phys.* **65**, 239 (2016).
- [10] G. De Palma, A. Serafini, V. Giovannetti, and M. Cramer, Necessity of Eigenstate Thermalization, *Phys. Rev. Lett.* **115**, 220401 (2015).
- [11] H.-H. Lai and K. Yang, Entanglement entropy scaling laws and eigenstate typicality in free fermion systems, *Phys. Rev. B* **91**, 081110(R) (2015).
- [12] J. R. Garrison and T. Grover, Does a single eigenstate encode the full Hamiltonian? [arXiv:1503.00729](https://arxiv.org/abs/1503.00729).
- [13] N. Lashkari, A. Dymarsky, and H. Liu, Eigenstate thermalization hypothesis in conformal field theory, [arXiv:1610.00302](https://arxiv.org/abs/1610.00302).
- [14] R. Haag, *Local Quantum Physics: Fields, Particles, Algebras* (Springer Science & Business Media, New York, 2012).
- [15] J. M. Deutsch, Thermodynamic entropy of a many-body energy eigenstate, *New J. Phys.* **12**, 075021 (2010).
- [16] J. Deutsch, H. Li, and A. Sharma, Microscopic origin of thermodynamic entropy in isolated systems, *Phys. Rev. E* **87**, 042135 (2013).
- [17] A. Ferraro, A. García-Saez, and A. Acín, Intensive temperature and quantum correlations for refined quantum measurements, *Europhys. Lett.* **98**, 10009 (2012).
- [18] M. Kliesch, C. Gogolin, M. J. Kastoryano, A. Riera, and J. Eisert, Locality of Temperature, *Phys. Rev. X* **4**, 031019 (2014).
- [19] S. Hernández-Santana, A. Riera, K. Hovhannisyanyan, M. Perarnau-Llobet, L. Tagliacozzo, and A. Acín, Locality of temperature in spin chains, *New J. Phys.* **17**, 085007 (2015).
- [20] H. Tasaki, From Quantum Dynamics to the Canonical Distribution: General Picture and a Rigorous Example, *Phys. Rev. Lett.* **80**, 1373 (1998).
- [21] H. Kim, T. N. Ikeda, and D. A. Huse, Testing whether all eigenstates obey the eigenstate thermalization hypothesis, *Phys. Rev. E* **90**, 052105 (2014).