

Successive Cambia: A Developmental Oddity or an Adaptive Structure?

Elisabeth M. R. Robert^{1,2*}, Nele Schmitz^{1,2}, Ilse Boeren^{2,3}, Tess Driessens¹, Kristof Herremans¹, Johan De Mey⁴, Elke Van de Castele^{5,6}, Hans Beeckman², Nico Koedam¹

1 Laboratory for Plant Biology and Nature Management (APNA), Vrije Universiteit Brussel (VUB), Brussels, Belgium, **2** Laboratory for Wood Biology and Xylarium, Royal Museum for Central Africa (RMCA), Tervuren, Belgium, **3** Institute of Botany (210), University of Stuttgart-Hohenheim, Stuttgart, Germany, **4** Radiology, Universitair Ziekenhuis Brussel, Brussels, Belgium, **5** SkyScan NV, Kontich, Belgium, **6** Vision Lab, Universiteit Antwerpen (UA), Wilrijk, Belgium

Abstract

Background: Secondary growth by successive cambia is a rare phenomenon in woody plant species. Only few plant species, within different phylogenetic clades, have secondary growth by more than one vascular cambium. Often, these successive cambia are organised concentrically. In the mangrove genus *Avicennia* however, the successive cambia seem to have a more complex organisation. This study aimed (i) at understanding the development of successive cambia by giving a three-dimensional description of the hydraulic architecture of *Avicennia* and (ii) at unveiling the possible adaptive nature of growth by successive cambia through a study of the ecological distribution of plant species with concentric internal phloem.

Results: *Avicennia* had a complex network of non-cylindrical wood patches, the complexity of which increased with more stressful ecological conditions. As internal phloem has been suggested to play a role in water storage and embolism repair, the spatial organisation of *Avicennia* wood could provide advantages in the ecologically stressful conditions species of this mangrove genus are growing in. Furthermore, we could observe that 84.9% of the woody shrub and tree species with concentric internal phloem occurred in either dry or saline environments strengthening the hypothesis that successive cambia provide the necessary advantages for survival in harsh environmental conditions.

Conclusions: Successive cambia are an ecologically important characteristic, which seems strongly related with water-limited environments.

Citation: Robert EMR, Schmitz N, Boeren I, Driessens T, Herremans K, et al. (2011) Successive Cambia: A Developmental Oddity or an Adaptive Structure? PLoS ONE 6(1): e16558. doi:10.1371/journal.pone.0016558

Editor: Wendy Peer, Purdue University, United States of America

Received: September 3, 2010; **Accepted:** December 27, 2010; **Published:** January 31, 2011

Copyright: © 2011 Robert et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The study was financially supported by the Institute for the Promotion of Innovation through Science and Technology in Flanders (IWT-Vlaanderen), the National Fund for Scientific Research (FWO, Belgium), the King Leopold III Fund for Nature Exploration and Conservation, Belgium, and the AFRODENDRO project of the Royal Museum for Central Africa (RMCA), Tervuren. The funders had no role in study design, content of the study, decision to publish or determination of the conclusions.

Competing Interests: Elke Van de Castele was, at the time the study was conducted, employed by Skyscan NV, Kontich, Belgium, which offered free micro-CT scanning. There is no other affiliation with Skyscan NV than scientific contacts. The input of the company was of a purely academic and technical nature. The company in no way obstructs publication and sharing of the data used for this publication. Herewith the authors confirm that this does not alter their adherence to all the PLoS ONE policies on sharing data and material, as detailed in the PLoS Policy on Declaration and Evaluation of Competing Interests.

* E-mail: erobert@vub.ac.be

Introduction

Expansion in girth in most vascular plant species is the result of the meristematic activity of one cylindrical vascular cambium, producing xylem towards the inner and phloem towards the outer part of the plant stem (Fig. 1a) [1]. However, some plant species (out of 34 families according to the taxonomy used [1]) have not one but several successive cambia causing the secondary growth and the resulting wood anatomy to be different from that of plants with only one vascular cambium (Fig. 1b). In the stem of plants with successive cambia a sequence of vascular cambia can be found, each responsible for the production of secondary xylem inwards and secondary phloem outwards (Fig. 1b) [2]. The cambia can literally develop successively but several cambia can also develop simultaneously [3]. This results in plants from which stem discs show a succession of dark coloured xylem tissue bands and pale coloured phloem tissue bands.

Special forms of secondary wood formation are typical for lianas and vines [e.g. 1,4,5,6,7], though the phenomenon of successive cambia is also found in a select group of herb, shrub and tree species [e.g. 1,8,9,10,11,12] coming from different phylogenetic clades [1,2,12]. The genus *Avicennia* is the only mangrove tree genus showing the developmental oddity of secondary growth by successive cambia. Xylem and phloem tissue in *Avicennia* are found to be non-concentric (Fig. 1c) [e.g. 3,13,14,15,16,17,18] and form an intricate three-dimensional network instead of separate concentric cylinders [3,13,15]. Detailed and four dimensional - the three spatial dimensions and time - observations of these xylem and phloem networks are lacking, though highly necessary to understand *Avicennia's* growth pattern and the secondary growth of trees with successive cambia in general.

From a study in two dimensions, the nature of secondary growth in *Avicennia marina* has been suggested to be patch-like, with active growth displacing around the stem circumference with time [3]. In




Figure 1. Spatial structure of the xylem and phloem tissue in *Avicennia* in comparison to other trees. Schematic view of a stem disc from (a) a tree with only one cambium, (b) a tree with successive cambia organised in concentric cylinders, giving rise to a stem disc with concentric circles of xylem tissue, phloem tissue and cambium circles and (c) *Avicennia*, having a reticulate organisation of its cambia and transport structure. The spatial structure of *Avicennia* is depicted by smoothed surface images after three-dimensional reconstruction in Reconstruct [42] (d,e). Only a small part of the xylem tissue has been visualised. It can be seen that *Avicennia* had a complex network of xylem patches that fused at certain heights of the tree and joined different patches at other parts of the stem. The connections between xylem patches changed rapidly with vertical distance. This is shown by the white arrowheads in the four serial micro-CT-scan images of an outside zone of an *Avicennia marina* stem disc, produced by a SkyScan 1172 scanner (g-k). Each section is 150 µm apart from the previous. On the micro-CT-images (g-k) the wide light grey bands are xylem tissue while the dark small strings are phloem tissue. The complexity of the internal structure can be expressed by the sum of the locations of free ending and branched phloem tissue (f) per stem disc surface area.
doi:10.1371/journal.pone.0016558.g001

Avicennia, cambium differentiation was found to be internally controlled [13,14,19], with indications for an interaction with the micro-environmental conditions such as soil water salinity [3]. What triggers the development of new vascular cambia however remains unclear. In the harsh and changing environmental conditions of the mangrove habitat, the secondary growth of *Avicennia marina* and the resulting pattern of vascular tissue have been proposed to offer a functional advantage for the water transport. While the non-lignified and more thin-walled phloem tissue in between the more rigid xylem tissue increases the mechanical flexibility of lianas [1,2,20], it can also provide water and photosynthate supply [3,13,21,22,23,24] for plants growing under water stress. The living phloem tissue in at least the outer part of the stem could play an important role in embolism repair [21,22,23,24,25,26]. In addition, beneficial growth conditions can

be exploited by simultaneous activity of more than one cambium or by alternating cambial activity around the stem disc. In this way, wood segments with different wood anatomical characteristics – and thus different physiological possibilities - can be created around the stem disc.

In this study we aimed at clarifying the three-dimensional structure of the xylem and phloem tissue of the mangrove genus *Avicennia* through (micro-)CT-scanning. X-ray (micro-)CT scanning allows a non-destructive visualization of the internal structures at high resolution. In order to differentiate between connected structures a difference in X-ray density is required. Since xylem is denser than phloem tissue, this technique is appropriate to study the 3D structure. We further investigated the effect of the environment on the spatial organisation of this structure to unveil aspects of the remarkable adaptation of

Avicennia to the mangrove ecosystem. In order to widen the scope of our conclusions, we tested the relation between habitats that are susceptible to water stress and the presence of woody species with concentric internal phloem. As successive cambia and the resulting internal phloem rings are thought to bring physiological advantage in harsh environmental conditions, we hypothesized that the major part of the woody species with internal concentric phloem are growing in habitats that are physiologically harsh either by drought or salt stress.

Results

The Three-Dimensional Structure of *Avicennia's* Transport System

A reticulate xylem structure, consisting of xylem patches joined in horizontal as well as in vertical direction (Fig. 1c–k), was found consistently in all species and tree samples studied. Within this reticulate system, the connections between xylem zones (patches) changed rapidly with height (Fig. 1h–k) creating a complex network of xylem patches that fused at certain heights of the tree but separated to join different patches in other parts of the tree stem (Fig. 1d–e). In small zones of wood, it could be observed that vessels, although in the same organisation with respect to each other, were more or less scattered depending on height. The remaining space was filled with fibre cells. Moreover, xylem volumes with their corresponding phloem could be found to belong to one growth patch at a certain tree height, while being part of another one at another height (Fig. 1h–k). Therefore phloem and xylem parts that were not connected and part of different growth patches at a certain place in the tree stem lined up lower or higher in the tree.

Phloem was observed to have the same reticulate structure as the xylem network: in most parts of the stem discs studied phloem did not form concentric rings but an assembly of free ending strings and branched circles (Fig. 1f–k). Considering this in three dimensions, we could recognize free ending sheets and branched cylinders, the three-dimensional equivalents of free ending tissue and branched circles respectively (Fig. 1d,e). At certain heights of

the tree stems, phloem was found to exist in very small portions that were not connected with the surrounding phloem network at that same height. In these parts of the tree a continuous circular band of phloem around the stem circumference was lacking. In a part of the stem of the *A. germinans* individual from Benin (Tw50689, Table 1) small isolated patches of xylem completely surrounded by phloem tissue were observed on the transverse CT-images. They were at least 4.56 cm in height.

Vessels were found to be rather straight at distances of less than 6.5 mm. However, we observed, at zones of the stem disc with a highly reticulate transport system, that some vessels were part of a certain wood patch at a certain height of the tree but were part of wood volumes that were non-existing at this height, only a few hundred micrometers higher in the tree.

The ratio of ending to branched phloem (Fig. 1f) depended on species ($H = 19.64$, $p < 0.001$, Kruskal-Wallis test) but not on the ecological conditions of the growing site (*A. marina* – Kenya; $H = 2.04$, $p > 0.05$, Kruskal-Wallis test). The species of the eastern biogeographic mangrove region (Indo-West Pacific and East Africa) had a significantly lower ratio than *A. germinans* of the western biogeographic mangrove region (America and West Africa) ($U = 32.50$, $p < 0.001$, Mann-Whitney U test, Fig. 2a). On the contrary, the level of branching of the xylem and phloem network, expressed as the number of points were the growth segments are not concentric per surface area, did not depend on species ($H = 1.99$, $p > 0.05$, Kruskal-Wallis test) but on growing site ($H = 16.67$, $p < 0.001$, Kruskal-Wallis test, Fig. 2b). The lowest values were found for trees growing in the study site with the lowest salinity and the highest inundation frequency (Table 2). There was a significant difference in ratio of phloem surface area to wood surface area in *A. officinalis* as compared to *A. marina* and *A. germinans* ($H = 21.49$, $p < 0.001$, Kruskal-Wallis test, Fig. 2c). Within *A. marina* the growing site had a significant effect on the phloem to xylem ratio ($H = 19.83$, $p < 0.001$, Kruskal-Wallis test, Fig. 2d).

Furthermore, tree height had a significant effect on the branching of the xylem and phloem tissue ($H = 38.63$, $p < 0.001$, Kruskal-Wallis test): from the base to the upper crown of the tree,

Table 1. Overview of the *Avicennia* samples used for the CT-analysis and micro-CT- analysis.

species	collection number	country	location	biogeographical region	analysis
<i>A. marina</i>	Tw58927-9	Kenya	Gazi Bay – site 1	eastern	CT
	Tw58916, 18, 19	Kenya	Gazi Bay – site 2	eastern	CT
	Tw58937, 38, 41	Kenya	Gazi Bay – site 3	eastern	CT
	Tw60819	Kenya	Gazi Bay – site 3	eastern	CT*
	Tw60820	Kenya	Gazi Bay – site 2	eastern	CT**
	Tw60821	Kenya	Gazi Bay	eastern	micro-CT
	Tw42907	Union of the Comoros	Ngasidja	eastern	CT
	Tw57257	Sri Lanka	Rekawa	eastern	CT
<i>A. officinalis</i>	Tw57246-8	Sri Lanka	Pambala	eastern	CT
	Tw57255	Sri Lanka	Rekawa	eastern	CT
<i>A. germinans</i>	Tw50689	Benin	Cotonou	western	CT
	Tw57679	Democratic Republic of Congo	Ile Bula	western	CT
	Tw55845	Democratic Republic of São Tomé and Príncipe	Lagoa Azul	western	CT

Samples were selected from the xylarium of the Royal Museum of Central Africa in Tervuren (Belgium) or collected in the field (Gazi Bay – Kenya).

* (root sample),

** (root sample and samples at different heights).

doi:10.1371/journal.pone.0016558.t001




Figure 2. Characteristics of the spatial network of the transport system in *Avicennia*. (a) Ratio of ending to branched phloem in function of the biogeographical mangrove regions; (b) Degree of branching of the transport network of *Avicennia marina* in function of the study site (Table 2); (c–d) Ratio of phloem surface area to wood surface area in function of the mangrove species (c) and in function of the study site (Table 2) for *A. marina* (d). Lines: medians. Different letters indicate significant differences. doi:10.1371/journal.pone.0016558.g002

the degree of branching of the xylem and phloem tissue was increasing (Fig. 3). In the roots however, the degree of branching was similar to that in the crown of the tree (Fig. 3). In contrast, no trend with tree height could be found in the ratio of ending to branched phloem or in the phloem to xylem ratio.

Distribution of concentric internal phloem species according to drought stress intensity

From the investigated woody species with concentric internal phloem, 39.6% could be found in habitats characterised with drought or salt stress while 60.4% could be found in rather wet habitats (Table 3). Then, again, the abundance of lianas was much higher in the non-stressed compared to the drought stressed category (49.5% versus 4.1% respectively) (Table 3). If lianas were excluded, 76.7% of the species was growing in dry conditions while 23.3% was growing in a wet, non-saline environment (Table 3). Ten of the fourteen shrub species growing without drought stress could also have a liana habitus. Excluding them was giving a balance of 84.9% of species in a dry and/or saline environment versus 15.1% in wet, non-saline conditions.

Discussion

The three-dimensional structure of *Avicennia* wood

Seen in three dimensions, the transport tissues in *Avicennia* had a network-like structure (Fig. 1c–k). This network of xylem and phloem tissue, extensively connected in horizontal and vertical direction, confirmed the existence of the structure proposed by Schmitz *et al.* [3,13] for *A. marina* as well as with the structure proposed by Zamski [15] who studied young *A. resinifera* and *A. germinans* trees. Our study however, added that a similar three-dimensional structure systematically occurred in stems of different *Avicennia* species from different origins, and in roots, branches and stems from *A. marina* (Table 1). Therefore, the structure could be considered a general characteristic of the mangrove genus *Avicennia*. Furthermore we added that the observed network was highly variable with tree height and position in the tree for all species studied as the connection pattern could change within a few micrometers only (Fig. 1d–e, h–k).

Similar reticulate three-dimensional structures have, as far as we are aware, been proposed (i) in small stems of *Anabasis articulata* and *Kochia indica* [11], (ii) in roots of *Beta vulgaris* [27] and (iii) in branches and stems of *Dalbergia paniculata* [28] having concentric

Table 2. Stand characteristics of *Avicennia marina* study sites in Gazi Bay, Kenya [3].

Location	°S	°E	Soil Water Nutrients ($10^{-3} \mu\text{mol}/\text{cm}^3$ soil)						P			LAI	Soil texture	Soil Water Salinity (‰)			Inundation class
			NO ₃ ⁻			NH ₄ ⁺			mean	min	max			mean	min	max	
			mean	min	max	mean	min	max	mean	min	max			mean	min	max	
site 1	4°24'54.0"	39°30'42.0"	0.5	0.2	0.6	3	0	5	1.4	0.9	1.8	1.42	silty clay	34.6	20.9	46.0	I
site 2	4°25'55.5"	39°30'34.5"	0.15	0.10	0.24	*	*	*	0.2	0.1	0.3	0.23	loamy sand	66.1	38.0	86.0	III
site 3	4°25'16.0"	39°30'27.0"	2	0	6	10	7	14	0.1	0	0.3	1.18	clay loam	63.3	40.0	79.9	III

Soil water for nutrient and salinity analyses was taken at about 25 cm depth. Inundation classes are defined according to Tomlinson [44].

P (soluble reactive phosphor), LAI (Leaf Area Index).

* (no data records).

doi:10.1371/journal.pone.0016558.t002

rings of successive cambia. Furthermore, this structure has been observed in branches of *Bougainvillea spectabilis* [29], in stems of *Phytolacca dioica* [30] and in small stems of *Atriplex halimus* [8,11] and *Haloxylon salicornicum* [11] having concentric rings of collateral vascular bundles. With the exception of Fahn *et al.* [8,11] and Zamski 1979 [15], three-dimensional visualisations and characterisations of these structures are lacking. In general, a three-dimensional way of interpreting internal phloem is not yet developed, although the articles of Fahn and Zamski were published in 1967, 1979 and 1986 already. This lack of spatial insight is obstructing in depth discussions on the ecological and developmental implications of the observed reticulate structures.

Observation of transversal sections of other species in which successive cambia have been described as concentric rings [e.g. 27,31,32] showed that many more species potentially have a complex network of transport tissues. Connections between rings could already be seen in two-dimensional observations. Furthermore, in the stem of *Combretum nigricans*, the diffuse internal phloem, seen as scattered patches on transversal stem sections, was also observed to be organised as a network if watched in three dimensions [33]. This only strengthened the demand for three-dimensional observations of plant anatomy and morphology and for wood anatomical studies with three-dimensional approaches of scientific questions.

Within the observed structure, vessels were criss-crossing different xylem patches along the tree stem. Since each xylem patch is formed by another (part of the) cambium, vessel elements of the same vessel were not always derived from one cambial zone only. To assure the longitudinal alignment of vessel elements, a xylogenetic signal might be given by the developing vessel to an undifferentiated cell in the cambial zone of a contiguous xylem patch via parenchyma-vessel pits [34,35,36].

The complex reticulate structure raised questions about its formation and functioning. What type of cambium can give rise to this kind of spatial structure? How does such a cambium originate? Is there a different system of molecular signalling in these non-continuous tissues than in a cylindrical cambium? Has this kind of organisation a functional significance? Answers to these questions touch upon our fundamental understanding of the hydraulic architecture and its ontogeny in plants and in trees in particular and may reveal unknown mechanisms of plant growth and functioning.

Looking at the visualized three-dimensional structure of the xylem and phloem tissue in *Avicennia*, we would expect the cambia in *Avicennia* to have the same reticulate structure as the xylem and phloem tissue, and, in that sense, be non-continuous, either in space or in time. Speculating about the nature of such cambia, we could define two possibilities: cambia that look like *fishnet stockings* or cambia that are *broken cylinders* (Fig. 4). In the first model (Fig. 4a), we imagined multiple layers of cambium sheets that are discontinuous along as well as around the tree stem and that are possibly connected, while in the second one (Fig. 4b), continuous cambium sheets have the ability to break up longitudinally and/or radially at certain locations, after formation. Both models have to be considered together with the possibility of cambia to line up radially and longitudinally once active and of different growth speeds in distinct parts of the tree as has been already proposed by Schmitz *et al.* [3].

The first as well as the second model on the three-dimensional structure and functioning of *Avicennia's* cambia are both compatible with the proposed parenchymatic origin of new cambia in *Avicennia* [3,13,14,15,18,30,37]. In the *fishnet stockings* model the parenchyma cells of the assumed cambial cylinder do not all dedifferentiate to meristematic cells, giving rise to the suggested




Figure 3. Degree of branching of the internal transport structure of *Avicennia* in function of tree height. Data are from one *Avicennia marina* tree from Gazi Bay (Kenya). Lines: medians; W: main root of the same tree; W*: main root of another *A. marina* tree from a different study site in the same mangrove forest. doi:10.1371/journal.pone.0016558.g003

reticulate structure. This has already been proposed for *Dalbergia paniculata*, a tree species with successive cambia [28]. In the *broken cylinders* model on the other hand, modifications of the cambia, initially organised in concentric cylinders, would occur after dedifferentiation. In reality, a combination of the two proposed models is not excluded. In that case, cambia would be reticulate while formed as well as able to break at certain locations. However, only growth experiments on high spatial and temporal resolution can illuminate the exact ontology of the transport system in *Avicennia*.

Several authors report a number of simultaneously active cambia in plant species with successive cambia [3,13,15,32,38]. We should not picture different cambial zones as separate unities only because they seem to be unconnected on transverse (thin) sections. With the three-dimensional structure in mind, different zones of the cambium in different rings might be part of one and the same connected cambial zone. In that sense, the cambia, as well as the resulting transporting tissues have to be considered as

co-functioning in a unit, as has been proposed for the rings of transport tissue in *B. spectabilis* by Zamski [29].

The Ecological Advantage of Successive Cambia

Within *Avicennia* we observed a higher degree of branching in the spatial structure of the transport system in trees growing in ecological conditions that are more demanding for the water transport, *i.e.* higher salt concentrations in the soil water (mean soil water salinity and soil water salinity range) and low inundation frequency [3,39] (Fig. 2b and Table 2). Furthermore, a higher

Table 3. Table showing the number of woody species with concentric internal phloem (Table S1) that occur in (periodically) dry and/or saline environments (stress) and in wet, non-saline environments (no stress) categorized by habit (lianas, shrubs and trees).

	Stress	No stress
Lianas	9 (4.1%)	110 (49.5%)
Shrubs	48 (21.6%)	14 (6.3%)
Trees	31 (14.0%)	10 (4.5%)
	88 (39.6%)	134 (60.4%)

Absolute numbers with percentage between brackets. doi:10.1371/journal.pone.0016558.t003




Figure 4. Nature of the successive cambia in *Avicennia*: two models. (a) Model of the *fishnet stockings*: multiple non-continuous cambial sheets give rise to the reticulate transport system. (b) Model of the *broken cylinders*: complete cambium shields have the ability to break at certain locations and can eventually line up in a later state of the development of the transport structure. Only one cambial layer is shown in both visualisations. doi:10.1371/journal.pone.0016558.g004

phloem to xylem tissue ratio has been measured in the trees of the ecologically most stressful study site (Fig. 2d and Table 2). These observations corroborated the current idea for a role of phloem tissue in embolism repair [21,22,23,24,25,26]. The more extensive branching of the phloem tissue, going with a higher absolute amount of internal phloem, as well as the higher phloem to xylem ratio might thus provide a safer water transport system, next to the enhanced water storage potential [21,22,23,24].

With increasing height aboveground, the degree of branching of the transport system within an *Avicennia* tree became higher in absolute value and bigger in range. In the roots, the degree of branching was similar to that of the crown (Fig. 3). This means that in the crown and in the roots, the phloem tissue was more entangled within the xylem, leading to closer contact between the two transport tissues. In the crown an increased tension on the water column exists, and thus putting a higher demand on the water transport system requiring a higher safety level. Although the tension is lowest in the roots, securing water flow is as essential at the start of the pipeline as it is in the higher parts of the crown.

Surprisingly, the ratio of ending to branched phloem differed between species, with a lower ratio in species from the eastern biogeographical mangrove region than in *A. germinans* from the western biogeographical mangrove region (Fig. 2a). This characteristic seemed to be species-specific and therefore part of the evolutionary background of the species rather than a flexible response to environmental drivers.

Supporting the findings in *Avicennia* spp. and the functional role for the phloem in safeguarding the water transport was that 84.9% of trees and shrubs with concentric internal phloem grow in dry or

saline habitats. This relationship, however, held only when lianas were excluded from the database since they were far more distributed in wet habitats (Table 3). Concerning the functional advantage concentric internal phloem - often deriving from successive cambia - might offer, a distinction has to be made between trees and shrubs on one side and lianas on the other side. In trees and shrubs, the main functional advantage of internal phloem is thought to be water and photosynthate storage as well as the role in embolism repair. For scandent lianas however, the succession of the rigid xylem tissue with the thin-walled phloem tissue provides the necessary flexibility [1,2,20]. Besides, the additional starch storage capability of the successive phloem layers offers an advantage for quick growth to the top of the canopy [2]. In terms of our hypothesis, it means that concentric internal phloem in lianas is not *per se* related to drought or salt stress conditions, whereas it is in trees and shrubs.

Apart from the majority of trees and shrubs with concentric internal phloem growing under drought stress conditions, there was still 15.1% related to habitats with adequate water supply. Although speculative, this can be explained by the wide range of habitats these species occur in, or by the evolutionary evolvement of these species from species in dryer habitats together with the neutral effect of growth via successive cambia on the survival of trees and shrubs growing under wet conditions. Another explanation is the higher flexibility and regeneration capacity offered by successive cambia. Flexibility can be advantageous for trees and shrubs confronted with storms and/or heavy wind while a good regeneration capacity can be due to the presence of more than one cambium. Lastly, the increased storage of photosynthates




Figure 5. From wood anatomy to ecology: overview of the study. The flow chart is showing the organisms involved (black boxes, capital text) and the results of the study of successive cambia and concentric internal phloem in these organisms (white boxes, normal text) establishing that internal phloem could offer advantages in ecological stressful conditions (grey box with black frame). In this way, the wood anatomical features of *Avicennia* can help in explaining its broad distribution compared to other mangrove genera. doi:10.1371/journal.pone.0016558.g005

is not only advantageous for lianas in their sprint to the canopy in order to receive light [2], but also for smaller trees growing in dense tropical forests.

Conclusions and perspectives

Growth via successive cambia offers an ecological advantage under water stress conditions. Woody species showing concentric internal phloem were predominant in salt or drought prone habitats, and the networking of the phloem tissue was more pronounced under harsher environmental conditions for the water transport (Fig. 5). Physiological experiments - *e.g.* those proving that the internal phloem tissue is still active [11] - should strengthen the hypothesis that the advantage is most probably related to the water storage capacity and the role in embolism repair of the internal phloem. Concentric internal phloem and thus growth via successive cambia was found to offer several ecological benefits for different groups of species growing in different habitats: under humid-wet conditions this anatomy offers ecological advantage for woody plants with a scandent growth, while under drought or salt stress conditions it can be beneficial for trees and shrubs. If this type of secondary growth is so beneficial, we can wonder why the commonly occurring growth through a single cambium is the main growth form of woody plants. The cost-benefit ratio of having more than one cambium might be the answer, although this has not been calculated to date, unless disadvantages as yet unknown to us would impede evolutionary spread of growth by more than one cambium.

The ecological benefits of successive cambia, in combination with the specialised vessel characteristics [39], help explaining the broad local and global distribution of *Avicennia* and its occurrence in extremely high salinity conditions. Generating data on the growth of species with network-like successive cambia at high temporal and spatial resolution with the laser-scanning method [40] or point dendrometers is needed to obtain more insight in the precise functioning of those cambia and the way the complex structure of the wood develops. The complexity of the water transport system structure that can originate from successive cambia is extending our current insight in plant secondary growth, as these types of structure can only be generated from cambia that are anatomically different from those generally observed in trees.

In conclusion, we can state that our study indicates that successive cambia are an important wood anatomical characteristic explaining ecological species distribution at least partially. It moreover stressed the importance of three-dimensional visualisations and interpretations of plant structures.

Materials and Methods

X-ray Computed Tomography (CT): CT-scanning

Wood Samples. In order to study the three-dimensional wood structure of the mangrove genus *Avicennia* L., we used (i) stems available in the xylarium of the Royal Museum of Central Africa in Tervuren (Belgium) and (ii) root samples and serial stem parts of one tree collected in the mangrove forest of Gazi Bay (Kenya).

Eighteen air-dried wood samples were selected from the Tervuren wood collection (Table 1). The samples were selected to allow comparison between (i) *Avicennia* species (*A. marina*, *A. officinalis* and *A. geminans*), (ii) biogeographical regions (eastern region: Indo-West Pacific and East Africa and western region: America and West Africa) and (iii) study sites mainly differing in degree of soil water salinity. All selected stems were at least 3 cm in diameter.

In addition to the stem samples out of the Tervuren wood collection, two cable root samples of *Avicennia marina* (Forssk.)

Vierh. were collected in the mangrove forest of Gazi Bay, Kenya, in February 2009 at two different study sites (Table 1). Study of the variation in wood structure by height was done with twelve samples of the same tree (distance between samples: 50 cm) sampled from study site 2 of which a root sample also has been taken (Table 1). All collected samples were been air-dried before analysis and now are part of the Tervuren wood collection (Tw60819-20 – Table 1).

Environmental data from the different study sites in Gazi Bay, Kenya, were determined during former studies (Table 2) [3,39].

Scanning Procedure. X-ray tomographical images have been made for all samples using a multi-slice spiral CT-scanner (CT-scan Brilliance 64 slice, Philips, The Netherlands) with the following characteristics: collimation: 20×0,625 mm; slice thickness: 0.7 mm; reconstructed slice interval: 0.35 mm; intensity: 330 mAs; tension: 140 kV; X-tube rotation time: 0.75 s; reconstruction filter: B-standard. The field of view and the pixel spacing were adapted to the object diameter, all with 512×512 image matrix. Radial and three-dimensional reconstructions were made on the Philips CT-scan with dedicated software.

Data Collection and Analysis. In order to investigate the degree of branching of the *Avicennia* vascular network, points of ending and branched phloem tissue (Fig. 2c–g) were counted on the CT-images using ImageJ 1.41k (Wayne Rasband, National Institute of Health, Bethesda, Maryland, USA). For every sample, measurements were made at three different heights at 30 mm distance. On every image the stem surface area was determined by the ellips tools in eFilm Lite 2.1.2 (Merge OEM, Mississauga, Canada) or after measuring the stem diameters in iQ-VIEW 2.5.0 (IMAGE Information Systems, London, UK). From these measurements we calculated (i) the degree of branching, defined as the sum of ending and branched phloem tissue *i.e.* the number of points where the growth segments are not concentric per surface area, giving an indication of the complexity of the network of transport tissue and (ii) the ratio of ending to branched phloem. On the same images the internal phloem surface area was determined semi-automatically (Color Range tool in Adobe Photoshop CS3 – version 10.0.1, Adobe Systems Inc., San Jose, California, USA). From these measurements, the xylem surface area (total stem surface area minus internal phloem surface area) as well as the ratio of internal phloem surface area to xylem surface area could be calculated.

Degree of branching, ratio of ending to branched phloem and ratio of phloem to xylem surface area were compared within and between species, biogeographical regions and study sites using Mann-Whitney U and Kruskal-Wallis tests. All statistical analyses were conducted using Statistica 7.0 (StatSoft Inc., Tulsa, OK, USA).

X-ray Computed Tomography (CT): Micro-CT-scanning

The samples used for micro-CT-scanning were representative *A. marina* samples of Gazi Bay, Kenya, out of the Tervuren wood collection. The pictures showed in this article are from the Tw60821 *A. marina* sample.

A high resolution desktop X-ray micro-CT-system (SkyScan 1172, SkyScan, Belgium) with closed X-ray micro-focus source, was used for non-destructive visualization of the internal structure of *A. marina*. It was possible to obtain an isotropic pixel resolution of 7.5 μm using a camera binning mode of 2 by 2 pixels resulting in a projection image of 1000×2000 pixels. A filter of 0.5 mm aluminium was chosen in order to get a better contrast and to reduce the beam hardening effect. The peak voltage of the source was set at 90 kV. Projection images were taken with a rotation step

of 0.4° over 180° and the signal to noise of the projection images was improved by using a frame averaging of 3. After acquiring the projection images the reconstruction was done using a modified Feldkamp cone-beam algorithm [41]. Two-dimensional cross-sectional images of the sample were obtained in consecutive slices throughout the object in order to obtain a three-dimensional dataset that can be viewed in any direction.

Three-dimensional smoothed surface images were made using the Reconstruct software (version 1.1.0.0 - John C. Fiala, Austin, Texas, USA) [42].

Database Analysis

In order to upscale and verify the hypothesis that concentric internal phloem is an adaptive characteristic of species growing under drought stress, either by a lack of water or by the presence of salt, we conducted a database analysis in which we analysed 198 woody species with concentric internal phloem. Species were selected from the modern wood database of the Inside Wood Database [12] (species with IAWA (International Association of Wood Anatomists) characteristic 133: included phloem, concentric) or from scientific literature. For each species the growth habit, the habitat and the (bio)geography were searched for in the Inside Wood Database or in scientific articles and websites (Table S1). In all, 194 (87 genera out of 25 families) could be taken into the analysis (Table S1).

For statistical analysis (descriptive statistics) habitats were classified according to Walter's zonobiomes (Table S1) [43] based on both habitat and (bio)geography of each species. Zonobiomes II – IV and VII – IX have been associated with water stress as a result of drought during at least one period of the year and also saline habitats were considered physiologically dry. In contrast, zonobiomes I, V and VI as well as mountain areas were considered to be without extensive drought stress. Zonobiome II has been divided into II a (semi-evergreen and wet season green forests) and II b (savannas, grassland and dry woodlands) according to Walters's vegetational zones [43]. Habitat descriptions were decisive to divide species into stress categories (drought and/or salt stress versus no drought stress). Species with two growth habits (tree/shrub or shrub/liana) were assigned to both groups.

References

1. Carlquist S (2001) Comparative Wood Anatomy: Systematic, Ecological, and Evolutionary Aspects of Dicotyledon Wood (Springer Series in Wood Sciences). Berlin - Heidelberg: Springer-Verlag. 448 p.
2. Carlquist S (2007) Successive cambia revisited: ontogeny, histology, diversity, and functional significance. *Journal of the Torrey Botanical Society* 134: 301–332.
3. Schmitz N, Robert EMR, Verheyden A, Kairo JG, Beekman H, et al. (2008) A patchy growth via successive and simultaneous cambia: key to success of the most widespread mangrove species *Avicennia marina*? *Annals of Botany* 101: 49–58.
4. Araujo GUC, Costa CG (2006) Cambial variant in the stem of *Serjania corrugata* (Sapindaceae). *Iawa Journal* 27: 269–280.
5. McDonald JA (1992) Evolutionary Implications of Typical and Anomalous Secondary Growth in Arborescent *Ipomoea* (Convolvulaceae). *Bulletin of the Torrey Botanical Club* 119: 262–267.
6. Nair MNB (1993) Structure of Stem and Cambial Variant in *Spatholobus roxburghii* (Leguminosae). *Iawa Journal* 14: 191–204.
7. Dobbins DR, Fisher JB (1986) Wound Responses in Girdled Stems of Lianas. *Botanical Gazette* 147: 278–289.
8. Fahn A, Zimmermann MH (1982) Development of the Successive Cambia in *Atriplex halimus* (Chenopodiaceae). *Botanical Gazette* 143: 353–357.
9. Rajput KS, Rao KS (2000) Secondary growth in the stem of some species of *Alternanthera* and *Achyranthes aspera* (Amaranthaceae). *Iawa Journal* 21: 417–424.
10. Rajput KS, Rao KS, Patil UG (2006) Stem anatomy of *Dolichos lablab* Linn (Fabaceae): Origin of cambium and reverse orientation of vascular bundles. *Flora* 201: 65–73.
11. Fahn A, Shchori Y (1967) The Organization of Secondary Conducting Tissues in Some Species of the Chenopodiaceae. *Phytomorphology* 17: 147–8.
12. InsideWood (2004-onwards) Available: <http://insidewood/lib.ncsu.edu/search>. Accessed: 2010 Sept 3.
13. Schmitz N, Verheyden A, Kairo JG, Beekman H, Koedam N (2007) Successive cambia development in *Avicennia marina* (Forssk.) Vierh. is not climatically driven in the seasonal climate at Gazi Bay, Kenya. *Dendrochronologia* 25: 87–96.
14. Zamski E (1981) Does Successive Cambia Differentiation in *Avicennia* Depend on Leaf and Branch Initiation. *Israel Journal of Botany* 30: 57–64.
15. Zamski E (1979) The mode of secondary growth and the three-dimensional structure of the phloem in *Avicennia*. *Botanical Gazette* 140: 67–76.
16. Baker RT (1915) The Australian "grey mangrove," (*Avicennia officinalis*, Linn.). *J Proc roy Soc NS Wales* 49: 257–281.
17. Chapman VJ (1947) Secondary thickening and lenticels in *Avicennia nitida* Jacq. *Proceedings of the Linnean Society* 158: 2–6.
18. Studholme WP, Philipson WR (1966) A comparison of the cambium in two woods with included phloem: *Heimerliodendron brunonianum* and *Avicennia resinifera*. *New Zealand Journal of Botany* 4: 355–365.
19. Gill AM (1971) Endogenous control of growth-ring development in *Avicennia*. *Forest Science* 17: 462–465.
20. Bhambe S (1972) Correlation between form, structure and habit in some lianas. *Proceedings of the Indian Academy of Sciences* 75: 246–256.
21. Salleo S, Lo Gullo MA, Trifilo P, Nardini A (2004) New evidence for a role of vessel-associated cells and phloem in the rapid xylem refilling of cavitated stems of *Laurus nobilis* L. *Plant Cell and Environment* 27: 1065–1076.
22. Salleo S, Trifilo P, Esposito S, Nardini A, Lo Gullo MA (2009) Starch-to-sugar conversion in wood parenchyma of field-growing *Laurus nobilis* plants: a component of the signal pathway for embolism repair? *Functional Plant Biology* 36: 815–825.

Supporting Information

Table S1 List of species with concentric internal phloem taken into account in the presented study.

Species names are according to the reference article while families are along the APG. Clear synonyms are removed from the list. For each species habit (L: liana, S: shrub or T: tree), habitat and (bio)geography have been searched for in scientific articles and websites. Based on this information, species have been categorised non salt tolerant (0) or at least salt tolerant (1) and have been classified to the different zonobiomes described in Walter's Vegetation of the Earth. Species from coastal areas have been classified as azonal (A), while species from mountain areas have been indicted with mountain (M). Genera that were found to have internal concentric phloem are not taken into account in the analysis but only mentioned in this list. Number of species according to Mabberley's Plant-Book are mentioned between brackets.

(PDF)

Acknowledgments

We thank James Gitundu Kairo of the Kenya Marine and Fisheries Research Institute for the logistic help during the field expedition, Abudhabi Jambia for the help during sampling in the field, Karel Robert for the development of the measuring method in Adobe Photoshop and the assistance with the figure creation, Maarten De Munter for his help during the CT-scanning and Wim Tavernier for his contribution concerning the Tervuren wood collection. We are very grateful to the Department of Radiology of the Universitair Ziekenhuis Brussel for offering free CT-scanning and to an anonymous reviewer for the critical comments on an earlier version of the manuscript.

Author Contributions

Conceived and designed the experiments: EMMR NS NK. Performed the experiments: EMRR JDM EVdC. Analyzed the data: EMRR NS IB HB NK. Contributed reagents/materials/analysis tools: JDM EVdC. Wrote the paper: EMRR. Data collection: EMRR TD KH.

23. Salleo S, Trifilo P, Lo Gullo MA (2006) Phloem as a possible major determinant of rapid cavitation reversal in stems of *Laurus nobilis* (laurel). *Functional Plant Biology* 33: 1063–1074.
24. Zwieniecki MA, Holbrook NM (2009) Confronting Maxwell's demon: biophysics of xylem embolism repair. *Trends in Plant Science* 14: 530–534.
25. Mauseth JD, Plemons-Rodriguez BJ (1997) Presence of paratracheal water storage tissue does not alter vessel characters in cactus wood. *American Journal of Botany* 84: 815–822.
26. Scholz FG, Bucci SJ, Goldstein G, Meinzer FC, Franco AC, et al. (2007) Biophysical properties and functional significance of stem water storage tissues in Neotropical savanna trees. *Plant Cell and Environment* 30: 236–248.
27. Krishnamurthy KV, Venugopal N (1984) Studies on successive cambia of a few plants. *Indian J Bot* 7: 183–189.
28. Nair MNB, Ram HYM (1990) Structure of Wood and Cambial Variant in the Stem of *Dalbergia paniculata* Roxb. *Iawa Bulletin* 11: 379–391.
29. Zamski E (1980) Vascular Continuity in the Primary and Secondary Stem Tissues of *Bougainvillea* (Nyctaginaceae). *Annals of Botany* 45: 561–8.
30. Wheat D (1977) Successive Cambia in Stem of *Phytolacca dioica*. *American Journal of Botany* 64: 1209–1217.
31. Bhambe S, Sharma A (1985) Ontogeny of cambium in *Amaranthus caudatus* L. and *Achyranthes aspera* L. *Proceedings of the Indian Academy of Sciences* 95: 295–301.
32. Terrazas T (1991) Origin and Activity of Successive Cambia in *Cycas* (Cycadales). *American Journal of Botany* 78: 1335–1344.
33. den Outer RW, van Veenendaal WLH (1995) Development of Included Phloem in the Stem of *Combretum nigricans* (Combretaceae). *IAWA Journal* 16: 151–158.
34. Kitin P, Fujii T, Abe H, Takata K (2009) Anatomical features that facilitate radial flow across growth rings and from xylem to cambium in *Cryptomeria japonica*. *Annals of Botany* 103: 1145–1157.
35. Kitin P, Takata K (2007) Structure and interfaces between secondary xylem, cambium and phloem. *International Conference on Plant Vascular Biology - Program and Abstract Book, Academia Sinica Taipei, Taiwan*. 20 p.
36. Murakami Y, Funada R, Sano Y, Ohtani J (1999) The differentiation of contact cells and isolation cells in the xylem ray parenchyma of *Populus maximowiczii*. *Annals of Botany* 84: 429–435.
37. Philipson WR, Ward JM, Butterfield BG (1971) Anomalous cambia. In: Chapman & Hall LTD. *The vascular cambium. Its development and activity*. 184 p.
38. Carlquist S (2007) Successive cambia in Aizoaceae: products and process. *Botanical Journal of the Linnean Society* 153: 141–155.
39. Robert EMR, Koedam N, Beeckman H, Schmitz N (2009) A safe hydraulic architecture as wood anatomical explanation for the difference in distribution of the mangroves *Avicennia* and *Rhizophora*. *Functional Ecology* 23: 649–657.
40. Dunisch O, Schulte M, Kruse K (2003) Cambial growth of *Swietenia macrophylla* King studied under controlled conditions by high resolution laser measurements. *Holzforschung* 57: 196–206.
41. Feldkamp LA, Davis LC, Kress JW (1997) Practical Cone beam algorithm. *Journal of Microscopy* 185: 67–75.
42. Fiala JC (2005) Reconstruct: a free editor for serial section microscopy. *Journal of Microscopy-Oxford* 218: 52–61.
43. Breckle SW (2002) *Walter's vegetation of the earth. The ecological systems of the Geo-Biosphere*. Berlin - Heidelberg: Springer-Verlag. 547 p.
44. Tomlinson PB (1994) *The Botany of Mangroves*. Cambridge: Cambridge University Press. 433 p.

Table S1: List of species with concentric internal phloem taken into account in the presented study. Species names are according to the reference article while families are along the APG [1]. Clear synonyms are removed from the list. For each species habit (L: liana, S: shrub or T: tree), habitat and (bio)geography have been searched for in scientific articles and websides. Based on this information, species have been categorised non salt tolerant (0) or at least salt tolerant (1) and have been classified to the different zonobiomes described in Walter’s Vegetation of the Earth (see below table) [2]. Species from coastal areas have been classified as azonal (A), while species from mountain areas have been been indicted with mountain (M). Genera that were found to have internal concentric phloem are not taken into account in the analysis but only mentionend in this list. Number of species accoring to Mabberley’s Plant-Book [3] are mentioned between brackets.

GENUS	SPECIES	FAMILY	HABIT	HABITAT	(BIO)GEOGRAPHY	SALT	WALTER	REFERENCE
<i>Abuta</i>	<i>brevifolia</i>	Menispermaceae	L	Rainforest	Middle and South America	0	I	[4-6]
<i>Abuta</i>	<i>colombiana</i>	Menispermaceae	L	Rainforest	Meso and South America	0	I	[4-6]
<i>Abuta</i>	<i>fluminum</i>	Menispermaceae	L	Amazonian forest	Peru, Amazonian area	0	I	[4-8]
<i>Abuta</i>	<i>grandifolia</i>	Menispermaceae	L/S	Humid, tropical areas, near and far away from water, in secondary forest and degraded meadows	Tropical South America	0	I	[4, 5, 9]
<i>Abuta</i>	<i>grisebachii</i>	Menispermaceae	L	Rainforest	Meso and South America	0	I	[4-6, 10]
<i>Abuta</i>	<i>imene</i>	Menispermaceae	L	Rainforest	Brazil	0	I	[4, 5, 7, 11, 12]
<i>Abuta</i>	<i>rufescens</i>	Menispermaceae	L	Non-flooded moist forest	Tropical South America	0	I	[4, 5, 13]
<i>Abuta</i>	<i>sandwithiana</i>	Menispermaceae	L	Dense, submountainous ombrophilous forest	Brazil	0	I	[4, 5, 7, 14]
<i>Albertisia</i>	<i>exelliana</i>	Menispermaceae	L	Rainforest, near lake forest	Tropical Africa	0	I	[4, 15]
<i>Albertisia</i>	<i>papuana</i>	Menispermaceae	L	Rainforest	South East Asia	0	I	[4, 16]
<i>Albertisia</i>	<i>villosa</i>	Menispermaceae	L	Dense humid forest	Tropical Africa	0	I	[4, 17]
<i>Allenrolfea</i>	<i>vaginata</i>	Amaranthaceae	T	Desert with saline soils	Temperate South America including Argentina, Chile, Uruguay and South Paraguay	1	III-VII	[5, 18]
<i>Anabasis</i>	<i>articulata</i>	Amaranthaceae	S	Shrub steppe, desert	Mediterranean including Northern Africa and Middle East	1	III-IV	[5, 19]
<i>Anamirta</i>	<i>cocculus</i>	Menispermaceae	L/S	Tropical evergreen forest	India, Indonesia, Sri-Lanka	0	I	[4, 5, 20, 21]
<i>Anisocycla</i>	<i>cymosa</i>	Menispermaceae	L	Sandy riversides and fringing forest	Congo	0	I	[4, 22]
<i>Anomospermum</i>	<i>chloranthum</i>	Menispermaceae	L	Moist forest	Meso and South America	0	I	[4, 5, 23]
<i>Anomospermum</i>	<i>reticulatum</i>	Menispermaceae	L	Dense tropical evergreen forest	Costa Rica, Southern America	0	I	[4, 24, 25]
<i>Arcangelisia</i>	<i>flava</i>	Menispermaceae	L	Wet forest, near river banks	Temperate China, tropical Asia	0	I-II-V-VI	[3, 4, 25]

GENUS	SPECIES	FAMILY	HABIT	HABITAT	(BIO)GEOGRAPHY	SALT	WALTER	REFERENCE
<i>Arthroa</i>	<i>leubnitziae</i>	Amaranthaceae	S	Desert and sand dunes	Namibia	0	III	[26]
<i>Arthrocnemum</i>	<i>macrostachyum</i>	Amaranthaceae	S	Saline habitats: coasts and banks of saline water pools	Mediterranean including Northern Africa and Middle East	1	IV	[5, 27]
<i>Arthrocnemum</i>	<i>perenne</i>	Amaranthaceae	S	Muddy salt marshes near the sea	Mediterranean including Northern Africa and Middle East	1	IV	[5, 15]
<i>Atriplex</i>	<i>halimus</i>	Amaranthaceae	S	Dry and saline habitats, desert	Mediterranean including Northern Africa and Middle East	1	IV	[5, 28]
<i>Atriplex</i>	<i>nummularia</i>	Amaranthaceae	S	Arid and semi arid regions, usually on saline soils	Australia	1	III	[29, 30]
<i>Avicennia</i>	<i>alba</i>	Acanthaceae	S/T	Mangrove forests	Indo-Pacific ocean	1	A	[31-34]
<i>Avicennia</i>	<i>bicolor</i>	Acanthaceae	S/T	Mangrove forests	West coast of Meso America	1	A	[31-34]
<i>Avicennia</i>	<i>germinans</i>	Acanthaceae	S/T	Mangrove forests	Meso America (up to Florida) and West Africa	1	A	[5, 31-37]
<i>Avicennia</i>	<i>integra</i>	Acanthaceae	S/T	Mangrove forests	North Australia	1	A	[31, 33, 34, 38]
<i>Avicennia</i>	<i>marina</i>	Acanthaceae	S/T	Mangrove forests	East Africa and Indo-Pacific	1	A	[5, 31-34, 38-40]
<i>Avicennia</i>	<i>officinalis</i>	Acanthaceae	S/T	Mangrove forests	Indo-Pacific	1	A	[31-34, 41]
<i>Avicennia</i>	<i>rumphiana</i>	Acanthaceae	S/T	Mangrove forests	Indo-Malaysia	1	A	[31, 33, 34]
<i>Avicennia</i>	<i>schauriana</i>	Acanthaceae	S/T	Mangrove forests	East coast of South America	1	A	[31-34]
<i>Azima</i>	<i>tetracantha</i>	Salvadoraceae	S	Tropical mainland, on saline or alluvial soils near rivers and coast	Tropical mainland Africa and adjacent islands (Madagascar, Mauritius, Réunion, and Comores)	1	I-IIa-IIb-III	[5, 15, 42]
<i>Barbeuia</i>	<i>madagascariensis</i>	Barbeuiaceae	L	Subhumid to humid forest	Madagascar	0	I-II a	[43-45]
<i>Beirnaertia</i>	<i>cabindensis</i>	Menispermaceae	L	Tropical wet forest	Tropical Africa	0	I	[3, 4, 46]
<i>Bosea</i>	<i>yervamora</i>	Amaranthaceae	S	Dry areas	Madeira and the Canary Islands	0	IV	[5, 26]
<i>Bougainvillea</i>	<i>spectabilis</i>	Nyctaginaceae	L/S	Rainforest	Brazil, naturalised elsewhere	0	I	[5, 25, 47-50]
<i>Bredemeyera</i>	sp.	Polygalaceae	S	Caatinga, forest patches in a savannah biome	Neotropics and temperate Brazil, tropical South American, New Guinea, tropical Australia	0	II b	[3, 5, 51]
<i>Cadaba</i>	<i>glandulosa</i>	Capparaceae	S	Deciduous bush land and grassland, dry places on clay or sandy soil and dry riverbeds	Mediterranean including Northern Africa and Middle East, Tropical mainland Africa and adjacent islands (Madagascar, Mauritius, Réunion, and Comores)	1	II b - IV	[5, 27]
<i>Cadaba</i>	<i>rotundifolia</i>	Capparaceae	S	Deciduous bush land, semi-desert and dry riverbeds	Mediterranean including Northern Africa and Middle East, tropical	1	II b - IV	[5, 27]

GENUS	SPECIES	FAMILY	HABIT	HABITAT	(BIO)GEOGRAPHY	SALT	WALTER	REFERENCE
					Africa and adjacent islands			
<i>Caryomene</i>	<i>olivascens</i>	Menispermaceae	L	Tropical rainforest	Meso and South America	0	I	[4, 5, 52]
<i>Casimirella</i>	<i>ampla</i>	Icacinaceae	L	Moist deciduous and semi-evergreen seasonal forest, near riverbanks	Tropical South America	0	II a	[5, 53]
<i>Celosia</i>	<i>floribunda</i>	Amaranthaceae	T	Dry land, scrubland	Mexico (Baja California)	0	III	[26]
<i>Ceratoides</i>	<i>lanata</i>	Amaranthaceae	S	Steppe and grassland, on saline, clay or calcareous soils	North America, north of Mexico	1	VII	[5, 54]
<i>Charpentiera</i>	<i>densiflora</i>	Amaranthaceae	T	Mesic forest	Hawaiian Islands	0	II a	[26, 55]
<i>Charpentiera</i>	<i>elliptica</i>	Amaranthaceae	T	Lowland wet or mesic forest	Hawaiian Islands	0	II a	[26, 56]
<i>Charpentiera</i>	<i>obovata</i>	Amaranthaceae	S	Mesic to occasionally wet rain forests	Pacific Islands (including New Caledonia, Samoa, Hawaii and Fiji)	0	II a	[5, 26]
<i>Cheiloclinium</i>	spp. (11)	Celastraceae	L	Campinarana forest	Mexico and Central America, tropical South America	0	II	[5, 57]
<i>Cheiloclinium</i>	<i>anomalum</i>	Celastraceae	L	Rainforest	Amazonian area	0	I	[5, 7, 58, 59]
<i>Cheiloclinium</i>	<i>belizense</i>	Celastraceae	L	Riparian forest	Belize, Panama, Brazil, Venezuela	0	I – II	[5, 13, 58, 60]
<i>Cheiloclinium</i>	<i>hippocrateoides</i>	Celastraceae	L	Lowland tropical to subtropical moist forest	South America	0	II a	[5, 13, 58, 61]
<i>Cheiloclinium</i>	<i>serratum</i>	Celastraceae	L	Riverine habitats	Middle and South America	0	I-II-III-V-VII	[5, 58, 62, 63]
<i>Chenopodium</i>	<i>sandwicheum</i>	Amaranthaceae	S	Dry land habitats from coastal to subalpine, adapted to drought	Pacific Islands (including New Caledonia, Samoa, Hawaii, and Fiji)	1	II b	[5, 64]
<i>Chondrodendron</i>	<i>microphyllum</i>	Menispermaceae	L	Rainforest	Meso and South America	0	I	[4, 65]
<i>Chondrodendron</i>	<i>tomentosum</i>	Menispermaceae	L	Rainforest	Meso and South America	0	I	[4, 65]
<i>Cissampelos</i>	<i>pareira</i>	Menispermaceae	L/S	Riverine and lowland forest	Tropical regions worldwide	0	I	[4, 5, 17, 49]
<i>Cocculus</i>	<i>hirsutus</i>	Menispermaceae	L/S	Lowland and medium-altitude bush land and woodland	India, Pakistan, Tropical Africa	0	II a - II b	[45, 47, 66, 67]
<i>Cocculus</i>	<i>laurifolius</i>	Menispermaceae	L	Mountain region, dry evergreen forest between 1200 and 1600 m	Temperate Asia (China, Japan and Russia)	0	M	[4, 5, 68]
<i>Cocculus</i>	<i>pendulus</i>	Menispermaceae	L	Savannah	Tropical and subtropical India, Pakistan and Africa	0	IIb	[4, 5, 17, 45, 69]
<i>Colignonia</i>	<i>scandens</i>	Nyctaginaceae	S	Arid regions, mountains > 3000 m	Tropical South America	0	M	[5, 70]
<i>Combretum</i>	<i>nigricans</i>	Combretaceae	S/T	Savannah	Tropical Africa and adjacent islands	0	II b	[5, 71]

GENUS	SPECIES	FAMILY	HABIT	HABITAT	(BIO)GEOGRAPHY	SALT	WALTER	REFERENCE
<i>Curarea</i>	<i>candicans</i>	Menispermaceae	L	Tropical lowland forest	Neotropics	0	I	[4, 5, 72]
<i>Curarea</i>	<i>tecunarum</i>	Menispermaceae	L	Tropical lowland forest	Neotropics	0	I	[4, 5, 72]
<i>Curarea</i>	<i>toxicofera</i>	Menispermaceae	L	Tropical lowland forest	Neotropics	0	I	[4, 5, 72]
<i>Cycas</i>	<i>circinalis</i>	Cycadaceae	T	Along the seashore, inland and occasionally in mountainous areas, open grass lands, shrub-woodlands in hilly areas to dense forests	India, South-East Asia	1	II a - II b – (M)	[73-77]
<i>Cycas</i>	<i>rumphii</i>	Cycadaceae	T	Tropical woodland or forest, mainly along coastal areas, generally on sandy soil and calcareous substrate	Moluccan island group, Papua, Indonesia	1	II a	[78-81]
<i>Cycas</i>	<i>thuarsii</i>	Cycadaceae	T	Near coast, behind the dunes, in open bush land	Madagascar, coast of East Africa	1	II a - II b	[15, 78, 81]
<i>Dalbergia</i>	<i>lanceolaria</i>	Fabaceae	T	Tropical dry thorn forest, tropical moist deciduous forest	Central South Asia	0	II a - II b	[5, 73, 82]
<i>Dalbergia</i>	<i>paniculata</i>	Fabaceae	T	Primary moist deciduous forest, mesomorphic habitat	India, Indo-China (tropical Asia)	0	II a	[25, 83]
<i>Dicranostyles</i>	<i>guianensis</i>	Convolvulaceae	L	Evergreen broadleaf rainforest with high rainfall	Tropical South America	0	I	[5, 53]
<i>Dicranostyles</i>	<i>mildbraediana</i>	Convolvulaceae	L	Evergreen broadleaf rainforest with high rainfall	Tropical South America	0	I	[5, 53]
<i>Dicranostyles</i>	<i>villosus</i>	Convolvulaceae	L	Evergreen broadleaf rainforest with high rainfall	Mexico and Central America	0	I	[5, 53]
<i>Dioscoreophyllum</i>	<i>cumminsii</i>	Menispermaceae	L	Evergreen tropical rainforest	Tropical Africa	0	I	[4, 72, 84]
<i>Diploclisia</i>	<i>glaucescens</i>	Menispermaceae	L	Tropical rainforest	Tropical Asia	0	I	[4, 45]
<i>Doliocarpus</i>	spp. (ca. 40)	Dilleniaceae	L	Transition gallery forest, cerrado	Mexico and Central America, tropical South America	0	I-II a	[5, 57]
<i>Doliocarpus</i>	<i>coriaceus</i>	Dilleniaceae	L	Wet lowland forest	Mexico and Central America, tropical South America	0	I	[5, 85]
<i>Elephantomene</i>	<i>eburnea</i>	Menispermaceae	L	Tropical rainforest	Meso and South America	0	I	[4, 5, 53]
* <i>Forchhammeria</i>	<i>longifolia</i>	Capparaceae	S		Mexico and Central America	0		[5, 86]
<i>Gallesia</i>	<i>integrifolia</i>	Phytolaccaceae	T	Semi-deciduous forest, upland forest with dense closed canopy, tropical lowland dry deciduous forest	Tropical South America	0	II a	[5, 87-89]
<i>Gallesia</i>	<i>scorododendrum</i>	Phytolaccaceae	T	Atlantic moist forest	Temperate South America including	0	I	[5, 90]

GENUS	SPECIES	FAMILY	HABIT	HABITAT	(BIO)GEOGRAPHY	SALT	WALTER	REFERENCE
					Argentina, Chile, Uruguay and South Paraguay			
<i>Gnetum</i>	<i>ula</i>	Gnetaceae	L	Moist deciduous rainforest	India (Eastern and Western Ghats)	0	II a	[47, 73, 91-95]
<i>Haematocarpus</i>	<i>subpeltatus</i>	Menispermaceae	L	Tropical forest along streams at low altitudes	Southeast Asia	0	II a	[4, 96]
<i>Haloxylon</i>	<i>persicum</i>	Amaranthaceae	S/T	Sand hills, deserts and sand ridges	Mediterranean including Northern Africa and Middle East	1	III-IV-VI-VII-VIII	[5, 97]
<i>Haloxylon</i>	<i>salicornicum</i>	Amaranthaceae	S	Desert and semi-desert areas in soils containing much salt	Northern Africa, Arabian peninsula, Western Asia, Indian subcontinent	1	III - IV – VII	[25, 98]
<i>Hyperbaena</i>	<i>domingensis</i>	Menispermaceae	L	Seasonal humid Amazonian rainforest, flooded grassland forest	Tropical South America	0	I - II a	[4, 5, 99]
<i>Hyperbaena</i>	<i>winzerlingii</i>	Menispermaceae	L	Flooded grassland forests	Tropical South America	0	I - II a	[5, 100]
<i>Hypserpa</i>	<i>nitida</i>	Menispermaceae	L	Tropical rainforest, forest margins	Southeast Asia	0	I	[4, 5, 101]
<i>Ipomoea</i>	<i>arborescens</i>	Convolvulaceae	T	Dry tropical forest	Mexico (Sonora)	0	II a	[102, 103]
<i>Ipomoea</i>	<i>murucoides</i>	Convolvulaceae	L	Tropical deciduous moist forest	Mexico, South America, tropical Africa and Southeast Asia. Australia	0	II a	[5, 102]
<i>Ipomoea</i>	<i>pauciflora</i>	Convolvulaceae	T	Dry coastal forests	Central Mexico, Meso America, Western South America	0	II a – III – IV	[25, 102, 104]
<i>Ipomoea</i>	<i>pentaphylla</i>	Convolvulaceae	L	Tropical beach, sand dunes, in forests on calcareous rock	Circumtropical	1	A	[47, 59]
<i>Ipomoea</i>	<i>pes-caprae</i>	Convolvulaceae	L	Sand dunes and beaches	Circumtropical	1	A	[47, 105]
<i>Ipomoea</i>	<i>praecana</i>	Convolvulaceae	L	Dry thicket or forest	Southern Mexico, Honduras, Nicaragua	0	II a	[85, 92, 102]
<i>Ipomoea</i>	<i>populina</i>	Convolvulaceae	L	Xeric shrub land	Mexico, Meso America	0	II a	[102, 106, 107]
<i>Ipomoea</i>	<i>wolcottiana</i>	Convolvulaceae	T	Tropical dry forest	Pacific slope of southern Mexico / Mexico (Northern America)	0	II a	[5, 86, 102, 103, 108]
<i>Iresine</i>	sp.	Amaranthaceae	S	Subtropical or tropical dry forest	Tropical South America, temperate South America including Argentina, Chile, Uruguay and South Paraguay, West Africa, South Japan	0	II b	[3, 5, 92]
<i>Kochia</i>	<i>sedifolia</i>	Amaranthaceae	S	Dry, open shrub land	Australia	1	II b –III – IV	[29, 98]
<i>Koompassia</i>	<i>malaccensis</i>	Fabaceae	T	Mixed swamp forest, Kerangas forest, moist deciduous and semi-evergreen seasonal forest	Indochina and Indomalaysia	0	II a	[5, 109]

GENUS	SPECIES	FAMILY	HABIT	HABITAT	(BIO)GEOGRAPHY	SALT	WALTER	REFERENCE
<i>*Legnephora</i>	<i>minutiflora</i>	Menispermaceae	L		Asia	0		[4]
<i>Machaerium</i>	<i>cobanense</i>	Fabaceae	L	Humid forest	Neotropics	0	I	[110-112]
<i>Machaerium</i>	<i>floribundum</i>	Fabaceae	L	Wet tropical rainforest	Mesoamerica	0	I	[5, 110, 113]
<i>Macrococculus</i>	<i>pomiferus</i>	Menispermaceae	L	Tropical rainforest	Southeast Asia (Indonesia)	0	I	[4, 53]
<i>Maerua</i>	<i>angolensis</i>	Capparaceae	S/T	Very dry areas, savannah	Tropical mainland Africa and adjacent islands (Madagascar, Mauritius, Réunion, and Comores), Southern Africa (south of the Tropic of Capricorn)	0	II b, III, IV	[5, 15]
<i>Maerua</i>	<i>filliformis</i>	Capparaceae	T	Very dry areas, savannah	Madagascar	0	II b	[5, 15, 45]
<i>Maerua</i>	<i>oblonifolia</i>	Capparaceae	S	Savannah woodland	Mediterranean including Northern Africa and Middle East, tropical Africa and adjacent islands	0	II b, III, IV	[5, 15]
<i>Maerua</i>	<i>rigida</i>	Capparaceae	S/T	Very dry areas, savannah	Tropical mainland Africa and adjacent islands (Madagascar, Mauritius, Réunion, and Comores), Southern Africa (south of the Tropic of Capricorn)	0	II b	[5, 15]
<i>Maerua</i>	<i>rosmarinoides</i>	Capparaceae	S/T	Woodland and bush land	Southern Africa (south of the Tropic of Capricorn)	0	II b	[5, 114]
<i>Mestoklema</i>	<i>tuberosum</i>	Aizoaceae	S	Shrub land	Cape (South Africa)	0	IV	[3, 112, 115]
<i>Nototrichum</i>	<i>sandwicense</i>	Amaranthaceae	S	Open dry forest	Hawaiian Islands	0	II a - II b	[26, 116]
<i>Orthomene</i>	<i>schomburgkii</i>	Menispermaceae	L	Riparian rainforest	Meso and South America	0	I	[4, 117]
<i>Pachygone</i>	<i>dasycarpa</i>	Menispermaceae	L	Rainforest	Southeast Asia	0	I	[4, 118]
<i>Pera</i>	<i>bicolor</i>	Euphorbiaceae	T	Moist deciduous and semi-evergreen seasonal forest	Tropical South America	0	II a	[5, 53]
<i>Peritassa</i>	spp. (14)	Celastraceae	L/S	Lowland, non-flooded, tropical rainforest	South America, Tobago	0	I	[5, 7, 53]
<i>Peritassa</i>	<i>bullata</i>	Celastraceae	L	Tropical forest	Ecuador	0	I	[5, 58, 119, 120]
<i>Peritassa</i>	<i>calypsoides</i>	Celastraceae	S	Non-flooded open bush	South Brazil	0	II a	[5, 58, 121, 122]
<i>Peritassa</i>	<i>huanucana</i>	Celastraceae	L	Riparian forest	Range includes Peru and Suriname	0	I (II b - III)	[5, 13, 58, 60]
<i>Peritassa</i>	<i>pruinosa</i>	Celastraceae	L	Dense and high Terra Firme lowland forest at low altitudes	Meso-America, northern South America	0	I	[13, 58, 123]

GENUS	SPECIES	FAMILY	HABIT	HABITAT	(BIO)GEOGRAPHY	SALT	WALTER	REFERENCE
<i>Petiveria</i>	<i>alliacea</i>	Phytolaccaceae	S	Wet forest	Tropical Africa, South America and Mesoamerica, Caribbean	0	I-II a	[87, 92, 124-126]
<i>Pfaffia</i>	<i>grandiflora</i>	Amaranthaceae	S	Lowland tropical to subtropical moist forest, Terra Firme forest	Temperate South America including Argentina, Chile, Uruguay and South Paraguay	0	II a	[5, 61]
<i>Phytolacca</i>	<i>dioica</i>	Phytolaccaceae	T	Pampas grassland	Brazil, Peru, Paraguay, Uruguay, North of Argentina	1	II b - VII	[50, 87, 127]
<i>Phytolacca</i>	<i>dodecandra</i>	Phytolaccaceae	L	Wide range of habitats, often riverine, evergreen bush land	Tropical Africa, Madagascar	0	II a	[87, 128, 129]
<i>Phytolacca</i>	<i>weberbaueri</i>	Phytolaccaceae	T	Pampas grassland	Brazil, Peru, Paraguay, Uruguay, North of Argentina	0	II b – VII	[50, 127, 130]
<i>Pisonia</i>	<i>brunoniana</i>	Nyctaginaceae	S/T	Dry to mesic forest, gulches and occasionally wet areas, especially along the sides of streams	Southeast Asia and Pacific islands, Australia and New Zealand	0	II b	[5, 48, 131, 132]
<i>Polygala</i>	sp.	Polygalaceae	S	Coastal or inland habitats in xeric vegetation, from dry to moist habitats	Subcosmopolitan (not in New Zealand)	1	IIa-IIb -III	[3, 5, 133, 134]
<i>Pycnarrhena</i>	<i>cauliflora</i>	Menispermaceae	L	Tropical rainforest	Southeast Asia	0	I	[4, 53]
<i>Pycnarrhena</i>	<i>celebica</i>	Menispermaceae	L	Tropical rainforest	Southeast Asia	0	I	[4, 53]
<i>Pycnarrhena</i>	<i>lucida</i>	Menispermaceae	L	Tropical forest	Southeast Asia	0	I	[4, 53]
<i>Pycnarrhena</i>	<i>tumefacta</i>	Menispermaceae	L	Tropical rainforest	Southeast Asia	0	I	[4, 45]
<i>Rhabdodendron</i>	<i>amazonicum</i>	Rhabdodendraceae	T	Riparian forest, lowland forest, Terra Firme forest	Tropical South America	0	I	[5, 13, 57, 135]
<i>Salacia</i>	spp. (200)	Celastraceae	L/S/T	Subtropical or tropical moist lowland rainforest	Africa, Madagascar, Malaysia, neotropics	0	I	[5, 24, 92]
<i>Salacia</i>	<i>adolfo-friderici</i>	Celastraceae	L	Tropical moist semi-deciduous forest	Cameroon, Ghana	0	I - II a	[5, 58, 86, 136]
<i>Salacia</i>	<i>alwynii</i>	Celastraceae	L	Rainforest	Western South America	0	I	[5, 58, 86, 137, 138]
<i>Salacia</i>	<i>amplectens</i>	Celastraceae	L	Dense and high Terra Firme lowland forest at low altitudes	Brazil, ...	0	I	[5, 13, 58, 86]
<i>Salacia</i>	<i>cauliflora</i>	Celastraceae	L	Rainforest	Amazonian area	0	I	[53, 58, 86, 139]
<i>Salacia</i>	<i>cerasifera</i>	Celastraceae	L	Guineo-congololese forest	Tropical Africa	0	I	[5, 45, 58, 86]
<i>Salacia</i>	<i>chinensis</i>	Celastraceae	L/S	Hot and humid forest	Africa, Indochina, Australia	0	I	[45, 58, 140]
<i>Salacia</i>	<i>chlorantha</i>	Celastraceae	L/S	Evergreen forest and fringing	Tropical Africa	0	I	[15, 58, 86]

GENUS	SPECIES	FAMILY	HABIT	HABITAT	(BIO)GEOGRAPHY	SALT	WALTER	REFERENCE
				forest				
<i>Salacia</i>	<i>crassifolia</i>	Celastraceae	T	Cerrado	Southern America	0	II b	[5, 58, 86, 141]
<i>Salacia</i>	<i>debilis</i>	Celastraceae	L	Tropical forest	West Tropical Africa	0	I	[5, 15, 58, 86]
<i>Salacia</i>	<i>disepala</i>	Celastraceae	L	Rainforest	Australia, ...	0	II b	[58, 142]
<i>Salacia</i>	<i>duckei</i>	Celastraceae	L	Tropical rainforest	Southern America	0	I	[5, 58, 86, 143]
<i>Salacia</i>	<i>elegans</i>	Celastraceae	L/S	Evergreen and fringing forest and dense deciduous woodland	Tropical Africa	0	I-II a	[5, 15, 58, 86]
<i>Salacia</i>	<i>erecta</i>	Celastraceae	L/S	Evergreen forest or thicket, 50-1700 m	Tropical Africa	0	I	[58, 86, 144]
<i>Salacia</i>	<i>germainii</i>	Celastraceae	L	African rainforest	Congo	0	I	[5, 27, 53, 58, 86]
<i>Salacia</i>	<i>impressifolia</i>	Celastraceae	L	Neotropical rainforest	Central and South America	0	I	[5, 58, 145, 146]
<i>Salacia</i>	<i>juruaana</i>	Celastraceae	L	Lowland tropical to subtropical moist forest	Amazonian area	0	I	[5, 58, 61, 86]
<i>Salacia</i>	<i>kanukuensis</i>	Celastraceae	L	Dense tropical forest	Northern South America, Southern America	0	I	[5, 58, 86, 139, 147]
<i>Salacia</i>	<i>lateritia</i>	Celastraceae	L	Tropical rainforest	Tropical Africa	0	I	[15, 53, 58, 86]
<i>Salacia</i>	<i>letestui</i>	Celastraceae	L	West African rainforest	Congo	0	I	[5, 58, 86, 148]
<i>Salacia</i>	<i>macrantha</i>	Celastraceae	T	Lowland tropical to subtropical moist forest	Amazonian area	0	I	[5, 58, 61, 86, 149]
* <i>Salacia</i>	<i>miqueliana</i>	Celastraceae	L	Riparian forest		0		[13, 58, 86]
<i>Salacia</i>	<i>multiflora</i>	Celastraceae	L	Dense, humid evergreen forest		0	I	[5, 13, 58, 86, 145]
<i>Salacia</i>	<i>nitida</i>	Celastraceae	L	Tropical forest	Tropical Africa	0	I	[15, 58, 86]
<i>Salacia</i>	<i>opacifolia</i>	Celastraceae	L	Submontaneous dense rainforest	Neotropics	0	I	[24, 58, 150]
<i>Salacia</i>	<i>prinoides</i>	Celastraceae	L	Forest along seashore, riverbanks	India, ...	1	II a - II b	[5, 58, 151]
<i>Salacia</i>	<i>pynaertii</i>	Celastraceae	L	Evergreen and fringing forest	Tropical Africa	0	I	[5, 58, 86, 144]
<i>Salacia</i>	<i>reticulata</i>	Celastraceae	L	Subtropical or tropical moist lowland rainforest	Central South Asia (India, Pakistan, Sri Lanka)	0	I	[5, 58, 92]
<i>Salacia</i>	<i>solimoesensis</i>	Celastraceae	L	Old growth tropical to subtropical moist forest	Amazonian area	0	I	[5, 58, 61, 86]
<i>Salacia</i>	<i>staudtiana</i>	Celastraceae	L	Tropical rainforest	Tropical Africa	0	I	[5, 15, 58, 86]

GENUS	SPECIES	FAMILY	HABIT	HABITAT	(BIO)GEOGRAPHY	SALT	WALTER	REFERENCE
<i>Salacia</i>	<i>whytei</i>	Celastraceae	L	Rainforest	Tropical Africa	0	I	[15, 58, 86]
<i>Salsola</i>	<i>baryosma</i>	Amaranthaceae	S	Saline and waste sandy places, Sahara desert	Mediterranean including Northern Africa and Middle East	1	III-IV	[5, 15]
<i>Salvadora</i>	<i>persica</i>	Salvadoraceae	S/T	Thorn shrub land, grassy savannahs, coastal or riverine scrub on saline, sandy or alluvial soils, tropical semi-arid areas	Central South Asia, Mediterranean including Northern Africa and Middle East, Tropical Africa and adjacent islands	1	II b – III – IV - VII	[5, 15, 42, 152]
<i>Sarcobatus</i>	<i>vermiculatus</i>	Sarcobataceae	S	Strongly alkaline and saline soils, in semi-arid to arid plains, in badlands, silt dunes	North America, north of Mexico	1	III	[5, 153, 154]
<i>Sciadotenia</i>	<i>eichleriana</i>	Menispermaceae	L	Moist forest	Meso and South America	0	I	[4, 23]
<i>Sciadotenia</i>	<i>sprucei</i>	Menispermaceae	L	Neotropical rainforest	Meso and South America	0	I	[4, 5, 155]
<i>Sciadotenia</i>	<i>toxicofera</i>	Menispermaceae	L	Floodplain, Terra Firme forest	Meso and South America	0	I	[4, 5, 156]
<i>Securidaca</i>	<i>diversifolia</i>	Polygalaceae	T	Tropical dry to premontaneous wet forest	Mexico and Central America, tropical South America	0	II a	[5, 157]
<i>Securidaca</i>	<i>philippinensis</i>	Polygalaceae	L	Rainforest	Indomalaysia	0	I	[5, 158]
<i>Securidaca</i>	<i>virgata</i>	Polygalaceae	L/S	Subtropical wet and moist valley forest and hillsides	Caribbean	0	I-II a	[5, 159]
<i>Seguieria</i>	<i>americana</i>	Phytolaccaceae	S	Subtropical seasonal forest	South America	0	II a	[25, 87, 160, 161]
<i>Seguieria</i>	<i>paraguayensis</i>	Phytolaccaceae	T	Semi-evergreen seasonal forest, tropical moist forest, deciduous forest	Temperate South America including Argentina, Chile, Uruguay and South Paraguay	0	II a – IV	[5, 53, 162]
<i>Simmondsia</i>	<i>chinensis</i>	Simmondsiaceae	S	Desert and chaparral	Sonoran and Mojave Desert (southwest US, northwest Mexico)	1	III	[163-165]
<i>Sleumeria</i>	<i>auriculata</i>	Icacinaceae	L	Rainforest	North Borneo	0	I	[3, 166]
<i>Solenostemma</i>	<i>argel</i>	Apocynaceae	S	Dry sandy semi-desert areas	Mediterranean including Northern Africa and Middle East, Sahara (Arabia)	1	III	[5, 15]
<i>Spatholobus</i>	<i>roxburghii</i>	Fabaceae	L	Sal and teak forest, tropical forest with high rainfall	southeast Asia	0	II a	[167]
<i>Stegnosperma</i>	<i>cubense</i>	Phytolaccaceae	L/S	Tropical dry forest	Cuba, Dominican Republic, Jamaica, Pacific Coast and central Mexico	0	II a	[24, 168-171]
<i>Stegnosperma</i>	<i>halimifolium</i>	Phytolaccaceae	S	Beach, along permanent and ephemeral watercourses, to fifty miles from the coast	coastal Baja California, Mexico	1	A	[168, 169, 171, 172]

GENUS	SPECIES	FAMILY	HABIT	HABITAT	(BIO)GEOGRAPHY	SALT	WALTER	REFERENCE
<i>Stegnosperma</i>	<i>watsonii</i>	Phytolaccaceae	S	Along seacoast and about saline swales	Mexico, coastal Sonora, northern Sinaloa	1	A	[168, 169, 171, 173]
<i>Strychnos</i>	<i>glabra</i>	Loganiaceae	T	Semi-evergreen seasonal forest, tropical moist forest	Tropical South America	0	II a	[5, 53]
<i>Suaeda</i>	<i>fruticosa</i>	Amaranthaceae	S	Saline alluvial flats with clayey or sandy soils, on drier sites and coastal belts	Mediterranean including Northern Africa and Middle East	1	A	[5, 45]
<i>Suaeda</i>	<i>monoica</i>	Amaranthaceae	S	Coastal habitat, halophytic, able to tolerate frequent sea-water flooding	East Africa and Asia	1	A	[5, 15]
<i>Syrrheonema</i>	<i>fasciculatum</i>	Menispermaceae	L	Rainforest, forest regrowth and fallow land, at 1200–1400 m altitude	Tropical Africa	0	I -II a – M	[4, 156]
<i>Telitoxicum</i>	<i>glaziovii</i>	Menispermaceae	L	Tropical lowland rainforest	Meso and South America	0	I	[4, 52]
<i>Telitoxicum</i>	<i>krukovii</i>	Menispermaceae	L	Neotropical rainforest	Meso and South America	0	I	[4, 155]
<i>Telitoxicum</i>	<i>minutiflorum</i>	Menispermaceae	L	Neotropical rainforest	Meso and South America	0	I	[4, 155]
<i>Telitoxicum</i>	<i>peruvianum</i>	Menispermaceae	L	Rainforest	Meso and South America	0	I	[4, 155]
<i>Tetracera</i>	<i>volubilis</i>	Dilleniaceae	L	Shrubby forest, semi-deciduous tropical dry forest	Mexico and Central America, tropical South America	0	II b	[5, 126]
<i>Tetrastigma</i>	<i>voinierianum</i>	Vitaceae	L	Wet rainforest	Indo-China (Vietnam and Laos)	0	I	[5, 174-176]
<i>Tiliacora</i>	<i>acuminata</i>	Menispermaceae	L	Riverine, deciduous forests	China, Nepal, Ceylon, Burma	0	II a	[4, 45, 177]
<i>Tiliacora</i>	<i>chrysobotrya</i>	Menispermaceae	L	Rainforest	Angola, Congo	0	I	[4, 53, 178]
<i>Tiliacora</i>	<i>dielsiana</i>	Menispermaceae	L	Rainforest and fringing forest	Guinea, Liberia, Ivory Coast, Ghana	0	I	[4, 22]
<i>Tiliacora</i>	<i>funifera</i>	Menispermaceae	L	In riverine and other evergreen forest and moist areas in woodland	Tropical Africa	0	I -II a	[4, 5, 179]
<i>Tiliacora</i>	<i>glycosmantha</i>	Menispermaceae	L	Lowland and upland rainforest, riverine forest and moist places in woodland	Tropical mainland Africa and adjacent islands (Madagascar, Mauritius, Réunion, and Comores)	0	II a	[5, 15]
<i>Tiliacora</i>	<i>laurentii</i>	Menispermaceae	L	Forest, subtropical and tropical moist lowland	Tropical Africa	0	II a	[4, 180]
<i>Tontelea</i>	<i>corymbosa</i>	Celastraceae	L	Neotropical rainforest	Amazonian area	0	I	[5, 7, 58, 155]
<i>Tontelea</i>	<i>mauritioides</i>	Celastraceae	L	Forest on Terra Firme	Brazil	0	I	[5, 7, 58, 181]
<i>Tontelea</i>	<i>micrantha</i>	Celastraceae	S/T	Dry grassland	Brazil	0	II b	[5, 7, 58, 182]
<i>Tontelea</i>	<i>nectandrifolia</i>	Celastraceae	L	Slope forest with well drained soil	South America	0	II a	[5, 7, 13, 58]

GENUS	SPECIES	FAMILY	HABIT	HABITAT	(BIO)GEOGRAPHY	SALT	WALTER	REFERENCE
<i>Triclisia</i>	<i>dictyophylla</i>	Menispermaceae	L/S	Dense, humid lowland to medium-altitude forest	Tropical Africa	0	I – II a	[4, 22]
* <i>Triclisia</i>	<i>jumelleana</i>	Menispermaceae	L		Africa and Madagascar	0		[4, 183]
<i>Triclisia</i>	<i>patens</i>	Menispermaceae	L	Lowland rainforest	Tropical West Africa	0	I	[4, 15]
<i>Triclisia</i>	<i>sacleuxii</i>	Menispermaceae	L	Lowland rainforest and riverine forest	Tropical Africa	0	I	[4, 15]
<i>Wisteria</i>	<i>floribunda</i>	Fabaceae	L	Forest edges and disturbed areas at low altitudes, can tolerate a variety of soil and moisture types but it prefers loamy, deep, well drained soils	Temperate Asia (China), Japan and Russia	0	V	[5, 184]

* No appropriate data could be found so that the species has not been included in the analysis of the data.

Zonobiomes according to Breckle 2002 [2]:

I	Zonobiome of the Evergreen Tropical Rain Forest	(Zonobiome of the Equatorial Humid Diurnal Climate)
II	Zonobiome of Savannas and Deciduous Forests and Grasslands	(Zonobiome of the Humido-arid Tropical Summer Rain Region)
	II a semi-evergreen and wet season green forests	
	II b savannas, grassland and dry woodlands	
III	Zonobiome of Hot Deserts	(Zonobiome of Subtropical Arid Climates)
IV	Zonobiome of Sclerophyllic Woodlands	(Zonobiome of the Arido-Humid Winter Rain Region)
V	Zonobiome of Laurel Forests	(Zonobiome of the Warm-Temperate Humid Climate)
VI	Zonobiome of Deciduous Forests	(Zonobiome of the Temperate Nemoral Climate)
VII	Zonobiome of Steppes and Cold Deserts	(Zonobiome of the Arid-Temperature Climate)
VIII	Zonobiome of the Taiga	(Zonobiome of the Cold-Temperature Boreal Climate)
IX	Zonobiome of the Tundra	(Zonobiome of the Arctic Climate)

References:

1. Angiosperm Phylogeny Website. Version 9, June 2008 [and more or less continuously updated since] - <http://www.mobot.org/MOBOT/research/APweb/> - [Date accessed: 26th of January 2010]
2. Breckle, S.-W. (2002). Walter's Vegetation of the Earth. The Ecological Systems of the Geo-Biosphere. Berlin - Heidelberg - New York: Springer-Verlag.
3. Mabberley, D.J. (2008). Mabberley's Plant-Book: A Portable Dictionary of Plants, Their Classification and Uses Book Description. Cambridge: Cambridge University Press.
4. Jacques, F.M.B., and De Franceschi, D. (2007). Menispermaceae wood anatomy and cambial variants. Iawa Journal 28, 139-172.
5. Inside Wood Database. <http://insidewood.lib.ncsu.edu/search> - [Date accessed: 26th of January 2010]
6. Krukoff, B.A., and Barneby, R.C. (1973). Supplementary notes on American Menispermaceae. Phytologia 25 - Accessed from the World Wide Web on the 26th of January 2010: http://www.archive.org/stream/phytologia36glea/phytologia36glea_djvu.txt

7. The Wood Explorer - <http://www.thewoodexplorer.com/> - [Date accessed: 26th of January 2010]
8. Culot, L., Huynen, M.C., Gerard, P., and Heymann, E.W. (2009). Short-term post-dispersal fate of seeds defecated by two small primate species (*Saguinus mystax* and *Saguinus fuscicollis*) in the Amazonian forest of Peru. *Journal of Tropical Ecology* 25, 229-238.
9. Maya Ethnobotanics - <http://www.maya-ethnobotanics.com/> - [Date accessed: 26th of January 2010]
10. Vamosi, S.M., Mazer, S.J., and Cornejo, F. (2008). Breeding systems and seed size in a Neotropical flora: Testing evolutionary hypotheses. *Ecology* 89, 2461-2472.
11. Silverton, J.V., Kabuto, C., Buck, K.T., and Cava, M.P. (1977). Structure of Imerubrine, a Novel Condensed Tropolone-Isoquinoline Alkaloid. *Journal of the American Chemical Society* 99, 6708-6712.
12. van Dulmen, A. (2001). Pollination and phenology of flowers in the canopy of two contrasting rain forest types in Amazonia, Colombia. *Plant Ecology* 153, 73-85.
13. L'Institut de recherche pour le développement en Guyane - <http://www.cayenne.ird.fr/> - [Date accessed: 26th of January 2010]
14. Sasaki, D., Zappi, D., and Milliken, W. (2008). Vegetação do Parque Estadual Cristalino. *Flora Cristalino*.
15. Aluka - digital library of scholarly resources from and about Africa - <http://www.aluka.org/> - [Date accessed: 26th of January 2010]
16. Wilkinson, H.P. (1989). Leaf Anatomy of the Menispermaceae Tribe Tiliaceae Miers. *Botanical Journal of the Linnean Society* 99, 125-174.
17. Schmelzer, G.H., and Gurib-Fakim, A. (2008). Plant Resources of Tropical Africa. Medicinal Plants. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
18. Navone, S., Abraham, E., Bargiela, M., Dent, D.L., Espoz-Alsina, C., Maggi, A., Montana, E., Morrison, S., Pastor, G., Rosatto, H., et al. (2006). Global Deserts Outlook. Chapter 4: State and trends of the world's deserts. - Accessed from the World Wide Web on the 26th of January 2010: <http://www.unep.org/Geo/gdoutlook/>
19. Flowers in Israel - <http://www.flowersinIsrael.com/> - [Date accessed: 26th of January 2010]
20. Plant Biographies - <http://www.plantlives.com/> - [Date accessed: 26th of January 2010]
21. Muthuramkumar, S., and Parthasarathy, N. (2001). Tree-liana relationships in a tropical evergreen forest at Varagalaiar, Anamalais, Western Ghats, India. *Journal of Tropical Ecology* 17, 395-409.
22. Protabase - Webdatabase on useful plants of Tropical Africa - <http://database.prota.org/> - [Date accessed: 26th of January 2010]
23. Mori, S.A., and Boom, B.M. (1987). Chapter II. The Forest. *Mem. New York Botanical Gardens* 44, 9-29.
24. National Biodiversity Institute - Costa Rica - <http://www.inbio.ac.cr/> - [Date accessed: 26th of January 2010]
25. National Plant Germplasm System - United States Department Of Agriculture - <http://www.ars-grin.gov/npgs/> - [Date accessed: 26th of January 2010]
26. Carlquist, S. (2003). Wood and stem anatomy of woody Amaranthaceae s.s.: ecology, systematics and the problems of defining rays in dicotyledons. *Botanical Journal of the Linnean Society* 143, 1-19.
27. Conservatoire et jardin botaniques - Ville de Genève - Databases - <http://ville-ge.ch/musinfo/bd/cjb/> - [Date accessed: 26th of January 2010]
28. Ben Hassine, A., Ghanem, M.E., Bouzid, S., and Lutts, S. (2008). An inland and a coastal population of the Mediterranean xero-halophyte species *Atriplex halimus* L. differ in their ability to accumulate proline and glycinebetaine in response to salinity and water stress. *Journal of Experimental Botany* 59, 1315-1326.
29. Carlquist, S. (1977). Ecological Factors in Wood Evolution - Floristic Approach. *American Journal of Botany* 64, 887-896.
30. Growing Native Plants - Australian National Botanic Gardens - <http://www.anbg.gov.au/gnp/> - [Date accessed: 26th of January 2010]
31. Spalding, M., Blasco, F., and Field, C. (1997). *World Mangrove Atlas*. Okinawa: The International Society for Mangrove Ecosystems.
32. Tomlinson, P.B. (1994). *The Botany of Mangroves*. Cambridge: Cambridge University Press.
33. Philipson, W.R., Ward, J.M., and Butterfield, B.G. (1971). Anomalous cambia. In *The vascular cambium. Its development and activity.*, 1st Edition. (Chapman & Hall LTD).
34. Duke, N.C. (1991). A systematic revision of the mangrove genus *Avicennia* (Avicenniaceae) in Australasia. *Aust. Syst. Bot.* 4, 299-324.
35. Gill, A.M. (1971). Endogenous control of growth-ring development in *Avicennia*. *Forest Science* 17, 462-465.
36. Zamski, E. (1979). The mode of secondary growth and the three-dimensional structure of the phloem in *Avicennia*. *Botanical Gazette* 140, 67-76.
37. Zamski, E. (1981). Does Successive Cambia Differentiation in *Avicennia* Depend on Leaf and Branch Initiation. *Israel Journal of Botany* 30, 57-64.
38. Duke, N.C. (2006). *Australia's mangroves. The authoritative guide to Australia's mangrove plants*. Brisbane: University of Queensland.
39. Sun, Q., and Suzuki, M. (2000). Wood anatomy of mangrove plants in Iriomote Island of Japan: a comparison with mangrove plants from lower latitudes. *Acta Phytotax. Geobot.* 51, 37-55.

40. Sun, Q., and Suzuki, M. (2001). Quantitative character variations of cambial derivatives in mangroves and their functional significance. *Trees-Structure and Function* 15, 249-261.
41. Baker, R.T. (1915). The Australian "grey mangrove," (*Avicennia officinalis*, Linn.). *J. Proc. roy. Soc. NS Wales* 49, 257-281.
42. Carlquist, S. (2002). Wood and bark anatomy of Salvadoraceae: ecology, relationships, histology of interxylary phloem. *Journal of the Torrey Botanical Society* 129, 10-20.
43. Carlquist, S. (1999). Wood anatomy, stem anatomy, and cambial activity of *Barbeuia* (Caryophyllales). *Iawa Journal* 20, 431-440.
44. The families of flowering plants: descriptions, illustrations, identification, and information retrieval - <http://delta-intkey.com> - [Date accessed: 26th of January 2010]
45. eFloras - <http://efloras.org/> - [Date accessed: 26th of January 2010]
46. Kubitzki, K., Rohwer, J.G., and Bittrich, V. (1993). The Families and Genera of Vascular Plants. Volume VII: Flowering plants - Dicotyledons: Magnoliid, Hamamelid, and Caryophyllid families. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
47. Bhambie, S. (1972). Correlation between form, structure and habit in some lianas. *Proceedings of the Indian Academy of Sciences* 75, 246-256.
48. Carlquist, S. (2004). Lateral meristems, successive cambia and their products: a reinterpretation based on roots and stems of Nyctaginaceae. *Botanical Journal of the Linnean Society* 146, 129-143.
49. Plants Database - Natural Resources Conservation Service - United States Department of Agriculture - <http://www.plants.usda.gov/> - [Date accessed: 26th of January 2010]
50. Blue Planet Biomes - <http://www.blueplanetbiomes.org/> - [Date accessed: 26th of January 2010]
51. Lüdtkke, R., de Souza-Chies, T.T., and Miotto, S.T.S. (2008). *Bredemeyera* Willd. e *Securidaca* L. (Polygalaceae) na Região Sul do Brasil. *Revista Brasileira De Biociências* 6.
52. Hecklau, E.F., Mori, S.A., and Brown, J.L. (2005). Specific epithets of the flowering plants of central French Guiana. *Brittonia* 57, 68-87.
53. Peel, M.C., Finlayson, B.L., and McMahon, T.A. (2007). Updated world map of the Koppen-Geiger climate classification. *Hydrology and Earth System Sciences* 11, 1633-1644.
54. National Park Service - U.S. Department of the Interior - <http://www.nps.gov/> - [Date accessed: 26th of January 2010]
55. Heddle, M.L., Wood, K.R., Asquith, A., and Gillespie, R.G. (2000). Conservation status and research on the Fabulous Green Sphinx of Kaua'i, *Tinostoma smaragditis* (Lepidoptera: Sphingidae), including checklists on the vascular plants of the diverse mesic forests of Kaua'i, Hawai'i. *Pacific Conservation* 54, 1-9.
56. Federal Register Environmental Documents - U.S. Environmental Protection Agency - <http://www.epa.gov/fedrgstr/> - [Date accessed: 26th of January 2010]
57. SysTax - a Database System for Systematics and Taxonomy - <http://www.biologie.uni-ulm.de/systax/> - [Date accessed: 26th of January 2010]
58. Menega, A.M.W. (1997). Wood anatomy of the Hippocrateoideae (Celastraceae). *Iawa Journal* 18, 331-368.
59. Botanicus - <http://www.botanicus.org/> - [Date accessed: 26th of January 2010]
60. Neotropical Herbarium Specimens - Fieldmuseum - <http://fm1.fieldmuseum.org/vrrc/> - [Date accessed: 26th of January 2010]
61. Atrium - Biodiversity Information System - <http://atrium.andesamazon.org/> - [Date accessed: 26th of January 2010]
62. Pagina dos colecionadores de frutas - <http://frutasraras.sites.uol.com.br/> - [Date accessed: 26th of January 2010]
63. Gonçalves-Esteves, V., and Sant'Anna Melhem, T. (2004). Palynotaxonomy of the Brazilian species of *Cheilochlinium* Miers (Hippocrateaceae Juss.). *Acta Botanica Brasilica* 18.
64. Senock, R.S., and Taylor, S.A. (2003). *Chenopodium oahense* (College of Agriculture, Forestry and Natural Resource Management, University of Hawaii). - Accessed from the World Wide Web on the 26th of January 2010: <http://www.fs.fed.us/global/iitf/pdf/shrubs/Chenopodium%20oahuenseFinalEd2.pdf>
65. Metabolomics - <http://www.metabolomics.jp> - [Date accessed: 26th of January 2010]
66. Rajput, K.S., and Rao, K.S. (2003). Cambial variant and xylem structure in the stem of *Cocculus hirsutus* (Menispermaceae). *Iawa Journal* 24, 411-420.
67. Friis, I., and Vollesen, K. (2005). Flora of the Sudan-Uganda border area east of the Nile. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
68. Biodiversity Informatics and co-Operation in Taxonomy for Interactive shared Knowledge base (BIOTIK) - <http://www.biotik.org/> - [Date accessed: 26th of January 2010]
69. Iwu, M.M. (1993). Handbook of African medicinal plants. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
70. De Feo, V., Piacente, S., Pizza, C., and Soria, R.U. (1998). Saponins from *Colignonia scandens* Benth. (Nyctaginaceae). *Biochemical Systematics and Ecology* 26, 251-253.
71. Simon, G., Dewelle, J., Nacoulma, O., Guissou, P., Kiss, R., Daloze, D., and Braekman, J.C. (2003). Cytotoxic pentacyclic triterpenes from *Combretum nigricans*. *Fitoterapia* 74, 339-344.
72. Ortiz-Gentry, R.d.C. (1999) Systematic revision of *Curarea* Barneby & Krukoff (Menispermaceae). Missouri, St Louis M. Sc.
73. Krishnamurthy, K.V., and Venugopal, N. (1984). Studies on successive cambia of a few plants. *Indian J. Bot.* 7, 183-189.
74. Yang, M.G., and Mickelse, O. (1968). Cycad Husk from Guam - Its Toxicity to Rats. *Economic Botany* 22, 149-&.

75. ARKive - Images of Life on Earth - <http://www.arkive.org/> - [Date accessed: 26th of January 2010]
76. Palm and Cycad Societies of Australia (PACSOA) - <http://www.pacsoa.org.au/> - [Date accessed: 26th of January 2010]
77. The Gymnosperm Database - <http://www.conifers.org/> - [Date accessed: 26th of January 2010]
78. Terrazas, T. (1991). Origin and Activity of Successive Cambia in *Cycas* (Cycadales). American Journal of Botany 78, 1335-1344.
79. Flowers of India - <http://www.flowersofindia.net/> - [Date accessed: 26th of January 2010]
80. Whitelock, L.M. (2002). The Cycads. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
81. New South Wales Flora Online - Botanic Gardens Trust - Sydney Australia - <http://plantnet.rbgsyd.nsw.gov.au/> - [Date accessed: 26th of January 2010]
82. Kodandapani, N., Cochrane, M.A., and Sukumar, R. (2008). A comparative analysis of spatial, temporal, and ecological characteristics of forest fires in seasonally dry tropical ecosystems in the Western Ghats, India. Forest Ecology and Management 256, 607-617.
83. Nair, M.N.B., and Ram, H.Y.M. (1990). Structure of Wood and Cambial Variant in the Stem of *Dalbergia paniculata* Roxb. Iawa Bulletin 11, 379-391.
84. Okafor, J. (1999). The use of farmer knowledge in non-wood forest product research. In: Current research issues and prospects for conservation and development. - Accessed from the World Wide Web on the 26th of January 2010: <http://www.fao.org/documents/>
85. Standley, P.C., Williams, L.O., and Gibson, D.N. (1970). Flora of Guatemala. Fieldiana: Botany 24.
86. The International Plant Names Index - <http://www.ipni.org/> - [Date accessed: 26th of January 2010]
87. Carlquist, S. (2000). Wood and stem anatomy of phytolaccoid and rivinoid Phytolaccaceae (Caryophyllales): ecology, systematics, nature of successive cambia. Aliso 19, 13-29.
88. dos Anjos, L., Volpato, G.H., Lopes, E.V., Serafini, P.P., Poletto, F., and Aleixo, A. (2007). The importance of riparian forest for the maintenance of bird species richness in an Atlantic Forest remnant, southern Brazil. Revista Brasileira De Zoologia 24, 1078-1086.
89. Markesteijn, L., Poorter, L., and Bongers, F. (2007). Light-dependent leaf trait variation in 43 tropical dry forest tree species. American Journal of Botany 94, 515-525.
90. Petit Lobão, E.V., and Valeri, S.V. (2007). Agroecossistema cacauceiro da bahia: cacauceira e fragmentos florestais na conservação de espécies arbóreas. Jaboticabal.
91. Augustine, A.C., and DSouza, L. (1997). Somatic embryogenesis in *Gnetum ula* Brongn (*Gnetum edule*) (Willd) Blume. Plant Cell Reports 16, 354-357.
92. Wikipedia - The Free Encyclopedia - <http://wikipedia.org/> - [Date accessed: 26th of January 2010]
93. Endlicher, S.L.F. (1847). Synopsis Coniferarum. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
94. Wild World - <http://www.worldwildlife.org/wildworld/> - [Date accessed: 26th of January 2010]
95. Reddy, S.M. (2003). University Botany 2. Gymnosperms, Plant Anatomy, Genetics, Ecology. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.be/>
96. Merrill, E.D. (1918). New or noteworthy Philippine plants. The Philippine Journal of Science 13(1), January.
97. Al-Khalifah, N.S., and Shanavaskhan, A.E. (2007). On the distribution, status and phenology of Ghada (*Haloxylon persicum* Bunge) in the Arabian Peninsula. Tropical Ecology 48, 51-60.
98. Warren, A. (1981). Arid-Land Ecosystems - Structure, Functioning and Management, Vol 1 - Goodall, Dw, Perry, Ra. Geographical Journal 147, 215-217.
99. Flora de la Region del Parque Nacional Amoro - <http://www.nybg.org/botany/nee/> - [Date accessed: 26th of January 2010]
100. ParksWatch - <http://www.parkswatch.org/> - [Date accessed: 26th of January 2010]
101. DeWalt, S.J., Ickes, K., Nilus, R., Harms, K.E., and Burslem, D.F.R.P. (2006). Liana habitat associations and community structure in a Bornean lowland tropical forest. Plant Ecology 186, 203-216.
102. McDonald, J.A. (1992). Evolutionary Implications of Typical and Anomalous Secondary Growth in Arborescent *Ipomoea* (Convolvulaceae). Bulletin of the Torrey Botanical Club 119, 262-267.
103. Center for Sonoran Desert Studies - Arizona-Sonora Desert Museum - <http://www.desertmuseum.org/center/> - [Date accessed: 26th of January 2010]
104. Llamas, K. (2003). Tropical Flowering Plants: A guide to identification and cultivation. - Accessed from the World Wide Web on the 26th of January 2010: <http://ebook30.com/>
105. Floridata - <http://www.floridata.com> - [Date accessed: 26th of January 2010]
106. Pérez-García, E.A., Meave, J., and Gallardo, C. (1988). Vegetación y Flora de la región de Nizanda, istmo de Tehuantepec, Oaxaca, México. Acta Botanica Mexicana 56.
107. NewCROP - Center for New Crops & Plant Products - Purdue University - <http://www.hort.purdue.edu/newcrop/> - [Date accessed: 26th of January 2010]

108. Parra-Tabla, V., and Bullock, S.H. (2005). Ecological and selective effects of stigma-anther separation in the self-incompatible tropical tree *Ipomoea wolcottiana* (Convolvulaceae). *Plant Systematics and Evolution* 252, 85-95.
109. Forest Department Sarawak - <http://www.forestry.sarawak.gov.my/> - [Date accessed: 26th of January 2010]
110. Leon-Gomez, C., and Monroy-Ata, A. (2005). Seasonality in cambial activity of four lianas from a Mexican lowland tropical rainforest. *Iawa Journal* 26, 111-120.
111. Discover Life - <http://www.discoverlife.org/> - [Date accessed: 26th of January 2010]
112. Carlquist, S. (2007). Successive cambia in Aizoaceae: products and process. *Botanical Journal of the Linnean Society* 153, 141-155.
113. Croat, T.B. (1978). Flora of Barro Colorado Island. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
114. Schmitz, E., Lötter, M., and McClelland, W. (2002). Trees and shrubs of Mpumalanga and Kruger National Park. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
115. Palmer, T. (2004). Vegetation of Makana. Makana LEAP: Comprehensive Environmental Audit - Accessed from the World Wide Web on the 26th of January 2010: <https://www.ru.ac.za/documents/Environment/b%20Vegetation%20of%20Makana.pdf>
116. Hawaiian Native Plant Propagation Database - College of Tropical Agriculture and Human Resources - University of Hawai at Manoa - <http://pdc.ctahr.hawaii.edu:591/hawnprop/> - [Date accessed: 26th of January 2010]
117. Gillespie, L.J., Hollowell, T., Funk, V.A., and Kelloff, C.L. (2003). Smithsonian Plant Collections, Guyana: 1989-1991, Lynn J. Gillespie. Contributions from the United States National Herbarium 44, 1-104.
118. Kabir, M.E., and Webb, E.L. (2006). Saving a forest: the composition and structure of a deciduous forest under community management in northeast Thailand. *Nat. Hist. Siam Soc.* 54, 239-260.
119. MacBride, J.F. (1961). Flora of Peru. Botanical Series XIII (1). - Accessed from the World Wide Web on the 26th of January 2010: http://www.archive.org/stream/floraofperueuphofimacb/floraofperueuphofimacb_djvu.txt
120. León, B., Pitman, N., and Roque, J. (2006). Introducción a las plantas endémicas del Perú. *Rev. Peru. Biol.* Numero especial 13, 9-22.
121. Lombardi, J.A., and Temponi, L.G. (1999). A new species of *Peritassa* Miers (Hippocrateaceae) from Southern Brazil, and notes on two confused species. *Novon* 9, 221-226.
122. de Araújo, D.S.D., and contributors, a. (2009). Área de proteção ambiental de Massambaba, Rio de Janeiro: caracterização fitofisionômica e florística. *Rodriguésia* 60, 67-96.
123. Smithsonian Tropical Research Institute Herbarium - <http://biogeodb.stri.si.edu/herbarium/> - [Date accessed: 26th of January 2010]
124. Carlquist, S. (1998). Wood and stem anatomy of *Petiveria* and *Rivina* (Caryophyllales); systematic implications. *Iawa Journal* 19, 383-391.
125. Garcia-Mateos, M.R., Sanchez, E.E., Espinosa-Robles, P., and Alvarez-Sanchez, M.E. (2007). Toxicity of *Petiveria alliacea* L. on Greenhouse Whitefly (*Trialeurodes vaporariorum* West.). *Interciencia*.
126. Gargiullo, M.B., Magnuson, B.L., and Kimball, L.D. (2008). A field guide to plants of Costa Rica. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
127. Wheat, D. (1977). Successive Cambia in Stem of *Phytolacca dioica*. *American Journal of Botany* 64, 1209-1217.
128. Esser, K.B., Semagn, K., and Wolde-Yohannes, L. (2003). Medicinal use and social status of the soap berry endod (*Phytolacca dodecandra*) in Ethiopia. *Journal of Ethnopharmacology* 85, 269-277.
129. Friis, I., and Vollesen, K. (1998). Flora of the Sudan-Uganda Border Area East of the Nile: I. Catalogue of vascular plants, 1st part. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
130. Wayne's Word - An On-line Textbook of Natural History - <http://waynesword.palomar.edu/> - [Date accessed: 26th of January 2010]
131. Studholme, W.P., and Philipson, W.R. (1966). A comparison of the cambium in two woods with included phloem: *Heimerliodendron brunonianum* and *Avicennia resinifera*. *New Zealand Journal of Botany* 4, 355-365.
132. Hawaiian Ethnobotany Online Database - <http://www2.bishopmuseum.org/ethnobotanydb/index.asp> - [Date accessed: 26th of January 2010]
133. Rodríguez, R.R. (2001). Variation in *Polygala guantanamana* (Polygalaceae), a cuban endemic species. *Willdenowia* 31.
134. Google Directory / Flora and Fauna - http://directory.google.com/Top/Science/Biology/Flora_and_Fauna/ - [Date accessed: 26th of January 2010]
135. Carlquist, S. (2001). Wood and stem anatomy of Rhabdodendraceae is consistent with placement in Caryophyllales *sensu lato*. *Iawa Journal* 22, 171-181.

136. Addo-Fordjour, P., Anning, A.K., Larbi, J.A., and Akyeampong, S. (2009). Liana species richness, abundance and relationship with trees in the Bobiri forest reserve, Ghana: Impact of management systems. *Forest Ecology and Management* 257, 1822-1828.
137. TropHort - Life and Earth Sciences - <http://trophort.com/> - [Date accessed: 26th of January 2010]
138. Plotkin, M.J., and Famolare, L. (1992). Sustainable harvest and marketing of rain forest products. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
139. Smith, A.C. (1940). The American species of Hippocrateaceae. *Brittonia* 3, 341-555.
140. Janick, J., and Paull, R.E. (2008). The Encyclopedia of Fruit & Nuts. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
141. Levey, D.J., Silva, W.R., and Galetti, M. (2002). Seed dispersal and frugivory: ecology, evolution and conservation -Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
142. Stanton, J.P., and Fell, D. (1940). The rainforests of Cape York Peninsula In Cooperative Research Centre for Tropical Rainforest Ecology and Management. p. 177.
143. Martins, V.L.C. (2005). Tipos das plantas vasculares do herbario do museu nacional (R) -XXIV: Gentianaceae, Haloragaceae, Hippocrateaceae, Humiriaceae, Icacinaceae, Verbenaceae, Violaceae e Vitaceae. *Boletim do museu nacional. Botânica* 126.
144. Flora Zambesiaca - Kew Gardens - <http://apps.kew.org/efloras/> - [Date accessed: 26th of January 2010]
145. Bongers, F., Charles-Dominique, P., Forget, P.-M., and Théry, M. (2001). Nouragues. Dynamics and plant-animal interactions in a neotropical rainforest. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.be>
146. Hanelt, P., Büttner, R., and Mansfeld, R. (2001). Mansfeld's Encyclopedia of Agricultural and Horticultural Crops (except ornamentals). - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
147. Virtual Herbarium Berlinense - Freie Universität Berlin - <http://ww2.bgbm.org/herbarium/> - [Date accessed: 26th of January 2010]
148. Poorter, L., Bongers, F., Koeamé, F.N., and Hawthorne, W.D. (2004). Biodiversity of West African Forests. An Ecological Atlas of Woody Plant Species -Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
149. Missouri Botanical Garden - <http://www.mobot.org> - [Date accessed: 26th of January 2010]
150. Oliveira-Filho, A.T., and Ratter, J.A. (1994). Database: woody flora of 106 forest areas of eastern tropical South America. Occasional Monographs from the Royal Botanic Garden Edinburgh - Accessed from the World Wide Web on the 26th of January 2010: <http://www.icb.ufmg.br/treetlan/Downloads/t02.pdf>
151. Khare, C.P. (2004). Indian Herbal Remedies. Rational Western Therapy, Ayurvedic and Other Traditional Usage, Botany. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.com/>
152. AgroForestryTree Database - <http://www.worldagroforestrycentre.org/SEA/Products/AFDbases/AF/> - [Date accessed: 26th of January 2010]
153. NatureServe Explorer - An Online Encyclopedia of Life - <http://www.natureserve.org/explorer/> - [Date accessed: 26th of January 2010]
154. Plants For A Future - Edible, medicinal and useful plants for a healthier world - <http://www.pfaf.org/> - [Date accessed: 26th of January 2010]
155. Nabe-Nielsen, J. (2001). Diversity and distribution of lianas in a neotropical rain forest, Yasuni National Park, Ecuador. *Journal of Tropical Ecology* 17, 1-19.
156. Burnham, R.J. (2004). Alpha and beta diversity of Lianas in Yasuni, Ecuador. *Forest Ecology and Management* 190, 43-55.
157. Graham, A., and Dilcher, D.L. (1998). Studies in neotropical paleobotany. XII. A palynoflora from the Pliocene Rio Banano Formation of Costa Rica and the Neogene vegetation of Mesoamerica. *American Journal of Botany* 85, 1426-1438.
158. Fischer, A.F. (1921). Minor products from Philippine forests. *Bulletin Nr 22 III* - Accessed from the World Wide Web on the 26th of January 2010: http://www.archive.org/stream/minorproductsofp03brow/minorproductsofp03brow_djvu.txt
159. Acevedo-Rodríguez, P., and Axelrod, F.S. (1999). Annotated Checklist for the Tracheophytes of Río Abajo Forest Reserve, Puerto Rico. *Caribbean Journal of Science* 35, 265-285.
160. Oliveira-Filho, A.T., Jarenkow, L.A., and Rodal, M.J.N. (2006). Floristic relationships of seasonally dry forests of Eastern South America based on tree species distribution patterns. In: *Neotropical Savannas and Seasonally Dry Forests: Plant Diversity, Biogeography, and Conservation*. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.be/>
161. Crabb, G. (1823). Universal technological dictionary on familiar explanation of the terms used in all arts and sciences. - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.be>

162. da Cunha, C.N., Junk, W.J., and Leitao-Filho, H.F. (2007). Woody vegetation in the Pantanal of Mato Grosso, Brazil: a preliminary typology. *Amazoniana-Limnologia Et Oecologia Regionalis Systemae Fluminis Amazonas* 19, 159-184.
163. Fire Effects Information System - USDA Forest Service - <http://www.fs.fed.us/database/feis/> - [Date accessed: 26th of January 2010]
164. Bailey, D.C. (1980). Anomalous Growth and Vegetative Anatomy of *Simmondsia chinensis*. *American Journal of Botany* 67, 147-161.
165. The International Jojoba Export Council - <http://www.ijec.net/> - [Date accessed: 26th of January 2010]
166. Utteridge, T., Nagamasu, H., Teo, S.P., White, L.C., and Gasson, P. (2005). *Sleumeria* (Icacinaceae): A new genus from northern Borneo. *Systematic Botany* 30, 635-643.
167. Nair, M.N.B. (1993). Structure of Stem and Cambial Variant in *Spatholobus roxburghii* (Leguminosae). *Iawa Journal* 14, 191-204.
168. Horak, K.E. (1981). Anomalous Secondary Thickening in *Stegnosperma* (Phytolaccaceae). *Bulletin of the Torrey Botanical Club* 108, 189-197.
169. AllExperts - <http://www.allexperts.com/> - [Date accessed: 26th of January 2010]
170. Sagebud - The Social Network For Gardeners - <http://sagebud.com/> - [Date accessed: 26th of January 2010]
171. Carlquist, S. (1999). Wood and stem anatomy of *Stegnosperma* (Caryophyllales); phylogenetic relationships; nature of lateral meristems and successive cambial activity. *Iawa Journal* 20, 149-163.
172. Ives, R.L. (1964). Vegetation and Flora of the Sonoran Desert - Shreve, F., Wiggins, I.L. *Journal of Geography* 63, 288-290.
173. Shreve, F., and Wiggins, I.L. (1964). Vegetation and Flora of the Sonoran Desert. *Journal of Geography* 63, 288-290.
174. Pacific Island Ecosystems at Risk (PIER) - Institute of Pacific Islands Forestry - <http://www.hear.org/pier/> - [Date accessed: 26th of January 2010]
175. Dobbins, D.R., and Fisher, J.B. (1986). Wound Responses in Girdled Stems of Lianas. *Botanical Gazette* 147, 278-289.
176. BugwoodWiki - <http://wiki.bugwood.org/> - [Date accessed: 26th of January 2010]
177. Tripathi, K.P., and Singh, B. (2009). Species diversity and vegetation structure across various strata in natural and plantation forests in Katerniaghat Wildlife Sanctuary, North India. *Tropical Ecology* 50, 191-200.
178. Behrens, J.W., and Uhlworm, O. (2010). *Botanisches Centralblatt; Referirendes Organ.* - Accessed from the World Wide Web on the 26th of January 2010: <http://books.google.be>
179. Flora of Zimbabwe - <http://www.zimbabweflora.co.zw/> - [Date accessed: 26th of January 2010]
180. ZipcodeZoo - <http://zipcodezoo.com/> - [Date accessed: 26th of January 2010]
181. University of Florida Herbarium Collections Catalog - <http://www.flmnh.ufl.edu/herbarium/cat/> - [Date accessed: 26th of January 2010]
182. Tannus, J.L.S., and Assis, M.A. (2005). Vascular species composition of dry and swamp grasslands in a savanna area, Itirapina - SP, Brazil. *Revista Brasileira De Botânica* 27.
183. Global Biodiversity Information Facility - <http://www.gbif.net/> - [Date accessed: 26th of January 2010]
184. Global Invasive Species Database - <http://www.invasivespecies.net/database/> - [Date accessed: 26th of January 2010]