

Published in final edited form as:

*Psychiatry Res.* 2013 August 30; 209(1): 60–65. doi:10.1016/j.psychres.2013.03.017.

## Suicidal ideation and risk factors in primary care patients with anxiety disorders

Jessica Bomyea<sup>a</sup>, Ariel J. Lang<sup>b,c</sup>, Michelle G. Craske<sup>d</sup>, Denise Chavira<sup>b</sup>, Cathy D. Sherbourne<sup>e</sup>, Raphael D. Rose<sup>d</sup>, Daniela Golinelli<sup>e</sup>, Laura Campbell-Sills<sup>b</sup>, Stacy S. Welch<sup>f</sup>, Greer Sullivan<sup>e,g,h</sup>, Alexander Bystritsky<sup>i</sup>, Peter Roy-Byrne<sup>f</sup>, and Murray B. Stein<sup>b,j</sup>

<sup>a</sup>SDSU/UCSD Joint Doctoral Program in Clinical Psychology, University of California, San Diego

<sup>b</sup>Department of Psychiatry, University of California, San Diego

<sup>c</sup>VA San Diego Health Care System Center of Excellence for Stress and Mental Health, San Diego, California

<sup>d</sup>Department of Psychology, University of California, Los Angeles

<sup>e</sup>RAND Corporation, Santa Monica, California

<sup>f</sup>Department of Psychiatry and Behavioral Sciences, University of Washington School of Medicine and Harborview Center for Healthcare Improvement for Addictions, Mental Illness, and Medically Vulnerable Populations (CHAMMP), Seattle, Washington

<sup>g</sup>Department of Psychiatry University of Arkansas for Medical Sciences, Little Rock

<sup>h</sup>VA South Central Mental Illness Research, Education, and Clinical Center University of Arkansas for Medical Sciences, Little Rock

<sup>i</sup>Psychiatry and Biobehavioral Sciences, David Geffen School of Medicine, University of California, Los Angeles

<sup>j</sup>Family and Preventive Medicine, University of California, San Diego

### Abstract

© 2013 Elsevier Ireland Ltd. All rights reserved.

Correspondence to: Jessica Bomyea.

Jessica Bomyea, MS and Ariel J. Lang, Ph.D., MPH, 8939 Villa La Jolla Dr. Suite 200, San Diego, CA 92037, Phone: 858-534-6438, Bomyea@hotmail.com, Fax: 858-534-6460

Michelle Craske, UCLA Anxiety Disorders Research Center, Department of Psychology, Franz Hall - Box 951563, Los Angeles, CA 90094-1563, Phone: 310-206-9191

Denise Chavira, 8939 Villa La Jolla Dr. Suite 200, San Diego, CA 92037, Phone: 858-246-0623

Cathy Sherbourne, 1776 Main Street, Santa Monica, CA, USA 90401, Phone: 310-393-0411

Raphael Rose, UCLA Anxiety Disorders Research Center, Department of Psychology, Franz Hall - Box 951563, Los Angeles, CA 90094-1563, Phone: 310-206-9191

Danielle Golinelli, 1776 Main Street, Santa Monica, CA, USA 90401, Phone: 310-393-0411

Laura Campbell-Sills, 8939 Villa La Jolla Dr. Suite 200, San Diego, CA 92037, Phone: 858-534-6448

Stacy Shaw Welch, 1200 5<sup>th</sup> Avenue Suite 800, Seattle WA 92101, Phone: 206-374-0109

Greer Sullivan, 4301 W. Markham # 554, Little Rock, AR 72205, Phone: 501-526-8100

Alexander Bystritsky, UCLA Psych and Behav Sciences, Box 956969 200 Medical Plaza Ste 2200, Los Angeles CA 90095, Phone: 310-206-5133

Peter Roy-Byrne, 325 9<sup>th</sup> Ave PSB – 5020, Box 359911, Seattle WA 98104, Phone: 206-897-4201

Murray B. Stein M.D., MPH, UCSD Department of Psychiatry, 8939 Villa La Jolla Dr. Suite 200, San Diego, CA 92037, Phone: 858-534-6451

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The presence of an anxiety disorder is associated with greater frequency of suicidal thoughts and behaviors. Given the high personal and societal costs of suicidal behaviors, suicide prevention is a priority. Understanding factors present within individuals with anxiety disorders that increase suicide risk may inform prevention efforts. The aims of the present study were to examine the prevalence of suicidal ideation and behaviors, as well as factors associated with suicide risk in patients with anxiety disorders in primary care. Data from a large scale randomized controlled study were analyzed to assess prevalence of suicidal thoughts and behaviors, as well as factors associated with suicide risk. Results revealed that suicidal ideation and behaviors were relatively common in this group. When examining mental and physical health factors jointly, presence of depression, mental health-related impairment, and social support each uniquely accounted for variance in suicide risk score. Methodological limitations include cross-sectional data collection and lack of information on comorbid personality disorders. Moreover, patients included were from a clinical trial with exclusion criteria that may limit generalizability. Results highlight the complex determinants of suicidal behavior and the need for more nuanced suicide assessment in this population, including evaluation of comorbidity and general functioning.

### Keywords

suicidality; depression; functional impairment; PTSD; GAD; Social anxiety; panic disorder

---

## 1. Introduction

Global estimates suggest that each year there are 10–20 million suicide attempts and one million completed suicides (World Health Organization, 1999). Suicide attempts are costly in terms of occupational and interpersonal disruption (American Foundation for Suicide Prevention, 2009). Moreover, substantial financial costs are associated with the intensive psychiatric resources devoted to these patients (Rissmiller et al., 2004). Thus, understanding features and correlates of suicide is a critical public health matter.

One challenge in studying suicide is the relatively low base rate of attempted and completed suicides observed in most populations. Thus, valuable information may be gained by studying factors that are associated with increased risk of suicide. Suicidal ideation and past suicide attempts are associated with risk for future suicide attempts and completions (Harris and Barraclough, 1997; Kessler et al., 1999; Kuo et al., 2001; Joiner et al., 2003; ten Have et al., 2009). Studying such thoughts and behaviors, therefore, may increase our understanding of who is likely to attempt or complete suicide, potentially informing suicide risk management and prevention efforts.

The presence of current or lifetime anxiety disorders, including panic disorder (PD), social phobia (SP), generalized anxiety disorder (GAD), and posttraumatic stress disorder (PTSD), is associated with increased suicide risk (Khan et al., 2002; Cogle et al., 2009a, b; Nepon et al., 2010; Nock et al., 2010; Lopez-Castroman et al., 2011). Individuals with anxiety disorders demonstrate increased suicidal ideation (Sareen et al., 2005a) and rates of self injury (Chartrand et al., in press) and more frequent suicide attempts (Bolton et al., 2008; Sareen et al., 2005a, b) than those without mental health disorders. For this reason, understanding factors associated with suicide risk within individuals with anxiety disorders is critical to managing this population.

In spite of the increased risk of suicide within anxiety disordered patients, the specific determinants of risk within this group are poorly understood. Studies of individuals with specific anxiety disorders including PD, SP, and PTSD indicate that the additional presence of depression and substance use disorders increases suicide risk (Noyes et al., 1991; Lepine

et al., 1993; Warshaw et al., 1995; Warshaw et al., 2000). Severity and aspects of functional impairment are also related to suicide risk in patients with anxiety disorders. For example, anxiety symptom severity is associated with increased suicidal ideation and attempts in PD and PTSD patients (Noyes et al., 1991; Freeman and Moore, 2000; Huang et al., 2010). Associations between suicide risk and impairment in general functioning in PTSD (Tarrier and Gregg, 2004; Panagioti et al., 2011) and social functioning in both PTSD and PD (Noyes et al., 1991; Huang et al., 2001; Panagioti et al., 2011) are also documented. However, few studies in patients with anxiety disorders comprehensively account for mental *and* physical health conditions and impairment that may be associated with suicide risk (although see Freeman and Moore, 2000 for an exception). The high comorbidity of anxiety disorders and physical health problems (Maier and Falkai, 1999; Levinson et al., 2008; Castro et al., 2009; Nicolson et al., 2009) suggests that the relationship between physical health and suicide in this population merits consideration. Physical health factors generally associated with suicide risk include pain (Braden and Sullivan, 2008; Ilgen et al., 2008), chronicity of medical conditions (Goodwin et al., 2003; Bartels et al., 2007; Robson et al., 2010), and functional limitation due to physical conditions (Kaplan et al., 2007; Park et al., 2010). Thus, our aim was to examine the potential contribution of these factors, along with other established predictors, to suicidality in individuals with anxiety disorders.

Primary care is an important setting to examine correlates of suicide risk in patients with anxiety disorders for a number of reasons. Anxiety and comorbid physical health conditions often present in primary care settings (Serrano-Blanco et al., 2010), and patients with anxiety disorders are often diagnosed and treated in primary care settings (Schulberg, 1995; Bijl and Ravelli, 2000; Price et al., 2000; Shear and Weisberg et al., 2007). Moreover, many individuals who eventually commit suicide present to primary care within weeks or days of making a suicide attempt (Luoma et al., 2002). Knowledge of potential risk factors could improve risk management in this setting by alerting providers to critical features of patients who may be likely to attempt suicide.

Given the great social and economic burden associated with suicide and its relatively high prevalence in primary care, examination of the features of suicide risk behaviors in this setting is warranted. The present study examined suicide risk variables in a sample of individuals with one or more anxiety disorders referred from primary care as part of the Coordinated Anxiety Learning and Management (CALM) study (Roy-Byrne et al., 2010). The first aim of the present study was to first examine the prevalence of suicidal ideation and behaviors in a large group of individuals recruited from the primary care setting. Second, we sought to examine which clinical factors were associated with suicide risk in these patients. To do so, we examined correlational and regression analyses of factors previously documented to be associated with suicide (including mental and physical health variables, as well as social support), and suicide risk score.

## 2. Methods

### 2.1 Participants

Participants were 1620 individuals who completed an initial eligibility assessment in a randomized controlled effectiveness trial comparing the CALM intervention to usual care conducted between June 2006 and August 2008 (UC; Roy-Byrne et al., 2010; [clinicaltrials.gov](http://clinicaltrials.gov) Identifier NCT00347269). Study procedures were reviewed and approved by the institutional review boards at all study sites. Eligible participants were between the ages of 18 and 75, met DSM-IV criteria for at least 1 anxiety disorder (SP, GAD, PTSD or PD), and indicated a score of 8 or greater (moderate anxiety symptoms on a scale ranging from 0–20) on the Overall Anxiety Severity and Impairment Scale (OASIS; Campbell-Sills et al., 2009). All participants who completed the eligibility screening are reported for

analyses of prevalence of suicide risk variables. Because excluded participants did not complete a comprehensive assessment battery, only those meeting inclusion criteria (N = 1002) were considered for regression analyses of suicide risk. Including the larger initial sample of individuals who conducted the screening allowed for examination of baseline rates *before* individuals who met exclusion criteria (including suicidality) were removed from the trial. Information on both the full sample and the screened sample are reported in the prevalence data for suicide risk variables. Exclusion criteria included the presence of life-threatening medical conditions, marked cognitive impairment, active suicidal or homicidal intent or plan, Bipolar I disorder, current substance dependence (except alcohol and marijuana), ongoing treatment in a cognitive behavioral intervention program, and inability to speak either English or Spanish. A total of 10 individuals were removed from the study for suicide or homicide-related reasons. Potential participants were referred from primary care physicians in 17 primary care clinics from 4 U.S. regions. Table 1 presents demographic data for the sample.

## 2.2 Assessments

**2.2.1. Diagnostic status and suicide risk score**—A trained study clinician administered the Mini International Neuropsychiatric Interview (MINI; Sheehan et al., 1998) to participants to determine diagnostic status (including presence of current SUDs and MDD) and assess for study eligibility. As part of this assessment, individuals completed a suicide risk module in which they were asked about the presence and severity of risk factors for future suicidal action. Specifically, participants were asked about a history of accidents with intention of self harm, passive suicidal ideation (“thinking you were better off dead”), desire for self harm, thoughts of suicide including frequency and intensity, and history of suicide attempts. Participants endorsing suicidal ideation were also asked about suicide plan and/or action upon plan, self injury, current intention for self harm, and self-reported impulse control. Consistent with prior studies, a suicide risk score (ranging from 0–33) was calculated by weighting and summing items from this module (e.g., Tang et al., 2010; Quevedo et al., 2011; Pompilli et al., in press). Items that are more strongly associated with suicide are given higher weight.

**2.2.2. Mental and health variables**—Participants were contacted via telephone by the RAND Corporation to complete a baseline assessment that included assessment of physical and mental health disability and social support. Symptom severity was operationalized using psychic and somatic anxiety symptoms assessed using 12 items from the anxiety and somatization subscales of the 18-item Brief Symptom Inventory (BSI-12; Derogatis and Savitz, 2000). Mental health-related disability was assessed using the Sheehan Disability Scale (SDS; Sheehan et al., 1996), which asks participants to rate how much their anxiety and depression interferes with occupational, social, and family/home responsibility domains.

**2.2.3. Physical health variables**—Physical health-related functioning (i.e., inability to complete daily tasks of living and work due to physical limitations) was assessed using the physical health composite score of the 12-item short form Heath Survey version 2 (PCS; Ware et al., 2002). Pain was assessed using one item from the EQ-5D (EQ-5D-pain; EuroQol Group, 1990) that asks participants to rate their level of pain on a three-point scale (“I have no/moderate/extreme pain or discomfort”). To assess total number of medical conditions, participants were read a list of serious medical conditions (e.g., asthma, arthritis, cancer) and asked if a doctor had ever told them they had such a condition.

**2.2.4. Social Support**—Four items from the MOS Social Support Survey (MOS-SSS; Sherbourne and Stewart, 1991) were used to assess perceived social support. These items ask about the level of support received by participants during relaxing activities, availability

of help for daily chores if sick, availability of help for dealing with personal problems, and whether or not others are available to exchange loving emotions. All items were rated on a 1–5 point scale ranging from “none of the time” to “all of the time”.

### 2.3 Statistical analysis

We first examined the frequency of suicide risk variables including passive suicidal ideation, thoughts of suicide in the past month, and history of suicide attempts. Each potential explanatory variable was also correlated with total suicide risk score. A hierarchical multiple regression analysis was utilized to describe the association between suicide risk variables and potential explanatory variables. Mental health variables were entered in the first step, followed by physical health variables, and social support was entered in the final step. To adjust for multiple comparisons the critical value was set to  $p < .01$ . For this regression, continuous explanatory variables were dichotomized by comparing those who were one standard deviation above the mean to the remainder of the group (because social support is scored in the reverse direction to other measures, individuals one standard deviation below the mean were compared to the others).

## 3. Results

### 3.1 Frequency of suicidal thoughts and behaviors

Table 2 presents the percentage of individuals endorsing each suicide risk variable question. Results revealed that approximately 26% of patients endorsed passive suicidal ideation while 16% endorsed thoughts of suicide in the last month. Eighteen percent endorsed a history of suicide attempts; number of lifetime attempts in the total sample ranged from 0 to 30 with an average of 1.9 ( $SD = 2.5$ ).

### 3.2 Bivariate associations with suicide risk score

Table 3 presents bivariate correlations between presence of MDD and SUDs, continuous measures of symptom severity, SDS, PCS, EQ-5D pain, and medical conditions and suicide risk score. With the exception of SUDs, all were significantly associated with the suicide risk score. The magnitude of these correlations indicated small associations between each of these variables and the suicide risk score (based on the guidelines of small  $r = .1$ , medium  $r = .30$ , large  $r = .5$  established by Cohen, 1988; Hemphill, 2003).

### 3.3 Regression analysis of suicide risk score

Hierarchical linear regression was used to examine the relationship between suicide risk score and our set of explanatory variables. Table 4 presents results from these analyses. Significant mental health variables included MDD and mental health-related disability. Addition of the physical health variables did not explain additional variance in the model. Social support significantly and independently accounted for variance in suicide risk score.

## 4. Discussion

Results of the present study suggest that suicidal thoughts and behaviors are common among patients with anxiety disorders referred from primary care settings. Passive suicidal ideation was most common among these patients, with over one quarter of the sample endorsing thoughts of death in the past month. Moreover, 16% endorsed thoughts of suicide within the last month and 18% had attempted suicide at some point in their lifetime. These percentages are significantly higher than those observed in population studies (Kessler et al., 1999) and closely approximate those from studies of other anxious samples (Weissman et al., 1989; Cox et al., 1994; Tarrier and Greg, 2004; Huang et al., 2010). Although individuals who were screened for participation in the randomized controlled trial had a relatively lower

prevalence of suicidal thoughts and behaviors, prevalence rates for many of the suicide related items were very similar. Thus, although patients were referred from a primary care setting, the level of presenting suicidal ideation and behavior appears similar to individuals in outpatient mental health care settings.

Examination of individual explanatory variables indicated that although all variables but SUDs were significantly correlated with suicide risk score, the magnitude of these correlations was small. Thus, it appears that these variables individually are relatively weak predictors of suicide risk score. When examining independent variables simultaneously within a linear regression model, mental health variables including presence of MDD and greater mental health-related disability predicted suicide risk score. Physical health factors were not associated with suicide risk score when controlling for mental health, but social support was a significant independent variable in the model. However, the incremental variance explained by adding social support to the model was small (1%). It is also important to note that overall these variables accounted for a relatively small proportion of the variance in overall suicide risk score (approximately 10%). In conjunction with the relatively weak associations found in the correlational analysis, this data suggests that other factors that were not assessed contribute substantially to overall suicide risk score. Nonetheless, these relationships are generally consistent with an established body of literature indicating that mental health factors, including comorbid conditions and disability, are related to suicide risk in individuals with anxiety disorders (Noyes et al., 1991; Lepine et al., 1993; Warshaw et al., 1995; Freeman and Moore, 2000; Warshaw et al., 2000; Huang et al., 2010). The replication of these relationships in the present sample suggests that these patterns generalize to individuals with an anxiety disorder presenting in primary care.

In contrast to prior literature (e.g., Braden et al., 2008, Ilgen et al., 2008), the hypothesized physical health factors did not emerge as independently associated with suicide risk score. Comparing bivariate correlations to the regression analysis suggests that mental health comorbidity may partially account for associations observed between physical health variables and suicide risk score. Alternatively, the measurement of physical health problems may have led to the discrepant findings. Prior studies have found that specific disorders (Goodwin et al, 2003; Maclean et al., 2011) are predictive of suicide risk. If these associations are specific to certain disorders then summing the number of disorders or assessing health status more generally might not adequately capture these relationships. Given that only one published study has examined physical conditions and suicide within individuals with an anxiety disorder (Freeman and Moore, 2000), further study is needed to determine association between these conditions and suicide risk in individuals with anxiety disorders, particularly in those referred from primary care settings. In addition, SUDs were not significantly associated with suicide risk scores in this sample. This is likely due to the exclusion of individuals with addiction problems other than alcohol or marijuana, suggesting that relatively mild use of marijuana and alcohol (below the level of dependence) may not be associated with increased suicide risk.

A number of limitations warrant considering in interpreting these results. First, the study did not assess all potentially relevant comorbid conditions. For example, the present analyses did not include any assessment of personality disorders. Extant literature indicates that these disorders increase suicide risk in patients with anxiety disorders and should thus be included in future research in this area (Nepon et al., 1991). Second, although descriptive data on suicide risk factors was available for a large sample of participants, examination of suicide risk score explanatory variables was done only with individuals enrolled in the clinical trial who met inclusion criteria. Thus, generalizability of this part of the results is limited to this type of population. In addition, data was collected cross-sectionally. Consequently, conclusions about directionality of these relationships or the temporal course of suicidal

thoughts and behaviors and mental and physical health, disability, and social support variables cannot be made. Finally, suicide risk score was utilized as a proxy for suicidal behaviors, but risk score did not translate into attempted or completed suicide in this sample. Given that the current study is limited by the lack of serious suicidal behaviors, future work examining factors associated with suicide attempts and completions in a longitudinal design would provide further information about key predictors to target in suicide prevention efforts.

The present study highlights the relatively high rates of suicidal thoughts and past suicidal behaviors in individuals referred from a primary care setting. Risk may be particularly high in the context of comorbid MDD and functional impairment and low social support. However, future work is needed to determine other potential variables that may impact suicide risk score given that general mental health, physical health, and social support factors appear to play a relatively small role in overall suicide risk scores. While suicide assessment for individuals with MDD has become more commonplace, findings point to the additional need for suicide screening and intervention for individuals with anxiety disorders, particularly those with low occupational and social functioning. Thus, this type of clinical evaluation should also consider level of disability rather than simply diagnostic evaluation. Consideration of impairments in daily activities due to mental health symptoms, as well as the degree of isolation and social support available to patients, may provide critical information about level of risk in primary care samples.

In summary, suicidal ideation and past suicidal behavior were relatively common in this sample of primary care patients with one or more anxiety disorders. Findings are consistent with literature in specific anxious populations suggesting that individuals with greater burden of mental illness in terms of mood disorder comorbidity and impairment as well as relatively low social support are more likely to endorse suicidal ideation and behavior. Findings highlight the importance of suicide risk assessments in this type of patient, and suggest that consideration of depression, disability, and social support may be important for risk management.

## Acknowledgments

This work was supported by grants U01 MH057858 & K24 MH065324 (Dr. Roy-Byrne), U01 MH070022 (Dr. Sullivan), U01 MH058915 (Dr. Craske), U01 MH070018 (Dr. Sherbourne), and U01 MH057835 & K24 MH64122 (Dr. Stein) from the National Institute of Mental Health

## References

- American Foundation for Suicide Prevention. [Accessed 15 October, 2010] Suicide prevention priorities for health reform. 2009. <http://healthyamericans.org/assets/files/Suicideprevention.pdf>
- Bartels SJ, Coakley E, Oxman TE, Constantino G, Oslin D, Chen H, Zubritsky C, Cheal K, Durai UN, Llorente M, Sanchez H. Suicidal and death ideation in older primary care patients with depression, anxiety, and at-risk alcohol use. *The American Journal of Geriatric Psychiatry Special Issue: Suicidal Behaviors in Older Adults*. 2002; 10:417–427.
- Bijl RV, Ravelli A. Psychiatric morbidity, service use, and need for care in the general population: Results of the Netherlands mental health survey and incidence study. *American Journal of Public Health*. 2000; 90:602–607. [PubMed: 10754976]
- Bolton JM, Cox BJ, Afifi TO, Enns MW, Bienvenu OJ, Sareen J. Anxiety disorders and risk for suicide attempts: Findings from the Baltimore epidemiologic catchment area follow-up study. *Depression and Anxiety*. 2008; 25:477–481. [PubMed: 17541978]
- Braden JB, Sullivan MD. Suicidal thoughts and behavior among adults with self-reported pain conditions in the national comorbidity survey replication. *The Journal of Pain*. 2008; 9:1106–1115. [PubMed: 19038772]

- Campbell-Sills L, Norman SB, Craske MG, Sullivan G, Lang AJ, Chavira DA, Bystrisky A, Sherbourne C, Roy-Byrne P, Stein MB. Validation of a brief measure of anxiety related severity and impairment: The overall anxiety severity and impairment scale (OASIS). *Journal of Affective Disorders*. 2009; 112:92–101. [PubMed: 18486238]
- Castro M, Kraychete D, Daltro C, Lopes J, Menezes R, Oliveira I. Comorbid anxiety and depression disorders in patients with chronic pain. *Arquivos De Neuro-Psiquiatria*. 2009; 67:982–985. [PubMed: 20069205]
- Chartrand H, Sareen J, Toews M, Bolton JM. Suicide attempts versus nonsuicidal self-injury among individuals with anxiety disorders in a nationally representative sample. *Depression and Anxiety*. in press.
- Cohen, J. *Statistical power analysis for the behavioral sciences*. Hillsdale, NJ: Lawrence Erlbaum Associations; 1988. p. 77-81.
- Cougle JR, Keough ME, Riccardi CJ, Sachs-Ericsson N. Anxiety disorders and suicidality in the national comorbidity survey replication. *Journal of Psychiatric Research*. 2009a; 43:825–829. [PubMed: 19147159]
- Cougle JR, Resnick H, Kilpatrick DG. PTSD, depression, and other comorbidity in relation to suicidality: Cross-sectional and prospective analyses of a national probability sample of women. *Depression and Anxiety*. 2009b; 26:1151–1157. [PubMed: 19842171]
- Cox BJ, Dorenfeld DM, Swinson RP, Norton GR. Suicidal ideation and suicide attempts in panic disorder and social phobia. *American Journal of Psychiatry*. 1994; 151:882–887. [PubMed: 8184998]
- Derogatis, LR.; Savitz, KL. The SCL-90–R and brief symptom inventory (BSI) in primary care. In: Maruish, ME., editor. *Handbook of psychological assessment in primary care settings*. Mahwah, NJ: Lawrence Erlbaum Associates Publishers; 2000. p. 297-334.
- EuroQol Group. EuroQol—a new facility for the measurement of health related quality of life. *Health Policy*. 1990; 16:199–208. [PubMed: 10109801]
- Freeman TW, Moore WM. A comparison of chronic combat-related posttraumatic stress disorder (PTSD) patients with and without a history of suicide attempt. *Journal of Nervous and Mental Disease*. 2000; 188:460–463. [PubMed: 10919706]
- Goodwin RD, Marusic A, Hoven CW. Suicide attempts in the united states: The role of physical illness. *Social Science & Medicine*. 2003; 56:1783–1788. [PubMed: 12639594]
- Harris EC, Barraclough B. Suicide as an outcome for mental disorders: A meta-analysis. *British Journal of Psychiatry*. 1997; 170:205–228. [PubMed: 9229027]
- Huang M, Yen C, Lung F. Moderators and mediators among panic, agoraphobia symptoms, and suicidal ideation in patients with panic disorder. *Comprehensive Psychiatry*. 2010; 51:243–249. [PubMed: 20399333]
- Ilgen MA, Zivin K, McCammon RJ, Valenstein M. Pain and suicidal thoughts, plans and attempts in the united states. *General Hospital Psychiatry*. 2008; 30:521–527. [PubMed: 19061678]
- Joiner TE, Steer RA, Brown G, Beck AT, Pettit JW, Rudd MD. Worst-point suicidal plans: A dimension of suicidality predictive of past suicide attempts and eventual death by suicide. *Behaviour Research and Therapy*. 2003; 41:1469–1480. [PubMed: 14583414]
- Kaplan MS, McFarland BH, Huguet N, Newsom JT. Physical illness, functional limitations, and suicide risk: a population-based study. *American Journal of Orthopsychiatry*. 2007; 77:56–60. [PubMed: 17352585]
- Kessler RC, Berglund P, Borges G, Nock M, Wang PS. Trends in suicide ideation, plans, gestures, and attempts in the united states, 1990–1992 to 2001–2003. *JAMA: Journal of the American Medical Association*. 2005; 293:2487–2495.
- Kessler RC, Borges G, Walters EE. Prevalence of and risk factors for lifetime suicide attempts in the national comorbidity survey. *Archives of General Psychiatry*. 1999; 56:617–626. [PubMed: 10401507]
- Khan A, Leventhal RM, Khan S, Brown WA. Suicide risk in patients with anxiety disorders: A meta-analysis of the FDA database. *Journal of Affective Disorders*. 2002; 68:183–190. [PubMed: 12063146]



- Kuo W, Gallo JJ, Tien AY. Incidence of suicide ideation and attempts in adults: The 13-year follow-up of a community sample in Baltimore, Maryland. *Psychological Medicine: A Journal of Research in Psychiatry and the Allied Sciences*. 2001; 31:1181–1191.
- Lepine JP, Chignon JM, Teherani M. Suicide attempts in patients with panic disorder. *Archives of General Psychiatry*. 1993; 50:144–149. [PubMed: 8427555]
- Levinson D, Karger CJ, Haklai Z. Chronic physical conditions and use of health services among persons with mental disorders—results from the israel national health survey. *General Hospital Psychiatry*. 2008; 30:226–232. [PubMed: 18433654]
- Lopez-Castroman J, Perez-Rodriguez MDIM, Jaussent I, Alegria AA, Artes-Rodriguez A, Freed P, Gillaumes S, Jollant F, Leiva-Murillo JM, Malafosse A, Oquendo MA, de Prado-Cumplido M, Saiz-Ruiz J, Cava-Garcia E, Courtet P. European Research Consortium for Suicide (EURECA). Distinguishing the relevant features of frequent suicide attempters. *Journal of Psychiatric Research*. 2011; 45:619–625. [PubMed: 21055768]
- MacLean J, Kinley DJ, Jacobi F, Bolton JM, Sareen J. The relationship between physical conditions and suicidal behavior among those with mood disorders. *Journal of Affective Disorders*. 2011; 130:245–250. [PubMed: 21078525]
- Maier W, Falkai P. The epidemiology of comorbidity between depression, anxiety disorders and somatic diseases. *International Clinical Psychopharmacology*. 1999; 14:S1–S6. [PubMed: 10471166]
- Malone KM, Haas GL, Sweeney JA, Mann JJ. Major depression and the risk of attempted suicide. *Journal of Affective Disorders*. 1995; 34:173–185. [PubMed: 7560545]
- Nepon J, Belik S, Bolton J, Sareen J. The relationship between anxiety disorders and suicide attempts: Findings from the national epidemiologic survey on alcohol and related conditions. *Depression and Anxiety*. 2010; 279:791–798. [PubMed: 20217852]
- Nicolson SE, Caplan JP, Williams DE, Stern TA. Comorbid pain, depression, and anxiety: Multifaceted pathology allows for multifaceted treatment. *Harvard Review of Psychiatry*. 2009; 17:407–420. [PubMed: 19968455]
- Nock MK, Hwang I, Sampson NA, Kessler RC. Mental disorders, comorbidity and suicidal behavior: Results from the national comorbidity survey replication. *Molecular Psychiatry*. 2010; 15:868–876. [PubMed: 19337207]
- Noyes R, Christiansen J, Clancy J, Garvey MJ. Predictors of serious suicide attempts among patients with panic disorder. *Comprehensive Psychiatry*. 1991; 32:261–267. [PubMed: 1884606]
- Panagioti M, Gooding PA, Dunn G, Tarrier N. Pathways to suicidal behavior in posttraumatic stress disorder. *Journal of Traumatic Stressm*. 2011; 24:137–145.
- Park S, Cho S, Moon S. Factors associated with suicidal ideation: Role of emotional and instrumental support. *Journal of Psychosomatic Research*. 2010; 69:389–397. [PubMed: 20846540]
- Price D, Beck A, Nimmer C, Bensen S. The treatment of anxiety disorders in a primary care HMO setting. *Psychiatric Quarterly*. 2000; 71:31–45. [PubMed: 10736815]
- Pompili M, Innamorati M, Rihmer Z, Gonda X, Serafini G, Akiskal H, Amore M, Niolu C, Sher L, Tatarelli R, Perugi G, Girardi P. Cyclothymic–depressive–anxious temperament pattern is related to suicide risk in 346 patients with major mood disorders. *Journal of Affective Disorders*. 2012; 13:405–411. [PubMed: 22177743]
- Quevedo L, da Silva RA, Coelho F, Tavares Pinheiro KA, Horta BL, Kapczinski F, Pinheiro RT. Risk of suicide and mixed episode in men in the postpartum period. *Journal of Affective Disorders*. 2011; 132:243–246. [PubMed: 21277023]
- Rissmiller DJ, Steer R, Ranieri WF, Rissmiller F. Factors complicating cost containment in the treatment of suicidal patients. *Hospital & Community Psychiatry*. 1994; 45:782–788. [PubMed: 7982693]
- Robson A, Scrutton F, Wilkinson L, MacLeod F. The risk of suicide in cancer patients: A review of the literature. *Psycho-Oncology*. 2010; 19:1250–1258. [PubMed: 20213857]
- Roy-Byrne P, Craske MG, Sullivan G, Rose RD, Edlund MJ, Lang AJ, Bystritsky A, Welch SS, Chavira D, Golinelli D, Campbell-Sills L, Sherbourne CD, Stein MB. Delivery of evidence-based treatment for multiple anxiety disorders in primary care: A randomized controlled trial. *JAMA: Journal of the American Medical Association*. 2010; 303:1921–1928.

- Sareen J, Cox BJ, Afifi TO, de Graaf R, Asmundson GJG, ten Have M, Stein MB. Anxiety disorders and risk for suicidal ideation and suicide attempts: A population-based longitudinal study of adults. *Archives of General Psychiatry*. 2005a; 62:1249–1257. [PubMed: 16275812]
- Sareen J, Houlahan T, Cox BJ, Asmundson GJG. Anxiety disorders associated with suicidal ideation and suicide attempts in the national comorbidity survey. *Journal of Nervous and Mental Disease*. 2005b; 193:450–454. [PubMed: 15985839]
- Serrano-Blanco A, Palao DJ, Luciano JV, Pinto-Meza A, Luján L, Fernández A, Roura P, Bertsch J, Mercader M, Haro JM. Prevalence of mental disorders in primary care: Results from the diagnosis and treatment of mental disorders in primary care study (DASMAP). *Social Psychiatry and Psychiatric Epidemiology*. 2010; 45:201–210. [PubMed: 19452110]
- Shear MK, Schulberg HC. Anxiety disorders in primary care. *Bulletin of the Menninger Clinic*. 1995; 59:A73–A85. [PubMed: 7795573]
- Sheehan DV, Harnett-Sheehan K, Raj BA. The measurement of disability. *International Clinical Psychopharmacology*. 1996; 11:89–95. [PubMed: 8923116]
- Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, Hergueta T, Baker R, Dunbar GC. The mini-international neuropsychiatric interview (M.I.N.I.): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry*. 1998; 59:22–33. [PubMed: 9881538]
- Sherbourne CD, Stewart AL. The MOS social support survey. *Social Science & Medicine*. 1991; 32:705–714. [PubMed: 2035047]
- Sokero TP, Melartin TK, Rytsälä HJ, Leskelä US, Lestelä-Mielonen PS, Isometsä ET. Suicidal ideation and attempts among psychiatric patients with major depressive disorder. *Journal of Clinical Psychiatry*. 2003; 64:1094–1100. [PubMed: 14628986]
- Tang T, Yen C, Cheng C, Yang P, Chen CS, Yang R, Huang MS, Jong YJ, Yu HS. Suicide risk and its correlate in adolescents who experienced typhoon-induced mudslides: A structural equation model. *Depression and Anxiety*. 2010; 27:1143–1148. [PubMed: 21049526]
- Tarrier N, Gregg L. Suicide risk in civilian PTSD patients: Predictors of suicidal ideation, planning and attempts. *Social Psychiatry and Psychiatric Epidemiology*. 2004; 39:655–661. [PubMed: 15300376]
- ten Have M, de Graaf R, van Dorsselaer S, Verdurmen J, van't Land H, Vollebergh W, Beekman A. Incidence and course of suicidal ideation and suicide attempts in the general population. *The Canadian Psychiatric Association Journal / La Revue De l'Association Des Psychiatres Du Canada*. 2009; 54:824–833.
- Ware, JE., Jr; Kosinski, M.; Turner-Bowker, DM.; Gandek, B. How to Score Version 2 of the SF-12™ Health Survey (With a Supplement Documenting Version 1). Lincoln (RI): QualityMetric Incorporated; 2002. 2002
- Warshaw MG, Dolan RT, Keller MB. Suicidal behavior in patients with current or past panic disorder: Five years of prospective data from the Harvard/Brown anxiety research program. *The American Journal of Psychiatry*. 2000; 157:1876–1878. [PubMed: 11058491]
- Warshaw MG, Massion AO, Peterson LG, Pratt LA. Suicidal behavior in patients with panic disorder: Retrospective and prospective data. *Journal of Affective Disorders*. 1995; 34:235–247. [PubMed: 7560552]
- Weisberg RB, Dyck I, Culpepper L, Keller MB. Psychiatric treatment in primary care patients with anxiety disorders: A comparison of care received from primary care providers and psychiatrists. *The American Journal of Psychiatry*. 2007; 164:276–282. [PubMed: 17267791]
- Weissman MM, Klerman GL, Markowitz JS, Ouellette R. Suicidal ideation and suicide attempts in panic disorder and attacks. *New England Journal of Medicine*. 1989; 321:1209–1214. [PubMed: 2797086]
- World Health Organization (WHO). *Figures and facts about suicide*. Geneva: 1999.

**Table 1**

## Baseline demographic variables

	Number of participants (percentage)
Race:	
Hispanic	195 (19%)
Black	115 (12%)
White	568 (57%)
Other	124 (12%)
Years of education †:	
Less than 12 years	54 (5.4)
12 years	165 (16.5)
More than 12 years	781 (77.9)
Gender:	
Female	713 (71.2)
Male	289 (28.8)
Mean Age:	43.5 (13.5)

† Education data missing for one patient.

*Note:* Table represents N in each cell for race, mean years of education, and gender. Cells represent means and standard deviation for age. Table represents data from the sample of 1002 eligible study participants

**Table 2**

Percent of sample endorsing items from MINI Suicide Risk module

Item	% of total sample endorsing (N =1620)	% of enrolled sample endorsing (N = 1002)
Think that you were better off dead or wish you were dead?	26.2	25.3
Want to harm yourself or to hurt or injure yourself?	6.4	5.1
Think about suicide?	15.6	13.4
Have a suicide plan?	2.5	1.3
Take any active steps to prepare to injure yourself or to prepare for a suicide attempt in which you expected or intended to die?	< 1	<1
Deliberately harm or injure yourself?	4.1	2.2
Attempt suicide (past month)?	< 1	<1
In your lifetime did you ever make a suicide attempt?	18.3	17.8

*Note.* Table represents data from the full sample of 1620 participants assessed for study eligibility as well as the sample of 1002 eligible study participants.

**Table 3**  
Bivariate correlations between mental health, physical health, and social support variables and suicide risk score

Measure	1	2	3	4	5	6	7	8	9
1. SUD	1.00								
2. MDD	0.04	1.00							
3. BSI	0.04	0.29***	1.00						
4. SDS	0.02	0.41***	0.54***	1.00					
5. PCS	0.07	-0.20***	-0.19***	-0.20***	1.00				
6. EQ5D-pain	-0.03	0.24***	0.26***	0.25***	-0.57***	1.00			
7. Medical conditions	-0.06	0.16***	0.16***	0.12***	-0.51***	0.40***	1.00		
8. MOS-SSS	0.003	-0.18***	-0.14***	-0.21***	0.19***	-0.17***	-0.17***	1.00	
9. Suicide risk score	0.07	0.21***	0.21***	0.22***	0.17***	0.17***	0.11**	-0.14***	1.00

Note. SUD/MDD: Presence of a substance use disorder or major depressive disorder; BSI-12: Brief Symptom Inventory-12 (Derogatis & Savitz, 2000), anxiety and somatization subscales, SDS: Sheehan Disability Scale (Sheehan et al., 1996); PCS: 12-item short form Health Survey version 2 (Ware et al., 2002); EQ5D-pain: pain item of the EQ-5D (EuroQol, 1990); MOS-SSS (MOS-Social Support Scale; Sherbourne & Stewart, 1991). For dichotomous variables (SUD, MDD) correlations represent point-biserial correlations. Table represents data from the sample of 1002 eligible study participants.

\*\*  
\* $p < .01$ .

\*\*\*  
 $p < .001$

Table 4

Hierarchical regression analyses predicting suicide risk score from mental health, physical health, and social support variables

Suicide Risk Score				
Variable	<i>B</i>	<i>SE B</i>	$\beta$	$\Delta R^2$
<u>Step 1</u>				0.08 ***
SUD diagnosis	.95	.42	.07	
MDD diagnosis	1.18	.25	.15 ***	
BSI-12	.83	.32	.09 **	
SDS	1.43	.34	.14 ***	
<u>Step 2</u>				.01
SUD diagnosis	1.04	.42	.08	
MDD diagnosis	1.08	.25	.14 ***	
BSI-12	.71	.32	.07	
SDS	1.29	.34	.13 ***	
PCS	-.07	.32	-.01	
GD5D-Pain	.85	.39	.07	
Medical Conditions	.08	.06	.05	
<u>Step 3</u>				0.01 **
SUD diagnosis	1.05	0.42	0.08	
MDD diagnosis	1.02	0.25	0.13 ***	
BSI-12	0.70	0.32	0.07	
SDS	1.25	0.34	0.12 ***	
PCS	-0.03	0.32	-0.003	
GD5D-Pain	0.84	0.38	0.07	
Medical Conditions	0.06	0.06	0.03	
MOS-SSS	-0.83	0.28	-0.09 **	

*Note.* SUD/MDD: Presence of a substance use disorder or major depressive disorder; BSI-12: Brief Symptom Inventory-12 (Derogatis & Savitz, 2000), anxiety and somatization subscales, SDS: Sheehan Disability Scale (Sheehan et al., 1996); PCS: 12-item short form Health Survey version 2 (Ware et al., 2002); EQ5D-pain: pain item of the EQ-5D (EuroQol, 1990); MOS-SSS (MOS-Social Support Sale; Sherbourne & Stewart, 1991). BSI-12, SDS, PCS, EQ-5D and MOS-SSS are dichotomous variables representing those above 1 standard deviation above the mean and those below (1 SD below the mean vs. above for MOS-SSS). Table represents data from the sample of 1002 eligible study participants.

\*\*  
 $p < .01$ .

\*\*\*  
 $p < .001$