

## Sulfate Reduction and Anaerobic Glycerol Degradation by a Mixed Microbial Culture

Abdel-Allah Qatibi,<sup>1,2</sup> André Bories,<sup>1</sup> and Jean-Louis Garcia<sup>2</sup>

<sup>1</sup>Laboratory of Biotechnology of A.A.I. Environment, Narbonne; <sup>2</sup>Laboratory of Microbiology ORSTOM, University of Provence, Marseilles, France

**Abstract.** Anaerobic glycerol degradation by a mixed microbial culture from a fermenter fed with industrial alcohol distillation waste water, was investigated in the absence or presence of sulfate, at 37°C and at a constant pH of 7.2. In the absence of sulfate, glycerol utilization was found to be characterized by the transient formation of 1,3-propanediol prior to propionate and acetate accumulation. In the presence of sulfate, 1,3-propanediol production was minor, and the carbon balance reflected a considerable accumulation of intermediate(s). A study of the role of sulfate reduction and methanogenesis on anaerobic 1,3-propanediol degradation showed that consumption of this substrate by the mixed microbial culture required a terminal electron acceptor. The number of fermentative and sulfate-reducing bacteria with glycerol or 1,3-propanediol as carbon and energy source revealed that sulfate-reducing bacteria outcompete fermentative bacteria for these substrates. The possible ecological role of sulfate-reducing bacteria in the metabolism of these reduced substrates is discussed.

The classical *Desulfovibrio* species utilize a rather limited range of electron donors for sulfate reduction. Lactate, pyruvate, ethanol, malate, formate, and hydrogen are the common energy substrates for these bacteria [27]. Recently, new types of *Desulfovibrio* species were isolated that utilize fructose [6, 26], sucrose [16], and amino acids [35].

It has also been recently reported that glycerol can be used as energy and carbon source by some *Desulfovibrio* species in pure culture [10, 17, 24-26, 28, 30, 31]. Little is known, however, about the utilization of glycerol by sulfate-reducing bacteria [31]. Glycerol is a major component of waste water from bioethanol production plants (distillery stillage), which also contain sulfate [2, 3]; and anaerobic digestion is one of the main purification processes used on these wastes [2, 3].

The purpose of the present study was to examine whether sulfate-reducing bacteria play a role in anaerobic glycerol and 1,3-propanediol digestion, and to quantify the populations involved in this process.

### Materials and Methods

**Sources of microorganisms.** The microbial mixed culture came from a continuous fermenter (20-liter Biolaftite reactor; agitation

rate, 50 rpm, temperature, 37°C; sequential feeding, 1 liter every 3 days with a mean organic load of 0.4 g COD L<sup>-1</sup> d<sup>-1</sup> and hydrolic residence time of 60 days) fed with waste water from an industrial distillery (INRA-Narbonne, France). The main substrates present in the waste were glycerol and lactic acid. Concentrations of sulfate varied between 6 and 12 mM. The inoculum was from an anaerobic lagoon of distillery waste waters [2, 3].

**Media.** The basal sulfate-free salt solution mineral-medium used for batch culture experiments had the following composition (g/L): K<sub>2</sub>HPO<sub>4</sub>, 0.4; KH<sub>2</sub>PO<sub>4</sub>, 0.4; NH<sub>4</sub>Cl, 0.9; NaCl, 0.9; MgCl<sub>2</sub> · 6H<sub>2</sub>O, 0.1; CaCl<sub>2</sub> · 2H<sub>2</sub>O, 0.1; cysteine-HCl · H<sub>2</sub>O (as a reducing agent), 0.4; resazurin, 0.001. A basal bicarbonate buffered and sulfide-reduced medium with vitamins and trace elements was composed and prepared as described by Widdel and Pfennig [39] and modified by Widdel [37]; it was supplemented with 0.01% yeast extract (Difco) for fermentative and sulfate-reducing bacterial counts.

**Growth conditions.** Batch culture experiments were carried out at 37°C either in 1-L magnetically stirred reactors with pH control or in 60-ml serum bottles closed with butyl-rubber stoppers, incubated in a gyratory shaker, without pH control. Prior to the batch culture experiments, the inoculum was anaerobically diluted to 50% (vol/vol) with the sterilized sulfate-free salt solution under O<sub>2</sub>-free N<sub>2</sub>, with a Hungate technique [15] as modified for the use of syringes [20]. The diluted samples were incubated under 100% N<sub>2</sub>, at 37°C, for 3 or 4 days to stimulate bacterial growth and to ensure that the sulfate derived from the samples was completely consumed. Samples were then distributed among reactors and

Address reprint requests to: Dr. J.L. Garcia, ORSTOM Laboratoire de Microbiologie, Université de Provence, 3, place V-Hugo, 13331 Marseille cedex 3, France.

O. R. S. T. O. M. Fonds Documentaire

N° : 34549, ex 1

Cote : B M p 57

30 OCT. 1991

bottles under anaerobic conditions. Organic substrates, sulfate, and chloroform (an inhibitor of methanogenesis) were added from separately sterilized stock solutions.

**Bacterial counts.** Fermentative and sulfate-reducing bacteria were counted by the most probable number (MPN) method with the following substrates: glycerol (20 mM) and 1,3-propanediol (20 mM). Sulfate-reducing bacteria were counted in the presence of sulfate (20 mM). Total flora was estimated by the acridine orange epifluorescence method [14]. The MPN tubes were incubated at 37°C for 4 weeks; growth of fermentative and sulfate-reducing bacteria was determined from optic density measurements at 580 nm in Bausch and Lomb Spectronic 70 spectrophotometer and sulfide production respectively. Metabolites produced from the various substrates were also determined.

**Analytical methods.** Sulfide production could not be demonstrated under these conditions by spectrophotometric methods because of the strong color of the waste used in the experiments. The sulfate reduction activity was, therefore, estimated from sulfate depletion in the culture medium. Sulfate was analyzed by the HPLC technique under the following conditions: pump, Shimadzu LC 6A; eluant, phthalic acid (3 mM) neutralized with Na-tetraborate at pH 4.9; flow rate, 2 ml min<sup>-1</sup>; injection loop, 100  $\mu$ l; column, Vydac anion exchange phase 302 IC (25 cm long); room temperature; conductivity detection was performed with a Shimadzu model CDD 6A detector; recorder integrator, Shimadzu chromatopak CR 3A; external standard. Sulfide was determined spectrophotometrically in the form of colloidal CuS, as described by Cord-Ruwisch [5]. Glycerol was determined by the enzymatic method developed by Batlle and Collon [1]. 1,3-propanediol was measured by means of FID gas chromatography (THEED 10% column on carbopack [SUPELCO, Inc.] 1 m long, 2 mm inner diameter; column temperature, 125°C; carrier gas, nitrogen; internal standard, ethyleneglycol). Volatile fatty acids were analyzed by means of FID gas chromatography (Glass column, 2 m long; 4 mm inner diameter; filled with Chromosorb (SUPELCO, Inc.), coated with 10% SP-1200 and 1% H<sub>3</sub>PO<sub>4</sub>; carrier gas, nitrogen; column temperature, 105°C; internal standard, ethyl 2-butyric acid. Alcohol and nonvolatile fatty acids were analyzed by HPLC measurements (column, Aminex HPX-87H (BIORAD, Richmond); detector, differential refractometer (Knauer); external standard). Methane and hydrogen were measured with TCD gas chromatography as described by Qatibi and Bories [29]. Total organic carbon (TOC) was measured on a Dohrmann DC 80 analyzer [18].

## Results

**Anaerobic glycerol degradation.** In the absence of sulfate, with pH control at 7.2, glycerol was degraded with transient 1,3-propanediol formation before propionate and acetate production occurred; the 1,3-propanediol accumulated was slowly degraded into propionate and acetate (Fig. 1a). The kinetics of the total organic carbon measured and calculated revealed the presence of unknown intermediate(s) (Fig. 1b). In the presence of sulfate (Fig. 2a), with pH maintained at 7.2, glycerol was degraded and a parallel sulfate depletion occurred, co-




Fig. 1. Time course of anaerobic glycerol degradation by (a) the mixed microbial culture and (b) total organic carbon (TOC) calculated and measured. The experiment was performed in the absence of sulfate at constant temperature and pH (37°C and 7.2 respectively). Symbols: ■, glycerol; □, 1,3-propanediol; ▲, propionate; △, acetate; ×, TOC measured; +, TOC calculated.

inciding with a transient formation of 1,3-propanediol and other intermediate(s), before propionate (then converted into acetate by sulfate reduction) and acetate production took place. A little 1,3-propanediol was produced, and the carbon balance (Fig. 2b) showed that considerable accumulation of the intermediate(s) had occurred.

**Effects of sulfate reduction and methanogenesis on anaerobic 1,3-propanediol degradation.** To determine the relationship between sulfate reduction and methanogenesis on anaerobic 1,3-propanediol degradation, experiments were carried out in which chloroform was added to inhibit methanogenesis (final concentration, 0.003% vol/vol) in the absence or presence of sulfate. As shown in Table 1, when




Fig. 2. Time course of anaerobic glycerol degradation by (a) the mixed microbial culture and (b) total organic carbon (TOC) calculated and measured. The experiment was performed in the presence of sulfate at constant temperature and pH (37°C and 7.2 respectively). Symbols: ■, glycerol; □, 1,3-propanediol; ▲, propionate; △, acetate; ◆, sulfate; ×, TOC measured; +, TOC calculated.

chloroform was added to the waste with or without sulfate, methane production from 1,3-propanediol as carbon and energy source was entirely inhibited. In the absence of sulfate, and in the presence of chloroform, hydrogen accumulated; it was consumed when sulfate was added, and the amount of 1,3-propanediol degraded was higher in the presence than in the absence of sulfate. This suggests that 1,3-propanediol degradation required either a terminal electron acceptor such as sulfate or syntrophic cooperation with methanogenic hydrogenotrophic bacteria.

**Bacterial counts.** Table 2 shows results from bacterial counts. In total, approximately  $10^{10}$  bacterial

cells were counted per ml by the epifluorescence technique with acridine orange. In low-sulfate glycerol medium,  $2.5 \times 10^9$  fermentative bacteria were estimated per ml by MPN counts. In the last positive tubes, propionate, acetate, methane, and presumably carbon dioxide were produced. *Propionobacterium* sp., *Desulfovibrio* sp., and *Methanibacterium* sp. were observed by microscopic examination. In high-sulfate glycerol medium,  $1.4 \times 10^9$  sulfate-reducing bacteria were counted per ml; acetate, sulfide, and presumably carbon dioxide accumulated. Species of *Desulfovibrio* were identified as the main glycerol degraders, from their morphology and fermentation products;  $4.0 \times 10^7$  fermentative bacteria were counted per ml with 1,3-propanediol as sole substrate in the absence of sulfate. Putative species of *Pelobacter* and *Desulfovibrio* were observed in the last positive tubes; propionate, acetate, methane, and presumably carbon dioxide were produced, whereas in the presence of sulfate,  $2.5 \times 10^7$  sulfate-reducing bacteria were estimated per ml. Species of *Desulfovibrio* were identified as the main 1,3-propanediol degraders in the last positive tubes with accumulation of acetate, sulfide, and presumably carbon dioxide.

## Discussion

Anaerobic degradation of glycerol requires the reduction of electron acceptors such as fumarate, as occurs in *Escherichia coli* [32] or *Streptococcus faecalis* [13] or  $\text{CO}_2$  in the case of homoacetogenic bacteria [8, 9]. The production of 1,3-propanediol as a reduced end product from glycerol has been observed in *Clostridium pasteurianum* [23], *Clostridium butylicum* [12], *Klebsiella pneumoniae* [11], *Lactobacillus brevis*, and *Lactobacillus buchneri* [33], *Citrobacter freundii* [22], and *Klebsiella aerogenes* [36]. In this study, it was demonstrated that in the absence of sulfate, anaerobic glycerol degradation by the mixed microbial culture studied was characterized by the transient formation of 1,3-propanediol prior to propionate and acetate accumulation. No lactate [19], ethanol and butyrate [12], acrolein [34], succinate, or pyruvate were produced. The presence of sulfate as a terminal electron acceptor seems to diminish the production of 1,3-propanediol by the mixed microbial culture. Thus, in the presence of sulfate, the problem of excess electron release during glycerol degradation is solved. These results suggest that sulfate-reducing bacteria may be involved in glycerol degradation with sulfate as terminal electron acceptor. The results of our enu-

Table 1. Effects of sulfate reduction and methanogenesis on anaerobic 1,3-propanediol degradation by the mixed microbial culture. Incubation at 37°C, for 4 weeks

Substrates	Initial concentration (mM)				Final concentration (mM)					
	1,3-OH <sup>a</sup>	C3	C2	sul	1,3-OH	C3	C2	sul	CH <sub>4</sub>	H <sub>2</sub>
1,3-OH	17.7	0	0	0	11	0.3	0	0	6.4	0
1,3-OH + sul	20.6	0	0	16.3	10.3	0.1	0	11.6	9.2	0
1,3-OH + chl	18.4	0	0	0	17.4	0.3	0.7	0	0	0.9
1,3-OH + sul + chl	18.6	0	0	15.9	9.2	0.1	0.3	6.7	0	0

<sup>a</sup> Symbols: 1,3-OH, 1,3-propanediol; C3, propionate; C2, acetate; sul, sulfate; chl, chloroform.

Table 2. MPN bacterial counts of fermentative and sulfate-reducing bacteria involved in anaerobic glycerol and 1,3-propanediol degradation by a mixed microbial culture.

Substrates	Bacterial counts (viable cells/ml)	Products of fermentation
- sulfate		
Glycerol	$2.5 \times 10^9$	propionate + acetate + methane
1,3-propanediol	$4.0 \times 10^7$	propionate + acetate + methane
+ sulfate		
Glycerol	$1.4 \times 10^9$	sulfide + acetate <sup>a</sup>
1,3-propanediol	$2.5 \times 10^7$	sulfide + acetate <sup>a</sup>
Total flora <sup>b</sup>	$10^{10}$	nd

<sup>a</sup> Traces of propionate were also detected.

<sup>b</sup> Estimated by the acridine epifluorescence method [15].

merations are consistent with this hypothesis. Sulfate-dependent glycerol oxidation was only recently unequivocally proved [10, 17, 24–26, 28, 30, 31, 35]. Furthermore, the values obtained in the absence of sulfate (53% carbon recovery when glycerol has disappeared, versus 91% at the end) and in the presence of sulfate (12.5% carbon recovery when glycerol has disappeared, versus 100% approximately at the end) are quite similar to those reported on pure culture of *Desulfovibrio carbinolicus* growing on glycerol with or without sulfate [24]; they strongly suggest the possibility that the intermediate produced during glycerol degradation by mixed microbial culture was 3-hydroxypropionate, although it is difficult to compare the results obtained because the media used were different, and the behavior of bacteria in pure culture differs greatly from that in mixed cultures.

Anaerobic oxidation of 1,3-propanediol has been reported with *D. carbinolicus* in the presence of sulfate [25] and with a strain of *Pelobacter carbinolicus* that oxidizes 1,3-propanediol into acetate only in the presence of a methanogenic hydrogen-

otrophic bacterium [7]. Our finding that 1,3-propanediol degradation did not occur when methanogenesis was inhibited in the absence of sulfate (Table 1) suggests that 1,3-propanediol degradation requires a terminal electron acceptor such as sulfate or a methanogenic hydrogenotrophic bacterium.

It is interesting to note that in low-sulfate medium tubes from enumerations with glycerol and 1,3-propanediol, *Desulfovibrio* species were observed by microscopic examination. It has, in fact, been demonstrated that *Desulfovibrio* species utilize glycerol and diols such as 1,2- and 1,3-propanediol by interspecies hydrogen transfer [28, 30, 31]. Our results on glycerol and 1,3-propanediol degradation by the mixed microbial culture agree with the concept that anaerobic degradation of reduced products of classical fermentations such as alcohols or fatty acids usually requires the reduction of electron acceptors such as sulfate [38], or a syntrophic association of fermenting and methanogenic bacteria, for example [4, 21]. Our results also show that glycerol and 1,3-propanediol (two substrates recently described as potential donors of reducing equivalents

for sulfate reduction) may be used by sulfate-reducing bacteria in the methanogenic ecosystem studied, possibly thanks to the presence of glycerol and sulfate in the original fermenter.

The isolation and characterization of the main sulfate-reducing bacteria using glycerol and 1,3-propanediol as energy and carbon sources from the positive tubes of enumerations will be described elsewhere [28].

#### ACKNOWLEDGMENTS

The authors wish to thank J.L. Batlle for glycerol assays, INRA experimental station (Pesh-Rouge-Narbonne, France), and J. Blanc for revising the manuscript. A.I. Qatibi's work was supported by ORSTOM and I.N.R.A grants.

#### Literature Cited

- Batlle JL, Collon Y (1979) Dosage enzymatique en flux continu du glycérol dans les vins. *Conn Vigne Vin* 1:45-50
- Bories A (1981) Méthanisation des eaux résiduaires de distilleries. *Trib Cebedeau*, 456:475-483
- Bories A, Raynal J, Jover JP (1982) Fixed film reactor with plastic media for methane fermentation of distilleries. In: Strub A, Chartier P, Schleser G (eds) *Energy from biomass*. London: Applied Science Publisher, pp 567-571
- Bryant MP, Wolin EA, Wolin MJ, Wolfe RS (1967) *Methanobacillus omelianskii*, a symbiotic association of two species of bacteria. *Ark Mikrobiol* 59:20-31
- Cord-Ruwisch R (1985) A quick method for the determination of dissolved and precipitated sulfides in cultures of sulfate-reducing bacteria. *J Microbiol Methods* 4:33-36
- Cord-Ruwisch R, Ollivier B, Garcia JL (1986) Fructose degradation by *Desulfovibrio* sp. in pure culture and in coculture with *Methanospirillum hungatei*. *Curr Microbiol* 13:285-289
- Dubourguier HC, Samain E, Prensier G, Albagnac G (1986) Characterisation of two strains of *Pelobacter carbinolicus* isolated from anaerobic digesters. *Arch Microbiol* 145:248-253
- Eichler B, Schink B (1984) Oxidation of primary aliphatic alcohols by *Acetobacterium carbinolicum* sp. nov., a homoacetogenic anaerobe. *Arch Microbiol* 140:147-152
- Emde R, Schink B (1987) Fermentation of triacetin and glycerol by *Acetobacterium* sp. No energy is conserved by acetate excretion. *Arch Microbiol* 149:142-148
- Esnault G, Caumette P, Garcia JL (1988) Characterization of *Desulfovibrio giganteus* sp. nov., a sulfate-reducing bacterium isolated from a brackish coastal lagoon. *System Appl Microbiol* 10:147-151
- Forage RG, Foster MA (1982) Glycerol fermentation in *Klebsiella pneumoniae*: functions of the coenzyme B12-dependent glycerol and diol dehydratases. *J Bacteriol* 149:413-419
- Forsberg CW (1987) Production of 1,3-propanediol from glycerol by *Clostridium acetobutylicum* and other *Clostridium* species. *Appl Environ Microbiol* 53:639-643
- Gunsalus IC (1947) Products of anaerobic glycerol fermentation by *Streptococcus faecalis*. *J Bacteriol* 54:239-244
- Hobbie JE, Daley RJ, Jasper S (1977) Use of nucleopore filters for counting bacteria by fluorescence microscopy. *Appl Environ Microbiol* 33:1225-1228
- Hungate RE (1960) Microbial ecology of rumen. *Bacteriol Rev* 24:353-364
- Joubert WA, Britz TJ (1987) Isolation of saccharolytic dissimilatory sulfate-reducing bacteria. *FEMS Microbiol Lett* 48:35-40
- Kremer DR, Hansen TA (1987) Glycerol and dihydroxyacetone dissimilation in *Desulfovibrio* strains. *Arch Microbiol* 147:249-256
- Landre J (1983) Détermination du carbone organique total dans les eaux potables par oxydation U.V. *Eau Industries Nuisances* 71:25-28
- Laroche M (1983) Métabolisme intermédiaire des acides gras volatils en fermentation méthanique. Thèse doctorat, Institut National Des Sciences Appliquées, Toulouse, France
- Macy JM, Snellen JE, Hungate RE (1972) Use of syringe methods for anaerobiosis. *Am J Clin Nutr* 25:1318-1323
- McInerney MJ, Bryant MP, Pfennig N (1979) Anaerobic bacterium that degrades fatty acids in syntrophic association with methanogens. *Arch Microbiol* 122:129-135
- Mickelson MN, Werkman CH (1940) The dissimilation of glycerol by *Coli-aerogenes* intermediates. *J Bacteriol* 39:709-715
- Nakas JP, Chaedle MS, Parkinson CM, Coonley CE, Tannenbaum SW (1983) System development of linked fermentation production of solvents from algal biomass. *Appl Environ Microbiol* 46:1017-1023
- Nanninga HJ, Gottschal JC (1986) Isolation of a sulfate-reducing bacterium growing with methanol. *FEMS Microbiol Eco* 38:125-130
- Nanninga HJ, Gottschal JC (1987) Properties of *Desulfovibrio carbinolicus* sp. nov. and other sulfate-reducing bacteria isolated from an anaerobic-purification plant. *Appl Environ Microbiol* 51:572-579
- Ollivier B, Cord-Ruwisch R, Hatchikian EC, Garcia JL (1988) Characterization of *Desulfovibrio fructosovorans* sp. nov. *Arch Microbiol* 149:447-450
- Postgate JR (1984) *The sulfate reducing bacteria*, 2nd ed London: Cambridge University Press
- Qatibi AI (1990) Fermentation du lactate, du glycérol et des diols par les bactéries sulfato-réductrices du genre *Desulfovibrio*. Thèse doctorat, Univer. Aix-Marseille I, France
- Qatibi AI, Bories A (1988) Glycerol fermentation and sulfate utilization during the anaerobic digestion process. In: Tilche A, Rozzi A (eds) *Fifth Intern. Symp. Anaerobic Digestion*, Monduzzi Editore, Bologna, pp 69-73
- Qatibi AI, Garcia JL (1989) Glycerol degradation by *Desulfovibrio* sp., in pure culture and in coculture with *Methanospirillum hungatei*. In: *FEMS Symp. microbiology and biochemistry of strict anaerobes involved in interspecies hydrogen transfer*. New York: Plenum Publ. Corp., in press
- Qatibi AI, Cayol JL, Garcia JL (1989) 1,2- and 1,3-propanediol degradation by *Desulfovibrio alcoholovorans* sp. nov., in pure culture or through interspecies transfer. In: *FEMS Symp. microbiology and biochemistry of strict anaerobes involved in interspecies hydrogen transfer*. New-York: Plenum Publ. Corp., in press
- Quastel JH, Stephenson M, Whetham MD (1925) Some reactions of resting bacteria in relation to anaerobic growth. *Biochem J* 19:304-317
- Schutz H, Radler F (1984) Anaerobic reduction of glycerol to propanediol-1,3 by *Lactobacillus brevis* and *Lactobacillus buchneri*. *Syst Appl Microbiol* 5:169-178

34. Serjak WC, Day WH, Van Lanen JM, Boruff CS (1954) Acrolein production by bacteria found in distillery grain mashes. *Appl Microbiol* 2:14-20
35. Stams A, Hansen TA, Skyring GW (1985) Utilisation of amino acids as energy substrates by two marine *Desulfovibrio* strains. *FEMS Microbiol Lett* 31:11-15
36. Thiamann KV (1955) *The life of bacteria*. New York: MacMillan Publishing Co
37. Widdel F (1980) Anaerober Abbau von Fettsäuren und Benzoesäure durch neu isolierte Arten Sulfat-reduzierender Bakterien. Doctoral thesis, Univ Göttingen, FRG
38. Widdel F (1988) Microbiology and ecology of sulfate- and sulfur-reducing bacteria. In: Zehnder AJB (ed) *Biology of anaerobic microorganisms*. New York: John Wiley, pp 469-585
39. Widdel F, Pfennig N (1977) A new anaerobic sporing acetate-oxidizing, sulfate-reducing bacterium, *Desulfotomaculum* (emend) *acetoxidans*. *Arch Microbiol* 112:119-122