

Errata to

A Sum Rule for Deep Inelastic Electroproduction
from Polarized Protons

John Ellis and Robert Jaffe

Page 4, Eq. 2 and following equation, replace $\sqrt{2}$ by $\frac{1}{\sqrt{2}}$.

Page 4, bottom equation, replace $\frac{1}{\sqrt{2}}$ by $\sqrt{2}$.

A SUM RULE FOR DEEP INELASTIC ELECTROPRODUCTION
FROM POLARIZED PROTONS*

John Ellis†

Lauritsen Laboratory of Physics, California Institute of Technology
Pasadena, California 91109

and

Stanford Linear Accelerator Center
Stanford University, Stanford, California 94305

Robert Jaffe††

Laboratory for Nuclear Science and Department of Physics
Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

and

Stanford Linear Accelerator Center
Stanford University, Stanford, California 94305

ABSTRACT

A sum rule is derived for the asymmetry in deep inelastic scattering of polarized electrons from polarized protons:

$$\int_0^1 d\xi g^{\text{ep}}(\xi) \approx 0.15 g_A$$

The result follows from the quark light-cone algebra and the assumption that strange quarks do not contribute to the asymmetry. The latter is justified by conventional parton model arguments.

*Work supported in part by the U. S. Atomic Energy Commission.

†Address after 1st September 1973, Theory Division, CERN, 1211 Geneva 23, Switzerland.

††Supported in part through funds provided by the U. S. Atomic Energy Commission under contract AT(11-1)-3069.

Measurements of the asymmetry in the inelastic scattering of polarized electrons from polarized protons will begin soon.¹ Several years ago Bjorken² derived a sum rule for the difference in polarization asymmetry in deep inelastic scattering from protons and neutrons. The sum rule has been rederived recently using the parton model³ and the quark light-cone algebra.⁴ Unfortunately, testing the Bjorken sum rule would require data on scattering from polarized deuterons which will be unavailable for some time. Here we derive a sum rule for the asymmetry in scattering from polarized protons alone. We use the standard quark light-cone algebra, the usual parton model assumptions that the only isosinglet (λ -type) quarks in the proton are in the "sea" (Pomeranchukon) and that the spins of partons in the "sea" are paired. These are discussed further later.

The structure functions for scattering polarized electrons from polarized protons are defined as follows:

$$\begin{aligned} \frac{1}{2} \left[W_{\mu\nu} - W_{\nu\mu} \right] &= i \epsilon_{\mu\nu\lambda\sigma} \frac{q^\lambda n^\sigma}{M^2} G_1(\nu, q^2) \\ &+ i \epsilon_{\mu\nu\lambda\sigma} \frac{q^\lambda}{M^4} (\nu n^\sigma - n \cdot q p^\sigma) G_2(\nu, q^2) \end{aligned}$$

where $\nu \equiv p \cdot q$, n^σ is the proton's polarization vector ($n^2 = -M^2$, $p \cdot n = 0$) and

$$W_{\mu\nu}(q, p) = \frac{1}{4\pi} \int d^4x e^{iq \cdot x} \langle p, n | \left[J_\mu(x), J_\nu(0) \right] | p, n \rangle$$

It is expected that νG_1 and $\nu^2 G_2$ should scale in the Bjorken limit ($\nu, q^2 \rightarrow \infty$, $\xi \equiv -q^2/2\nu$ fixed):

$$\text{Lim}_{\text{Bj}} \frac{\nu}{M^2} G_1(q^2, \nu) \equiv g_1(\xi)$$

$$\lim_{\text{Bj}} \frac{\nu^2}{M^4} G_2(q^2, \nu) \equiv g_2(\xi)$$

Conventional quark light-cone algebra⁵ yields the following expression for $g_1(\xi)$:

$$g_1^{(\text{ep})}(\xi) = \frac{1}{4\pi} \int_{-\infty}^{\infty} d(x \cdot p) e^{i\xi x \cdot p} \left[\pm \frac{1}{6} S_3^5(x \cdot p) + \frac{1}{6\sqrt{3}} S_8^5(x \cdot p) + \sqrt{\frac{2}{27}} S_0^5(x \cdot p) \right] \quad (1)$$

where the $S_a^5(x \cdot p)$ are defined from the bilocal operators:

$$\langle p, n | S_{a\sigma}^5(x|0) | p, n \rangle \Big|_{x^2=0} = n_\sigma S_a^5(x \cdot p) + \dots$$

where $S_{a\sigma}^5(x|0)$ is the axial vector bilocal which is symmetric under interchange of x and 0 , and has the following form in free field theory:

$$S_{a\sigma}^5(x|0) = \bar{\psi}(x) \frac{\lambda_a}{2} \gamma_\sigma \gamma_5 \psi(0) + \bar{\psi}(0) \frac{\lambda_a}{2} \gamma_\sigma \gamma_5 \psi(x)$$

Bjorken's sum rule follows from inverting the Fourier transform in Eq. (1), setting $x \cdot p$ to zero, taking the ep, en difference and noting that

$$\langle p, n | S_{3\sigma}^5(0|0) | p, n \rangle = 2g_A n_\sigma$$

since the local limit of the bilocal is just the neutral isovector axial current.

In trying to derive a sum rule for scattering off polarized protons it is necessary to know $\langle p, n | S_{8\sigma}^5(0|0) | p, n \rangle$ and $\langle p, n | S_{0\sigma}^5(0|0) | p, n \rangle$. The first of these can be estimated using SU(3) for the baryon matrix elements of the axial charges. Hyperon decays are known to agree well with SU(3) and the Cabibbo theory, and the F/D ratio for axial charges is well determined.⁶ However, nothing is known a priori about $\langle p, n | S_{0\sigma}^5(0|0) | p, n \rangle$. To evaluate it we argue

on the basis of the parton model that the matrix elements of the combination

$$\sqrt{2} S_{0\sigma}^5(x|0) - S_{8\sigma}^5(x|0) \quad (2)$$

between nucleons are expected to be small.

The combination (2) gets contributions only from λ and $\bar{\lambda}$ type quarks. Data from neutrino reactions⁷ already suggest that the nucleons' wave functions have a small $(\lambda, \bar{\lambda})$ components, and that it is reasonable to assume they only occur in the "sea" associated with the Pomeranchukon. The Pomeranchukon does not contribute to the spin-dependent structure functions G_1 and G_2 ,⁸ so one might suspect that the $(\lambda, \bar{\lambda})$ contribution to them should be small. In fact

$$\langle p, n | (\sqrt{2} S_{0\sigma}^5(0|0) - S_{8\sigma}^5(0|0)) | p, n \rangle \propto \int_0^1 d\xi (\lambda_+(\xi) + \bar{\lambda}_+(\xi) - \lambda_-(\xi) - \bar{\lambda}_-(\xi))$$

where $\lambda_{\pm}(\xi)$ are the distributions of λ quarks in the nucleon with helicities parallel (antiparallel) to the helicity of the nucleon itself,³ and $\bar{\lambda}_{\pm}(\xi)$ are defined similarly. The usual spin and charge conjugation properties of the Pomeranchukon imply

$$\lambda_+(0) = \bar{\lambda}_+(0) = \lambda_-(0) = \bar{\lambda}_-(0)$$

Our assumption is that this remains at least approximately valid for $\xi \neq 0$ for the "sea" of $q\bar{q}$ pairs:

$$\lambda_+(\xi) \approx \bar{\lambda}_+(\xi) \approx \lambda_-(\xi) \approx \bar{\lambda}_-(\xi) \quad (3)$$

Assuming that the "sea" has the property (3) it follows that⁹

$$\langle p, n | S_{0\sigma}^5(x|0) | p, n \rangle \approx \frac{1}{\sqrt{2}} \langle p, n | S_{8\sigma}^5(x|0) | p, n \rangle$$

and separate sum rules for the proton and neutron asymmetries follow immediately:

$$\int_0^1 g_1^{\text{ep}}(\xi) d\xi = \frac{g_A}{12} \left(1 + \frac{5}{3} \frac{3 \frac{F}{D} - 1}{\frac{F}{D} + 1} \right) \quad (4)$$

$$\int_0^1 g_1^{\text{en}}(\xi) d\xi = \frac{g_A}{12} \left(-1 + \frac{5}{3} \frac{3 \frac{F}{D} - 1}{\frac{F}{D} + 1} \right) \quad (5)$$

The difference of Eqs. (4) and (5) is just Bjorken's sum rule.² In writing (4) and (5) we have used SU(3) to re-express the ratio

$$\frac{\langle p, n | S_{3\sigma}^5(0|0) | p, n \rangle}{\langle p, n | S_{8\sigma}^5(0|0) | p, n \rangle}$$

in terms of the F/D ratio of the axial charges. Since the axial charges seem to form an excellent octet⁶ we regard this as a minor assumption. Using the currently accepted value⁶

$$\frac{F}{D} = .581 \pm .029$$

we obtain

$$\int_0^1 d\xi g_1^{\text{ep}}(\xi) = \frac{g_A}{12} (1.78) \quad (6)$$

$$\int_0^1 d\xi g_1^{\text{en}}(\xi) = \frac{g_A}{12} (-0.22) \quad (7)$$

where $g_A = 1.248 \pm .010$.

We do not expect the sum rules Eqs. (6), (7) to be exact because of SU(3) breaking, the uncertainty in the F/D ratio, and the parton model assumption about the $(\lambda, \bar{\lambda})$ distributions. However, as long as the momentum of strange quarks extracted from neutrino experiments (see footnote 7) remains small,

we expect Eqs. (6) and (7) to be quite accurate estimates of polarization asymmetries in scattering off protons and neutrons separately. The proton sum rule (6) is particularly interesting because experiments using polarized proton targets will soon be performed, and because (following Bjorken¹⁰) it predicts that a large mean asymmetry of the order of 35% will be observed.

Acknowledgements

We thank F. Gilman for useful conversations and SLAC for its generous support and hospitality while this work was performed.

REFERENCES

1. Yale-SLAC-MIT-NSF Collaboration, SLAC Experiment No. E-80.
2. J. D. Bjorken, Phys. Rev. 148, 1467 (1966).
3. J. Kuti and V. F. Weisskopf, Phys. Rev. D 4, 3418 (1971).
R. P. Feynman, Photon-Hadron Interactions (W. A. Benjamin, Reading, 1972).
4. A. J. G. Hey and J. E. Mandula, Phys. Rev. D 5, 2610 (1972).
D. Wray, Nuovo Cimento 9A, 463 (1972).
5. H. Fritzsche and M. Gell-Mann, "Broken scale invariance and the light-cone," 1971 Coral Gables Conference on Fundamental Interactions at High Energy, edited by M. Dal Cin, G. J. Iverson and A. Perlmutter (Gordon and Breach, New York, 1971).
6. H. Ebenhoh et al., Z. Physik 241, 473 (1971).
7. In the quark model

$$\int_0^1 d\xi \xi (\lambda(\xi) + \bar{\lambda}(\xi)) = 9 \int_0^1 d\xi F_2^{\gamma N}(\xi) - \frac{5}{2} \int_0^1 d\xi F_2^{\nu N}(\xi) \Big|_{\Delta S=0}$$

where $\lambda(\xi)$ and $\bar{\lambda}(\xi)$ are the distributions of λ and $\bar{\lambda}$ quarks in the nucleon, and $F_2^{\gamma N}(\xi)$ and $F_2^{\nu N}(\xi)$ are structure functions averaged over neutrons and protons. D. Perkins (Proceedings of the XVI International Conference on High Energy Physics, National Accelerator Laboratory, 1972; Vol. 4, p. 189) has estimated

$$\int_0^1 d\xi F_2^{\gamma N}(\xi) = 0.14 \pm .02, \quad \int_0^1 d\xi F_2^{\nu N}(\xi) \Big|_{\Delta S=0} = 0.49 \pm .07$$

implying that

$$\int_0^1 d\xi \xi (\lambda(\xi) + \bar{\lambda}(\xi)) = 0.02 \pm 0.25 .$$

8. See for example, Kuti and Weisskopf (Ref. 3) and references therein.
9. M. Kugler informs us that this relation can also be obtained using the Melosh transformation from current to constituent quarks.
10. J. D. Bjorken, Phys. Rev. D 1, 1376 (1970).