

SHORT COMMUNICATION

Summer aggregations of the common eagle ray, *Myliobatis aquila*

PEDRO AFONSO & NUNO VASCO-RODRIGUES



Afonso, P. & N. Vasco-Rodrigues 2015. Summer aggregations of the common eagle ray, *Myliobatis aquila*. *Arquipelago*. Life and Marine Sciences 32.

Pedro Afonso (email: afonso@uac.pt): IMAR-UAz - Institute of Marine Research at the University of the Azores. Dept. of Oceanography and Fisheries, PT-9901-862 Horta; MARE – Marine and Environmental Sciences Centre, University of the Azores, Dept. of Oceanography and Fisheries, 9901-862 Horta, Açores, Portugal; Nuno Vasco-Rodrigues, MARE – Marine and Environmental Sciences Centre, ESTM, Polytechnic Institute of Leiria, 2520-641 Peniche, Portugal; Flying Sharks, Rua Jorge Castilho 1613, 7C, 1900-272 Lisboa, Portugal.

The aggregation of multiple individuals is a long known behaviour in a variety of elasmobranchs, including reef sharks (Speed et al. 2011), hammerhead sharks (Klimley & Nelson 1984), whale sharks (Hoffmayer et al. 2007), devilrays (Ward-Page et al. 2013; Sobral & Afonso 2014), sting-rays (Semeniuk & Rotley 2008), eagle rays (Silliman & Gruber 1999) and cownose rays (Smith et al. 1987). Various functions have been hypothesised to explain these aggregations, including feeding (e.g. Wilson et al. 2001; Rohner et al. 2013), courtship or mating (Whitney et al. 2004; Dudgeon et al. 2008), and cleaning stations (Dewar et al. 2008). However, apart from the obvious cases whenever individuals directly engage in reproductive activities, the social functions that these aggregations might serve are not understood or even described.

During the summer of 2014, while conducting regular fish visual surveys, an aggregation of up to at least 30 common eagle rays was observed at Radares point, a site located on eastern point of the Monte da Guia marine protected area (MPA), in the Azorean island of Faial, mid-north Atlantic (Fig. 1 and 2). Individuals were identified based on morphological characteristics (Fisher et al. 1981) and previous citations for the region (Santos et al. 1997; Barreiros & Gadig 2011).




Fig. 1. Location of the aggregation site (dark cloud) on the perimeter of the Monte da Guia Marine Reserve, Faial Island, Azores (dashed line).



Fig. 2. Common eagle rays aggregating at Radares point, Faial Island.

The aggregation was observed in four consecutive dives spanning eight days (Table 1). Upon the first sighting, another six dedicated dives were made until November. Dives were conducted at various times of the day, from dawn to dusk. Individuals were observed almost daily during the week around the full moon of August (08, 10, 11 and 14-08-2014). The aggregation held 20 to 30 observable individuals at one given dive, which measured an estimated 30 to 90 cm disc width (DW) (Fig. 2). The rays were always encountered within a localized area (ca. 1 ha) of the reef (Fig. 1), in spite of the prospective transects done in the areas surrounding the aggregation site. The site lies at the transition from the rocky to the sandy bottom at 40 m depth just off the point, but the animals were swimming in mid-water (10-30 m depth). All individuals appeared to be females as no claspers were seen sticking out of the fins

Table 1. Details of observations on the aggregation site; * indicates full moon.

Date	Time	N° indiv.	Size range (DW, cm)
08/08/2014	16:00	25	30-90
10/08/2014*	18:30	30	30-90
11/08/2014	07:30	20	30-90
14/08/2014	10:00	20	30-90
03/09/2014	14:00	0	-
08/09/2014*	08:00	0	-
25/11/2014	09:00	0	-

trailing edge, which would be expected in sub-adult and adult males (Capapé et al. 2007). There was no apparent close interaction between individuals, feeding, or cleaning activity. Right after this period, the aggregation disappeared.

The common eagle ray is a relatively rare elasmobranch in coastal areas of the Azores, where it was only observed in 1.6% of over 500 transects for fish counts across the archipelago and over 15 years, and there was never an occasion when more than one individual had been observed (P. Afonso, personal observation). McEachran & Séret (1990) state in their review of the family that the species "is often found in groups" with no further data on locations, numbers or behaviour. Barreiros & Gadig (2011) also refer that "this species can be observed in reproductive aggregations in the summers, in one specific cave, in Ilhéu das Cabras, Terceira Island" but provide no information to ascertain the function of the aggregation. Similar aggregations were also observed in the summer at an offshore reef in Graciosa Island, apparently also constituted only by females of various sizes (Rolando Oliveira, pers. comm.). The fact that no particular behaviour was observed makes it impossible to even speculate about the nature of the aggregation, although it seems that there are no obvious feeding or parasite cleaning functions. As such, it remains possible that these aggregations serve social functions. It is particularly intriguing that only females were confirmedly observed, although it remains possible that immature males were in the group but went unnoticed. Thus, the aggregation might serve reproductive functions, namely related to egg deposition. Capapé et al. (2007) refer that this species breeds in August and September in the northwestern Mediterranean. Indeed, recruits of this species (ca. 10-15 cm DW) were observed from July to the end of summer in the neighbour nursery area of Porto Pim, just around the perimeter of Monte da Guia (Figure 1; P. Afonso unpublished data), so it appears that the reproductive schedule in the Azores matches that of the Mediterranean, as in most coastal fishes. This would explain why the aggregation is exclusively composed of mature females.

The nature of the aggregation and its potential significance for the conservation of this species should be investigated given its status as 'data deficient' and vulnerable population characteristics.

ACKNOWLEDGEMENTS

The authors would like to thank C. Moura and M. Schmiing for dive support and M. Simões for the help given in photo treatment. This paper is a contribution to the MoniZec project (DRCT/M212/I/018/2011).

REFERENCES

- Barreiros, J. P. & O.B.F. Gadig 2011. *Sharks and Rays from the Azores: An illustrated catalogue*. Instituto Açoriano de Cultura, Angra do Heroísmo, Terceira, Azores. 187pp.
- Capapé, C., O. Guélorget, Y. Vergne & J.P. Quignard 2007. Reproductive biology of the common Eagle Ray *Myliobatis aquila* (Chondrichthyes: Myliobatidae) from the Coast of Languedoc (Southern France, Northern Mediterranean). *Vie milieu: Life Environment* 57 (3): 1-6.
- Dewar, H., P. Mous, M. Domeier, A. Muljadi, J. Pet & J. Whitty 2008. Movements and site fidelity of the giant manta ray, *Manta birostris*, in the Komodo Marine Park, Indonesia. *Marine Biology* 155: 121–133.
- Dudgeon, C.L., M.J. Noad & J.M. Lanyon 2008. Abundance and demography of a seasonal aggregation of zebra sharks *Stegostoma fasciatum*. *Marine Ecology Progress Series* 368: 269–281.
- Fisher, W., G. Bianchi & W.B. Scott (Eds) 1981. *FAO species identification sheets for fisheries purposes*. Eastern Central Atlantic; Fishing areas 34, 47 (in part). Canada Funds-in-Trust. Ottawa, Department of Fisheries and Oceans Canada, by arrangement with the Food and Agriculture Organization of the United Nations, vol. 5.
- Hoffmayer, E.R., J.S. Franks, W.B. Driggers, K.J. Oswald, K.J. & J.M. Quattro 2007. Observations of a feeding aggregation of whale sharks, *Rhincodon typus*, in the north central Gulf of Mexico. *Gulf and Caribbean Research* 19(2): 69–73.
- Klimley, A.P. & D.R. Nelson 1984. Diel movement patterns of the scalloped hammerhead shark (*Sphyrna lewini*) in relation to El Bajo Espiritu Santo: a refuging central-position social system. *Behavioural Ecology and Sociobiology* 15: 45-54.
- McEachran, J.D. & B. Séret 1990. Myliobatidae. Pp.67-70 in: Quero, J.C., J.C. Hureau, C. Karrer, A. Post. & L. Saldanha (Eds) *Check-list of the fishes of the eastern tropical Atlantic (CLOFETA)*. JNICT, Lisbon; SEI, Paris; and UNESCO, Paris. Vol. 1. 519 pp.

- Rohner, C.A., S.J. Pierce, A.D. Marshall, S.J. Weeks, M.B. Bennett. & A.J. Richardson 2013. Trends in sightings and environmental influences on a coastal aggregation of manta rays and whale sharks. *Marine Ecology Progress Series* 482: 153-168.
- Santos, R.S., F.M. Porteiro & J.P. Barreiros 1997. Marine Fishes of the Azores: annotated checklist and bibliography. *Arquipelago. Life and Marine Sciences*. Supplement 1. 244 pp.
- Semeniuk, C. & K. Rothley 2008. Costs of group-living for a normally solitary forager: effects of provisioning tourism on southern stingrays *Dasyatis americana*. *Marine Ecology Progress Series* 357: 271-282.
- Silliman, W.R. & S.H. Gruber 1999. Behavioral biology of the spotted eagle ray, *Aetobatus narinari* (Euphrasen, 1790), in Bimini, Bahamas; an interim report. *Bahamas Journal of Science* 7:13–20.
- Sobral, A.F. & P. Afonso 2014. Occurrence of mobulids in the Azores, central North Atlantic. *Journal of the Marine Biological Association of the United Kingdom* 94 (8): 1671-1675.
- Speed, C.W., M.G. Meekan, I.C. Field, C.R. McMahon, J.D. Stevens, F. McGregor, C. Huvaneer, Y. Berger & C.J.A. Bradshaw 2011. Spatial and temporal movement patterns of a multi-species coastal reef shark aggregation. *Marine Ecology Progress Series* 429: 261-275.
- Ward-Paige, C.A., B. Davis & B.Worm, 2013. Global Population Trends and Human use Patterns of *Manta* and *Mobula* Rays. *Plos One* 8(9): e74835.
- Whitney, N.M., J.H.L. Pratt & J.C. Carrier 2004. Group courtship, mating behaviour and siphon sac function in the whitetip reef shark, *Triaenodon obesus*. *Animal Behaviour* 68: 1435–1442.
- Wilson, S.G., J.G. Taylor & A.F. Pearce 2001. The Seasonal Aggregation of Whale Sharks at Ningaloo Reef, Western Australia: Currents, Migrations and the El Niño/Southern Oscillation. *Environmental Biology of Fishes* 61: 1-11.

Received 24 Apr 2015. Accepted 26 Jun 2015,

Published online 13 Jul 2015.