

Supersymmetric quantum mechanics and new potentials

Elso Drigo Filho

Instituto de Física Teórica, Universidade Estadual Paulista, Rua Pamplona 145, São Paulo, 01405, SP, Brasil

and

Instituto de Biociência Letras e Ciências Exatas, Universidade Estadual Paulista, Rua Cristóvão Colombo 1165, São José do Rio Preto, 15055, SP, Brasil

Received May 30, 1990

Abstract Using supersymmetric quantum mechanics we generalize some exactly solvable potentials: the particle in the box, Pöschl-Teller and Rosen-Morse. We evaluate the new potentials and indicate their eigenfunctions and spectra.

1. INTRODUCTION

We know that the number of Schrödinger equations that have analytic solutions is quite small. In recent years some works have tried to increase this number, starting from potentials whose solutions are known (e.g. Abraham and Moses¹ and Pursey²). Supersymmetric quantum mechanics (SQM) has also been used for that purpose. The superalgebra is used to construct a hierarchy of Hamiltonians³ and to build new Hamiltonians from a Riccati equation^{4,5,6}.

The method to construct new potentials from known potentials using SQM, which we use in this paper, was proposed by Nieto⁴ and Alves and Drigo Filho⁵. It is based on the factorisation method which was applied by Mielnik⁷ to the harmonic oscillator and by Fernandez⁸ to the Coulomb potential. This method is also applicable to spatially limited potentials. We will see it through the example of the particle in the box.

Work supported by FAPESP and CNPq, Brasil.

Supersymmetric quantum mechanics and new potentials

Here, we use the superalgebra to construct new potentials from exactly solvable potentials. This construction was used to generalize the Coulomb and the harmonic oscillator potentials⁵, as well as the Morse potential⁶. Firstly, we present the method for a general potential in sec. 2. Then, we apply it to the simple potential of a particle in a box (sec. 3). We treat the Poschl-Teller and Rosen-Morse potentials in sec. 4 and we comment on the results in sec. 5.

2. Generalization method

In SQM^{9,10} we have two supersymmetric charges Q and Q^+ ; they satisfy the anticommutation relations

$$\{Q, Q^+\} = H_{ss}; \quad (Q, Q) = 0 \quad , \quad (Q^+, Q^+) = 0 \tag{1}$$

A simple realization of this algebra is

$$Q = \begin{pmatrix} 0 & 0 \\ d^- & 0 \end{pmatrix} \quad \text{and} \quad Q^+ = \begin{pmatrix} 0 & q^+ \\ 0 & 0 \end{pmatrix} \tag{2}$$

and we have

$$H_{ss} = \begin{pmatrix} H_+ & 0 \\ 0 & H_- \end{pmatrix} = \begin{pmatrix} d^+d^- & 0 \\ 0 & d^-d^+ \end{pmatrix} \tag{3}$$

H_- is called the supersymmetric partner of H_+ they have the same spectrum except for the zero-energy ground state which belongs to H_+ only. We note that

$$Q \begin{bmatrix} \psi_+ \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ d^-\psi_+ \end{bmatrix} \quad \text{and} \quad Q^+ \begin{bmatrix} 0 \\ \psi_- \end{bmatrix} = \begin{bmatrix} d^+\psi_- \\ 0 \end{bmatrix} \tag{4}$$

i.e., Q and Q^+ induce transformations between the "bosonic" sector ($|\psi_+\rangle$) and the "fermionic" sector ($|\psi_-\rangle$). Then, the H_- eigenfunctions can be written in terms of H_+ eigenfunctions ($\psi_- \propto d^-\psi_+$). The reciprocal is also true, i.e. $|\psi_+\rangle \propto d^+\psi_-$ are the eigenfunctions of H_+ with the exception of the ground-state. With operators d^\pm written in the usual form

$$d^\pm = \mp \frac{d}{dx} + \frac{dW}{dx}(x) \tag{5}$$

the supersymmetric Hamiltonian is written as

Elso Drigo Filho

$$H_{ss} = -\frac{d^2}{dx^2} + \left[\frac{dW}{dx}(x) \right]^2 + \sigma_3 \frac{d^2}{dx^2} W(x) \quad (6)$$

where σ_3 is the **Pauli** matrix, $W(x)$ is called superpotential and is **associated** with the H_+ ground state eigenfunctions $\psi_{+,0}$

$$W(x) = -\ln \psi_{+,0}(x) \quad (7)$$

We can construct the new potentials from a generalization of the d^\pm operators

$$D^* = \mp \frac{d}{dx} + F(x) \quad (8)$$

The function $F(x)$ is determined when we impose that

$$H_- = D^- D^+ \quad (9)$$

and we obtain the **Riccati** equation

$$F^2(x) + \frac{d}{dx} F(x) = \left[\frac{d}{dx} W(x) \right]^2 + \frac{d^2}{dx^2} W(x) \quad (10)$$

The **commutator** of the new operators is

$$[D^-, D^+] = 2 \frac{d}{dx} F(x) \quad (11)$$

that defines a new Hamiltonian

$$\mathcal{H}_+ = D^+ D^- = D^- D^+ - [D^-, D^+] = D^- D^+ - 2 \frac{d}{dx} F(x) \quad (12)$$

\mathcal{H}_+ gives a new potential which is different from the H_+ and H_- potentials. However, from the supersymmetric algebra we know that the \mathcal{H}_+ spectrum is the **same** as that of H_- and the \mathcal{H}_+ eigenfunctions (Ψ_+) are

$$\Psi_+ = D^+ \psi_- = D^+ d^- \psi_+ \quad (13)$$

This map is not complete, because it excludes the \mathcal{H}_+ ground state. It is obtained by the equation

Supersymmetric *quantum* mechanics and new *potentials*

$$D\Psi_{+,0} = 0 \quad (14)$$

3. Particle in the box

The Hamiltonian of the one-dimensional **particle** in the **box**¹ is

$$H_+ = -\frac{d^2}{dx^2} - 1 \quad ; \quad -\frac{1}{2}\pi \leq x \leq \frac{1}{2}\pi \quad (15)$$

H_+ has eigenfunctions and eigenvalues given by

$$\psi_{+,n}(x) = \begin{cases} \sqrt{\frac{2}{\pi}} \sin(nx) & n \text{ even} \\ \sqrt{\frac{2}{\pi}} \cos(nx) & n \text{ odd} \end{cases} \quad (16)$$

$$E_n = n^2 - 1 \quad n = 1, 2, 3 \quad (17)$$

The constant term in (15) **only** displace the spectrum. It sets the eigenvalue of the ground state to zero, $E_1 = 0$.

We note that eq.(15) can be factorized, $H_+ = d^+d^-$, by

$$d^\pm = \mp \frac{d}{dx} + \frac{d}{dx}W(x) = \mp \frac{d}{dx} - \frac{d}{dx} \ln \psi_{+,1}(x) = \mp \frac{d}{dx} + \tan x \quad (18)$$

Thus, the **supersymmetric** partner of H_+ is

$$H_- = d^-d^+ = -\frac{d^2}{dx^2} + \frac{1 + \sin^2 x}{\cos^2 x} \quad (19)$$

Defining new operators D^\pm , as in eqs. (8), we obtain $F(x)$ given by

$$F(x) = \tan x + \frac{4 \cos^2 x}{\sin 2x + 2x + 4\Gamma} \equiv \tan x + \phi(x) \quad (20)$$

Thus, the new Hamiltonian is

$$\mathcal{H}_+ = D^+D^- = D^-D^+ - [D^-, D^+] = -\frac{d^2}{dx^2} - 1 - \frac{32 \cos x [(x + 2\Gamma) \sin x + \cos x]}{(\sin 2x + 2x + 4\Gamma)^2} \quad (21)$$

which corresponds to the generalized potential

Elso Drigo Filho

$$\mathcal{V}_+(x) = -1 - \frac{32 \cos x [(x + 2\Gamma) \sin x + \cos x]}{(\sin 2x + 2x + 4\Gamma^2)} \quad (22)$$

The constant Γ is arbitrary and is chosen to avoid singularities; here we choose: $\Gamma < -\frac{\pi}{4}$ or $\Gamma > \frac{\pi}{4}$. The spectrum of this Hamiltonian is the same of the particle in the box (17). Its eigenfunctions are

$$\Psi_{+,n}(x) = D^+ \psi_{-,n}(x) = D^+ d^- \psi_{+,n}(x) \quad (23)$$

with $\psi_{+,n}$ given by (16) and the ground state ($\tilde{\Psi}_+$) is evaluated by using (14)

$$\tilde{\Psi}_+ \propto \cos^{-1} x \mp \left[\int_{-\frac{\pi}{2}}^x \phi(\bar{x}) d\bar{x} \right] \quad (24)$$

4. Other potentials

Using the Pöschl-Teller potential¹¹ we write the Hamiltonian

$$H_+^{(PT)} = -\frac{d^2}{dz^2} + \frac{k(k-1)}{\sin^2 z} + \frac{A(A-1)}{\cos^2 z} - (k-\lambda)^2, \quad z = \alpha x \quad (25)$$

where α, k and λ are constants. The eigenfunctions and eigenvalues are¹²

$$\psi_{+,n}^{PT} = N(\alpha, k, \lambda, n) (\sin z)^k (\cos z)^\lambda P_n^{(k-\frac{1}{2}, \lambda-\frac{1}{2})} (1 - 2 \sin^2 z) \quad (26)$$

$$E_n = (k + \lambda + 2n)^2 - (k + \lambda)^2 \quad (27)$$

where

$$P_n^{(k-\frac{1}{2}, \lambda-\frac{1}{2})} (1 - 2 \sin^2 z)$$

are the Jacobi polynomials and $N(\alpha, k, \lambda, n)$ is the normalization constant. The factor $[-(k+\lambda)^2]$ in (25) sets the ground state eigenvalue to zero.

The Hamiltonian (25) is factorized by

$$d_{PT}^\mp = \pm \frac{d}{dz} + \frac{d}{dz} W_{PT}(z) = \mp \frac{d}{dz} - k \cot z + \lambda \tan z \quad (28)$$

and the supersymmetric partner is

Supersymmetric quantum mechanics and new potentials

$$\begin{aligned}
 H_-^{(PT)} &= d_{PT}^- d_{PT}^+ = d_{PT}^+ d_{PT}^- - [d_{PT}^-, d_{PT}^+] = \\
 &= -\frac{d^2}{dz^2} + \frac{k(k+1)}{\sin^2 z} - \frac{\lambda(\lambda+1)}{\cos^2 z} - (\lambda+k)^2
 \end{aligned} \tag{29}$$

The generalization of the operators (28), as indicated in (8), leaves us with

$$\begin{aligned}
 F_{PT}(z) &= -k \cot g z + X \operatorname{tg} z + \frac{[(\sin z)^{2k} + (\cos z)^{2\lambda}]^{-1}}{\Gamma_{PT} + \int_0^z [(\sin \bar{z})^{2k} + (\cos \bar{z})^{2\lambda}]^{-1} d\bar{z}} \\
 &\equiv -k \cot z + \lambda \tan z + \phi_{PT}(z)
 \end{aligned} \tag{30}$$

We choose $\Gamma_{PT} > 0$ to avoid singularities . Thus, the new Hamiltonian is

$$\begin{aligned}
 \mathcal{H}_{PT} &= D_{PT}^+ D_{PT}^- = D_{PT}^- D_{PT}^+ - [D_{PT}^-, D_{PT}^+] = \\
 &= -\frac{d^2}{dz^2} + \lambda(\lambda-1) \frac{1}{\cos^2 z} + \frac{k(k-1)}{\sin^2 z} - (\lambda+k)^2 - 2 \frac{d}{dz} \phi_{PT}(z)
 \end{aligned} \tag{31}$$

and the potential is

$$\mathcal{V}_{PT}(z) = \frac{\lambda(\lambda+1)}{\cos^2 z} + \frac{k(k+1)}{\sin^2 z} + (\lambda+k)^2 - 2 \frac{d}{dz} \phi_{PT}(z) \tag{32}$$

This Hamiltonian has the spectrum given by (27) and its eigenfunctions are

$$\Psi_n^{PT}(z) = D_{PT}^+ d_{PT}^- \Psi_n^{PT} \tag{33}$$

and the ground state ($\tilde{\Psi}_{PT}(z)$) is

$$\tilde{\Psi}_{PT}(z) \propto (\sin z)^{-k} (\cos z)^{-\lambda} \exp \left\{ \int_0^z \phi_{PT}(\bar{z}) d\bar{z} \right\} \tag{34}$$

The other potential that we treat is the Rosen-Morse one¹³. It was recently studied by Nieto¹² and Aragão de Carvalho¹⁴. Its Hamiltonian be written in the form

$$H_+^{RM} = k^2 \varphi_0^2 \left\{ -\frac{d^2}{dz^2} + \beta \tanh z - \gamma_+ \operatorname{sech}^2 z \right\} + \mu^2 + \varphi_0 \tag{35}$$

Elso Drigo Filho

where $z = k\varphi_0 x$; $\beta = 2\mu/k^2\varphi_0$; $\gamma_+ = (1+k)/k^2$; k, φ_0 and μ are constants. The eigenvalues and the eigenfunctions of this Hamiltonian are

$$\psi_{+,n}^{(\text{RM})} = N(k, \varphi_0, \mu; n) e^{az} \cosh^{-b}(z) F(-n, (4\gamma_+ + 1)^{1/2} - n; a + b + 1; 1/2[1 + \tanh z]) \quad (36)$$

$$E_n = -K^2\varphi_0^2(a^2 + b^2) + \mu^2 + \varphi_0^2; \quad a = \frac{-\beta}{(4\gamma_+ + 1)^{1/2} - 2n - 1}$$

and

$$b = \frac{1}{2}[(4\gamma_+ + 1)^{1/2} - 2n - 1] \quad (37)$$

We can factorize the Hamiltonian (35) by the operators

$$d_{\text{RN}}^\pm = \mp k\varphi_0 \frac{d}{dz} \mp \varphi_0 \tanh z \pm \mu \quad (38)$$

that satisfy the commutation relation

$$[d_{\text{RM}}^-, d_{\text{RM}}^+] = 2k\varphi_0^2 \text{sech}^2 z \quad (39)$$

The supersymmetric partner of (35) is

$$H_-^{\text{RM}} = d_{\text{RM}}^+ d_{\text{RM}}^- = k^2\varphi_0^2 \left\{ -\frac{d^2}{dz^2} + \beta \tanh z - \gamma_- \text{sech}^2 z \right\} + \mu^2 + \varphi_0^2 \quad (40)$$

where

$$\gamma_- = \frac{1-k}{k^2}$$

From the generalized operators (8) we obtain

$$F_{\text{RM}}(z) = \varphi_0 \tanh z \pm \mu \pm \frac{e^{2\mu} (\cosh z)^{-2\varphi_0}}{\Gamma_{\text{RM}} + \int_0^z e^{-2\mu z} (\cosh z)^{-2\varphi_0} dz} \mp \equiv \varphi_0 \tanh z + \mu + \phi_{\text{RM}}(z) \quad (41)$$

To avoid singularities we choose $\Gamma_{\text{RM}} > 0$. Then, the new Hamiltonian is

Supersymmetric quantum mechanics and new potentials

$$\begin{aligned} \mathcal{H}_{\text{RM}} &= D_{\text{RM}}^+ D_{\text{RM}}^- = D_{\text{RM}}^- D_{\text{RM}}^+ [D_{\text{RM}}^-, D_{\text{RM}}^+] = \\ &= k^2 \varphi_0^2 \left\{ -\frac{d^2}{dz^2} + \beta \tanh z - \gamma_+ \operatorname{sech} z - \frac{2}{k\varphi_0} - \frac{d}{dz} \phi_{\text{RM}}(z) \right\} + \mu^2 + \varphi_0^2 \end{aligned} \quad (42)$$

whose potential is

$$\mathcal{V}_{\text{RM}} = 2\varphi_0\mu \tanh z - \varphi_0(k+1)\operatorname{sech}^2 z - 2k\varphi_0 \frac{d}{dz} \phi_{\text{RM}}(z) \quad (43)$$

We note that the new Hamiltonian (42) has the spectrum given by (37) and its eigenfunctions are given in terms of the functions $\psi_{+,n}^{(\text{RM})}$ (36):

$$\Psi_n^{(\text{RM})}(z) = D_{\text{RM}}^+ d_{\text{RM}}^- \psi_{+,n}^{(\text{RM})} \quad (44)$$

with ground state

$$\tilde{\Psi}_{\text{RM}}(z) \alpha (\cosh z)^{1/k} e^{\mu z/k\varphi_0} \exp \left\{ \int_0^z \frac{\phi(\bar{z})}{k\varphi_0} d\bar{z} \right\} \quad (45)$$

5. Conclusion

From the potentials studied (particle in the box, Pöschl-Teller and Rosen-Morse) we obtained new potentials (eq. (22), (32) and (43)), which are different from the original ones, but whose spectra and eigenfunctions are known. The relation between the old system and the new one is established through the SQM.

As the spectrum of one potential is the same as that of its generalized version, some papers have appeared trying to distinguish these systems through the scattering produced by them. Cooper et al¹⁵ and Kare and Sukhatme¹⁶ have worked in this direction but they use a generalization method different from the one we used here.

Nieto⁴ explored the link between the generalization of the potential from supersymmetry and from the inverse scattering method. Using this result, the potential obtained in (22) should be the same (up to integration constants) as the potential

Elsó Drigo Filho

found by Abraham and Moses¹ for the particle in the box. Unfortunately, these potentials are different. However, we can see that in ref. 1 eq. (44) is not a solution of eq. (10) and also that eq. (45) is not derived from eq. (44); these mistakes justify the difference between the results.

The author wishes to thank Dr. M. A. Manna for helpful discussions about the paper of AM and Drs. R. M. Ricotta e V. Pleitez for their useful suggestions to this paper.

References

1. P. B. Abraham and H. E. Moses, Phys. Rev. **A22**, 1333 (1980).
2. D. L. Pursey, Phys. Rev. **D33**, 1049 (1986).
3. C. V. Sukumar, J. Phys. **A18**, L57 (1988).
4. M. M. Nieto, Phys. Lett. **145D**, 208 (1984).
5. N. A. Alves and E. Drigo Filho, J. Phys. **A21**, 3215 (1988).
6. E. Drigo Filho, J. Phys. A41, L1025 (1988).
7. B. Mielnik, J. Math. Phys. 25, 3387 (1984).
8. D. J. Fernandez C., Lett. Math. Phys. 8, 337 (1984).
9. F. Cooper and B. Freedman, Ann. Phys., NY 146, 262 (1983).
10. R. W. Haymaker and A. R. P. Rau, Am. J. Phys. 54, 928 (1986).
11. G. Pöschl and E. Teller, Z. Phys. 83, 143 (1933).
12. M. M. Nieto, Phys. Rev. A17, 1273 (1978).
13. N. Rosen and P. M. Morse, Phys. Rev. 42, 210 (1932).
14. C. Aragão de Carvalho, "From Polymers to Supersymmetry" Proceedings "1a. Escuela Latino-Americana de Física de Altas Energias", Valparaiso (1987).
15. F. Cooper, J. N. Ginocchio and W. Wipf, Phys. Lett. **129A**, 145 (1988).
16. A. Khare and U. Sukhatme, J. Phys. **A21**, L501 (1988)

Resumo

Usando a mecânica quântica supersimétrica vamos generalizar os potenciais: da partícula em uma caixa, Pöschl-Teller e Rosen-Morse. Calculamos os novos potenciais e indicamos suas respectivas autofunções e espectro.