

DORIANA

Supplemento agli
ANNALI DEL MUSEO CIVICO DI STORIA NATURALE "G. DORIA"

GENOVA

Vol. VI - N. 271

15 - XI - 1990

BERNHARD SEIFERT *)

SUPPLEMENTATION TO THE REVISION OF EUROPEAN
SPECIES OF THE ANT SUBGENUS *CHTHONOLASIUS*
RUZSKY, 1913

(HYMENOPTERA: FORMICIDAE)

INTRODUCTION

In a revision of the European of ant subgenus *Chthonolasius* (SEIFERT, 1988) I have demonstrated the existence of 11 morpho-ecological entities: 10 good species plus one entity which is either a well-defined subspecies or a parapatric species. Weak points of that revision were the low number of investigated specimens from the Mediterraneans and an incomplete consideration of synonymies. The investigations of material seen after the finishing of the manuscript confirmed the general conception of these 11 entities. However, one further good species *Lasius viehmeyeri* EMERY, 1922 from Greece has to be added and a replacement name for the invalid name *Lasius affinis* (SCHENCK, 1852) has to be published. Further, this paper wants to give supplementary notes on types and synonymies.

ACKNOWLEDGEMENTS

I wish to thank Valter Raineri from Museo Civico di Storia Naturale Genova (MCSNG) for the kind loan of several types of *C. EMERY* and help in supply with literature. Further I am deeply indebted to Claude Besuchet/Musée d'Histoire Naturelle Genève (MHNG)

*) Staatliches Museum für Naturkunde Görlitz, Am Museum 1 8900 Görlitz, E. Germany

for the loan of FOREL types and with many thanks I acknowledge the loan of BONDROIT types by Paul Dessart from Institute Royal des Sciences Naturelles de Belgique (I.R. Sc. N.B.) Bruxelles. I acknowledge further the friendly help of W. Dorow and J. -P. Kopelke in search for material of the Schenck collection at Senckenberg Museum Frankfurt and loan of specimens.

INVESTIGATION METHODS AND TERMINOLOGY

The investigation methods are explained elsewhere (SEIFERT, 1988) and I repeat here only the character terminology:

AW	- maximum alitrunk width before the tegulae
GHL	- length of longest hair on dorsal face of first gaster tergite.
HL	- maximum head length measured from midpoint of occipital border of anterior clypeal border; the head has to be tilted carefully until maximum length is in measuring plane.
HTMAX	- maximum diameter of hind tibia at midpoint; to measure the real cuticular surface, transmitted-light was used.
HTMIN	- minimum diameter of hind tibia at midpoint; transmitted-light was used.
HW	- maximum measurable head width; this is the outer distance of eyes (in males and queen) or the maximum width of head capsule (in workers normally across or slightly behind the eyes).
nHHT	- numbers of standing hairs projecting more than 11 μm from cuticular surface of hind tibia extensor profile, except the hairs on distal apex which are always present.
F2	- median line length of second funiculus segment in dorsal visual position (looking at the plane of scape).
IF2	- ratio F2/maximum width of second funiculus segment; the width is measured in dorsal view and transmitted-light conditions.

- PDF - average pubescence distance in front of midocellus; the number of pubescence hairs crossing a straight, transversal measuring line of 250-500 μm is counted and the average distance between two hairs is calculated.
- PDG - average pubescence distance on dorsum of first tergite obtained as in PDF.
- SL - maximum straight line length of scape excluding articular bulb.
- SMAX - maximum diameter of scape at midpoint; transmitted-light was used to measure real cuticular surface.
- SMIN - minimum diameter of scape at midpoint measured as in SMAX.

Redescription of *Lasius viehmeyeri* Emery, 1922

In the collection of MCSNG still exists one queen of Carlo Emery's type material found at Erymanthos/Greece. I have labelled this specimen as lectotype, det. Seifert 1989. The pin carries the original labels « Morea sept. Erymth. H. Vlasis, 800-1000 m, VIII.1901. Holtz » and « *L. umbratus* var. *viehmeyeri*, Type Emery ».

Description of the lectotype queen:

In many of absolute measurements clearly larger than the upper extremes of all other European species, particularly HW (1931 μm against 1807 μm in the largest *umbratus* queen) and HTMAX (388 against 310 μm in the largest *meridionalis* or *longiceps* queen). Scape flattening extreme, comparable to *jensi*. Flattening of hind tibia extreme, HTMAX/HTMIN as large as in upper extremes of *jensi* or *longiceps*. Second funiculus segment as much elongated as in upper extremes of *jensi*. Head, alitrunk and gaster with a dilute subdecumbent pubescence and with clearly shining surfaces (but somewhat less brilliant than in *rabaudi*). Pilosity on tibiae very sparse: hind tibiae on proximal part of extensor profile each with 4 suberect to erect hair the longest of which projects 46 μm from cuticular surface (Fig. 2). Dorsal surface of scape with subdecumbent pubescence projecting 15-20 μm from cuticular surface and with 1-2 weak erect hairs projecting up to 30 μm (Fig. 1).


As a consequence, scape and tibiae appear more smooth than in other *Chthonolasius* taxa with flat appendages (*rabaudi*, *meridionalis*, *jensi*, *longiceps*). Petiolar scale in caudal view with straight or slightly convex sides, converging a little towards dorsal crest that is notably emarginated (Fig. 3). Erect hairs on whole dorsal surface of gaster tergites sparse and as short as in *mixtus* but on dorsal face of first tergite 100-125 μm long. Genae each with 5-10 rather short, standing hairs which project at the most 45 μm from cuticular surface. Mesonotum with many erected hairs of 70-108 μm length. Length of longest hair on scutellum 158 μm . Morphological data:

HL 1591 μm , HW 1931 μm , SL 1424 μm , SMAX 181 μm , SMIN 84 μm , SMAX/SMIN 2.17, F2 194 μm , IF2 2.08, AW 1558 μm , PDF 12.9 μm , PDG 13.5 μm , GHL 125 μm , HTMAX 388 μm , HTMAX/HTMIN 2.94, n_{HHT} 4, HL/HW 0.824, SL/HL 0.895, SL/HW 0.737, overall length from mandibular tip to acidopore 8900 μm , maximum petiole width 704 μm .

Comments

Taxonomic judgements from single specimens are often very problematic but it seems to be, in this case, no risk to regard *viehmeyeri* as good species because its character combination is outstanding and the specimen shows no signs of morphological malformations or abnormalities. The *Lasius viehmeyeri* queen is not to confuse with any other European species: It is a combination of extremely flattened scape and hind tibia, strongly elongated second funiculus segment, extremely large body size, very weak tibial and scape pilosity and notably emarginated petiolar scape. A little similar is the *rabaudi* queen but the *viehmeyeri* lectotype is much larger (HW 1931 against 1720 μm in the largest *rabaudi* queen), the standing hairs on whole body are much shorter in *viehmeyeri*, the appendage pilosity is much less developed, the flattening of hind tibia is stronger and the scape length indices are below the range of *rabaudi* (lower extremes in *rabaudi* 0.925 for SL/HL and 0.812 for SL/HW).

From the detailed description of A. Starcke, it is most probable that his *Lasius viehmeyeri* var. *dalmatica* STARCKE, 1937 from the Dinaric Alps belongs indeed to *viehmeyeri*. In Starcke's description of the *dalmatica* queen we find no contradictions to the above description of the *viehmeyeri* lectotype in statements on pubescence, pilosity, cuticular surface, petiolar scale, body shape and measurements. Star-



Figs. 1-3: lectotype queen of *Lasius viehmeyeri* EMERY, 1922; 1: scape in frontal view with minimum midpoint diameter in visual plane and seen in transmitted-light; 2: hind tibia with maximum midpoint diameter in visual plane and seen in transmitted light; 3: petiolar scale in frontal view.

not to see the Emery material and suspected that he possibly was going to produce a synonym (STÄRCKE 1937), gives for *dalmatica*: HTMAX 369 μm , maximum petiole width 730 μm , SL 1405 μm .

The worker and the male of *Lasius viehmeyeri* are unknown.

Lectotype Fixation for **Lasius distinguendus** Emery, 1916

The date of publication of Emery's original description was February 1916. He gave no date of collecting and reported as collecting site simply « Bologna ». According to the present knowledge (SEIFERT 1988), only queens guarantee a safe distinction of *Lasius distinguendus* from related species. I could examine a total of 9 queens from the original material of Emery. 4 queens were labelled with « Bologna,

1916, Emery », 2 queens with « Borgo Panigale, Estate 1914 » and three others with « Bologna, 1912, Emery ». The queens of 1916 can not be regarded as types because they must have been collected after the date of publication (the first young queens are found in the nests June). As a consequence, I have selected as lectotype a queen with a label « Bologna, 1912, Emery » and labelled it as « *Lasius distinguendus* Emery, 1916; Lectotype (det. Seifert 1989) ». This lectotype is stored in MCSNG.

The examination of 3 type queens of *Formicina umbrata* v. *nuda* BONDROIT 1917 (stored in coll. of I.R. Sc. N.B. Bruxelles) showed clearly the synonymy with *L. distinguendus* EMERY, 1917. It should be emphasized that all the Emery and Bondroit specimens and all further queen material I have examined after the finishing of my revision confirmed the general validity of the *distinguendus* diagnosis stated by SEIFERT (1988) and there remains no doubt that it is a good morpho-species different from *L.umbratus* NYLANDER.

There are few queen examples from Central and East Germany with a somewhat doubtful identity but their interpretation as lower pilosity extremes of *umbratus* seems reasonable.

In my revision is suggested that workers of *Lasius distinguendus* and *balcanicus* could be separated in a number of cases by morphometry of nest samples. In the Emery collection there are two workers accompanied with *distinguendus* queens under the same label (« Borgo Panigale ») which show characters typical for *balcanicus* (longer scape, flatter appendages). If these workers were really from the same nest as the queens, they completely disproved my earlier suggestion that SL/HL, HTMAX/HTMIN and SMAX/SMIN could be used in many nest samples for separation of *balcanicus* and *distinguendus* workers. So I can not tell just a single objective criterion for separation of workers while the distinction of the two species' queens has proved as very easy in each case.

Lasius mixtoumbratus FOREL, 1974 can not be interpreted

The original paper of Forel from 1874 contains no figure, no statements on collecting date and locality, and the short verbal description is absolutely insufficient to permit a taxonomic interpretation in such a difficult ant group. From the MHNG collection I got a loan of 3 queens and 6 workers labelled apparently by Forel himself with « Burghölzli, 23 V 91, L. mixto-umbratus For., Coll. Forel, Typus ». The collecting date definitely shows that this series can not be

recognized as type material since it was collected 17 years after the first description. The absence of types and the useless original description make it clear that the name *Lasius mixtroumbratus* FOREL, 1874 can not be referred safely to any species and its use should be avoided. On the other hand, it would be unreasonable to apply for an opinion of the International Commission on Zoological Nomenclature to give the Burghölzli series the status of neotypes; there is not to defend any stability of the name *mixtroumbratus* which was used very rarely during the last 100 years and was forgotten in the last 50 years. The correct name for Forels Burghölzli series must be *Lasius sabularum* (BONDROIT, 1918) - a species which is widely distributed in Central Europe and of which we have a confirmed type queen.

Lasius citrinus Emery, 1922 - a replacement name for the invalid taxon **Lasius affinis** (SCHENCK, 1852)

I was informed by Barry Bolton/BMNH London that the name *Formica affinis* SCHENCK, 1852 was preoccupied by *Formica affinis* LEACH, 1825 (Formicidae: *incertae sedis*) and *Formica affinis* LE GUILLOU, 1841 (now *Polyrhachis affinis* (Le G., 1841)). The international Code of Zoological Nomenclature requires the replacement of the name *Lasius affinis* (SCHENCK) by the first available junior synonym. Such formal taxonomic acts are disputed if they refer to a long-established, widely used name that has been transferred to another genus soon after its introduction and which has been « virtually » no homonym for several generations of myrmecologists. *Formica affinis* (SCHENCK, 1852) was removed from *Formica* by MAYR already in 1861 and did not reappear under *Formica* in the myrmecological literature of the last 120 years. As a consequence, it seems that the replacement of *Lasius affinis* by a forgotten name could produce irritations among a majority of myrmecologists. However, in case of *Lasius affinis*, I have no practical objections to follow the International Code because the « *Lasius affinis* » conception of the last 120 years is already confused and de facto invalidated by the many false subjective interpretations of species. So we have many « *Lasius affinis* » which are in reality *Lasius distinguendus* EMERY or *Lasius balcanicus* SEIFERT.

Schenck's description of *Formica affinis* is unusually detailed in terms of his time and in Senckenberg Museum Frankfurt exist two queens and two worker specimens ex coll. Schenck (all without collecting data or type labels) which characters are in very good agreement with the description of *Lasius affinis* (SCHENCK, 1852), sensu

SEIFERT, 1988. So we have a good notion which species is meant the more it has an outstanding character combination and can be identified in single workers. After *L. viehmeyeri*, the species is the rarest *Chthonolasius* in European ant collections and I have found only one name among 26 *Chthonolasius* taxa I know from Europe which can serve as replacement name: *Lasius bircornis* var. *citrina* EMERY, 1922 of which exists a lectotype in the collection of MCSNG. This worker specimen is labelled « Monte Gargano, 1907, legit. M. Hilf, coll. O. Leonhard » and « lectotype, designated by E.O. Wilson ». All characters of this specimen are typical for the species *Lasius affinis* sensu SEIFERT (1988). So I can state the following list of synonymies on the basis of investigated specimens (without having seen their specimens, I can not say which species Mayr, the early Emery, Forel, Bondroit, Baroni Urbani and Kutter determined as *Lasius affinis*):

Lasius citrinus EMERY, 1922 **stat nov.**

Lasius bicornis var. *citrina*, EMERY, 1922

Formica affinis SCHENCK, 1852:62. Junior primary homonym of

Formica affinis LEACH, 1825 [Formicidae, *incertae sedis*], and of

Formica affinis [LE GUILLOU, 1841 [now *Polyrhachis affinis* (LE GUILLOU, 1841)].

Lasius affinis (SCHENCK) sensu VAN BOVEN, 1955

Lasius affinis (SCHENCK) sensu COLLINGWOOD, 1963

Lasius affinis (SCHENCK) sensu SEIFERT, 1988

The following table gives data of the *Lasius citrinus* lectotype and West Palaearctic population data of *citrinus*, *balcanicus* and *distinguendus* workers.

	<i>citrinus</i> lectotype	<i>citrinus</i> (n = 33)		<i>balcanicus</i> (n = 35)		<i>distinguendus</i> (n = 26)	
		mean	SD	mean	SD	mean	SD
HL	1236	1211.5 ±	50.1	1068.5 ±	55.2	1062.4 ±	62.4
SMAX	107	107.8 ±	5.78	97.7 ±	8.29	93.4 ±	8.86
GHIL	117	116.4 ±	7.50	105.7 ±	8.83	105.9 ±	7.94
PDG	5.7	6.10 ±	1.01	11.52 ±	3.11	11.64 ±	2.88
PDF	8.2	7.72 ±	1.43	10.50 ±	1.97	9.91 ±	1.90
HL/HW	1.036	1.035 ±	0.0167	1.045 ±	0.0161	1.047 ±	0.0186
SL/HL	0.856	0.847 ±	0.0145	0.879 ±	0.0141	0.873 ±	0.0186
SMAX/SMIN	1.384	1.461 ±	0.0859	1.471 ±	0.0688	1.382 ±	0.0967
nHHT	0	0.12 ±	0.28	6.68 ±	1.79	4.56 ±	2.84
nGen	0	0.66 ±	1.05	10.26 ±	2.96	7.21 ±	3.16

(nGen = number standing hairs on genae in front of eye which project, in dorsal view of head, more than 11 µm from cuticular surface; the value is the arithmetic mean of both genae).

These data show that even single *citrinus* workers can be separated from *balcanicus*/*distinguendus* using the character combination nHHT--nGen-PDG. The *citrinus* worker's body size is frequently above the upper extremes of these species and its appendages appear much more smooth due to more appressed pubescence and the lacking standing pilosity (SEIFERT, 1988).

Our knowledge on biology of *Lasius citrinus* is poor. It typically nests in hollow trees, rotten logs or stumps at sunny wood stands. The reports for occurrence of *citrinus* in treeless xerothermous grasslands or steppes refer surely in most cases to either *distinguendus* or *balcanicus*. Alates occur on average earlier than in many other *Chthonolasius* species even in the northern parts of its distributional range which indicates that elder sexual larvae, pupae or even imagines hibernate. Sure data are: May 2, 13; June 2, 7, 11, 17; July 7, 8, 17, 21; August 18. VAN BOVEN (1955) reports a large nest area with a numerous population and as inner nest structure a brown-yellow carton nest. A dealate queen with a *Lasius brunneus* worker attached on its antenna (Rüppurr near Karlsruhe, June 11 1951, stored in Naturkundemuseum Karlsruhe) and the preferred nest sites indicate that *brunneus* is a probable host species.

Lasius citrinus is more rarely collected than *bicornis* and I give here the *citrinus* sites of which I have checked specimens:

- | | |
|-------------|---|
| Belgium: | 1. Mechelen, leg. Van Boven 4.V.1954 |
| | 2. Chevremont, Liège, 21.VII.1921, in coll. I.R.Sc. N.B. |
| Germany: | 3. Nassau (most probable site but not label), coll. Schenck |
| | 4. Rottleben/Bad Frankenhausen, leg. Pisarski 2. VI.1964 |
| | 5. Rüppurr near Karlsruhe, 11.VI.1951 |
| | 6. Karlsruhe, 17.VI.1955 |
| Spain: | 7. Huesca, Torla-Ordesa, 13.V.1953, leg. Yarrow |
| Italy: | 8. Monte Gargano, leg. M. Hilf 1907 |
| | 9. Tirol, Bolzano, July 1919, leg. C. Minozzi |
| Yugoslavia: | 10. Dalmatia, leg. Köller 20.IV. 1932 |
| Bulgaria: | 11. 30 km N of Karnobat, leg. Krampfl 8.VII.1981 |
| S Russia: | 12. Tatyankovka/Donetz River, 18.VIII.1938 |
| | 13. Svyatogorsk near Donetzk, 7.VI.1939 |
| | 14. Smiyev near Kharkov, 20.VIII.1938 |
| | 15. Ubinskaya/Kuban River, 22.IX.1953 |

- Caucasus: 16. Kazbegi (1800 m), 28.VI.1963
 Azerbaidzhan: 17. Alazapin/Talysh Mountains, leg. Arnoldi July 1929

Commented synonymic list of European **Chthonolasius** species

1. ***Lasius citrinus*** EMERY, 1922; good species, lectotype seen
Formica affinis SCHENCK, 1852; invalid preoccupied name, original material investigated
Lasius umbratus (NYLANDER) sensu WILSON, 1955, partim; material seen
2. ***Lasius balcanicus*** SEIFERT, 1988; good species, types seen; species often named by different authors as *Lasius affinis*
Lasius affinis (SCHENCK) sensu STÄRCKE, 1937; synonymy clear from detailed description and investigation of original material
3. ***Lasius bicornis*** (FÖRSTER, 1850); good species, identity clear from description
Formica incisa SCHENCK, 1852; synonymy can be assumed from description, two worker specimens ex coll. Schenck were studied
Lasius bicornis subsp. *oertzeni* FOREL, 1910; lectotype studied, no subspecies
Formicina microgyna BONDROIT, 1918; synonymy clear from description of the queen
Lasius bicornis var. *neapolitana* EMERY, 1922; type studied: in all characters a typical *bicornis* queen
4. ***Lasius distinguendus*** EMERY, 1916; good species, types seen
Lasius bicornis distinguendus var. *hybrida* EMERY, 1916; original material seen
Formicina umbrata var. *nuda* BONDROIT, 1917; type queen seen
Lasius umbratus distinguendus var. *cereomicans* STÄRCKE, 1937; synonymy with *distinguendus* very probable from description
Lasius hybridus (EMERY) sensu ARNOLDI and DLUSSKY, 1978; material seen
Lasius umbratus (NYLANDER) sensu WILSON, 1955, partim; material seen
5. ***Lasius jensi*** SEIFERT, 1982; good species, types seen
Lasius meridionalis (BONDROIT) sensu PISARSKI, 1975, partim; material seen

- Lasius rabaudi* (BONDROIT) sensu ARNOLDI and DLUSSY, 1978, partim; material seen
6. *Lasius jensi longiceps* SEIFERT, 1988; type seen, unclear whether parapatric species or well-defined subspecies
7. *Lasius meridionalis* (BONDROIT, 1919; good species, type queen seen
Lasius meridionalis (BONDROIT) sensu STÄRCKE, 1937; the identity of Stärcke's Material is clear from his detailed descriptions
Lasius rabaudi (BONDROIT) sensu WILSON, 1955, partim; material seen
Lasius rabaudi (BONDROIT) sensu COLLINGWOOD, 1963; synonymy clear from description and geographic origin of material
Lasius rabaudi (BONDROIT) sensu KUTNER, 1977, partim; material seen
Lasius rabaudi (BONDROIT) sensu ARNOLDI and DLUSSKY, 1978, partim; material seen
8. *Lasius mixtus* (NYLANDER, 1846); good species; identity clear from detailed description of type (STÄRCKE, 1937) and type locality
Lasius umbratus (NYLANDER) sensu WILSON, 1955, partim; material seen
9. *Lasius rabaudi* (BONDROIT, 1917); good species, type queen seen
10. *Lasius sabularum* (BONDROIT, 1918); good species, type queen seen
Lasius umbratus (NYLANDER) sensu WILSON, 1955, partim; material seen
11. *Lasius umbratus* (NYLANDER, 1846); good species, identity clear from detailed description of type (STÄRCKE, 1937) and type locality
Formicina belgarum BONDROIT, 1918; type queen seen
Lasius umbratus var. *hirtiscapus* STÄRCKE, 1937; synonymy concluded from Stärcke's detailed description
12. *Lasius viehmeyeri* EMERY, 1922; good species, type queen seen
Lasius viehmeyeri var. *dalmatica* STÄRCKE, 1937; synonymy can be concluded from Stärcke's detailed description
Lasius umbratus (NYLANDER) sensu WILSON, 1955, partim; material seen

Incertae sedis

Lasius mixtoumbratus FOREL, 1874; insufficient description, no types available

Lasius umbratus var. *exacutus* RUZSKY, 1902; types not got, description insufficient

Lasius affinoumbratus DONISTHORPE, 1927; concluded from type locality it must be a synonym of either *mixtus*, *sabularum* or *meridionalis*

Chthonolasius [!] *affinis* var. *nyaradi* RÖSZLER, 1943; insufficient description, types not available

Synonyms of ***Lasius flavus*** (FABRICIUS, 1871)

Lasius umbratus ibericus SANTSCHI, 1925; type seen

Lasius umbratus ibericus var. *sancho* SANTSCHI, 1925; types seen

REFERENCES

- BONDROIT, J. (1917): Notes sur quelques Formicidae de France. - *Bull. Soc. Ent. France*, **86**: 174-177.
- COLLINGWOOD, C. (1963): The *Lasius* (Chthonolasius) *umbratus* species complex in North Europe. - *The Entomologist* **96**: 145-158.
- EMERY, C. (1916): Fauna entomologica italiana. I. Hymenoptera-Formicidae. - *Boll. Soc. Ent. Ital.* **47**: 79-275.
- (1922): Il genere *Lasius* (F.) Mayr e particolarmente le forme mediterranee del gruppo *umbratus* Nyl. - *Boll. Soc. Ent. Ital.* **54**: 9-15.
- FOREL, A. (1874): Les fourmis de la Suisse. - *Neue Denkschriften der allg. schweizerischen Gesellschaft für die gesamten Naturwissenschaften*, Zürich, **26**: 48.
- (1910): Fourmis de Palestine et de Syrie. - *Ann. Soc. Ent. Belgique* **54**: 6-32.
- MAYR, G.L. (1861): Die europäischen Formiciden. Wien 1861, pp. 1-80
- SANTSCHI, F. (1925): Fourmis d'Espagne et autres espèces palearctiques (Hymenopt.) - *EOS*, Inst. Espagnol de Entomologia Madrid **1** (4): 339-360.
- SCHENCK, C.F. (1852): Beschreibung nassauischer Ameisenarten. - *Jahrb. Ver. Naturkunde Nassau*, **8**, Wiesbaden: 3-149.
- SEIFERT, B. (1988): A revision of the European species of the ant subgenus *Chthonolasius*. - *Entomol. Abh. Mus. Tierkunde. Dresden* **51** (8): 143-180.
- STÄRCKE, A. (1937): Retouches sur quelques fourmis d'Europe. II *Lasius* groupe *umbratus* Nylander. - *Tijdschrift voor Entomologie* **80**: 37-72.
- VAN BOVEN, J.K.A. (1917): *Lasius* (Chthonolasius) *affinis* Schenck - Fauna Neerl. nov. spec. (Hymenoptera, Formicidae). - *Nat. His. Maandblad* **44**: 6-10.

SUMMARY

A redescription of *Lasius viehmeyeri* EMERY, 1922 is given showing that it is a good species with outstanding characters. *Lasius citrinus* EMERY, 1922 is redescribed and published as replacement name for the invalid name *Lasius affinis* (SCHENCK, 1852) which is a junior primary homonym. 17 proved sites of *Lasius citrinus* are reported for Europe and Caucasia. A lectotype of *Lasius distinguendus* EMERY, 1916 is fixed. The taxon *Lasius mixtoumbratus* FOREL, 1874 can not be interpreted because types are nor available. A commented synonymic list of 26 taxa referred to *Chthonolasius* is given for the territory of Europe.

RIASSUNTO

Viene descritta la specie *L. viehmeyeri* Emery, 1922 dimostrando che è una specie valida, *L. citrinus* Emery, 1922 è ridescritto e pubblicato per rimpiazzare il nome non valido di *L. affinis* (Schenck, 1852) che è più recente omonimo primario. Sono segnalate 17 località per l'Europa ed il Caucaso dove è stato trovato *L. citrinus*. È stabilito il lectotipo di *L. distinguendus* Emery, 1916. Il taxon *L. mixtoumbratus* Forel, 1874 non può essere interpretato per la mancanza di tipi. È fornita una lista sinonimica di 26 taxa di *Chthonolasius* d'Europa.
