## Downloaded from http://aacrjournals.org/cancerres/article-pdf/79/19/5125/2783177/5125.pdf by guest on 24 August 2022

## Editor's Note: Suppression of Tumor Cell Growth Both in Nude Mice and in Culture by n-3 Polyunsaturated Fatty Acids: Mediation through Cyclooxygenase-Independent Pathways



Mary D. Boudreau, Kyung Hee Sohn, Sang Hoon Rhee, Sam W. Lee, Jay D. Hunt, and Daniel H. Hwang

The editors are publishing this note to alert readers to a concern about this article (1). The editors were made aware of duplications of Fig. 5 in this article, with Fig. 3 in ref. 2 and Fig. 5 in ref. 3 depicting the results from soft-agar colony assays. Because satisfactorily corrected figures could not be provided, the editors are publishing this note to alert readers to these concerns.

## References

- Boudreau MD, Sohn KH, Rhee SH, Lee SW, Hunt JD, Hwang DH. Suppression of tumor cell growth both in nude mice and in culture by n-3 polyunsaturated fatty acids: mediation through cyclooxygenase-independent pathways. Cancer Res 2001;61:1386–91.
- Lee SW, Reimer CL, Oh P, Campbell DB, Schnitzer JE. Tumor cell growth inhibition by caveolin reexpression in human breast cancer cells. Oncogene 1998;16:1391–7.
- 3. Lee SW, Reimer CL, Fang L, Iruela-Arispe ML, Aaronson SA. Overexpression of kinase-associated phosphatase (KAP) in breast and prostate cancer and inhibition of the transformed phenotype by antisense KAP expression. Mol Cell Bio 2000;20:1723–32.

Published first October 1, 2019. Cancer Res 2019;79:5125 doi: 10.1158/0008-5472.CAN-19-2362 ©2019 American Association for Cancer Research.