

Surface modifications of silica nanoparticles are crucial for their inert versus proinflammatory and immunomodulatory properties

Viviana Marzaioli¹
 Juan Antonio Aguilar-Pimentel²
 Ingrid Weichenmeier¹
 Georg Luxenhofer³
 Martin Wiemann⁴
 Robert Landsiedel⁵
 Wendel Wohlleben⁵
 Stefanie Eiden⁶
 Martin Mempel⁷
 Heidrun Behrendt¹
 Carsten Schmidt-Weber¹
 Jan Gutermuth^{1,8}
 Francesca Alessandrini¹

¹Center of Allergy and Environment (ZAUM), Technische Universität and Helmholtz Zentrum München, Member of the German Center for Lung Research (DZL), Munich, Germany; ²Department of Dermatology and Allergy Biederstein, Technische Universität München (TUM) and German Mouse Clinic, Helmholtz Zentrum München, German Research Center for Environmental Health, Munich-Neuherberg, Germany; ³Institute of Developmental Genetics, Helmholtz Zentrum München, Neuherberg, Germany; ⁴IBE R&D gGmbH, Münster, Germany; ⁵BASF, Ludwigshafen, Germany; ⁶Bayer Technology Services, Leverkusen, Germany; ⁷Department of Dermatology, Venereology and Allergology, Universitätsmedizin Göttingen (UMG), Göttingen, Germany; ⁸Department of Dermatology, Vrije Universiteit Brussel, Brussels, Belgium

Correspondence: Francesca Alessandrini Institute for Allergy Research (IAF), Helmholtz Zentrum München/ZAUM Center of Allergy and Environment, TU München, Ingolstädter Landstr. 1, D-85764 Neuherberg, Germany
 Tel +49 89 3187 2524
 Fax +49 89 3187 2540
 Email franci@helmholtz-muenchen.de

Background: Silica (SiO₂) nanoparticles (NPs) are widely used in diverse industrial and biomedical applications. Their applicability depends on surface modifications, which can limit potential health problems.

Objective: To assess the potential impact of SiO₂ NP exposure and NPs chemical modifications in allergic airway inflammation.

Methods: Mice were sensitized by five repetitive intraperitoneal injections of ovalbumin/aluminum hydroxide (1 µg) over 42 days, then intratracheally instilled with plain or modified SiO₂ NPs (50 µg/mouse), and subsequently aerosol challenged for 20 minutes with ovalbumin. One or 5 days later, allergic inflammation was evaluated by cell differentiation of bronchoalveolar lavage fluid, lung function and gene expression and histopathology, as well as electron and confocal microscopy of pulmonary tissue.

Results: Plain SiO₂ NPs induced proinflammatory and immunomodulatory effects in vivo, highlighted by enhanced infiltration of inflammatory cells in the bronchoalveolar lavage fluid, induction of a pulmonary T helper type 2 (Th2) cytokine pattern, differentiation of type 2 macrophages, and by morphological changes in the lung of sensitized mice. These effects were dramatically attenuated using surface-functionalized NPs with amino and phosphate groups, but not with polyethylene glycol. The role of macrophages in taking up SiO₂ NPs was confirmed by flow cytometry, confocal microscopy, and gene expression analysis.

Conclusion: Our data suggest that amino and phosphate surface modifications, but not polyethylene glycol (PEG), mitigate the proinflammatory and immunomodulatory effect of SiO₂ NPs in allergic airway inflammation, paving the way for new strategies in the production of nanomaterials with lower health impact for humans.

Keywords: immunomodulation, allergic inflammation, health impact

Introduction

Engineered nanoparticles (NPs) are frequently used in commercial products due to their unique physical and chemical characteristics, which depend on their specific dimensions (100 nm or less in one or more dimension).¹ Silicon dioxide (silica [SiO₂]) nanomaterials are of great importance in the fabrication of electric and thermal insulators, catalyst supports, and drug carriers, as well as in gene delivery, media for coating processes, and filler materials.² They are also produced on an industrial scale as additives for cosmetics, drugs, printer toners, varnishes, and food packing.³ The same properties that mediate the advantages of nanotechnology also confer their potential toxic effects, resulting in a major health concern regarding the use of nanomaterials.^{1,4}

NPs may enter the body via different routes, such as the gastrointestinal tract,⁵ skin,^{6,7} and lung.^{8,9} Due to their small size, NPs can translocate from entry portals into

the circulatory and lymphatic systems, causing long-lasting cell damage.^{10,11} Several studies show that SiO₂ NPs induce dose-dependent cytotoxicity in *in vitro* systems, including human lung epithelial cells¹² and bronchoalveolar carcinoma-derived cells¹³ and human alveolar barrier,¹⁴ as well as murine macrophages.¹⁵ Short-term instillations or inhalation exposure to amorphous SiO₂ NPs showed that these particles possess the potential to cause toxic effects in the lung.^{16–18} Immunomodulatory effects by particle exposure during the sensitization phase have also been demonstrated.^{19,20} Recently, surface modifications of SiO₂ NPs were synthesized to study their influence on the biological activity of SiO₂ NPs. *In vitro* studies using various mammalian cells have demonstrated that unmodified SiO₂ NPs exhibit higher cytotoxic and genotoxic properties compared to surface-modified NPs.^{18,21,22}

The aim of this study was to investigate the effects of amorphous SiO₂ NPs and their surface modifications in the elicitation phase in a murine model of ovalbumin-specific allergic airway inflammation. SiO₂ and polyethylene glycol (PEG) SiO₂ (SiO₂-PEG) exerted proinflammatory and immunomodulatory activity in allergic airway inflammation, while modifying SiO₂ NPs with amino and phosphate groups markedly decreased these effects. Alternative activation of pulmonary macrophages that take up SiO₂ NPs and distinct chemokine patterns in the lung were identified as underlying mechanisms.

Materials and methods

Animals

Female 6- to 10-week-old BALB/c mice were obtained from Charles River (Wilmington, MA, USA), housed under specific

pathogen-free conditions in individually ventilated cages (VentiRack; BioZoneGlobal Ltd, Ramsgate, UK) and fed a standard diet and water *ad libitum*. The study was conducted under federal guidelines for the use and care of laboratory animals and was approved by the Government of the District of Upper Bavaria and the Animal Care and Use Committee of the Helmholtz Center Munich, Munich, Germany.

NP preparation

Plain SiO₂ (SiO₂), PEGylated SiO₂ (SiO₂-PEG), phosphate-coated SiO₂ (SiO₂-P), and amino-coated SiO₂ (SiO₂-NH₂) NPs were obtained from BASF SE, Ludwigshafen, Germany, and fluorescein isothiocyanate (FITC)-conjugated SiO₂ (SiO₂-FITC) NPs from Bayer Technology Services GmbH, Leverkusen, Germany. All NPs were amorphous with a single particle/agglomerate size between 5 and 50 nm, with the exception of SiO₂-FITC (23–30 nm). NPs were freshly diluted in water (*aqua ad iniectionem*; B Braun Melsungen AG, Melsungen, Germany) and buffered with phosphate-buffered saline (PBS) to physiological pH, to the final concentration of 1 mg/mL, just before intratracheal instillation. Supernatant controls (SUP) were obtained by hard sedimentation (24,000 rpm, 15 hours). Additional NP characteristics are disclosed in the “Supplementary materials” section.

Study design

To evaluate the immunomodulatory effect of NPs, a protocol of mild allergic inflammation in the lung was used, as previously described (Figure 1A).²³ Briefly, mice were sensitized by repetitive intraperitoneal injections of 1 µg ovalbumin (OVA) (grade VI; Sigma-Aldrich, St Louis, MO, USA) in PBS adsorbed to 2.5 mg aluminum




Figure 1 Experimental protocol.

Notes: (A) BALB/c mice were intraperitoneally sensitized (S) with OVA/aluminum hydroxide or non-sensitized (NS) with phosphate-buffered saline/aluminum hydroxide (black arrows). On day 52, S and NS mice were intratracheally instilled with SiO₂ nanoparticles or with SUP (white arrow) and subsequently challenged with OVA aerosol (arrowhead). Sacrifice was on day 53 or 57, according to the analysis performed (stars). (B) Experimental groups.

Abbreviations: FITC, fluorescein isothiocyanate; IP, intraperitoneal; IT, intratracheal; OVA, ovalbumin; PEG, polyethylene glycol; SiO₂, silicon dioxide; SiO₂-NH₂, amino-coated SiO₂; SiO₂-P, phosphate-coated SiO₂; SiO₂-PEG, PEGylated SiO₂; SUP, supernatant controls.

hydroxide (alum) (Thermo Fisher Scientific, Waltham, MA, USA) on days 0, 7, 14, 28, and 42. Blood samples were taken before and after sensitization. OVA/alum-sensitized mice (S mice), compared to non-sensitized mice (NS mice), were characterized by high titers of OVA-specific immunoglobulin E (8.05 ± 1.64 versus 0.1 ± 0.03 $\mu\text{g/mL}$). At day 52, mice were intratracheally instilled with 50 μg (1 mg/mL) of SiO_2 NPs, or with their correspondent SUP (Figure 1B). For dose-response studies, S mice were instilled with 12.5, 25, or 50 μg of SiO_2 and SiO_2 -PEG NPs. NP instillation was followed by OVA aerosol challenge for 20 minutes with 1% OVA in PBS delivered by a PARI BOY nebulizer (PARI GmbH, Starnberg, Germany). Lung function, flow cytometry, and confocal analysis were performed on day 53. Bronchoalveolar lavage (BAL), histology, scanning electron microscopy (SEM), and messenger (m)RNA expression analysis were performed 5 days after OVA challenge (day 57).

Flow cytometry

On day 53, single cell suspensions were generated from lungs of NS and S mice exposed to SiO_2 -FITC or to correspondent SUP ($n=5-7$), dissociated with a lung dissociation kit (kit number 130-065-927; Miltenyi Biotec GmbH, Bergisch Gladbach, Germany), and analyzed by flow cytometry, monitoring the FITC conjugate. Untreated mice were used as negative controls. To investigate the interaction of SiO_2 -FITC with different lung cell populations, cells were stained and gated as described previously.²⁴ anti-mouse CD45 PE-Cyanine7, 7AAD, and anti-mouse F4/80 PE were obtained from eBioscience (San Diego, CA, USA), anti-mouse CD11b APC-CyTM7 from BD Biosciences (San Jose, CA, USA), anti-mouse CD11c APC from BioLegend (San Diego, CA, USA), and GR1 (LY6C/G) Pacific OrangeTM from Invitrogen (Life Technologies, Carlsbad, CA, USA). Samples were measured in a BD FACSCanto II flow cytometer and data were analyzed using a FlowJo graphics application V9.7.2 (TreeStar Inc., Ashland, OR, USA).

Interstitial macrophages were defined as CD45^+ , Gr-1^{Low} , F4/80^+ , CD11b^+ but CD11c^- ; alveolar macrophages as CD45^+ , Gr-1^{Low} , F4/80^+ , CD11c^+ but CD11b^- ; neutrophils as CD45^+ , $\text{Gr-1}^{\text{High}}$, and CD11b^+ ; and lymphocytes as CD45^+ , but CD11b^- and SSC^{Low} .

Confocal microscopy

On day 53, lungs were embedded in, Tissue-TekTM CRYO-OCT Compound (Thermo Fisher Scientific), removed, and snap frozen in liquid nitrogen ($n=2/\text{group}$). Pulmonary tissue sections (20 μm) were fixed in 4% paraformaldehyde

(Merck KGaA, Darmstadt, Germany) and subsequently blocked for 2 hours with 3% bovine serum albumin/PBS. The primary antibody for detecting macrophages (anti-CD68 marker, 1:100; AbD Serotec, Kidlington, UK) was incubated overnight and the secondary antibody ([anti-rat immunoglobulin G] Alexa Fluor[®] 555, 1:7000; Cell Signaling Technology, Inc., Danvers, MA USA) for 1 hour. Antibodies were diluted in 1% bovine serum albumin/PBS. Nuclei were stained with 4',6-diamidino-2-phenylindole (DAPI) (0.1 $\mu\text{g/mL}$ in PBS; Sigma-Aldrich) for 5 minutes. Sections were embedded in Moviol 4-88 (10% polyvinyl alcohol, 20% glycerol in PBS; Merck). Images were acquired with an inverted confocal laser scanning microscope (LSM 510 Meta; Carl Zeiss Meditec AG, Jena, Germany) with 40 \times objective, Z-Stack size 6 μm , and collapsed two stacks. For detailed images, a 100 \times objective was used, with Z-Stack size 2 μm . Orthogonal sections of single cells were generated with the LSM Image Browser software (Carl Zeiss Meditec AG).

Histology and SEM

On day 57, after BAL, the lungs were excised and the left lobe fixed in 4% buffered formalin and embedded in paraffin ($n=4-5/\text{group}$). Sections of 3 μm thickness were stained with hematoxylin and eosin and periodic acid-Schiff. For electron microscopy, lungs were processed as previously described.²⁵

Lung function analysis

Lung function analysis was performed on day 53, 24 hours after allergen challenge in intubated, mechanically ventilated animals ($n=5-8/\text{group}$; Buxco[®] Research Systems, Wilmington, NC, USA). Anesthesia was induced by an intraperitoneal injection of ketamine (100 mg/kg) and xylazine (5 mg/kg) in PBS. After cannulation of the trachea, the animals were ventilated with room air at rates of 130 breaths/minute. A protocol for measuring airway hyperreactivity was initiated by challenging the mice with increasing methacholine concentrations by inhalation, using an in-line nebulizer (5 μL methacholine solution in PBS delivered for 30 seconds at the following concentrations: 0, 1.25, 2.5, 5, 10, 20, and 40 mg/mL). Data were recorded using FinePoint software version 1.0 (Buxco[®] Research Systems). The highest value of respiratory system resistance and the lowest value of dynamic compliance were recorded every 5 seconds during the data-recording interval, which was set at 3 minutes after each methacholine level. The heart rate of each animal was continuously monitored using an electrocardiography device connected with three subcutaneous electrodes throughout the entire experiment (Buxco[®] Research Systems).

Polymerase chain reaction arrays and real-time polymerase chain reaction

Total RNA was extracted from snap-frozen lung tissue as previously described.²⁵ For polymerase chain reaction (PCR) arrays (n=4/group of experiment) (SABiosciences, Hilden, Germany), 0.4 µg RNA was converted into complementary (c)DNA with the RT² First Strand Kit (Qiagen) and real-time PCR was performed with RT² SYBR Green ROX qPCR Mastermix (Qiagen) and ViiATM7 thermocycler (Applied Biosystems; Life Technologies). Relative expression levels were calculated using the 2^{ΔΔct} method, normalized to the arithmetic mean of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and beta-actin (ACTB). For the dose-response experiment, 0.4 µg of RNA was converted to cDNA following the manufacturer's instructions (RevertAid First Strand cDNA Synthesis Kit #K1622; Fermentas, Thermo Fisher Scientific). Real-time PCR was performed with SYBR Green and ViiATM7 thermocycler (n=3/group) (Applied Biosystems). Relative expression levels were calculated using the 2^{ΔΔct} method, normalized to ACTB. Primer sequences are listed in Table S1. Data were considered significant with a *P*-value ≤0.05 with a confidence interval of 95%.

Data analysis

Results are shown as boxplots indicating minimum, 25th percentile, median, 75th percentile, and maximum, or as mean ± standard deviation. Statistical significance among groups was determined by the Mann–Whitney *U*-test for boxplots, two-way analysis of variance with post hoc least significant difference test for lung function studies, and one-way analysis of variance with Bonferroni post hoc test for PCR and flow cytometry (GraphPad Prism; GraphPad Software, Inc., La Jolla, CA, USA). Statistically significant values were categorized as follows: *P*≤0.05, *P*≤0.01, and *P*≤0.001.

Results

SiO₂ NPs exert proinflammatory and Th2 immunomodulatory activity in vivo

In order to investigate possible proinflammatory and immunomodulatory effects of SiO₂, increasing concentrations of SiO₂ NPs (0, 12.5, 25, and 50 µg/mouse) were intratracheally instilled in the lungs of S mice prior to OVA challenge. Bronchoalveolar lavage fluid (BALF) was analyzed for differential cell count 5 days after OVA challenge. The lowest NP concentration used (12.5 µg) did not evoke a significant increase of inflammatory cells into the BALF compared to the respective SUP. Increasing NP dose to 25 µg led to a significant increase of macrophages (*P*<0.01)

and neutrophils (*P*<0.01) (Figure 2A). The highest dose of 50 µg SiO₂ NPs led to a significant increase of macrophages, lymphocytes, neutrophils, and eosinophils (*P*<0.001) in the BALF compared to sensitized SUP (Figure 2A). Similar effects were shown following SiO₂-PEG instillation (Figure S1A). Intratracheal instillation of 50 µg/mouse SiO₂ NPs in NS mice exerted a proinflammatory effect, leading to significantly increased neutrophils and lymphocytes (*P*<0.001) and to a slight increase of eosinophils (*P*<0.05) compared to non-sensitized SUP (Figure 2B). These results suggest that SiO₂ NPs exert both a strong proinflammatory and immunomodulatory effect in allergic airway inflammation.

Pulmonary macrophages take up SiO₂ NPs

To determine whether OVA sensitization affects pulmonary retention of NPs, mice were instilled with SiO₂-FITC or respective SUP prior to OVA challenge. Twenty-four hours later, a minor, but not significant, increase in the percentage of FITC⁺ cells was observed in the lung of S mice, compared to NS mice. FITC⁺ cells in the negative control and in S and NS mice instilled with SUP were not higher than background level (Figure 3A).

Flow cytometry of lung cell subtypes showed that macrophages displayed the highest percentage of FITC positivity. In detail, in the lungs of NS mice, 15.3% of alveolar macrophages and 8.2% of interstitial macrophages were positive for SiO₂-FITC NPs. In the lungs of S mice, a significantly increased percentage of alveolar macrophages (23.2%, *P*<0.01 versus NS mice) and a reduced percentage of interstitial macrophages (4.3%) were positive for SiO₂-FITC (Figure 3A). Pulmonary suspensions of negative controls displayed minimal FITC autofluorescence, comparable to the lungs instilled with FITC-SUP. A minimal percentage of T-cells and neutrophils were FITC positive (data not shown).

To evaluate whether SiO₂ NPs interacted only with the surface of macrophages or were internalized by these cells, confocal analysis was performed 24 hours after OVA challenge. Figure 3B shows representative lung sections of NS and S mice stained with the macrophage marker CD68 (red), and with DAPI for the nuclei (blue), suggesting that pulmonary macrophages take up SiO₂ NPs (green) in both NS and S mice. The 100× orthogonal sections of single cells show that NPs were present inside the macrophages.

Specific surface modifications of SiO₂ NPs significantly reduce their immunomodulatory activity

In order to evaluate the possible reduction of immunomodulatory effects on the allergic response by the




Figure 2 SiO₂ NPs exert proinflammatory and immunomodulatory activity.

Notes: Inflammatory cell infiltration in bronchoalveolar lavage fluid of sensitized (A) and non-sensitized (B) mice 5 days after intratracheal instillation with increasing concentrations of SiO₂ NPs (12.5–50 µg/mouse, (A) or 50 µg/mouse (B) SiO₂ NPs [pattern and red filled bar] or with respective SUP [white bars]). Data are displayed in boxplots (five to ten mice per bar). * $P \leq 0.05$; ** $P \leq 0.01$; *** $P \leq 0.001$ versus SUP (0 µg).

Abbreviations: NP, nanoparticle; ns, not significant; SiO₂, silicon dioxide; SUP, supernatant controls.

functionalization of SiO₂ NPs, we evaluated the BAL cell infiltration in NS and S mice after intratracheal instillation of SiO₂ NPs with PEG, amino, and phosphate surface modifications.

Similarly to SiO₂ plain NPs, SiO₂-PEG NPs exerted proinflammatory effects, with a significant increase of neutrophils and lymphocytes ($P < 0.001$) and a minor increase of eosinophils ($P < 0.05$), in NS mice. Moreover, functionalizing SiO₂ plain NPs with PEG did not impair the immunomodulatory effect of SiO₂ NPs in the BALF of S mice. Lymphocyte,

neutrophil, and eosinophil total cell numbers were still significantly increased in the BALF of S mice ($P < 0.001$) (Figure 4A).

In contrast, mice treated with SiO₂-P or SiO₂-NH₂ (Figure 4B and C) showed minimal proinflammatory effects in NS mice, which reached significance only for neutrophil and eosinophil counts in SiO₂-NH₂-treated mice ($P < 0.05$). In S mice, no significant effects on the BALF total cell count were shown compared to control, apart from a slight increase in the macrophage cell number after SiO₂-NH₂ instillation ($P < 0.05$).




Figure 3 Macrophage populations take up nanoparticles.

Notes: (A) Percentage of FITC⁺ cells in lung (a) and within AMs (b) and IMs (c) of NS and S mice 24 hours after 50 µg SiO₂-FITC and OVA challenge. Neg CTL: untreated mouse; SUP: instillation of SiO₂-FITC-supernatant. Mean ± standard deviation (n=5–7/group). **P≤0.01 versus SUP. (B) NS (a) and S (b) lungs 24 hours after SiO₂-FITC (green) and OVA challenge. Macrophages: red; nuclei: blue. Alexa 555: Alexa Fluor® 555 (1:7000; Signaling Technology, Inc., Danvers, MA USA). Left: 40× and right: 100× orthogonal sections of single cell (arrowed cells 1 and 2).

Abbreviations: AM, alveolar macrophage; DAPI, 4',6-diamidino-2-phenylindole; FITC, fluorescein isothiocyanate; IM, interstitial macrophage; Neg CTL, negative controls; NS, non-sensitized; OVA, ovalbumin; S, sensitized; SiO₂, silicon dioxide; SUP, supernatant controls.

SiO₂ and SiO₂-PEG increase pulmonary inflammation and goblet cell necrosis

To further elucidate the effects of SiO₂ NPs and their surface coatings, we evaluated pulmonary inflammation in the lung. Intratracheal instillation of the supernatants of all four NP preparations did not affect pulmonary histopathology (Figure 5A). In S mice, SiO₂ and SiO₂-PEG NPs increased the inflammatory lung infiltrate and, although in higher degree SiO₂-

PEG, mucus hypersecretion when compared to sensitized SUP mice (Figure 5A). In contrast, mice treated with SiO₂-P and SiO₂-NH₂ NPs showed minimal pulmonary inflammation when compared with sensitized SUP mice; these observations were in accordance with the effects observed in the BALF differential cell count in S mice. Irrespective of functionalization, SiO₂ NPs did not induce pulmonary pathological alterations in NS mice (data not shown).




Figure 4 Surface modifications of SiO₂ nanoparticles influence their proinflammatory and immunomodulatory effects in vivo.

Notes: Bronchoalveolar lavage fluid inflammatory infiltrate of NS and S mice 5 days after intratracheal instillation of 50 µg SiO₂-PEG (A), SiO₂-P (B), or SiO₂-NH₂ (C), or respective SUP. Data are displayed as boxplots (n=6–10/bar). *P≤0.05; **P≤0.01; ***P≤0.001.

Abbreviations: ns, not significant; NS, non-sensitized; PEG, polyethylene glycol; S, sensitized; SiO₂, silicon dioxide; SiO₂-NH₂, amino-coated SiO₂; SiO₂-P, phosphate-coated SiO₂; SiO₂-PEG, PEGylated SiO₂; SUP, supernatant controls.

To evaluate structural changes of the airways following SiO₂ NP instillation, SEM was performed. NS mice instilled with the four different particles and respective supernatants showed no significant ultrastructural modifications of the lung (data not shown), apart from sporadic hypotrophy of Clara cells present in all samples (Figure 5B). S mice instilled with the four different sensitized SUP developed the classical pulmonary alterations due to OVA sensitization and challenge, ie, metaplasia of Clara cells to mucus-secreting goblet cells (Figure 5B). Lungs of S mice instilled with SiO₂ and SiO₂-PEG NPs showed morphological alterations of goblet cells, including the perforation of cell membrane, indicating cell death (Figure 5B). In contrast, no additional effects due to particle

exposure were observed in S mice instilled with SiO₂-NH₂ or SiO₂-P (Figure 5B).

SiO₂ and SiO₂-PEG NPs, but not SiO₂-P and SiO₂-NH₂ NPs, induce moderate alterations in lung function

Lung function tests performed 24 hours after intratracheal instillation of SiO₂ NPs and OVA challenge in S mice showed different results for the four particles tested. SiO₂ plain did not have an effect on respiratory system resistance when compared to SUP; however, dynamic compliance was significantly decreased in S mice instilled with SiO₂ already, before methacholine challenge (P<0.05), maintaining the lower trend at 1.25 and 2.5 mg mL




Figure 5 SiO₂ and SiO₂-PEG NPs exacerbate goblet cell hyperplasia and induce necrosis.

Notes: Representative NS and S lungs 5 days after SUP (a and b, respectively) or 50 μg nanoparticles (c: S SiO₂; d: S SiO₂-PEG; e: S SiO₂-P; f: S SiO₂-NH₂) and ovalbumin challenge. **(A)** Periodic acid–Schiff staining: mucus hypersecretion (arrowheads) and inflammatory infiltrate (arrows). Bar: 100 μm. **(B)** Scanning electron microscopy of bronchiolar epithelium. (a) Clara cells (arrowheads); (b) goblet cells (arrowheads); (c and d) necrosis of goblet cells (insets, arrows). Bar: 5 μm (insets: 3 μm).

Abbreviations: NS, non-sensitized; PEG, polyethylene glycol; S, sensitized; SiO₂, silicon dioxide; SiO₂-NH₂, amino-coated SiO₂; SiO₂-P, phosphate-coated SiO₂; SiO₂-PEG, PEGylated SiO₂; SUP, supernatant controls.

methacholine concentrations ($P=0.05$ and $P=0.06$, respectively) (Figure 6A and B). The instillation of SiO₂-PEG in S mice led to an increase of resistance following the highest methacholine concentration, namely 40 mg/mL ($P<0.01$). Dynamic compliance was lower in SiO₂-PEG with respect to sensitized SUP, but this difference was not statistically significant (Figure 6A and B). On the contrary, SiO₂-P and SiO₂-NH₂ NPs had no effect on either lung resistance or compliance in S mice (Figure 6 A and B). The effect of the instillation of NPs in NS mice was minimal for all NPs tested (Figure 6).

SiO₂ and SiO₂-PEG NPs, but not SiO₂-P and SiO₂-NH₂ NPs, augment the Th2 proinflammatory milieu in the lungs of OVA-sensitized mice

To characterize the distinct effects of the different surface modification of SiO₂ NPs on immune responses, we performed PCR arrays on total RNA from lung homogenates.

A distinct increase of T helper type 2 (Th2) cytokines (interleukin [*IL*]-4/*IL*-13) was observed in S mice instilled with SiO₂-PEG and, to a lesser extent, by SiO₂, with no effect on Th1 (interferon-γ/tumor necrosis factor-α) cytokines (Figure 7). Different Th2 chemokines (*MCP-1/CCL2*, *MIP-1/CCL3*, *Tarc/CCL17*, *KC/CXCL1*) involved in recruitment of macrophages, eosinophils, lymphocytes, and neutrophils were exclusively upregulated by SiO₂-PEG and SiO₂, and downregulated or unregulated by SiO₂-NH₂ and SiO₂-P. Moreover, SiO₂ and SiO₂-PEG only increased markers of Th2 alternative activation of macrophage (*Arg1* and *Retnla*) and eosinophil (*Ear11*) activation, and had no effect on the markers for classical activation (*CXCL9*, *Nos2*) (Figure 7 and Figure S2). These effects of NPs were observed also for other chemotactic chemokines (*MIP-3/CCL20*, *MDC/CCL22*, *MIP-2/CXCL2*) as well as for asthma markers (*Retnlg*, *Chi314*). *Muc5ac* was only upregulated by SiO₂-PEG in S mice (Figure S2). The effects of SiO₂ and SiO₂-PEG NP instillation on selective Th2 gene expression levels were dose dependent, and significant upregulation of




Figure 6 SiO₂ and SiO₂-PEG nanoparticles induce moderate alterations in pulmonary function.

Notes: Lung resistance (**A**) and dynamic compliance (**B**) of NS and S mice exposed to increasing concentrations of methacholine (1.25–40 mg/mL) 24 hours after 50 µg SiO₂ plain (a), SiO₂-PEG (b), SiO₂-P (c), or SiO₂-NH₂ (d) (solid lines) or respective SUP (dashed lines) and ovalbumin challenge. Mean ± standard deviation (n=6–8/group). *P≤0.05; **P≤0.01 versus S SUP.

Abbreviations: NS, non-sensitized; PBS, phosphate-buffered saline; PEG, polyethylene glycol; S, sensitized; SiO₂, silicon dioxide; SiO₂-NH₂, amino-coated SiO₂; SiO₂-P, phosphate-coated SiO₂; SiO₂-PEG, PEGylated SiO₂; SUP, supernatant controls.

IL-13, *MIP-1/CCL3*, and *Arg1* was detected at the highest concentration (50 µg/mouse) (Figure S1B).

Discussion

In this study, we used a well-characterized, murine model of mild allergic airway inflammation²³ to investigate SiO₂

NP-mediated, surface-dependent proinflammatory and immunomodulatory effects in the lung. SiO₂ plain and SiO₂-PEG exerted potent Th2-immunomodulatory activities that involved alternatively activated alveolar macrophages that presumably phagocytized SiO₂ NPs. Moreover, functionalization of SiO₂ NPs with amino or phosphate groups conferred




Figure 7 SiO₂ nanoparticles differentially modulate inflammatory genes.

Notes: Representative gene regulations in NS and S lungs 5 days after intratracheal instillation with 50 µg SiO₂ plain (a), SiO₂-PEG (b), SiO₂-P (c), or SiO₂-NH₂ (d), or respective SUP and ovalbumin challenge. Mean ± standard deviation (n=4/group). *P≤0.05; **P≤0.01; ***P≤0.001 versus respective SUP.

Abbreviations: AAM, alternatively activated macrophages; EOS, eosinophils; IFN, interferon; IL, interleukin; NS, non-sensitized; PEG, polyethylene glycol; S, sensitized; SiO₂, silicon dioxide; SiO₂-NH₂, amino-coated SiO₂; SiO₂-P, phosphate-coated SiO₂; SiO₂-PEG, PEGylated SiO₂; SUP, supernatant controls; TNF, tumor necrosis factor; Th1, T helper type 1; Th2, T helper type 2.

a higher NP tolerance, with a minimal inflammatory milieu in OVA-sensitized mice.

Only a few studies have investigated the modulation of allergic disease by NPs.^{19,20,26} Our results showed significant immunomodulatory effects exerted by SiO₂ NPs in OVA-sensitized mice, with a significant increase in BALF inflammatory infiltrate, mainly comprising macrophages and eosinophils. Our data are in line with a previous study testing titanium dioxide and gold NPs in the elicitation phase of the allergic response.²⁶ We focused our read-out to day 5 after challenge. The reason for this choice was that this time point is representative for the elicitation phase of the allergic response in our mouse model, since it is characterized by strong recruitment of inflammatory cells into the lung.²⁷ Since the Th2-adjuvant effects of SiO₂ NPs applied during the sensitization phase have been previously demonstrated,^{19,20} we

now can assume that SiO₂ NPs also exacerbate the elicitation phase of allergic airway inflammation.

SiO₂ NPs showed a strong proinflammatory effect in NS mice, leading to neutrophil and lymphocyte recruitment after intratracheal instillation, which is in agreement with a previous work¹⁶ showing neutrophil recruitment in the early inflammation stages (up to 1 week after instillation) following intratracheal instillation of SiO₂ NPs. We also demonstrated an increase of eosinophils in NS mice exposed to SiO₂ NPs, a phenomenon also observed in rats after titanium NP exposure.¹⁰

The observation that, 24 hours after intratracheal instillation, the percentage of FITC⁺ cells in S mice was higher compared to NS mice, is in agreement with our earlier study.²⁸ NP uptake occurred mainly by pulmonary macrophages, as has been observed in healthy animals.²⁹ Confocal analysis

confirmed that NPs were located inside the phagocytes, and did not just interact with the cell surface. Moreover, we demonstrated that, in S mice, more alveolar macrophages phagocytize NPs compared to in NS mice and, consequently, particle translocation and retention in the interstitium by interstitial macrophages occurs to a higher extent in NS mice. Since these cells exert immunoregulatory properties, this could explain the short-term NP-induced inflammation in NS mice.¹⁶ The higher percentage of phagocytizing alveolar macrophages in S mice relates to our observation of SiO₂-dependent enhancement of expression of markers for alternatively activated macrophages, like arginase 1 and mannose receptors in NP-treated S mice. These cells show increased phagocytic activity³⁰ and aggravate allergic airway inflammation.³¹ The SiO₂-dependent increase in alternatively activated macrophages is of high clinical importance and could account, at least in part, for the differential localization of SiO₂ NPs in S and NS mice.

A few groups have investigated the effects of surface modifications of SiO₂ NPs in vitro and in vivo;^{21,22,32,33} however, to our knowledge, this is the first study that explores the effect of SiO₂ NP functionalization on their proinflammatory and immunomodulatory effects in allergic airway inflammation in vivo. Our data showed that coating NPs with short-chain PEG was not a successful strategy for modifying biological properties of SiO₂ NPs: in fact, both NS and S mice treated with SiO₂-PEG responded similarly in terms of inflammation to those treated with SiO₂ plain. Moreover, both SiO₂ plain and SiO₂-PEG enhanced mucus production and exerted cytotoxic effects. The ultrastructural damages are very likely mediated by SiO₂ NP-induced oxidative stress and apoptosis.³⁴ Possibly, the 500 Da PEG chain used to coat the NPs was not long enough to mask the surface of the NPs. Lipka et al suggested that the PEG chain length is essential for determining the biodistribution of NPs.¹¹ Moreover, both molecular weight and chain density of PEG influence mesoporous SiO₂ NP binding to serum proteins and phagocytosis from THP-1 macrophages, proposing 10 kDa as the minimal molecular weight to inhibit phagocytosis.³⁵ In light of these studies, our PEG chain might have been too small to mask the effect of SiO₂ NPs. In contrast, our results showed that coating with amino or phosphate groups successfully mitigated the toxic and immunomodulatory effects of SiO₂ NPs. These effects were observed at histopathological and functional levels, with less BALF cell infiltrate, absence of goblet cell necrosis, and normal lung function in both NS and S mice instilled with SiO₂-P and SiO₂-NH₂ versus SiO₂ plain or SiO₂-PEG. It has been demonstrated that coating of NPs can affect their interaction with biological fluids.³⁶ Proteins

bind to the surface of NPs and form a coating known as the “protein corona,” which can critically affect the interaction of the NPs with living systems. Speculation can be made as to whether our surface modifications of SiO₂ NPs affected the thermodynamic characteristics of lipid monolayers, including lung surfactant or cell membrane, as previously observed in 1,2-dipalmitoyl-sn-glycerol-3-phosphocholine-cholesterol Langmuir monolayers.^{37,38}

Different cellular uptake mechanisms may be involved when NPs are functionalized; in fact, it has been previously shown that SiO₂ NPs coated with amine or carboxyl groups display a different intracellular distribution compared to those that are uncoated.³⁹ In our model, both amino and phosphate functionalization failed to regulate markers of macrophage activation, suggesting a different compartmentalization of SiO₂-P and SiO₂-NH₂ within macrophages. Further studies should investigate these aspects.

The successful mitigation of effects of SiO₂ NPs coated with amino groups is in agreement with the results of Yamashita et al,³² which showed less fetotoxicity of amino-coated NPs when compared to plain NPs. To the best of our knowledge, this is the first study showing mitigation of side effects of SiO₂ NPs after phosphate coating.

Our findings on the expression of key genes of inflammation support the hypothesis that functionalization of NPs modify their biological activity in vivo. The immunomodulatory NPs (SiO₂ plain and SiO₂-PEG), but not the counterpart NPs (SiO₂-P and SiO₂-NH₂), increased the inflammatory milieu in the pulmonary tissue, with upregulation of Th2 cytokines, as well as sensitization-induced chemokines, the latter responsible for chemoattraction of granulocytes (*MIP-1/CCL3* and *MIP-2/CXCL2*, *KC/CXCL1*), lymphocytes (*Tarc/CCL17*, *MIP-3/CCL20*, and *MDC/CCL22*), and macrophages (*MCP-1/CCL2*).⁴⁰ These data support the concept of recruitment of inflammatory cells observed in the BALF of OVA-sensitized mice by immunomodulatory NPs only. Furthermore, our gene expression analysis showed that only SiO₂ plain and SiO₂-PEG NPs could lead to increased marker of eosinophils activation (*Ear11*)⁴¹ and alternative activation of macrophages, with no modulation of markers of classical activation of macrophages (*Arg1/Retnla* versus *Nos2/CXCL9*).³¹ The asthmatic profile present in the lungs of S mice was further confirmed by the upregulation of genes involved in allergic airway inflammation: Chitinase 3-like 4 (*Chi3l4*)⁴² was modulated in our experiments by both SiO₂ and SiO₂-PEG NPs in S mice. Among the resistin family, *FIZZ1* (*Retnla*) and *FIZZ2* (*Retnlb*), but not *FIZZ3* (*Retnlg*), mRNAs were

found to be upregulated in the lungs of S mice 6 hours after the OVA challenge;⁴³ in this study we still found modulation of *FIZZ1* and *FIZZ3* in S mice 5 days after OVA challenge, which was significantly enhanced by both SiO₂ and SiO₂-PEG NPs and downregulated by SiO₂-NH₂. The prolongation of *FIZZ1* and *FIZZ3* expression after challenge may have been due to our relatively long sensitization protocol.

In most of the genes evaluated in our array, SiO₂-PEG had a greater effect than SiO₂. A clear example regards the regulation of the mucin *Muc5ac* gene, responsible for mucus production,⁴⁴ which was selectively upregulated by SiO₂-PEG NPs but not by SiO₂ NPs. Since *IL-13* is a known modulator of *Muc5ac*,⁴⁵ the lower expression of *IL-13* mRNA upon SiO₂ compared to SiO₂-PEG could explain this finding.

Conclusion

Our observations indicate the need for selecting surface modifications of engineered NPs for commercial purposes, in order to improve their biological activity and reduce or abolish their side effects. Moreover, our results can inform and help in the choice of drug delivery strategy, as the selection of the surface modification of NPs is crucial for targeted application.

Acknowledgments

We thank Johanna Grosch, Benjamin Schnautz, Christine Weil, Katja Haslauer, and Alexandra Seisenberger for excellent technical assistance. We thank Claudia Traidl-Hoffmann, Tanja Seher, Antje Vennemann, and Sebastian Öder for fruitful discussions. We thank Thomas Kuhlbusch for the excellent coordination of the NanoGEM Project. This study was supported by the German Federal Ministry of Education BMBF, NanoGEM Project, FKZ 03X0105, and CK Care, Christine Kühne Center for Allergy Research and Education.

Disclosure

The authors report no conflicts of interest in this work.

References

- Arora S, Rajwade JM, Paknikar KM. Nanotoxicology and in vitro studies: the need of the hour. *Toxicol Appl Pharmacol*. 2012;258:151–165.
- Napierska D, Thomassen LC, Lison D, Martens JA, Hoet PH. The nano-silica hazard: another variable entity. *Part Fibre Toxicol*. 2010;7:39.
- Hansen SF, Michelson ES, Kamper A, Borling P, Stuer-Lauridsen F, Baun A. Categorization framework to aid exposure assessment of nanomaterials in consumer products. *Ecotoxicology*. 2008;17:438–447.
- Nel A, Xia T, Mädler L, Li N. Toxic potential of materials at the nanolevel. *Science*. 2006;311:622–627.
- Fröhlich E, Roblegg E. Models for oral uptake of nanoparticles in consumer products. *Toxicology*. 2012;291:10–17.
- Mortensen LJ, Oberdörster G, Pentland AP, Delouise LA. In vivo skin penetration of quantum dot nanoparticles in the murine model: the effect of UVR. *Nano Lett*. 2008;8:2779–2787.
- Yu KO, Grabinski CM, Schrand AM, et al. Toxicity of amorphous silica nanoparticles in mouse keratinocytes. *J Nanopart Res*. 2009;11:15–24.
- Kreyling WG, Semmler-Behnke M, Seitz J, et al. Size dependence of the translocation of inhaled iridium and carbon nanoparticle aggregates from the lung of rats to the blood and secondary target organs. *Inhal Toxicol*. 2009;21 Suppl 1:55–60.
- Ho M, Wu KY, Chein HM, Chen LC, Cheng TJ. Pulmonary toxicity of inhaled nanoscale and fine zinc oxide particles: mass and surface area as an exposure metric. *Inhal Toxicol*. 2011;23:947–956.
- Gustafsson Å, Lindstedt E, Elfsmark LS, Bucht A. Lung exposure of titanium dioxide nanoparticles induces innate immune activation and long-lasting lymphocyte response in the Dark Agouti rat. *J Immunotoxicol*. 2011;8:111–121.
- Lipka J, Semmler-Behnke M, Sperling RA, et al. Biodistribution of PEG-modified gold nanoparticles following intratracheal instillation and intravenous injection. *Biomaterials*. 2010;31:6574–6581.
- Berg JM, Romoser AA, Figueroa DE, Spencer West C, Sayes CM. Comparative cytological responses of lung epithelial and pleural mesothelial cells following in vitro exposure to nanoscale SiO₂. *Toxicol In Vitro*. 2013;27:24–33.
- Lin W, Huang YW, Zhou XD, Ma Y. In vitro toxicity of silica nanoparticles in human lung cancer cells. *Toxicol Appl Pharmacol*. 2006;217:252–259.
- Farcial LR, Uboldi C, Mehn D, et al. Mechanisms of toxicity induced by SiO₂ nanoparticles of in vitro human alveolar barrier: effects on cytokine production, oxidative stress induction, surfactant proteins A mRNA expression and nanoparticles uptake. *Nanotoxicology*. 2013;7(6):1095–1110.
- Park MDZ, Lynch I, Ramírez-García S, et al. In vitro evaluation of cytotoxic and inflammatory properties of silica nanoparticles of different sizes in murine RAW 264.7 macrophages. *J Nanopart Res*. 2011;13:6775–6787.
- Cho WS, Choi M, Han BS, et al. Inflammatory mediators induced by intratracheal instillation of ultrafine amorphous silica particles. *Toxicol Lett*. 2007;175:24–33.
- Klein CL, Wiench K, Wiemann M, Ma-Hock L, van Ravenzwaay B, Landsiedel R. Hazard identification of inhaled nanomaterials: making use of short-term inhalation studies. *Arch Toxicol*. 2012;86:1137–1151.
- Landsiedel R, Ma-Hock L, Kroll A, et al. Testing metal-oxide nanomaterials for human safety. *Adv Mater*. 2010;22:2601–2627.
- Han B, Guo J, Abrahaley T, et al. Adverse effect of nano-silicon dioxide on lung function of rats with or without ovalbumin immunization. *PLoS One*. 2011;6:e17236.
- Brandenberger C, Rowley NL, Jackson-Humbles DN, et al. Engineered silica nanoparticles act as adjuvants to enhance allergic airway disease in mice. *Part Fibre Toxicol*. 2013;10:26.
- Lankoff A, Arabski M, Wegierek-Ciuk A, et al. Effect of surface modification of silica nanoparticles on toxicity and cellular uptake by human peripheral blood lymphocytes in vitro. *Nanotoxicology*. 2013;7(3):235–250.
- Morishige T, Yoshioka Y, Inakura H, et al. The effect of surface modification of amorphous silica particles on NLRP3 inflammasome mediated IL-1beta production, ROS production and endosomal rupture. *Biomaterials*. 2010;31:6833–6842.
- Alessandrini F, Schulz H, Takenaka S, et al. Effects of ultrafine carbon particle inhalation on allergic inflammation of the lung. *J Allergy Clin Immunol*. 2006;117:824–830.
- Bedoret D, Wallemacq H, Marichal T, et al. Lung interstitial macrophages alter dendritic cell functions to prevent airway allergy in mice. *J Clin Invest*. 2009;119:3723–3738.
- Alessandrini F, Weichenmeier I, van Miert E, et al. Effects of ultra-fine particles-induced oxidative stress on Clara cells in allergic lung inflammation. *Part Fibre Toxicol*. 2010;7:11.

26. Hussain S, Vanoirbeek JA, Luyts K, et al. Lung exposure to nanoparticles modulates an asthmatic response in a mouse model. *Eur Respir J*. 2011;37:299–309.
27. Alessandrini F, Beck-Speier I, Krappmann D, et al. Role of oxidative stress in ultrafine particle-induced exacerbation of allergic lung inflammation. *Am J Respir Crit Care Med*. 2009;179:984–991.
28. Alessandrini F, Semmler-Behnke M, Jakob T, Schulz H, Behrendt H, Kreyling W. Total and regional deposition of ultrafine particles in a mouse model of allergic inflammation of the lung. *Inhal Toxicol*. 2008;20:585–593.
29. Morfeld P, Treumann S, Ma-Hock L, Bruch J, Landsiedel R. Deposition behavior of inhaled nanostructured TiO₂ in rats: fractions of particle diameter below 100 nm (nanoscale) and the slicing bias of transmission electron microscopy. *Inhal Toxicol*. 2012;24:939–951.
30. Alexis NE, Soukup J, Nierkens S, Becker S. Association between airway hyperreactivity and bronchial macrophage dysfunction in individuals with mild asthma. *Am J Physiol Lung Cell Mol Physiol*. 2001;280:L369–L375.
31. Moreira AP, Hogaboam CM. Macrophages in allergic asthma: fine-tuning their pro- and anti-inflammatory actions for disease resolution. *J Interferon Cytokine Res*. 2011;31:485–491.
32. Yamashita K, Yoshioka Y, Higashisaka K, et al. Silica and titanium dioxide nanoparticles cause pregnancy complications in mice. *Nat Nanotechnol*. 2011;6:321–328.
33. Yoshida T, Yoshioka Y, Matsuyama K, et al. Surface modification of amorphous nanosilica particles suppresses nanosilica-induced cytotoxicity, ROS generation, and DNA damage in various mammalian cells. *Biochem Biophys Res Commun*. 2012;427:748–752.
34. Ahamed M. Silica nanoparticles-induced cytotoxicity, oxidative stress and apoptosis in cultured A431 and A549 cells. *Hum Exp Toxicol*. 2013;32:186–195.
35. He Q, Zhang J, Shi J, et al. The effect of PEGylation of mesoporous silica nanoparticles on nonspecific binding of serum proteins and cellular responses. *Biomaterials*. 2010;31:1085–1092.
36. Monopoli MP, Aberg C, Salvati A, Dawson KA. Biomolecular coronas provide the biological identity of nanosized materials. *Nat Nanotechnol*. 2012;7(12):779–786.
37. Guzmán E, Ferrari M, Santini E, Liggieri L, Ravera F. Influence of silica nanoparticles on phase behavior and structural properties of DPPC – Palmitic acid Langmuir monolayers. *Colloids Surf A Physicochem Eng Asp*. 2012;413:280–287.
38. Guzmán E, Liggieri L, Santini E, Ferrari M, Ravera F. Mixed DPPC-cholesterol Langmuir monolayers in presence of hydrophilic silica nanoparticles. *Colloids Surf B Biointerfaces*. 2013;105:284–293.
39. Nabeshi H, Yoshikawa T, Arimori A, et al. Effect of surface properties of silica nanoparticles on their cytotoxicity and cellular distribution in murine macrophages. *Nanoscale Res Lett*. 2011;6:93.
40. Palmqvist C, Wardlaw AJ, Bradding P. Chemokines and their receptors as potential targets for the treatment of asthma. *Br J Pharmacol*. 2007;151:725–736.
41. Cormier SA, Yuan S, Crosby JR, et al. T(H)2-mediated pulmonary inflammation leads to the differential expression of ribonuclease genes by alveolar macrophages. *Am J Respir Cell Mol Biol*. 2002;27:678–687.
42. Zhang Q, Hitchins VM, Schrand AM, Hussain SM, Goering PL. Uptake of gold nanoparticles in murine macrophage cells without cytotoxicity or production of pro-inflammatory mediators. *Nanotoxicology*. 2011;5:284–295.
43. Stütz AM, Pickart LA, Trifilieff A, Baumruker T, Prieschl-Strassmayr E, Woisetschlager M. The Th2 cell cytokines IL-4 and IL-13 regulate found in inflammatory zone 1/resistin-like molecule alpha gene expression by a STAT6 and CCAAT/enhancer-binding protein-dependent mechanism. *J Immunol*. 2003;170:1789–1796.
44. Rose MC, Voynow JA. Respiratory tract mucin genes and mucin glycoproteins in health and disease. *Physiol Rev*. 2006;86:245–278.
45. Kanoh S, Tanabe T, Rubin BK. IL-13-induced MUC5AC production and goblet cell differentiation is steroid resistant in human airway cells. *Clin Exp Allergy*. 2011;41:1747–1756.

Supplementary materials

Nanoparticle characteristics

Covalent functionalization of nanoparticles (NPs) was obtained with three different low-molar-mass silanes:

- PEGylated silicon dioxide (silica [SiO₂]) NPs: polyethylene glycol of chain length molecular weight =500 g/mol.
- Amino-coated SiO₂ NPs: carry a positively charged amino end group on a flexible, but short C3 linker.
- Phosphate-coated SiO₂ NPs: carries a negatively charged PO₃ end group on the same C3 linker.

Successful functionalization was confirmed by transmission electron microscopy, which showed that NPs maintained a spherical structure with no aggregates,

agglomerates, or secondary nucleation. Size distribution in water as measured by analytical ultracentrifugation¹ was not significantly changed by the surface modification, with a peak diameter at 14 nm as shown in Figure S3A. Surface charge was measured by electrophoretic mobility with pH titration (Figure S3B), showing that the non-functionalized SiO₂ NPs (plain) retained their negative charge across the physiological pH range; phosphate-coated SiO₂ NPs was strongly negatively charged; and the charge was reduced in PEGylated SiO₂ and inverted to positive values in amino-coated SiO₂.

Moreover, surface chemistry was evaluated by secondary ion mass spectrometry measurements, showing the presence




Figure S1 Dose-dependent effects of SiO₂ and SiO₂-PEG nanoparticles.

Notes: (A) Bronchoalveolar lavage fluid cellular infiltration 5 days after intratracheal instillation with increasing concentrations of SiO₂-PEG and ovalbumin challenge. Data are displayed as boxplots (per bar =5 mice/group). (B) Pulmonary messenger RNA expression of sensitized lungs 5 days after intratracheal instillation with the same concentrations of SiO₂ plain (red) or SiO₂-PEG (green). Data are expressed as mean ± standard deviation (n=3/group). *P≤0.05; **P≤0.01; ***P≤0.001 versus SUP (0 μg). **Abbreviations:** IL, interleukin; ns, not significant; PEG, polyethylene glycol; SiO₂, silicon dioxide; SiO₂-NH₂, amino-coated SiO₂; SiO₂-P, phosphate-coated SiO₂; SiO₂-PEG, PEGylated SiO₂; SUP, supernatant controls.




Figure S2 SiO₂ nanoparticles differentially modulate inflammatory genes.

Notes: Representative gene regulations in NS and S lungs 5 days after intratracheal instillation with 50 μg SiO₂, SiO₂-PEG, SiO₂-P or SiO₂-NH₂, or respective SUP (white) and ovalbumin challenge. Mean ± standard deviation (n=4/group). *P≤0.05; **P≤0.01; ***P≤0.001 versus respective SUP.

Abbreviations: NS, non-sensitized; PEG, polyethylene glycol; S, sensitized; SiO₂, silicon dioxide; SiO₂-NH₂, amino-coated SiO₂; SiO₂-P, phosphate-coated SiO₂; SiO₂-PEG, PEGylated SiO₂; SUP, supernatant controls.

of free SiO_x groups and free functionalized groups, underlining a partial coverage of the NPs (Figure S3C shows an example of one of the NPs used in this study, phosphate-coated SiO₂ NPs). All NPs used in this study were tested for lipopolysaccharide concentration. This was <0.25 EU/mL NPs suspension, which resulted in <0.000055 EU/

intratracheal instillation for the highest NP concentrations as determined by limulus amoebocyte lysate assay (PyroGene rFC Endotoxin Detection system, product number 50-658U; Lonza Group Ltd, Basel, Switzerland).

Supernatants for each NP were obtained by hard sedimentation (24,000 rpm, 15 hours), with successful removal

Table S1 Murine primers for real-time polymerase chain reaction

Gene name	GenBank accession number	Primer forward 5'-3'	Primer reverse 5'-3'
<i>IL13</i>	NM_008355.3	CCTCTGACCCTTAAGGAGCTTAT	CGTTGCACAGGGGAGTCT
<i>MIP-1/Ccl3</i>	NM_011337.2	TTCTCTGTACCATGACACTCTGC	CGTGGAACTTCCGGCTGTAG
<i>Arg1</i>	NM_007482.3	GGAACCCAGAGAGAGCATGA	TTTTTCCAGCAGACCAGCTT
<i>ACTB</i>	NM_007393.3	TTCTTTGCAGCTCCTTCGTT	ATGGAGGGAATACAGCCC

Abbreviations: IL13, interleukin 13; MIP-1/CCL3, macrophage inflammatory protein 1/Chemokine (C-C motif) ligand 3; ARG1, arginase; ACTB, actin beta.




Figure S3 Physical-chemical characterization.

Notes: (A) Size distributions by analytical ultracentrifugation (AUC), plotted as differential distribution of concentration dC per diameter interval dD . The area under the curve is proportional to the mass within this diameter range; (B) Surface charge by electrophoretic mobility with pH titration; (C) Surface chemistry by SIMS (SiO_2 -P), showing in the top 1 nm layer of the particles both the added PO_3^- functionality and the intrinsic SiO_2 fragments of the intrinsic surface chemistry.

Abbreviations: SiO_2 , silicon dioxide; $\text{SiO}_2\text{-NH}_2$, amino-coated SiO_2 ; $\text{SiO}_2\text{-P}$, phosphate-coated SiO_2 ; $\text{SiO}_2\text{-PEG}$, PEGylated SiO_2 ; PEG, polyethylene glycol; dC/dD , distribution of concentration per diameter interval.




Figure S4 Size and agglomeration characterization of SiO_2 NPs by transmission electron microscopy.

Note: (A) Silicon dioxide (plain) and (B) silicon dioxide–fluorescein isothiocyanate nanoparticles.

Abbreviations: SiO_2 , silicon dioxide; NPs, nanoparticles.




Figure S5 Absorption and fluorescence spectra of silica nanoparticles.

of 95%–98% of NPs in solution. Table S2 summarizes the characteristics of the NPs.

SiO₂-FITC NPs were prepared as described previously.² The FITC was covalently bound to the SiO₂ core and an SiO₂ shell was then produced with the addition of tetraethoxysilane. This shell prevented the influence of the FITC dye on the charge of the NP cores. Successful synthesis of the fluorescent SiO₂ NPs (SiO₂-FITC) was confirmed by transmission electron microscopy, which showed spherical NPs with no aggregates, agglomerates, or secondary nucleation.

The size of the particles was determined by transmission electron microscopy and dynamic light scattering (Figure S4A: plain SiO₂; B: SiO₂-FITC). The surface charge was measured by electrophoretic mobility with pH titration. The results showed no significant difference between the plain SiO₂ and the fluorescent SiO₂ NPs. The fluorescent properties were confirmed by fluorescence spectroscopy (Figure S5). Figure S5 shows the absorption and the emission spectra of the SiO₂ particles excited at 460 nm.

Table S2

	SiO ₂ plain	SiO ₂ -PEG	SiO-NH ₂	SiO ₂ -P
Primary particle size (transmission electron microscopy)	15 nm	15 nm	15 nm	15 nm
Surface charge				
Isoelectric point	<1	4	7.2	<1
ζ-pot at pH 7	-38 mV	-26 mV	0 mV	-43 mV
Dispersability in water (analytical ultracentrifugation)				
D50	19 nm	21 nm	20 nm	20 nm
Av. Agglom. Number (AAN)	1	1	1	1
Dispersability in DMEM/FCS				
D50	420 nm	3,200 nm	1,350 nm	30 nm
AAN	28	213	90	2

Abbreviations: AAN, average agglomeration number; DMEM, Dulbecco's Modified Eagle's Medium; D50, median diameter; PEG, polyethylene glycol; FCS, fetal calf serum; SiO₂, silicon dioxide; SiO-NH₂, amino-coated SiO₂; SiO₂-P, phosphate-coated SiO₂; SiO₂-PEG, PEGylated SiO₂.

References

1. Wohlleben W. Validity range of centrifuges for the regulation of nanomaterials: from classification to as-tested coronas. *J Nanopart Res.* 2012;14:1300.
2. Herz E, Ow H, Bonner D, Burns A, Wiesner U. Dye structure–optical property correlations in near-infrared fluorescent core-shell silica nanoparticles. *J Mater Chem.* 2009;19:6341–6347.

International Journal of Nanomedicine**Dovepress****Publish your work in this journal**

The International Journal of Nanomedicine is an international, peer-reviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. This journal is indexed on PubMed Central, MedLine, CAS, SciSearch®, Current Contents®/Clinical Medicine,

Journal Citation Reports/Science Edition, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-nanomedicine-journal>