



Morbidity and Mortality Weekly Report (*MMWR*)

[MMWR](#)

Surveillance for Foodborne-Disease Outbreaks --- United States, 1998--2002

Surveillance Summaries

November 10, 2006 / 55(SS10);1-34

Michael Lynch, MD

John Painter, DVM

Rachel Woodruff, MPH

Christopher Braden, MD

Division of Foodborne, Bacterial, and Mycotic Diseases National Center for Zoonotic, Vector-Borne, and Enteric Diseases (proposed)

Corresponding author: Michael Lynch, MD, Division of Foodborne, Bacterial, and Mycotic Diseases, National Center for Zoonotic, Vector-Borne, and Enteric Diseases (proposed), 1600 Clifton Road, NE, MS A-38, Atlanta, GA 30333. Telephone: 404-639-2206; Fax: 404-639-2205; E-mail: mlynch1@cdc.gov.

Abstract

Problem/ Condition: Since 1973, CDC has maintained a collaborative surveillance program for collection and periodic reporting of data on the occurrence and causes of foodborne-disease outbreaks (FBDOs) in the United States.

Reporting Period Covered: 1998--2002.

Description of System: The Foodborne Disease Outbreak Surveillance System reviews data on FBDOs, defined as the occurrence of two or more cases of a similar illness resulting from the ingestion of a common food. State and local public health departments have primary responsibility for identifying and investigating FBDOs. State, local, and territorial health departments use a standard form to report these outbreaks to CDC. In 1998, CDC implemented enhanced surveillance for FBDOs by increasing communication with state, local, and territorial health departments and revising the outbreak report form. Since 2001, reports of FBDOs are submitted through a web application on the Internet called the electronic Foodborne Outbreak Reporting System (eFORS).

Results: During 1998--2002, a total of 6,647 outbreaks of foodborne disease were reported (1,314 in 1998, 1,343 in 1999, 1,417 in 2000, 1,243 in 2001, and 1,330 in 2002). These outbreaks caused a reported 128,370 persons to become ill. Among 2,167 (33%) outbreaks for which the etiology was determined, bacterial pathogens caused the largest percentage of outbreaks (55%) and the largest percentage of cases (55%). Among bacterial pathogens, *Salmonella* serotype Enteritidis accounted for the largest number of outbreaks and outbreak-related cases; *Listeria monocytogenes* accounted for the majority of deaths of any pathogen. Viral pathogens, predominantly norovirus, caused 33% of outbreaks and 41% of cases; the proportion of outbreaks attributed to viral agents increased from 16% in 1998 to 42% in 2002. Chemical agents caused 10% of outbreaks and 2% of cases, and parasites caused 1% of outbreaks and 1% of cases.

Interpretation: Following implementation of measures to enhance outbreak surveillance, the annual number of FBDOs reported to CDC increased during this period compared with previous years. Viral pathogens accounted for an increased proportion of outbreaks each year during this reporting period and a higher proportion of outbreaks of known etiology

during this reporting period than preceding reporting periods, probably reflecting the increased availability of improved viral diagnostic tests. *S. Enteritidis* continued to be a major cause of illness and *L. monocytogenes* was a major cause of death. In addition, multistate outbreaks caused by contaminated produce and outbreaks caused by *Escherichia coli* O157:H7 remained prominent.

Public Health Actions: Methods to detect FBDOs are improving, and several changes to improve the ease and timeliness of reporting FBDO data have been implemented (e.g., a revised form to simplify FBDO reporting by state health departments and improved electronic reporting methods). State and local health departments continue to investigate and report FBDOs as part of efforts to better understand and define the epidemiology of foodborne disease in the United States. At the regional and national levels, surveillance data provide an indication of the etiologic agents, vehicles of transmission, and contributing factors associated with FBDOs and help direct public health actions to reduce illness and death caused by FBDOs.

Introduction

The reporting of foodborne and waterborne diseases in the United States began approximately 80 years ago when state and territorial health officers, concerned about the high morbidity and mortality caused by typhoid fever and infantile diarrhea, recommended that cases of "enteric fever" be investigated and reported. The purpose of investigating and reporting these cases was to obtain information about the role of food, milk, and water in outbreaks of intestinal illness as the basis for public health action. Beginning in 1925, the U.S. Public Health Service (PHS) published summaries of outbreaks of gastrointestinal illness attributed to milk (1). In 1938, PHS added summaries of outbreaks caused by all foods. These early surveillance efforts led to the enactment of important public health measures (e.g., the Pasteurized Milk Ordinance) that resulted in decreased incidence of enteric diseases, particularly those transmitted by milk and water (2).

During 1951--1960, the National Office of Vital Statistics reviewed reports of outbreaks of foodborne illness and published annual summaries in *Public Health Reports*. In 1961, CDC assumed responsibility for publishing reports about foodborne illness. During 1961--1965, CDC stopped publishing annual reviews but reported pertinent statistics and detailed individual investigations in *MMWR*.

The current system of surveillance for outbreaks of foodborne and waterborne diseases began in 1966, when reports of enteric disease outbreaks attributed to microbial or chemical contamination of food or water were incorporated into an annual summary. Since 1966, the quality of investigative reports has improved greatly, with more active participation by state and federal epidemiologists in outbreak investigations. Outbreaks of waterborne diseases and foodborne diseases have been reported in separate annual summaries since 1978 because of increased interest and activity in surveillance for waterborne diseases. Previous summaries of data reported to the Foodborne Disease Outbreak Surveillance System were published for 1983--1987 (3), 1988--1992 (4), and 1993--1997 (5). Outbreak surveillance has served three purposes:

- **Disease prevention and control.** The investigation of foodborne disease outbreaks leads to prevention and control measures in the food industry. Public health officials identify critical control points in the path from farm to table that can be monitored to reduce contamination by foodborne pathogens. Changes at all levels of food production (e.g., farm, slaughterhouse, and production plant) have contributed to less contamination in the food supply. Summarizing these investigations illustrates the burden of the outbreaks and the efforts needed to control them.
- **Knowledge of disease causation.** Outbreak investigations are a critical means of identifying new and emerging pathogens and maintaining awareness about ongoing problems. However, the pathogen is not identified in many outbreaks because of delayed or incomplete laboratory investigation, inadequate laboratory capacity, or inability to recognize a pathogen as a cause of foodborne disease. Prompt and thorough investigations of foodborne outbreaks aid in the timely identification of etiologic agents and lead to appropriate prevention and control measures. Summarizing the results provides an index of the relative importance and impact of specific pathogens.
- **Administrative guidance.** By analyzing several years of data on foodborne disease outbreaks, public health authorities can monitor trends over time in the prevalence of outbreaks caused by specific etiologic agents, the food that is the vehicle for the agent, and common errors in food handling. This information provides the basis for regulatory and other changes to improve food safety. Analysis of specific subsets of outbreaks can illustrate the challenges associated with specific pathogens, food vehicles, and settings and has helped define linkages between specific pathogens and foods.

This report summarizes epidemiologic data on FBDOs reported to CDC during 1998--2002.

Methods

Sources of Data for the Foodborne Disease Outbreak Surveillance System

Agencies use a standard form (CDC form 52.13, Investigation of a Foodborne Outbreak) to report FBDOs to CDC. In 1998, CDC increased communication with state, local, and territorial health departments to enhance surveillance for FBDOs, including formal confirmation procedures to finalize reports from each state each year. This led to a substantial increase in the number of reports, resulting in a surveillance discontinuity during 1997--1998. A revised form became effective in 1999. The revised form expanded the range of food items, places, and contributing factors that could be

reported. In 2001, state, local, and territorial health departments began submitting reports through a web-based version of this form. This web-based outbreak surveillance system is called the Electronic Foodborne Outbreak Reporting System (eFORS). This report summarizes data collected with both the paper and web-based forms (Appendix A). The majority of reports are submitted by state, local, and territorial health departments; however, they also can be submitted by federal agencies and other sources. Reporting officials use published criteria to determine whether a specific etiologic agent has been confirmed for an outbreak (Appendix B) and submit reasons that reported food vehicles were implicated. Implicated food vehicles for all reasons are included in this report.

Definition of Terms

An FBDO is defined as the occurrence of two or more cases of a similar illness resulting from the ingestion of a food in common. Laboratory or clinical guidelines for confirming an etiology of a FBDO outbreak vary for bacterial, chemical, parasitic, and viral agents (Appendix B). An outbreak in which more than one etiologic agent was confirmed is categorized as attributable to multiple etiologies. Food vehicles identified in outbreak investigations that can be classified into a single commodity are classified into one of 12 major food commodity categories. Some reported food vehicles cannot be categorized in a single commodity category and are listed as unclassifiable. Outbreaks in which more than one implicated food is reported or the implicated food contains ingredients from multiple commodities are classified as attributable to complex food vehicles.

Exclusions from and Limitations of the Surveillance System

The findings in this report are subject to at least four limitations. First, several types of outbreaks are excluded from the Foodborne Disease Outbreak Surveillance System, such as outbreaks that occur on cruise ships (these are summarized and published periodically in scientific publications) (6); outbreaks in which the food was eaten outside the United States, even if the illness occurred within the United States; and outbreaks that are traced to water intended for drinking (these are reported to the Waterborne Disease Outbreak Reporting System). In addition, FBDOs are excluded from the surveillance system if the route of transmission from the contaminated food to the infected persons is indirect. For example, in 1988, chitterlings (pig intestines) were the ultimate source of a cluster of *Yersinia enterocolitica* infections among several infants; however, this outbreak was not included because the infants did not eat the chitterlings (7). Similarly, outbreaks that occur as result of direct contact with animals are excluded.

Second, for many reports, information on certain aspects of the outbreak, such as the etiology, the implicated food vehicle, or the factors that might have contributed to the outbreak, is missing or incomplete. The category of "unknown etiology" is broad. Outbreaks with some etiologic information might not meet guidelines for confirmation and are presented in this report as "unknown etiology." Clinical and descriptive epidemiologic information that suggests etiologic categories for outbreaks of unknown etiology have not been used in this report (8).

Third, food vehicles are reported by investigating agencies as individual food items in varying levels of details (e.g., milk, 2% milk, pasteurized 2% milk). A particular reported food item with multiple ingredients could be classified under several food commodity categories; however, in this surveillance summary, the reported food item for each outbreak is classified under only one food commodity category. Food items that cannot be classified under one food commodity category are counted as unclassifiable. As a result, the reported number of outbreaks attributed to one food vehicle category might not include all outbreaks attributable to a particular food ingredient in that food.

Finally, no standard criteria exist for classifying a death as being FBDO-related. This determination is made by the reporting agency.

How Data Are Presented

In this report, 1998--2002 data on foodborne-disease outbreaks are presented as follows:

- Reported outbreak reports, by years, 1993--2002 (Figure 1).
- Outbreaks, by state, for each of the 5 years (Figures 2--6).
- Outbreaks, cases, and deaths, by etiology, for the 5-year period combined (Table 1).
- Outbreaks, cases, and deaths, by etiology, for each of the 5 years (Tables 2--6).
- Outbreaks, by etiology and month of occurrence, for the 5-year period combined (Table 7).
- Outbreaks, by etiology and place where food was eaten, for the 5-year period combined (Table 8).
- Outbreaks, cases, and deaths, by vehicle of transmission, for each of the 5 years (Tables 9--13).
- Outbreaks, by etiology and vehicle of transmission, for each of the 5 years (Tables 14--18).
- Outbreaks, by etiology and contributing factors, for the 5-year period combined (Table 19).

Results

During 1998--2002, the annual number of reported outbreaks ranged from 1,243 to 1,417 (Tables 2--6). The average

annual number of outbreaks reported during this period (1,329) was substantially greater than the average annual number of outbreaks reported during 1993--1997 (550) ([Figure 1](#)). The average number of cases per outbreak during 1998--2002 (19) was lower than the average number of cases per outbreak during 1993--1997 (31). During 1998--2002, a total of 2,167 (33%) of the 6,647 outbreaks reported to CDC had a known etiology; these outbreaks accounted for 68,981 (54%) of 128,370 illnesses ([Table 1](#)). Of the 2,167 outbreaks with a known etiology, 55% (55% of illnesses) were caused by bacterial pathogens, 33% (41% of illnesses) by viruses, 10% (2% of illnesses) by chemical agents, and 1% (1% of illnesses) by parasites. The proportion of outbreaks with known etiology attributable to viruses increased from 16% in 1998 to 42% in 2002. In the majority (67%) of outbreaks, the etiology was not determined. However, the proportion of outbreaks for which an etiology was determined increased during the reporting period, from 28% in 1998 to 37% in 2002.

Local investigators might report factors they believe contributed to the outbreak. These factors are grouped into those that investigators believed led to contamination of the food, those that allowed proliferation of the pathogen in the food, and those that contributed to survival of the pathogen in the food. During 1998--2002, at least one contributing factor was reported in 3,072 (46%) outbreaks. The most commonly reported contamination factor that contributed to FBDOS was "bare-handed contact by handler/worker/preparer" ([Table 19](#)). For outbreaks caused by bacterial pathogens "raw product/ingredient contaminated by pathogens from animal or environment" was the most commonly reported contamination factor. The most commonly reported proliferation factor was "allowing foods to remain at room or warm outdoor temperature for several hours"; the most common survivability factor was "insufficient time and/or temperature during initial cooking/heat processing."

In the majority of foodborne outbreaks during this period, food was eaten outside the home ([Table 8](#)). Restaurants were the most commonly reported place where food was eaten. Many outbreaks caused by *Salmonella* or norovirus occurred at a school or nursing home. In outbreaks caused by ciguatoxin and *L. monocytogenes*, food was more commonly reported to have been eaten at a private home.

During this period, notable outbreaks were reported that were caused by ground beef contaminated with *E. coli* O157:H7 ([9](#)) and fresh produce contaminated with *Salmonella*, *E. coli* O157:H7, *Cyclospora cayetanensis*, or hepatitis A (Tables 14--18). Multidrug-resistant strains of *Salmonella* caused outbreaks linked to unpasteurized milk and ground beef. A large multistate outbreak of listeriosis caused by contaminated deli meat led to one of the largest food recalls in the United States ([10](#)). Scombrototoxin (fish-derived histaminic agent) caused the majority of outbreaks attributable to a chemical etiology. The majority of these outbreaks was associated with tuna, although several were associated with nonscombroid fish, including 10 outbreaks associated with escolar. Unexpected vehicles of transmission (e.g., dry cereal [[11](#)], parsley [[12](#)], and mangoes [[13](#)]) also were reported.

During 1998--2002, norovirus caused 657 (30%) of the 2,167 FBDOS with a known etiology and 39% of all outbreak-related cases in these outbreaks. *S. Enteritidis*, the most frequently reported bacterial cause of FBDOS, caused 204 outbreaks, accounting for 9% of outbreaks for which an etiology was determined. Eggs caused more *S. Enteritidis* outbreaks than any other food vehicle. *L. monocytogenes* resulted in 38 outbreak-related deaths among 256 cases, more deaths, and a higher case-fatality rate (15%) than any other pathogen.

Discussion

Foodborne-Disease Outbreaks, 1998--2002

The annual number of FBDOS reported to CDC increased during this period compared with previous years, following implementation of measures to enhance outbreak surveillance ([3--5](#)). Certain observations suggest that the increase in outbreak reports probably represents the effect of enhanced surveillance rather than a true increase in the occurrence of FBDOS. First, after a marked increase during 1997--1998 with implementation of enhanced surveillance, the number of reported outbreaks remained within a relatively narrow range. Second, the number of cases of foodborne infections identified through routine surveillance, of which outbreak cases are a part, decreased or remained stable during this period ([14](#)). Finally, the average size of reported outbreaks during 1998--2002 was smaller than the average size of outbreaks during 1993--1997, indicating that a substantial portion of the increase in reported outbreaks might be caused by smaller outbreaks that were not reported in previous years. Because of this increased reporting, comparisons of the number of reported FBDOS attributable to a specific etiology or vehicle of transmission between this period and previous reporting periods are difficult to make. Comparisons of the proportion of FBDOS related to specific causes are less likely to be influenced by the effect of enhanced surveillance but should be made with caution.

As in previous years, bacterial pathogens caused the majority of outbreaks and infections among outbreaks with a known etiology ([3--5](#)). Viral pathogens accounted for a much greater proportion of outbreaks and infections than in previous years, probably because of the increased availability of methods to diagnose viral agents. Although 67% of reported FBDOS during 1998--2002 were of unknown etiology, the proportion of outbreaks of unknown etiology decreased during 1998--2002. Much of this decrease is attributed to increased norovirus diagnostic capacity in state health department laboratories ([15](#)) and improved strategies to obtain diagnostic specimens ([16](#)). With continued improvements in epidemiologic and laboratory investigations, the proportion of outbreaks of unknown etiology might decrease further.

Of FBDOS with a known etiology, multistate outbreaks caused by contaminated produce and outbreaks caused by *E. coli*

O157:H7 remained prominent. Investigation of several multistate outbreaks attributed to *L. monocytogenes*, detected by linking information from molecular subtyping of isolates from several states, led to recalls of implicated products (10, 17, 18). Although *S. Enteritidis* continued to be a major cause of illness and death, it caused a much smaller proportion of outbreaks for which an etiology was known than in the past. The decrease in outbreaks attributed to *S. Enteritidis* parallels the decrease in *S. Enteritidis* infections reported to the National *Salmonella* Surveillance System and might reflect the role of Egg Quality Assurance Programs and other public health interventions in reducing the incidence of *S. Enteritidis* infection (19). Persons can decrease their risk for egg-associated infections caused by *S. Enteritidis* by not eating raw or undercooked eggs. Nursing homes, hospitals, and commercial kitchens should use pasteurized egg products for all recipes requiring pooled or lightly cooked eggs (20).

Interpretation of Data from the Foodborne Disease Outbreak Surveillance System

Foodborne diseases cause an estimated 76 million illnesses and 5,000 deaths in the United States each year (21). Although foodborne diseases are common, only a fraction of these illnesses are routinely reported to CDC because a complex chain of events must occur before a foodborne infection is reported; a break at any point in the chain will result in a case not being reported. In addition, the majority of reported foodborne illnesses are sporadic; only a small number are identified as being part of an outbreak and reported through the Foodborne Disease Outbreak Surveillance System. For example, *Salmonella* infection causes an estimated 1.4 million foodborne illnesses annually (22). However, during 1998--2002, a total of 164,044 *Salmonella* infections (approximately 32,000 annually) were reported through the National *Salmonella* Surveillance System (23--27), which is a passive, public health laboratory-based system. During the same period, 585 recognized outbreaks of *Salmonella* infection resulting in 16,821 illnesses were reported through the Foodborne Disease Outbreak Surveillance System, not all of which were necessarily culture-confirmed. Therefore, the system represents only a fraction of the burden of foodborne disease.

The number of outbreaks summarized in this report represents a small proportion of the outbreaks that actually occurred during the surveillance period. Some outbreaks are never recognized, and those that are recognized frequently go unreported. The likelihood that public health authorities are alerted about an outbreak depends on many factors, including its size and the severity of illnesses; consumer and physician awareness, interest, and motivation to report the incident; and the resources and disease surveillance activities of state and local public health and environmental agencies. Outbreaks that are most likely to be brought to the attention of public health authorities include those that are large, interstate, or restaurant-associated or that can cause serious illness, hospitalization, or death. The degree of underreporting might vary by etiology; therefore, this report provides limited information about the absolute or relative incidence of foodborne-disease outbreaks related to specific causes. For example, foodborne diseases characterized by short incubation periods (e.g., those caused by a chemical agent or staphylococcal enterotoxin) are more likely to be recognized as common source FBDOs than are diseases with longer incubation periods (e.g., hepatitis A). Outbreaks involving less commonly identified pathogens (e.g., *Bacillus cereus*, enterotoxigenic *E. coli*, or *Giardia intestinalis*) are less likely to have a confirmed etiology because these organisms are not always considered in clinical, epidemiologic, and laboratory investigations of FBDOs.

The objective of this report is to present simple analyses of the data on outbreaks of foodborne disease reported during 1998--2002. These data will continue to be analyzed in detail, along with other relevant data, to answer specific questions of public health importance, and findings will be published in the scientific literature. Specifically, a more detailed analysis of outbreak data to estimate the attribution of illness to specific food commodities would take into account the burden of illness attributed to specific etiologies and the attributable portion of those illnesses caused by particular food commodities. The simple frequencies of outbreaks caused by certain food commodities presented here do not, by themselves, provide a good measure of the burden of illness associated with one food commodity compared with another.

Future Directions

Methods to detect FBDOs continue to improve. For example, two tools that have enhanced detection of FBDOs are the Statistical Outbreak Detection Algorithm (SODA) and the National Molecular Subtyping Network for Foodborne Disease Surveillance (PulseNet). SODA applies a statistical algorithm to data reported through CDC's National *Salmonella*, *Shigella*, and *E. coli* Surveillance Systems to identify substantial increases over a historical baseline for any given serotype (28). This technology can be used to help identify clusters or outbreaks. PulseNet is a national network of public health laboratories that perform pulsed-field gel electrophoresis (PFGE) analysis on bacteria that might be foodborne (29). PulseNet was initiated in four states in 1996 and reached full participation of all 50 states and several large cities by 2001. This network permits rapid comparison of PFGE patterns through an electronic database at CDC; closely related PFGE patterns suggest a common source. PulseNet has helped in the detection and investigation of outbreaks, particularly those that involve multiple states. An assessment of the impact of introducing PulseNet PFGE subtyping in one state indicated that it increased the number of detected outbreaks of *E. coli* O157:H7 by 40% (30).

Several changes have improved the ease and timeliness of reporting. In October 1999, CDC issued a revised FBDO reporting form to simplify reporting by state health departments. In addition, eFORS was implemented in 2001 to help improve the timeliness of foodborne disease outbreak reporting. Upcoming versions of eFORS will include an automated search algorithm for more ready access to foodborne outbreak surveillance data. An annual listing of foodborne disease outbreaks reported to CDC is available at http://www.cdc.gov/foodborneoutbreaks/outbreak_data.htm.

The investigation and reporting of FBDOs by state and local health departments are important steps in efforts to better understand and define the epidemiology of foodborne disease in the United States. At the regional and national levels, surveillance data provide an indication of the etiologic agents, vehicles of transmission, and contributing factors associated with FBDOs and help direct public health actions.

Acknowledgments

The authors would like to thank all State and Territorial Epidemiologists and, in particular, members of the Foodborne Disease Outbreak reporting network, who contributed reports of foodborne outbreak investigations included in this summary.

References

1. Public Health Service. Annual report of the Surgeon General on the Public Health Service of the United States for the fiscal year 1924--1925. Washington, DC: Government Printing Office; 1925:34--5.
2. Potter ME, Kaufmann AF, Blake PA, Feldman RA. Unpasteurized milk: the hazards of a health fetish. *JAMA* 1984;252:2048--52.
3. [CDC. Foodborne-disease outbreaks, 5-year summary, 1983--1987. In: Surveillance Summaries. MMWR 1990;39\(No. SS-1\).](#)
4. [CDC. Surveillance for foodborne-disease outbreaks---United States, 1988--1992. In: Surveillance Summaries. MMWR 1996;45\(No. SS-5\).](#)
5. [CDC. Surveillance for foodborne-disease outbreaks---United States, 1993--1997. In: Surveillance Summaries. MMWR 2000;49\(No. SS-1\).](#)
6. Koo D, Maloney K, Tauxe R. Epidemiology of diarrheal disease outbreaks on cruise ships, 1986 through 1993. *JAMA* 1996;275:545--7.
7. Lee LA, Taylor J, Carter GP, et al. *Yersinia enterocolitica* O:3: an emerging cause of pediatric gastroenteritis in the United States. *J Infect Dis* 1991;163:660--3.
8. Hall J, Goulding J, Bean N, et al. Epidemiologic profiling: evaluating foodborne outbreaks for which no pathogen was isolated by routine laboratory testing: United States, 1982--9. *Epi and Infect* 2001;127:381--7.
9. [CDC. Multistate outbreak of *Escherichia coli* O157:H7 infections associated with eating ground beef---United States, June--July, 2002. MMWR 2002;51:637--9.](#)
10. [CDC. Outbreak of listeriosis---Northeastern United States, 2002. MMWR 2002;51:950--1.](#)
11. [CDC. Multistate outbreak of *Salmonella* Serotype Agona infections linked to toasted oats cereal---United States, April--May 1998. MMWR 1998;47:462--4.](#)
12. [CDC. Outbreaks of *Shigella sonnei* infection associated with eating fresh parsley---United States and Canada, July--August, 1998. MMWR 1999;48:285--9.](#)
13. Sivapalasingham S, Barret E, Kimura A, et al. A multistate outbreak of *Salmonella enterica* Serotype Newport infection linked to mango consumption: impact of water-dip disinfestation technology. *Clin Infect Dis* 2003;37:1585--90.
14. [CDC. Summary of notifiable diseases, United States, 2002. MMWR 2002;51:1--84.](#)
15. Widdowson M, Sulka A, Bulens S, et al. Norovirus and foodborne disease, United States, 1991--2000. *Emerg Infect Dis* 2005;11:95--102.
16. Jones T, Bulens S, Gettner S, et al. Use of stool collection kits delivered to patients can improve confirmation of etiology in foodborne disease outbreaks. *Clin Infect Dis* 2004;39:1454--9.
17. [CDC. Multistate outbreak of listeriosis---United States, 1998. MMWR 1998; 47:1085--6.](#)
18. [CDC. Multistate outbreak of listeriosis, United States, 2000. MMWR 2000;49:1129--30.](#)
19. Mumma G, Griffin P, Meltzer M, Braden C, Tauxe RV. Egg quality assurance programs and egg-associated *Salmonella* Enteritidis infections, United States. *Emerg Infect Dis* 2004;10:279--82.
20. Levine WC, Smart JF, Archer DL, Bean NH, Tauxe RV. Foodborne disease outbreaks in nursing homes, 1975 through 1987. *JAMA* 1991;266:2105--9.
21. Mead PS, Slutsker L, Dietz V, et al. Food-related illness and death in the United States. *Emerg Infect Dis* 1999;5:607--25.
22. Voetsch AC, Van Gilder TJ, AnguloFJ, et al. FoodNet estimate of the burden of illness caused by nontyphoidal *Salmonella* infections in the United States. *Clin Infect Dis* 2004;38:127--34.
23. CDC. *Salmonella* surveillance: annual tabulation summary, 1998. Atlanta, GA: US Department of Health and Human Services, Public Health Service, CDC; 1994.

24. CDC. *Salmonella* surveillance: annual tabulation summary, 1999. Atlanta, GA: US Department of Health and Human Services, Public Health Service, CDC; 1995.
25. CDC. *Salmonella* surveillance: annual tabulation summary, 2000. Atlanta, GA: US Department of Health and Human Services, Public Health Service, CDC; 1996.
26. CDC. *Salmonella* surveillance: annual tabulation summary, 2001. Atlanta, GA: US Department of Health and Human Services, Public Health Service, CDC; 1997.
27. CDC. *Salmonella* surveillance: annual tabulation summary, 2002. Atlanta, GA: US Department of Health and Human Services, Public Health Service, CDC; 1998.
28. Hutwagner LC, Maloney EK, Bean NH, Slutsker L, Martin SM. Using laboratory-based surveillance data for prevention: an algorithm for detecting *Salmonella* outbreaks. *Emerg Infect Dis* 1997;3:395--400.
29. Swaminathan B, Barrett TJ, Hunter SB, Tauxe RV. PulseNet: the molecular subtyping network for foodborne disease surveillance---United States. *Emerg Infect Dis* 2001;7:382--9.
30. Bender J, Hedberg C, Besser J, et al. Surveillance for *Escherichia coli* O157:H7 infections in Minnesota by molecular subtyping. *N Engl J Med* 1997;337:388--94.


Table 1**TABLE 1. Number of reported foodborne-disease outbreaks, cases, and deaths, by etiology — United States, 1998-2002**

Etiology	Outbreaks		Cases		Deaths	
	No.	(%)	No.	(%)	No.	(%)
Bacterial						
<i>Bacillus cereus</i>	37	(0.6)	571	(0.4)	0	(0.0)
<i>Brucella</i>	1	(0.0)	4	(0.0)	0	(0.0)
<i>Campylobacter</i>	61	(0.9)	1,440	(1.1)	0	(0.0)
<i>Clostridium botulinum</i>	12	(0.2)	52	(0.0)	1	(1.1)
<i>Clostridium perfringens</i>	190	(2.0)	6,724	(5.2)	4	(4.5)
<i>Escherichia coli</i> *	140	(2.1)	4,854	(3.8)	4	(4.5)
<i>Listeria monocytogenes</i>	11	(0.2)	256	(0.2)	38	(43.2)
<i>Salmonella</i>	585	(8.8)	16,821	(13.1)	20	(22.7)
<i>Shigella</i>	67	(1.0)	3,677	(2.9)	1	(1.1)
<i>Staphylococcus aureus</i>	101	(1.5)	2,766	(2.2)	2	(2.3)
<i>Streptococcus</i>	1	(0.0)	4	(0.0)	0	(0.0)
<i>Vibrio cholerae</i> †	3	(0.0)	12	(0.0)	0	(0.0)
<i>Vibrio parahaemolyticus</i>	25	(0.4)	613	(0.5)	0	(0.0)
<i>Vibrio</i> , other	1	(0.0)	2	(0.0)	0	(0.0)
<i>Yersinia enterocolitica</i>	8	(0.1)	87	(0.1)	0	(0.0)
Other bacterial	1	(0.0)	4	(0.0)	0	(0.0)
Total bacterial	1,184	(17.8)	37,887	(29.5)	70	(79.5)
Chemical						
Ciguatoxin	84	(1.3)	315	(0.2)	1	(1.1)
Heavy metals	2	(0.0)	23	(0.0)	0	(0.0)
Mushroom toxin	2	(0.0)	6	(0.0)	0	(0.0)
Scombrototoxin	118	(1.8)	463	(0.4)	0	(0.0)
Shellfish toxin	5	(0.1)	36	(0.0)	0	(0.0)
Other chemical	10	(0.2)	297	(0.2)	0	(0.0)
Total chemical	221	(3.3)	1,140	(0.9)	1	(1.1)
Parasitic						
<i>Anisakis</i>	1	(0.0)	14	(0.0)	0	(0.0)
<i>Cryptosporidium parvum</i>	4	(0.1)	139	(0.1)	0	(0.0)
<i>Cyclospora cayentanensis</i>	9	(0.1)	325	(0.3)	0	(0.0)
<i>Giardia intestinalis</i>	3	(0.0)	119	(0.1)	0	(0.0)
<i>Trichinella spiralis</i>	6	(0.1)	33	(0.0)	0	(0.0)
Total parasitic	23	(0.3)	630	(0.5)	0	(0.0)
Viral						
Astrovirus	1	(0.0)	14	(0.0)	0	(0.0)
Hepatitis A	50	(0.8)	981	(0.8)	4	(4.5)
Norovirus	657	(9.9)	27,171	(21.2)	1	(1.1)
Rotavirus	1	(0.0)	108	(0.1)	0	(0.0)
Total viral	709	(10.7)	28,274	(22.0)	5	(5.7)
Multiple etiologies	30	(0.5)	1,050	(0.8)	0	(0.0)
Confirmed etiology	2,167	(32.6)	68,981	(53.7)	76	(86.4)
Unknown etiology	4,480	(67.4)	59,389	(46.2)	12	(13.6)
Total 1998--2002	6,647	(100.0)	128,370	(100.0)	88	(100.0)

* Enterohemorrhagic (132 outbreaks), Enterotoxigenic (7), Enteroaggregative (1)

† Serotype O1 (1 outbreak), Serotype non-O1, non-O139 (1), serotype unspecified (1)

[Return to top.](#)[Figure 1](#)

FIGURE 1. Number of reported foodborne-disease outbreaks, 1993–2002

* Electronic Foodborne Outbreak Reporting System.

[Return to top.](#)


[Table 2](#)

TABLE 2. Number of reported foodborne-disease outbreaks, cases, and deaths, by etiology — United States, 1998

Etiology	Outbreaks		Cases		Deaths	
	No.	(%)	No.	(%)	No.	(%)
Bacterial						
<i>Bacillus cereus</i>	10	(0.8)	213	(0.8)	0	(0.0)
<i>Brucella</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Campylobacter</i>	12	(0.9)	483	(1.8)	0	(0.0)
<i>Clostridium botulinum</i>	3	(0.2)	8	(0.0)	0	(0.0)
<i>Clostridium perfringens</i>	24	(1.8)	1,328	(4.9)	0	(0.0)
<i>Escherichia coli</i>	32	(2.4)	1,613	(5.9)	0	(0.0)
<i>Listeria monocytogenes</i>	2	(0.2)	105	(0.4)	21	(65.6)
<i>Salmonella</i>	125	(9.5)	2,731	(10.0)	6	(18.8)
<i>Shigella</i>	17	(1.3)	1,266	(4.6)	0	(0.0)
<i>Staphylococcus aureus</i>	15	(1.1)	615	(2.3)	0	(0.0)
<i>Streptococcus</i>	1	(0.1)	4	(0.0)	0	(0.0)
<i>Vibrio cholerae</i>	1	(0.1)	6	(0.0)	0	(0.0)
<i>Vibrio parahaemolyticus</i>	13	(1.0)	532	(2.0)	0	(0.0)
<i>Vibrio, other</i>	1	(0.1)	2	(0.0)	0	(0.0)
<i>Yersinia enterocolitica</i>	1	(0.1)	9	(0.0)	0	(0.0)
Other bacterial	1	(0.1)	4	(0.0)	0	(0.0)
Total bacterial	258	(19.6)	8,919	(32.7)	27	(84.4)
Chemical						
Ciguatoxin	16	(1.2)	73	(0.3)	0	(0.0)
Heavy metals	0	(0.0)	0	(0.0)	0	(0.0)
Mushroom toxin	1	(0.1)	2	(0.0)	0	(0.0)
Scombrototoxin	27	(2.1)	124	(0.5)	0	(0.0)
Shellfish toxin	1	(0.1)	6	(0.0)	0	(0.0)
Other chemical	3	(0.2)	124	(0.5)	0	(0.0)
Total chemical	48	(3.7)	329	(1.2)	0	(0.0)
Parasitic						
<i>Anisakis</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Cryptosporidium parvum</i>	1	(0.1)	88	(0.3)	0	(0.0)
<i>Cyclospora cayentanensis</i>	1	(0.1)	17	(0.1)	0	(0.0)
<i>Giardia intestinalis</i>	1	(0.1)	3	(0.0)	0	(0.0)
<i>Trichinella spiralis</i>	0	(0.0)	0	(0.0)	0	(0.0)
Total parasitic	4	(0.3)	116	(0.4)	0	(0.0)
Viral						
Astrovirus	0	(0.0)	0	(0.0)	0	(0.0)
Hepatitis A	13	(1.0)	293	(1.1)	1	(3.1)
Norovirus	47	(3.6)	2,563	(9.4)	0	(0.0)
Rotavirus	0	(0.0)	0	(0.0)	0	(0.0)
Total viral	60	(4.6)	2,856	(10.5)	1	(3.1)
Multiple etiologies	2	(0.2)	31	(0.1)	0	(0.0)
Confirmed etiology	372	(28.3)	12,251	(44.9)	28	(87.5)
Unknown etiology	942	(71.7)	15,007	(55.1)	4	(12.5)
Total 1998	1,314	(100.0)	27,258	(100.0)	32	(100.0)

[Return to top.](#)

[Figure 2](#)

FIGURE 2. Number of reported foodborne-disease outbreaks, by state — United States,* 1998

* Includes Guam, Puerto Rico, and the U.S. Virgin Islands.

[Return to top.](#)

[Table 3](#)


TABLE 3. Number of reported foodborne-disease outbreaks, cases, and deaths, by etiology — United States, 1999

Etiology	Outbreaks		Cases		Deaths	
	No.	(%)	No.	(%)	No.	(%)
Bacterial						
<i>Bacillus cereus</i>	7	(0.5)	194	(0.8)	0	(0.0)
<i>Brucella</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Campylobacter</i>	5	(0.4)	85	(0.3)	0	(0.0)
<i>Clostridium botulinum</i>	1	(0.1)	3	(0.0)	0	(0.0)
<i>Clostridium perfringens</i>	22	(1.6)	1,166	(4.7)	1	(10.0)
<i>Escherichia coli</i>	28	(2.1)	842	(3.4)	0	(0.0)
<i>Listeria monocytogenes</i>	5	(0.4)	28	(0.1)	2	(20.0)
<i>Salmonella</i>	111	(8.3)	3,463	(13.9)	2	(20.0)
<i>Shigella</i>	14	(1.0)	221	(0.9)	0	(0.0)
<i>Staphylococcus aureus</i>	19	(1.4)	353	(1.4)	0	(0.0)
<i>Streptococcus</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Vibrio cholerae</i>	1	(0.1)	2	(0.0)	0	(0.0)
<i>Vibrio parahaemolyticus</i>	3	(0.2)	14	(0.1)	0	(0.0)
<i>Vibrio</i> , other	0	(0.0)	0	(0.0)	0	(0.0)
<i>Yersinia enterocolitica</i>	1	(0.1)	32	(0.1)	0	(0.0)
Other bacterial	0	(0.0)	0	(0.0)	0	(0.0)
Total bacterial	217	(16.2)	6,403	(25.7)	5	(50.0)
Chemical						
Ciguatera	12	(0.9)	47	(0.2)	1	(10.0)
Heavy metals	1	(0.1)	2	(0.0)	0	(0.0)
Mushroom toxin	0	(0.0)	0	(0.0)	0	(0.0)
Scombrototoxin	21	(1.6)	67	(0.3)	0	(0.0)
Shellfish toxin	0	(0.0)	0	(0.0)	0	(0.0)
Other chemical	1	(0.1)	2	(0.0)	0	(0.0)
Total chemical	35	(2.6)	118	(0.5)	1	(10.0)
Parasitic						
<i>Anisakis</i>	1	(0.1)	14	(0.1)	0	(0.0)
<i>Cryptosporidium parvum</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Cyclospora cayentanensis</i>	2	(0.1)	153	(0.6)	0	(0.0)
<i>Giardia intestinalis</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Trichinella spiralis</i>	0	(0.0)	0	(0.0)	0	(0.0)
Total parasitic	3	(0.2)	167	(0.7)	0	(0.0)
Viral						
Astrovirus	0	(0.0)	0	(0.0)	0	(0.0)
Hepatitis A	12	(0.9)	387	(1.6)	0	(0.0)
Norovirus	98	(7.3)	4,745	(19.1)	1	(10.0)
Rotavirus	0	(0.0)	0	(0.0)	0	(0.0)
Total viral	110	(8.2)	5,132	(20.6)	1	(10.0)
Multiple etiologies	5	(0.4)	267	(1.1)	0	(0.0)
Confirmed etiology	370	(27.6)	12,087	(48.6)	7	(70.0)
Unknown etiology	973	(72.4)	12,907	(51.4)	3	(30.0)
Total 1999	1,343	(100.0)	24,894	(100.0)	10	(100.0)

[Return to top.](#)

Figure 3

FIGURE 3. Number of reported foodborne-disease outbreaks, by state — United States,* 1999



* Includes Guam, Puerto Rico, and the U.S. Virgin Islands.

[Return to top.](#)


Table 4

TABLE 4. Number of reported foodborne-disease outbreaks, cases, and deaths, by etiology — United States, 2000

Etiology	Outbreaks		Cases		Deaths	
	No.	(%)	No.	(%)	No.	(%)
Bacterial						
<i>Bacillus cereus</i>	8	(0.6)	61	(0.2)	0	(0.0)
<i>Brucella</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Campylobacter</i>	15	(1.1)	205	(0.8)	0	(0.0)
<i>Clostridium botulinum</i>	2	(0.1)	5	(0.0)	1	(4.8)
<i>Clostridium perfringens</i>	22	(1.6)	791	(3.0)	0	(0.0)
<i>Escherichia coli</i>	32	(2.3)	1,392	(5.3)	2	(9.5)
<i>Listeria monocytogenes</i>	2	(0.1)	41	(0.2)	7	(33.3)
<i>Salmonella</i>	127	(9.0)	2,850	(10.9)	2	(9.5)
<i>Shigella</i>	12	(0.8)	866	(3.3)	1	(4.8)
<i>Staphylococcus aureus</i>	23	(1.6)	657	(2.5)	2	(9.5)
<i>Streptococcus</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Vibrio cholerae</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Vibrio parahaemolyticus</i>	4	(0.3)	37	(0.1)	0	(0.0)
<i>Vibrio</i> , other	0	(0.0)	0	(0.0)	0	(0.0)
<i>Yersinia enterocolitica</i>	0	(0.0)	0	(0.0)	0	(0.0)
Other bacterial	0	(0.0)	0	(0.0)	0	(0.0)
Total bacterial	247	(17.4)	6,905	(26.4)	15	(71.4)
Chemical						
Ciguatera toxin	12	(0.8)	46	(0.2)	0	(0.0)
Heavy metals	1	(0.1)	21	(0.1)	0	(0.0)
Mushroom toxin	0	(0.0)	0	(0.0)	0	(0.0)
Scombrototoxin	20	(1.4)	81	(0.3)	0	(0.0)
Shellfish toxin	3	(0.2)	9	(0.0)	0	(0.0)
Other chemical	2	(0.1)	36	(0.1)	0	(0.0)
Total chemical	38	(2.7)	193	(0.7)	0	(0.0)
Parasitic						
<i>Anisakis</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Cryptosporidium parvum</i>	1	(0.1)	8	(0.0)	0	(0.0)
<i>Cyclospora cayentanensis</i>	2	(0.1)	73	(0.3)	0	(0.0)
<i>Giardia intestinalis</i>	1	(0.1)	82	(0.3)	0	(0.0)
<i>Trichinella spiralis</i>	2	(0.1)	6	(0.0)	0	(0.0)
Total parasitic	6	(0.4)	169	(0.6)	0	(0.0)
Viral						
Astrovirus	0	(0.0)	0	(0.0)	0	(0.0)
Hepatitis A	12	(0.8)	135	(0.5)	1	(4.8)
Norovirus	163	(11.5)	6,969	(26.7)	0	(0.0)
Rotavirus	1	(0.1)	108	(0.4)	0	(0.0)
Total viral	176	(12.4)	7,212	(27.6)	1	(4.8)
Multiple etiologies	3	(0.2)	22	(0.1)	0	(0.0)
Confirmed etiology	470	(33.2)	14,501	(55.5)	16	(76.2)
Unknown etiology	947	(66.8)	11,621	(44.5)	5	(23.8)
Total 2000	1,417	(100.0)	26,122	(100.0)	21	(100.0)

[Return to top.](#)
[Figure 4](#)

FIGURE 4. Number of reported foodborne-disease outbreaks, by state — United States,* 2000




* Includes Guam, Puerto Rico, and the U.S. Virgin Islands.

[Return to top.](#)**Table 5****TABLE 5. Number of reported foodborne-disease outbreaks, cases, and deaths, by etiology — United States, 2001**

Etiology	Outbreaks		Cases		Deaths	
	No.	(%)	No.	(%)	No.	(%)
Bacterial						
<i>Bacillus cereus</i>	5	(0.4)	61	(0.2)	0	(0.0)
<i>Brucella</i>	1	(0.1)	4	(0.0)	0	(0.0)
<i>Campylobacter</i>	16	(1.3)	317	(1.3)	0	(0.0)
<i>Clostridium botulinum</i>	3	(0.2)	22	(0.1)	0	(0.0)
<i>Clostridium perfringens</i>	31	(2.5)	1,232	(4.9)	3	(27.3)
<i>Escherichia coli</i>	22	(1.8)	521	(2.1)	0	(0.0)
<i>Listeria monocytogenes</i>	1	(0.1)	28	(0.1)	0	(0.0)
<i>Salmonella</i>	111	(8.9)	3,141	(12.5)	7	(63.6)
<i>Shigella</i>	15	(1.2)	1,006	(4.0)	0	(0.0)
<i>Staphylococcus aureus</i>	23	(1.9)	646	(2.6)	0	(0.0)
<i>Streptococcus</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Vibrio cholerae</i>	1	(0.1)	4	(0.0)	0	(0.0)
<i>Vibrio parahaemolyticus</i>	3	(0.2)	19	(0.1)	0	(0.0)
<i>Vibrio, other</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Yersinia enterocolitica</i>	3	(0.2)	33	(0.1)	0	(0.0)
Other bacterial	0	(0.0)	0	(0.0)	0	(0.0)
Total bacterial	235	(19.9)	7,034	(29.0)	10	(90.9)
Chemical						
Ciguatoxin	24	(1.9)	81	(0.3)	0	(0.0)
Heavy metals	0	(0.0)	0	(0.0)	0	(0.0)
Mushroom toxin	0	(0.0)	0	(0.0)	0	(0.0)
Scombrototoxin	29	(2.3)	132	(0.5)	0	(0.0)
Shellfish toxin	0	(0.0)	0	(0.0)	0	(0.0)
Other chemical	1	(0.1)	15	(0.1)	0	(0.0)
Total chemical	54	(4.3)	228	(0.9)	0	(0.0)
Parasitic						
<i>Anisakis</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Cryptosporidium parvum</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Cyclospora cayentanensis</i>	2	(0.2)	42	(0.2)	0	(0.0)
<i>Giardia intestinalis</i>	1	(0.1)	34	(0.1)	0	(0.0)
<i>Trichinella spiralis</i>	2	(0.2)	14	(0.1)	0	(0.0)
Total parasitic	5	(0.4)	90	(0.4)	0	(0.0)
Viral						
Astrovirus	0	(0.0)	0	(0.0)	0	(0.0)
Hepatitis A	6	(0.5)	116	(0.5)	1	(9.1)
Norovirus	150	(12.1)	6,335	(25.2)	0	(0.0)
Rotavirus	0	(0.0)	0	(0.0)	0	(0.0)
Total viral	156	(12.6)	6,451	(25.7)	1	(9.1)
Multiple etiologies	9	(0.7)	190	(0.8)	0	(0.0)
Confirmed etiology	450	(36.9)	13,993	(55.7)	11	(100.0)
Unknown etiology	784	(63.1)	11,137	(44.3)	0	(0.0)
Total 2001	1,243	(100.0)	25,130	(100.0)	11	(100.0)

[Return to top.](#)**Figure 5**

FIGURE 5. Number of reported foodborne-disease outbreaks, by state — United States,* 2001

* Includes Guam, Puerto Rico, and the U.S. Virgin Islands.

[Return to top.](#)

[Table 6](#)


TABLE 6. Number of reported foodborne-disease outbreaks, cases, and deaths, by etiology — United States, 2002

Etiology	Outbreaks		Cases		Deaths	
	No.	(%)	No.	(%)	No.	(%)
Bacterial						
<i>Bacillus cereus</i>	7	(0.5)	42	(0.2)	0	(0.0)
<i>Brucella</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Campylobacter</i>	13	(1.0)	350	(1.4)	0	(0.0)
<i>Clostridium botulinum</i>	3	(0.2)	14	(0.1)	0	(0.0)
<i>Clostridium perfringens</i>	31	(2.3)	2,207	(8.8)	0	(0.0)
<i>Escherichia coli</i>	26	(2.0)	496	(1.9)	2	(14.3)
<i>Listeria monocytogenes</i>	1	(0.1)	54	(0.2)	8	(57.1)
<i>Salmonella</i>	111	(8.3)	4,636	(18.6)	3	(21.4)
<i>Shigella</i>	9	(0.7)	318	(1.3)	0	(0.0)
<i>Staphylococcus aureus</i>	21	(1.6)	495	(2.0)	0	(0.0)
<i>Streptococcus</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Vibrio cholerae</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Vibrio parahaemolyticus</i>	2	(0.2)	11	(0.0)	0	(0.0)
<i>Vibrio, other</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Yersinia enterocolitica</i>	3	(0.2)	13	(0.1)	0	(0.0)
Other bacterial	0	(0.0)	0	(0.0)	0	(0.0)
Total bacterial	227	(17.1)	8,626	(34.6)	13	(92.9)
Chemical						
Ciguatoxin	20	(1.5)	68	(0.3)	0	(0.0)
Heavy metals	0	(0.0)	0	(0.0)	0	(0.0)
Mushroom toxin	1	(0.1)	4	(0.0)	0	(0.0)
Scombrototoxin	21	(1.6)	59	(0.2)	0	(0.0)
Shellfish toxin	1	(0.1)	21	(0.1)	0	(0.0)
Other chemical	3	(0.2)	120	(0.5)	0	(0.0)
Total chemical	46	(3.5)	272	(1.1)	0	(0.0)
Parasitic						
<i>Anisakis</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Cryptosporidium parvum</i>	2	(0.2)	43	(0.2)	0	(0.0)
<i>Cyclospora cayentanensis</i>	2	(0.2)	40	(0.2)	0	(0.0)
<i>Giardia intestinalis</i>	0	(0.0)	0	(0.0)	0	(0.0)
<i>Trichinella spiralis</i>	1	(0.1)	5	(0.0)	0	(0.0)
Total parasitic	5	(0.4)	88	(0.4)	0	(0.0)
Viral						
Astrovirus	1	(0.1)	14	(0.1)	0	(0.0)
Hepatitis A	7	(0.5)	50	(0.2)	1	(7.1)
Norovirus	199	(15.0)	6,559	(26.3)	0	(0.0)
Rotavirus	0	(0.0)	0	(0.0)	0	(0.0)
Total viral	207	(15.6)	6,623	(26.5)	1	(7.1)
Multiple etiologies	11	(0.8)	540	(2.2)	0	(0.0)
Confirmed etiology	496	(37.3)	16,149	(64.7)	14	(100.0)
Unknown etiology	834	(62.7)	8,817	(35.3)	0	(0.0)
Total 2002	1,330	(100.0)	24,966	(100.0)	14	(100.0)

[Return to top.](#)

Figure 6

FIGURE 6. Number of reported foodborne-disease outbreaks, by state — United States,* 2002



* Includes Guam, Puerto Rico, and the U.S. Virgin Islands.

[Return to top.](#)

Table 7

TABLE 7. Number of reported foodborne-disease outbreaks, by etiology and month of occurrence — United States, 1998–2002

Etiology	Month of occurrence												Total
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	
Bacterial													
<i>Bacillus cereus</i>	3	3	2	5	5	7	4	3	1	1	1	2	37
<i>Brucella</i>	—	—	—	—	—	—	1	—	—	—	—	—	1
<i>Campylobacter</i>	4	5	5	4	7	13	3	9	4	1	5	1	61
<i>Clostridium botulinum</i>	3	—	—	1	—	1	3	1	1	1	—	1	12
<i>Clostridium perfringens</i>	3	9	6	17	14	8	8	11	8	13	15	18	130
<i>Escherichia coli</i>	—	1	3	8	19	18	24	21	15	19	10	2	140
<i>Listeria monocytogenes</i>	—	—	—	—	1	1	1	2	1	2	1	2	11
<i>Salmonella</i>	32	22	31	43	50	88	80	76	61	33	39	30	585
<i>Shigella</i>	8	3	2	5	4	11	3	6	11	6	5	3	67
<i>Staphylococcus aureus</i>	4	6	9	8	11	13	5	7	9	13	6	10	101
<i>Streptococcus</i>	—	—	1	—	—	—	—	—	—	—	—	—	1
<i>Vibrio cholerae</i>	1	—	—	—	—	1	—	—	1	—	—	—	3
<i>Vibrio parahaemolyticus</i>	1	—	1	2	2	8	5	5	1	—	—	—	25
<i>Vibrio, other</i>	—	—	—	—	—	—	—	—	—	—	1	—	1
<i>Yersinia enterocolitica</i>	2	1	—	—	—	—	—	—	1	—	3	1	8
Other bacterial	—	—	—	—	1	—	—	—	—	—	—	—	1
Total bacterial	61	50	60	93	114	169	137	141	114	89	86	70	1,184
Chemical													
Ciguatera	4	2	4	8	12	6	17	6	10	8	5	2	84
Heavy metals	—	—	—	1	—	—	—	—	—	—	—	1	2
Mushroom toxin	—	—	—	—	—	—	1	—	—	1	—	—	2
Scombrototoxin	7	6	5	14	11	12	11	12	10	12	7	11	118
Shellfish toxin	1	—	—	—	—	—	—	3	—	1	—	—	5
Other chemical	—	—	1	—	4	—	—	1	—	1	1	2	10
Total chemical	12	8	10	23	27	18	29	22	20	23	13	16	221
Parasitic													
<i>Anisakis</i>	—	1	—	—	—	—	—	—	—	—	—	—	1
<i>Cryptosporidium parvum</i>	—	—	—	1	—	—	—	—	2	1	—	—	4
<i>Cyclospora cayentanensis</i>	3	—	—	—	3	2	1	—	—	—	—	—	9
<i>Giardia intestinalis</i>	—	—	—	1	—	—	—	1	—	1	—	—	3
<i>Trichinella spiralis</i>	—	—	—	1	1	—	—	2	1	—	1	—	6
Total parasitic	3	1	0	3	4	2	1	3	3	2	1	0	23
Viral													
Astrovirus	1	—	—	—	—	—	—	—	—	—	—	—	1
Hepatitis A	4	2	9	4	—	5	3	3	4	10	2	4	50
Norovirus	46	51	64	60	70	49	43	42	28	47	66	91	657
Rotavirus	—	—	1	—	—	—	—	—	—	—	—	—	1
Total viral	51	53	74	64	70	54	46	45	32	57	68	95	709
Multiple etiologies	1	2	1	5	3	1	3	5	3	4	2	—	30
Confirmed etiology	128	114	145	188	218	244	216	216	172	175	170	181	2,167
Unknown etiology	329	355	422	425	462	394	349	334	267	316	377	450	4,480
Total 1998–2002	457	469	567	613	680	638	565	550	439	491	547	631	6,647

[Return to top.](#)[Table 8](#)

TABLE 8. Number of reported foodborne-disease outbreaks, by etiology and place where food was eaten* — United States, 1998–2002

Etiology	Place where food was eaten								
	Private residence	Restaurant or Delicatessen	Grocery	School	Daycare center	Workplace cafeteria	Picnic	Church	Camp
Bacterial									
<i>Bacillus cereus</i>	7	13	—	2	—	2	—	—	—
<i>Brucella</i>	1	—	—	—	—	—	—	—	—
<i>Campylobacter</i>	16	25	—	2	—	—	1	1	2
<i>Clostridium botulinum</i>	10	—	—	—	—	1	—	1	—
<i>Clostridium perfringens</i>	15	36	—	11	—	10	2	6	1
<i>Escherichia coli</i>	40	41	1	9	2	2	7	5	6
<i>Listeria monocytogenes</i>	9	2	2	—	—	—	—	—	—
<i>Salmonella</i>	169	271	2	21	5	7	12	20	9
<i>Shigella</i>	10	39	—	5	2	—	3	2	—
<i>Staphylococcus aureus</i>	20	26	1	10	2	8	4	9	3
<i>Streptococcus</i>	—	—	1	—	—	—	—	—	—
<i>Vibrio cholerae</i>	1	2	—	—	—	—	—	—	—
<i>Vibrio parahaemolyticus</i>	4	17	—	—	—	—	—	—	—
<i>Vibrio</i> , other	—	—	—	—	—	—	—	—	—
<i>Yersinia enterocolitica</i>	4	1	—	—	—	1	—	—	—
Other bacterial	—	—	—	—	—	—	—	—	—
Total bacterial	306	473	7	60	11	31	29	44	21
Chemical									
Ciguatera	73	7	—	—	—	—	1	—	2
Heavy metals	1	—	—	1	—	—	—	—	—
Mushroom toxin	2	—	—	—	—	—	—	—	—
Scombrototoxin	14	84	2	1	—	3	—	—	—
Shellfish toxin	4	—	—	—	—	—	—	—	—
Other chemical	2	3	—	2	—	1	—	—	—
Total chemical	96	94	2	4	—	4	1	0	2
Parasitic									
<i>Anisakis</i>	1	—	—	—	—	—	—	—	—
<i>Cryptosporidium parvum</i>	2	—	—	—	—	—	—	—	—
<i>Cyclospora cayentanensis</i>	2	2	—	—	—	—	—	—	—
<i>Giardia intestinalis</i>	1	1	—	—	—	—	—	—	—
<i>Trichinella spiralis</i>	3	—	—	—	—	—	1	—	1
Total parasitic	9	3	—	—	—	—	1	—	1
Viral									
Astrovirus	—	1	—	—	—	—	—	—	—
Hepatitis A	5	29	—	2	1	3	2	—	—
Norovirus	83	279	—	51	1	30	11	17	12
Rotavirus	—	—	—	1	—	—	—	—	—
Total viral	88	309	—	54	2	33	13	17	12
Multiple etiologies	7	8	1	2	1	—	2	—	1
Confirmed etiology	506	887	10	120	14	68	46	61	37
Unknown etiology	791	2,447	6	165	10	133	45	54	29
Total 1998–2002	1,297	3,334	16	285	24	201	91	115	66

* More than one place might be reported per outbreak.

TABLE 8. (Continued) Number of reported foodborne-disease outbreaks, by etiology and place where food was eaten* — United States, 1998–2002

Etiology	Place where food was eaten						Total	
	Fair or festival	Hospital	Nursing home	Prison	Other	Known place		Unknown place
Bacterial								
<i>Bacillus cereus</i>	—	—	—	—	8	33	4	37
<i>Brucella</i>	—	—	—	—	—	1	—	1
<i>Campylobacter</i>	1	—	—	1	13	57	4	61
<i>Clostridium botulinum</i>	—	—	—	—	—	11	1	12
<i>Clostridium perfringens</i>	—	4	—	10	33	125	5	130
<i>Escherichia coli</i>	3	—	1	7	20	127	13	140
<i>Listeria monocytogenes</i>	—	2	1	—	2	11	—	11
<i>Salmonella</i>	7	4	19	10	68	540	45	585
<i>Shigella</i>	—	—	—	—	4	64	3	67
<i>Staphylococcus aureus</i>	4	1	1	1	18	96	5	101
<i>Streptococcus</i>	—	—	—	—	—	1	—	1
<i>Vibrio cholerae</i>	—	—	—	—	—	3	—	3
<i>Vibrio parahaemolyticus</i>	—	—	—	—	2	23	2	25
<i>Vibrio, other</i>	—	—	—	—	—	—	1	1
<i>Yersinia enterocolitica</i>	—	—	—	—	1	7	1	8
Other bacterial	—	—	—	—	—	—	1	1
Total bacterial	15	11	22	29	169	1,099	85	1,184
Chemical								
Ciguatera	—	—	—	—	2	82	2	84
Heavy metals	—	—	—	—	—	2	—	2
Mushroom toxin	—	—	—	—	—	2	—	2
Scombrototoxin	—	1	1	—	8	113	5	118
Shellfish toxin	—	—	—	—	1	4	1	5
Other chemical	—	—	—	—	1	9	1	10
Total chemical	—	1	1	—	12	212	9	221
Parasitic								
<i>Anisakis</i>	—	—	—	—	—	1	—	1
<i>Cryptosporidium parvum</i>	—	—	—	—	2	4	—	4
<i>Cyclospora cayentanensis</i>	—	—	—	—	4	7	2	9
<i>Giardia intestinalis</i>	—	—	—	—	1	3	—	3
<i>Trichinella spiralis</i>	—	—	—	—	3	6	—	6
Total parasitic	—	—	—	—	10	21	2	23
Viral								
Astrovirus	—	—	—	—	—	1	—	1
Hepatitis A	—	—	—	—	4	42	8	50
Norovirus	5	7	21	6	148	636	21	657
Rotavirus	—	—	—	—	—	1	—	1
Total viral	5	7	21	6	152	680	29	709
Multiple etiologies	1	—	1	4	3	29	1	30
Confirmed etiology	21	19	45	39	346	2,041	126	2,167
Unknown etiology	23	32	23	20	576	4,222	258	4,480
Total 1998–2002	44	51	68	59	922	6,263	384	6,647

* More than one place might be reported per outbreak.

[Return to top.](#)[Table 9](#)**TABLE 9. Number of reported foodborne-disease outbreaks, cases, and deaths, by vehicle of transmission — United States, 1998**

Vehicle of transmission	Outbreaks		Cases		Deaths	
	No.	(%)	No.	(%)	No.	(%)
Beef	26	(2.0)	805	(3.0)	0	(0.0)
Dairy	18	(1.4)	492	(1.8)	0	(0.0)
Eggs	7	(0.5)	48	(0.2)	0	(0.0)
Game	2	(0.2)	13	(0.0)	0	(0.0)
Pork	29	(2.2)	610	(2.2)	0	(0.0)
Poultry	62	(4.7)	876	(3.2)	0	(0.0)
Vegetables	27	(2.1)	1,299	(4.8)	2	(6.3)
Fruits and nuts	17	(1.3)	586	(2.1)	0	(0.0)
Grains	9	(0.7)	306	(1.1)	0	(0.0)
Oils and sugars	1	(0.1)	4	(0.0)	0	(0.0)
Finfish	69	(5.3)	493	(1.8)	0	(0.0)
Shellfish	38	(2.9)	880	(3.2)	0	(0.0)
Unclassifiable vehicle	41	(3.1)	632	(2.3)	2	(6.3)
Complex vehicle	432	(32.9)	10,851	(39.8)	23	(71.9)
Known vehicle	778	(59.2)	17,896	(65.7)	27	(84.4)
Unknown vehicle	536	(40.8)	9,363	(34.3)	5	(15.6)
Total 1998	1,314	(100.0)	27,258	(100.0)	32	(100.0)

[Return to top.](#)[Table 10](#)

TABLE 10. Number of reported foodborne-disease outbreaks, cases, and deaths, by vehicle of transmission — United States, 1999

Vehicle of transmission	Outbreaks		Cases		Deaths	
	No.	(%)	No.	(%)	No.	(%)
Beef	62	(4.6)	1,332	(5.4)	0	(0.0)
Dairy	15	(1.1)	199	(0.8)	0	(0.0)
Eggs	25	(1.9)	762	(3.1)	0	(0.0)
Game	0	(0.0)	0	(0.0)	0	(0.0)
Pork	26	(1.9)	559	(2.2)	0	(0.0)
Poultry	74	(5.5)	947	(3.8)	0	(0.0)
Vegetables	43	(3.2)	1,273	(5.1)	0	(0.0)
Fruits and nuts	19	(1.4)	629	(2.5)	0	(0.0)
Grains	19	(1.4)	139	(0.6)	0	(0.0)
Oils and sugars	5	(0.4)	135	(0.5)	0	(0.0)
Finfish	64	(4.8)	322	(1.3)	1	(10.0)
Shellfish	28	(2.1)	253	(1.0)	0	(0.0)
Unclassifiable vehicle	37	(2.8)	1,513	(6.1)	0	(0.0)
Complex vehicle	372	(27.7)	7,189	(28.9)	5	(50.0)
Known vehicle	799	(58.7)	15,252	(61.3)	6	(60.0)
Unknown vehicle	554	(41.3)	9,642	(38.7)	4	(40.0)
Total 1999	1,343	(100.0)	24,894	(100.0)	10	(100.0)

[Return to top.](#)[Table 11](#)**TABLE 11. Number of reported foodborne-disease outbreaks, cases, and deaths, by vehicle of transmission — United States, 2000**

Vehicle of transmission	Outbreaks		Cases		Deaths	
	No.	(%)	No.	(%)	No.	(%)
Beef	43	(3.0)	696	(2.7)	1	(4.8)
Dairy	22	(1.6)	300	(1.1)	0	(0.0)
Eggs	25	(1.8)	620	(2.4)	0	(0.0)
Game	0	(0.0)	0	(0.0)	0	(0.0)
Pork	27	(1.9)	610	(2.3)	0	(0.0)
Poultry	61	(4.3)	829	(3.2)	7	(33.3)
Vegetables	41	(2.9)	872	(3.3)	2	(9.5)
Fruits and nuts	21	(1.5)	1,527	(5.8)	1	(4.8)
Grains	28	(2.0)	434	(1.7)	0	(0.0)
Oils and sugars	1	(0.1)	27	(0.1)	0	(0.0)
Finfish	63	(4.4)	267	(1.0)	0	(0.0)
Shellfish	25	(1.8)	134	(0.5)	0	(0.0)
Unclassifiable vehicle	51	(3.5)	959	(3.6)	0	(0.0)
Complex vehicle	455	(32.1)	9,525	(36.5)	9	(42.9)
Known vehicle	963	(60.9)	16,800	(64.3)	20	(95.2)
Unknown vehicle	554	(39.1)	9,322	(35.7)	1	(4.8)
Total 2000	1,417	(100.0)	26,122	(100.0)	21	(100.0)

[Return to top.](#)[Table 12](#)**TABLE 12. Number of reported foodborne-disease outbreaks, cases, and deaths, by vehicle of transmission — United States, 2001**

Vehicle of transmission	Outbreaks		Cases		Deaths	
	No.	(%)	No.	(%)	No.	(%)
Beef	33	(2.6)	525	(2.1)	1	(9.1)
Dairy	21	(1.7)	536	(2.1)	0	(0.0)
Eggs	12	(1.0)	465	(1.9)	0	(0.0)
Game	5	(0.4)	45	(0.2)	0	(0.0)
Pork	30	(2.4)	560	(2.2)	0	(0.0)
Poultry	73	(5.9)	1,010	(4.0)	0	(0.0)
Vegetables	37	(3.0)	1,997	(7.9)	0	(0.0)
Fruits and nuts	21	(1.7)	585	(2.3)	2	(18.2)
Grains	11	(0.9)	92	(0.4)	0	(0.0)
Oils and sugars	4	(0.3)	95	(0.4)	0	(0.0)
Finfish	75	(6.0)	330	(1.3)	0	(0.0)
Shellfish	33	(2.7)	291	(1.2)	0	(0.0)
Unclassifiable vehicle	51	(4.1)	1,182	(4.7)	1	(9.1)
Complex vehicle	384	(31.0)	8,112	(32.3)	1	(9.1)
Known vehicle	790	(63.6)	15,825	(63.0)	5	(45.5)
Unknown vehicle	453	(36.4)	9,305	(37.0)	6	(54.5)
Total 2001	1,243	(100.0)	25,130	(100.0)	11	(100.0)

[Return to top.](#)[Table 13](#)

TABLE 13. Number of reported foodborne-disease outbreaks, cases, and deaths, by vehicle of transmission — United States, 2002

Vehicle of transmission	Outbreaks		Cases		Deaths	
	No.	(%)	No.	(%)	No.	(%)
Beef	44	(3.3)	831	(3.3)	3	(21.4)
Dairy	16	(1.2)	704	(2.8)	0	(0.0)
Eggs	14	(1.1)	317	(1.3)	0	(0.0)
Game	3	(0.2)	33	(0.1)	0	(0.0)
Pork	26	(2.0)	360	(1.4)	0	(0.0)
Poultry	75	(5.6)	1325	(5.3)	8	(57.1)
Vegetables	44	(3.3)	1506	(6.4)	0	(0.0)
Fruits and nuts	9	(0.7)	169	(0.7)	0	(0.0)
Grains	14	(1.1)	177	(0.7)	0	(0.0)
Oils and sugars	1	(0.1)	4	(0.0)	0	(0.0)
Finfish	66	(5.0)	280	(1.1)	0	(0.0)
Shellfish	27	(2.0)	200	(0.8)	0	(0.0)
Unclassifiable vehicle	52	(3.9)	1049	(4.2)	0	(0.0)
Complex vehicle	436	(32.8)	9369	(37.5)	1	(7.1)
Known vehicle	827	(62.2)	16414	(65.7)	12	(85.7)
Unknown vehicle	503	(37.8)	8552	(34.3)	2	(14.3)
Total 2002	1330	(100.0)	24966	(100.0)	14	(100.0)

[Return to top.](#)

[Table 14](#)

TABLE 14. Number of reported foodborne-disease outbreaks, by etiology and vehicle of transmission — United States, 1998

Etiology	Vehicle of transmission								
	Beef	Dairy	Eggs	Game	Pork	Poultry	Vegetables	Fruits and nuts	Grains
Bacterial									
<i>Bacillus cereus</i>	—	—	—	—	—	—	—	—	—
<i>Brucella</i>	—	—	—	—	—	—	—	—	—
<i>Campylobacter</i>	—	2	—	—	—	1	1	—	—
<i>Clostridium botulinum</i>	—	—	—	—	—	—	—	—	—
<i>Clostridium perfringens</i>	4	—	—	—	3	2	—	—	—
<i>Escherichia coli</i>	1	2	—	—	—	—	2	1	—
<i>Listeria monocytogenes</i>	—	—	—	—	—	—	—	—	—
<i>Salmonella</i>	—	3	3	1	2	6	3	1	2
<i>Shigella</i>	—	—	—	—	—	1	1	—	—
<i>Staphylococcus aureus</i>	1	—	—	—	2	1	—	—	—
<i>Streptococcus</i>	—	—	—	—	—	1	—	—	—
<i>Vibrio cholerae</i>	—	—	—	—	—	—	—	—	—
<i>Vibrio parahaemolyticus</i>	—	—	—	—	—	1	—	—	—
<i>Vibrio, other</i>	—	—	—	—	—	—	—	—	—
<i>Yersinia enterocolitica</i>	—	—	—	—	—	—	—	—	—
Other bacterial	—	—	—	—	—	—	—	—	—
Total bacterial	6	7	3	1	7	13	7	2	2
Chemical									
Ciguatera	—	—	—	—	—	—	—	—	—
Heavy metals	—	—	—	—	—	—	—	—	—
Mushroom toxin	—	—	—	—	—	—	1	—	—
Scombrototoxin	—	—	—	—	—	—	—	—	—
Shellfish toxin	—	—	—	—	—	—	—	—	—
Other chemical	—	—	—	—	—	—	1	—	—
Total chemical	—	—	—	—	—	—	2	—	—
Parasitic									
<i>Anisakis</i>	—	—	—	—	—	—	—	—	—
<i>Cryptosporidium parvum</i>	—	—	—	—	—	—	—	—	—
<i>Cyclospora cayentanensis</i>	—	—	—	—	—	—	—	—	—
<i>Giardia intestinalis</i>	—	—	—	—	—	—	—	—	—
<i>Trichinella spiralis</i>	—	—	—	1	—	—	—	—	—
Total parasitic	—	—	—	1	—	—	—	—	—
Viral									
Astrovirus	—	—	—	—	—	—	—	—	—
Hepatitis A	—	—	—	—	—	—	—	2	—
Norovirus	—	1	—	—	1	—	1	2	—
Rotavirus	—	—	—	—	—	—	—	—	—
Total viral	—	1	—	—	1	—	1	4	—
Multiple etiologies	—	—	—	—	—	—	—	—	—
Confirmed etiology	6	8	3	2	8	13	10	6	2
Unknown etiology	20	10	4	—	21	49	17	11	7
Total 1998	26	18	7	2	29	62	27	17	9

TABLE 14. (Continued) Number of reported foodborne-disease outbreaks, by etiology and vehicle of transmission — United States, 1998

Etiology	Vehicle of transmission						Total	
	Oils and sugars	Finfish	Shellfish	Unclassifiable vehicle	Complex vehicle	Known vehicle		Unknown vehicle
Bacterial								
<i>Bacillus cereus</i>	—	—	1	—	7	8	2	10
<i>Brucella</i>	—	—	—	—	—	—	—	—
<i>Campylobacter</i>	—	—	1	—	2	7	5	12
<i>Clostridium botulinum</i>	—	2	—	—	1	3	—	3
<i>Clostridium perfringens</i>	—	—	—	—	12	21	3	24
<i>Escherichia coli</i>	—	—	—	—	12	18	14	32
<i>Listeria monocytogenes</i>	—	—	—	—	2	2	—	2
<i>Salmonella</i>	—	1	1	4	35	62	63	125
<i>Shigella</i>	—	1	—	—	4	7	10	17
<i>Staphylococcus aureus</i>	—	—	—	—	9	13	2	15
<i>Streptococcus</i>	—	—	—	—	—	1	—	1
<i>Vibrio cholerae</i>	—	—	1	—	—	1	—	1
<i>Vibrio parahaemolyticus</i>	—	—	11	—	—	12	1	13
<i>Vibrio, other</i>	—	—	1	—	—	1	—	1
<i>Yersinia enterocolitica</i>	—	—	—	—	1	1	—	1
Other bacterial	—	—	—	—	1	1	—	1
Total bacterial	—	4	16	4	86	158	100	258
Chemical								
Ciguatera	—	16	—	—	—	16	—	16
Heavy metals	—	—	—	—	—	—	—	—
Mushroom toxin	—	—	—	—	—	1	—	1
Scombrototoxin	—	27	—	—	—	27	—	27
Shellfish toxin	—	—	1	—	—	1	—	1
Other chemical	—	1	—	—	1	3	—	3
Total chemical	—	44	1	—	1	48	—	48
Parasitic								
<i>Anisakis</i>	—	—	—	—	—	—	—	—
<i>Cryptosporidium parvum</i>	—	—	—	—	—	—	1	—
<i>Cyclospora cayentanensis</i>	—	—	—	—	—	—	1	—
<i>Giardia intestinalis</i>	—	—	—	—	1	1	—	—
<i>Trichinella spiralis</i>	—	—	—	—	—	1	—	1
Total parasitic	—	—	—	—	1	2	2	1
Viral								
Astrovirus	—	—	—	—	—	—	—	—
Hepatitis A	—	—	—	—	—	2	11	13
Norovirus	—	—	1	3	17	26	21	47
Rotavirus	—	—	—	—	—	—	—	—
Total viral	—	—	1	3	17	28	32	60
Multiple etiologies	—	—	—	—	1	1	1	2
Confirmed etiology	—	48	18	6	107	237	135	372
Unknown etiology	1	21	20	35	325	541	401	942
Total 1998	1	69	38	41	432	778	536	1,314

[Return to top.](#)

[Table 15](#)

TABLE 15. Number of reported foodborne-disease outbreaks, by etiology and vehicle of transmission — United States, 1999

Etiology	Vehicle of transmission								
	Beef	Dairy	Eggs	Game	Pork	Poultry	Vegetables	Fruits and nuts	Grains
Bacterial									
<i>Bacillus cereus</i>	—	—	—	—	—	—	—	—	3
<i>Brucella</i>	—	—	—	—	—	—	—	—	—
<i>Campylobacter</i>	—	1	—	—	—	1	—	—	—
<i>Clostridium botulinum</i>	—	—	—	—	—	—	—	—	—
<i>Clostridium perfringens</i>	4	—	1	—	—	3	1	—	—
<i>Escherichia coli</i>	12	—	—	—	—	—	5	—	—
<i>Listeria monocytogenes</i>	1	—	—	—	—	—	—	—	—
<i>Salmonella</i>	3	2	19	—	3	15	7	5	—
<i>Shigella</i>	—	—	1	—	—	—	—	—	—
<i>Staphylococcus aureus</i>	2	—	—	—	6	—	1	—	—
<i>Streptococcus</i>	—	—	—	—	—	—	—	—	—
<i>Vibrio cholerae</i>	—	—	—	—	—	—	—	—	—
<i>Vibrioparahaemolyticus</i>	—	—	—	—	—	—	—	—	—
<i>Vibrio, other</i>	—	—	—	—	—	—	—	—	—
<i>Yersinia enterocolitica</i>	—	—	—	—	1	—	—	—	—
Other bacterial	—	—	—	—	—	—	—	—	—
Total bacterial	22	3	21	0	10	19	14	5	3
Chemical									
Ciguatoxin	—	—	—	—	—	—	—	—	—
Heavy metals	—	—	—	—	—	—	—	—	—
Mushroom toxin	—	—	—	—	—	—	—	—	—
Scombrototoxin	—	—	—	—	—	—	—	—	—
Shellfish toxin	—	—	—	—	—	—	—	—	—
Other chemical	—	—	—	—	—	—	—	—	—
Total chemical	—	—	—	—	—	—	—	—	—
Parasitic									
<i>Anisakis</i>	—	—	—	—	—	—	—	—	—
<i>Cryptosporidium parvum</i>	—	—	—	—	—	—	—	—	—
<i>Cyclospora cayentanensis</i>	—	—	—	—	—	—	1	1	—
<i>Giardia intestinalis</i>	—	—	—	—	—	—	—	—	—
<i>Trichinella spiralis</i>	—	—	—	—	—	—	—	—	—
Total parasitic	—	—	—	—	—	—	1	1	—
Viral									
Astrovirus	—	—	—	—	—	—	—	—	—
Hepatitis A	—	—	—	—	—	—	—	—	—
Norovirus	5	2	—	—	1	2	3	5	—
Rotavirus	—	—	—	—	—	—	—	—	—
Total viral	5	2	—	—	1	2	3	5	—
Multiple etiologies	—	—	—	—	—	1	—	—	1
Confirmed etiology	27	5	21	—	11	22	18	11	4
Unknown etiology	35	10	4	—	15	52	25	8	15
Total 1999	62	15	25	—	26	74	43	19	19

TABLE 15. (Continued) Number of reported foodborne-disease outbreaks, by etiology and vehicle of transmission — United States, 1999

Etiology	Vehicle of transmission						Total	
	Oils and sugars	Finfish	Shellfish	Unclassifiable vehicle	Complex vehicle	Known vehicle		Unknown vehicle
Bacterial								
<i>Bacillus cereus</i>	—	—	—	—	3	6	1	7
<i>Brucella</i>	—	—	—	—	—	—	—	—
<i>Campylobacter</i>	—	—	—	—	—	2	3	5
<i>Clostridium botulinum</i>	—	—	—	—	1	1	—	1
<i>Clostridium perfringens</i>	—	—	—	—	12	21	1	22
<i>Escherichia coli</i>	—	—	—	1	3	21	7	28
<i>Listeria monocytogenes</i>	—	—	—	—	3	4	1	5
<i>Salmonella</i>	—	—	—	1	21	76	35	111
<i>Shigella</i>	—	—	—	—	7	8	6	14
<i>Staphylococcus aureus</i>	—	—	—	—	7	16	3	19
<i>Streptococcus</i>	—	—	—	—	—	—	—	—
<i>Vibrio cholerae</i>	—	—	1	—	—	1	—	1
<i>Vibrioparahaemolyticus</i>	—	—	3	—	—	3	—	3
<i>Vibrio, other</i>	—	—	—	—	—	—	—	—
<i>Yersinia enterocolitica</i>	—	—	—	—	—	1	—	1
Other bacterial	—	—	—	—	—	—	—	—
Total bacterial	—	—	4	2	57	160	57	217
Chemical								
Ciguatera	—	12	—	—	—	12	—	12
Heavy metals	—	—	—	—	1	1	—	1
Mushroom toxin	—	—	—	—	—	—	—	—
Scombrototoxin	—	20	—	—	1	21	—	21
Shellfish toxin	—	—	—	—	—	—	—	—
Other chemical	1	—	—	—	—	1	—	1
Total chemical	1	32	—	—	2	35	—	35
Parasitic								
Anisakis	—	1	—	—	—	1	—	1
<i>Cryptosporidium parvum</i>	—	—	—	—	—	—	—	—
<i>Cyclospora cayentanensis</i>	—	—	—	—	—	2	—	2
<i>Giardia intestinalis</i>	—	—	—	—	—	—	—	—
<i>Trichinella spiralis</i>	—	—	—	—	—	—	—	—
Total parasitic	—	1	—	—	—	3	—	3
Viral								
Astrovirus	—	—	—	—	—	—	—	—
Hepatitis A	—	—	—	1	5	6	6	12
Norovirus	1	—	2	3	29	53	45	98
Rotavirus	—	—	—	—	—	—	—	—
Total viral	1	—	2	4	34	59	51	110
Multiple etiologies	—	—	—	1	—	3	2	5
Confirmed etiology	2	33	6	7	93	260	110	370
Unknown etiology	3	31	22	30	279	529	444	973
Total 1999	5	64	28	37	372	789	554	1,343

[Return to top.](#)

[Table 16](#)

TABLE 16. Number of reported foodborne-disease outbreaks, by etiology and vehicle of transmission — United States, 2000

Etiology	Vehicle of transmission								
	Beef	Dairy	Eggs	Game	Pork	Poultry	Vegetables	Fruits and nuts	Grains
Bacterial									
<i>Bacillus cereus</i>	—	—	—	—	—	—	—	—	2
<i>Brucella</i>	—	—	—	—	—	—	—	—	—
<i>Campylobacter</i>	—	10	—	—	—	—	1	—	—
<i>Clostridium botulinum</i>	—	—	—	—	—	—	2	—	—
<i>Clostridium perfringens</i>	3	—	—	—	2	4	—	—	1
<i>Escherichia coli</i>	11	—	—	—	—	2	1	2	—
<i>Listeria monocytogenes</i>	—	1	—	—	—	1	—	—	—
<i>Salmonella</i>	1	—	19	—	4	12	4	3	—
<i>Shigella</i>	—	—	—	—	1	—	—	—	—
<i>Staphylococcus aureus</i>	2	—	—	—	3	2	1	—	1
<i>Streptococcus</i>	—	—	—	—	—	—	—	—	—
<i>Vibrio cholerae</i>	—	—	—	—	—	—	—	—	—
<i>Vibrioparae hemolyticus</i>	—	—	—	—	—	—	—	—	—
<i>Vibrio, other</i>	—	—	—	—	—	—	—	—	—
<i>Yersinia enterocolitica</i>	—	—	—	—	—	—	—	—	—
Other bacterial	—	—	—	—	—	—	—	—	—
Total bacterial	17	11	19	—	10	21	9	5	4
Chemical									
Ciguatera	—	—	—	—	—	—	—	—	—
Heavy metals	—	—	—	—	—	—	—	1	—
Mushroom toxin	—	—	—	—	—	—	—	—	—
Scombrotxin	—	—	—	—	—	—	—	—	—
Shellfish toxin	—	—	—	—	—	—	—	—	—
Other chemical	—	—	—	—	—	—	—	1	—
Total chemical	—	—	—	—	—	—	—	1	—
Parasitic									
<i>Anisakis</i>	—	—	—	—	—	—	—	—	—
<i>Cryptosporidium parvum</i>	—	—	—	—	—	—	—	—	—
<i>Cyclospora cayentanensis</i>	—	—	—	—	—	—	—	1	—
<i>Giardia intestinalis</i>	—	—	—	—	—	—	—	—	—
<i>Trichinella spiralis</i>	—	—	—	—	—	—	—	—	—
Total parasitic	—	—	—	—	—	—	—	1	—
Viral									
Astrovirus	—	—	—	—	—	—	—	—	—
Hepatitis A	—	—	—	—	—	—	1	1	—
Norovirus	1	—	—	—	2	1	8	6	—
Rotavirus	—	—	—	—	—	—	—	—	—
Total viral	1	—	—	—	2	1	9	7	—
Multiple etiologies	—	—	—	—	1	—	—	—	2
Confirmed etiology	18	11	19	—	13	22	18	14	6
Unknown etiology	25	11	6	—	14	39	23	7	22
Total 2000	43	22	25	—	27	61	41	21	28

TABLE 16. (Continued) Number of reported foodborne-disease outbreaks, by etiology and vehicle of transmission — United States, 2000

Etiology	Vehicle of transmission						Total	
	Oils and sugars	Finfish	Shellfish	Unclassifiable vehicle	Complex vehicle	Known vehicle		Unknown vehicle
Bacterial								
<i>Bacillus cereus</i>	—	1	—	—	3	6	2	8
<i>Brucella</i>	—	—	—	—	—	—	—	—
<i>Campylobacter</i>	—	—	—	—	—	11	4	15
<i>Clostridium botulinum</i>	—	—	—	—	—	2	—	2
<i>Clostridium perfringens</i>	—	—	—	1	9	20	2	22
<i>Escherichia coli</i>	—	—	—	2	4	22	10	32
<i>Listeria monocytogenes</i>	—	—	—	—	—	2	—	2
<i>Salmonella</i>	—	2	—	4	37	86	41	127
<i>Shigella</i>	—	—	—	2	6	9	3	12
<i>Staphylococcus aureus</i>	—	—	—	1	9	19	4	23
<i>Streptococcus</i>	—	—	—	—	—	—	—	—
<i>Vibrio cholerae</i>	—	—	—	—	—	—	—	—
<i>Vibrioparahaemolyticus</i>	—	—	3	—	1	4	—	4
<i>Vibrio, other</i>	—	—	—	—	—	—	—	—
<i>Yersinia enterocolitica</i>	—	—	—	—	—	—	—	—
Other bacterial	—	—	—	—	—	—	—	—
Total bacterial	—	3	3	10	69	181	66	247
Chemical								
Ciguatoxin	—	12	—	—	—	12	—	12
Heavy metals	—	—	—	—	—	1	—	1
Mushroom toxin	—	—	—	—	—	—	—	—
Scombrototoxin	—	20	—	—	—	20	—	20
Shellfish toxin	—	—	3	—	—	3	—	3
Other chemical	—	—	—	—	2	2	—	2
Total chemical	—	32	3	—	2	38	—	38
Parasitic								
<i>Anisakis</i>	—	—	—	—	—	—	—	—
<i>Cryptosporidium parvum</i>	—	—	—	—	1	1	—	1
<i>Cyclospora cayentanensis</i>	—	—	—	—	1	2	—	2
<i>Giardia intestinalis</i>	—	—	—	1	—	1	—	1
<i>Trichinella spiralis</i>	—	—	—	—	2	2	—	2
Total parasitic	—	—	—	1	4	6	—	6
Viral								
Astrovirus	—	—	—	—	—	—	—	—
Hepatitis A	—	—	1	1	5	9	3	12
Norovirus	—	1	2	4	49	74	89	163
Rotavirus	—	—	—	1	—	1	—	1
Total viral	—	1	3	6	54	84	92	176
Multiple etiologies	—	—	—	—	—	3	—	3
Confirmed etiology	—	36	9	17	129	312	158	470
Unknown etiology	1	27	16	34	326	551	396	947
Total 2000	1	63	25	51	455	863	554	1,417

[Return to top.](#)

[Table 17](#)

TABLE 17. Number of reported foodborne-disease outbreaks, by etiology and vehicle of transmission — United States, 2001

Etiology	Vehicle of transmission								
	Beef	Dairy	Eggs	Game	Pork	Poultry	Vegetables	Fruits and nuts	Grains
Bacterial									
<i>Bacillus cereus</i>	—	—	—	—	—	—	1	—	—
<i>Brucella</i>	—	1	—	—	—	—	—	—	—
<i>Campylobacter</i>	—	2	—	1	—	2	—	1	—
<i>Clostridium botulinum</i>	—	—	—	1	—	—	—	—	—
<i>Clostridium perfringens</i>	10	—	—	—	—	4	1	—	—
<i>Escherichia coli</i>	—	1	—	—	—	—	1	1	—
<i>Listeria monocytogenes</i>	—	—	—	—	—	1	—	—	—
<i>Salmonella</i>	1	4	10	—	5	9	3	6	—
<i>Shigella</i>	1	—	—	—	—	1	2	—	—
<i>Staphylococcus aureus</i>	—	—	—	—	6	2	—	—	—
<i>Streptococcus</i>	—	—	—	—	—	—	—	—	—
<i>Vibrio cholerae</i>	—	—	—	—	—	—	—	—	—
<i>Vibrioparahaemolyticus</i>	—	—	—	—	—	—	—	—	—
<i>Vibrio, other</i>	—	—	—	—	—	—	—	—	—
<i>Yersinia enterocolitica</i>	—	—	—	—	2	—	—	—	—
Other bacterial	—	—	—	—	—	—	—	—	—
Total bacterial	12	8	10	2	13	19	8	8	—
Chemical									
Ciguatoxin	—	—	—	—	—	—	—	—	—
Heavy metals	—	—	—	—	—	—	—	—	—
Mushroom toxin	—	—	—	—	—	—	—	—	—
Scombrototoxin	—	—	—	—	—	—	—	—	—
Shellfish toxin	—	—	—	—	—	—	—	—	—
Other chemical	—	—	—	1	—	—	—	—	—
Total chemical	—	—	—	1	—	—	—	—	—
Parasitic									
Anisakis	—	—	—	—	—	—	—	—	—
<i>Cryptosporidium parvum</i>	—	—	—	—	—	—	—	—	—
<i>Cyclospora cayentanensis</i>	—	—	—	—	—	—	1	—	—
<i>Giardia intestinalis</i>	—	—	—	—	—	—	—	—	—
<i>Trichinella spiralis</i>	—	—	—	1	—	—	—	—	—
Total parasitic	—	—	—	1	—	—	1	—	—
Viral									
Astrovirus	—	—	—	—	—	—	—	—	—
Hepatitis A	—	1	—	—	—	—	1	—	—
Norovirus	—	2	—	1	1	3	7	5	1
Rotavirus	—	—	—	—	—	—	—	—	—
Total viral	—	3	—	1	1	3	8	5	1
Multiple etiologies	1	—	—	—	1	1	—	—	—
Confirmed etiology	13	11	10	5	15	23	17	13	1
Unknown etiology	20	10	2	—	15	50	20	8	10
Total 2001	33	21	12	5	30	73	37	21	11

TABLE 17. (Continued) Number of reported foodborne-disease outbreaks, by etiology and vehicle of transmission — United States, 2001

Etiology	Vehicle of transmission						Total	
	Oils and sugars	Finfish	Shellfish	Unclassifiable vehicle	Complex vehicle	Known vehicle		Unknown vehicle
Bacterial								
<i>Bacillus cereus</i>	—	—	—	—	4	5	—	5
<i>Brucella</i>	—	—	—	—	—	1	—	1
<i>Campylobacter</i>	—	—	—	—	5	11	5	16
<i>Clostridium botulinum</i>	—	1	—	—	1	3	—	3
<i>Clostridium perfringens</i>	—	—	—	—	11	26	5	31
<i>Escherichia coli</i>	—	—	—	1	6	10	12	22
<i>Listeria monocytogenes</i>	—	—	—	—	—	1	—	1
<i>Salmonella</i>	—	—	1	8	90	77	34	111
<i>Shigella</i>	—	—	2	—	4	10	5	15
<i>Staphylococcus aureus</i>	—	—	—	—	14	22	1	23
<i>Streptococcus</i>	—	—	—	—	—	—	—	—
<i>Vibrio cholerae</i>	—	—	1	—	—	1	—	1
<i>Vibrio parahaemolyticus</i>	—	—	1	—	2	3	—	3
<i>Vibrio, other</i>	—	—	—	—	—	—	—	—
<i>Yersinia enterocolitica</i>	—	—	—	—	—	2	1	3
Other bacterial	—	—	—	—	—	—	—	—
Total bacterial	—	1	5	9	77	172	63	235
Chemical								
Ciguatoxin	—	24	—	—	—	24	—	24
Heavy metals	—	—	—	—	—	—	—	—
Mushroom toxin	—	—	—	—	—	—	—	—
Scombrototoxin	—	29	—	—	—	29	—	29
Shellfish toxin	—	—	—	—	—	—	—	—
Other chemical	—	—	—	—	—	1	—	1
Total chemical	—	53	—	—	—	54	—	54
Parasitic								
<i>Anisakis</i>	—	—	—	—	—	—	—	—
<i>Cryptosporidium parvum</i>	—	—	—	—	—	—	—	—
<i>Cyclospora cayentanensis</i>	—	—	—	—	—	1	1	2
<i>Giardia intestinalis</i>	—	—	—	—	—	—	1	1
<i>Trichinella spiralis</i>	—	—	—	—	1	2	—	2
Total parasitic	—	—	—	—	1	3	2	5
Viral								
Astrovirus	—	—	—	—	—	—	—	—
Hepatitis A	—	—	—	2	—	4	2	6
Norovirus	1	1	8	10	40	80	70	150
Rotavirus	—	—	—	—	—	—	—	—
Total viral	1	1	8	12	40	84	72	156
Multiple etiologies	—	—	—	—	4	7	2	9
Confirmed etiology	1	55	13	21	122	321	199	459
Unknown etiology	3	20	20	30	262	470	314	784
Total 2001	4	75	33	51	384	791	453	1,243

[Return to top.](#)

[Table 18](#)

TABLE 18. Number of reported foodborne-disease outbreaks, by etiology and vehicle of transmission — United States, 2002

Etiology	Vehicle of transmission								
	Beef	Dairy	Eggs	Game	Pork	Poultry	Vegetables	Fruits and nuts	Grains
Bacterial									
<i>Bacillus cereus</i>	—	—	—	—	—	2	—	—	1
<i>Brucella</i>	—	—	—	—	—	—	—	—	—
<i>Campylobacter</i>	—	4	—	—	—	2	1	—	—
<i>Clostridium botulinum</i>	—	—	—	—	—	—	—	—	—
<i>Clostridium perfringens</i>	8	—	1	—	2	5	—	—	—
<i>Escherichia coli</i>	6	—	—	—	—	—	3	—	1
<i>Listeria monocytogenes</i>	—	—	—	—	—	1	—	—	—
<i>Salmonella</i>	4	2	6	2	3	8	8	3	1
<i>Shigella</i>	—	—	—	—	—	—	1	—	—
<i>Staphylococcus aureus</i>	1	—	—	—	4	2	—	—	—
<i>Streptococcus</i>	—	—	—	—	—	—	—	—	—
<i>Vibrio cholerae</i>	—	—	—	—	—	—	—	—	—
<i>Vibrioparahaemolyticus</i>	—	—	—	—	—	—	—	—	—
<i>Vibrio</i> , other	—	—	—	—	—	—	—	—	—
<i>Yersinia enterocolitica</i>	—	—	—	—	1	—	—	—	—
Other bacterial	—	—	—	—	—	—	—	—	—
Total bacterial	19	6	7	2	10	20	13	3	3
Chemical									
Ciguatoxin	—	—	—	—	—	—	—	—	—
Heavy metals	—	—	—	—	—	—	—	—	—
Mushroom toxin	—	—	—	—	—	—	1	—	—
Scombrototoxin	—	—	—	—	—	—	—	—	—
Shellfish toxin	—	—	—	—	—	—	—	—	—
Other chemical	—	1	—	—	—	1	—	—	—
Total chemical	—	1	—	—	—	1	1	—	—
Parasitic									
<i>Anisakis</i>	—	—	—	—	—	—	—	—	—
<i>Cryptosporidium parvum</i>	—	—	—	—	—	—	—	—	—
<i>Cyclospora cayentanensis</i>	—	—	—	—	—	—	—	1	—
<i>Giardia intestinalis</i>	—	—	—	—	—	—	—	—	—
<i>Trichinella spiralis</i>	—	—	—	1	—	—	—	—	—
Total parasitic	—	—	—	1	—	—	—	1	—
Viral									
Astrovirus	—	—	—	—	—	—	—	—	—
Hepatitis A	—	—	—	—	—	—	—	—	—
Norovirus	4	2	—	—	—	2	11	1	—
Rotavirus	—	—	—	—	—	—	—	—	—
Total viral	4	2	—	—	—	2	11	1	—
Multiple etiologies	1	—	—	—	1	—	—	—	—
Confirmed etiology	24	9	7	3	11	23	25	5	3
Unknown etiology	20	7	7	—	15	52	19	4	11
Total 2002	44	16	14	3	26	75	44	9	14

TABLE 18. (Continued) Number of reported foodborne-disease outbreaks, by etiology and vehicle of transmission — United States, 2002

Etiology	Vehicle of transmission							Total
	Oils and sugars	Finfish	Shellfish	Unclassifiable vehicle	Complex vehicle	Known vehicle	Unknown vehicle	
Bacterial								
<i>Bacillus cereus</i>	—	—	—	—	4	7	—	7
<i>Brucella</i>	—	—	—	—	—	—	—	—
<i>Campylobacter</i>	—	—	—	—	4	11	2	13
<i>Clostridium botulinum</i>	—	2	—	—	1	3	—	3
<i>Clostridium perfringens</i>	—	—	—	—	13	29	2	31
<i>Escherichia coli</i>	—	—	—	3	7	20	6	26
<i>Listeria monocytogenes</i>	—	—	—	—	—	1	—	1
<i>Salmonella</i>	—	1	—	3	34	75	36	111
<i>Shigella</i>	—	—	—	—	2	3	6	9
<i>Staphylococcus aureus</i>	—	—	—	—	13	20	1	21
<i>Streptococcus</i>	—	—	—	—	—	—	—	—
<i>Vibrio cholerae</i>	—	—	—	—	—	—	—	—
<i>Vibrio parahaemolyticus</i>	—	—	2	—	—	2	—	2
<i>Vibrio, other</i>	—	—	—	—	—	—	—	—
<i>Yersinia enterocolitica</i>	—	—	—	1	—	2	1	3
Other bacterial	—	—	—	—	—	—	—	—
Total bacterial	—	3	2	7	78	173	54	227
Chemical								
Ciguatoxin	—	20	—	—	—	20	—	20
Heavy metals	—	—	—	—	—	—	—	—
Mushroom toxin	—	—	—	—	—	1	—	1
Scombrototoxin	—	21	—	—	—	21	—	21
Shellfish toxin	—	1	—	—	—	1	—	1
Other chemical	—	—	—	1	—	3	—	3
Total chemical	—	42	—	1	—	46	—	46
Parasitic								
<i>Anisakis</i>	—	—	—	—	—	—	—	—
<i>Cryptosporidium parvum</i>	—	—	—	—	—	—	2	2
<i>Cyclospora cayentanensis</i>	—	—	—	—	1	2	—	2
<i>Giardia intestinalis</i>	—	—	—	—	—	—	—	—
<i>Trichinella spiralis</i>	—	—	—	—	—	1	—	1
Total parasitic	—	—	—	—	1	3	2	5
Viral								
Astrovirus	—	—	—	—	—	—	1	1
Hepatitis A	—	—	—	—	1	1	6	7
Norovirus	—	—	2	10	62	94	105	199
Rotavirus	—	—	—	—	—	—	—	—
Total viral	—	—	2	10	63	95	112	207
Multiple etiologies	—	—	—	—	5	7	4	11
Confirmed etiology	—	45	4	18	147	324	172	496
Unknown etiology	1	21	23	34	289	503	331	834
Total 2002	1	66	27	52	436	827	503	1,330

[Return to top.](#)
[Table 19](#)

TABLE 19. Number of reported foodborne-disease outbreaks, by etiology and contributing factors*† — United States, 1998–2002

Etiology	Contamination factors								
	C1	C2	C3	C4	C5	C6	C7	C8	C9
Bacterial									
<i>Bacillus cereus</i>	—	—	—	—	—	3	—	—	—
<i>Brucella</i>	—	—	—	—	—	1	—	—	—
<i>Campylobacter</i>	—	—	—	—	—	18	7	—	10
<i>Clostridium botulinum</i>	—	—	—	—	—	5	1	—	—
<i>Clostridium perfringens</i>	—	—	—	—	—	14	1	—	5
<i>Escherichia coli</i>	—	1	—	—	—	33	4	1	14
<i>Listeria monocytogenes</i>	—	—	—	—	—	1	1	—	—
<i>Salmonella</i>	1	—	—	—	4	97	42	3	85
<i>Shigella</i>	—	—	—	—	—	1	—	—	1
<i>Staphylococcus aureus</i>	—	—	—	—	—	—	—	—	1
<i>Streptococcus</i>	—	—	—	—	—	—	—	—	—
<i>Vibrio cholerae</i>	—	—	—	—	—	—	1	1	—
<i>Vibrio parahaemolyticus</i>	1	—	—	—	—	7	4	2	2
<i>Vibrio</i> , other	—	—	—	—	—	—	—	—	—
<i>Yersinia enterocolitica</i>	—	—	—	—	—	—	—	—	2
Other bacterial	—	—	—	—	—	—	—	—	—
Total bacterial	2	1	—	—	4	180	61	7	120
Chemical									
Ciguatera	54	—	—	—	—	—	—	4	—
Heavy metals	—	—	1	—	—	1	—	—	—
Mushroom toxin	1	—	—	—	—	—	—	—	—
Scombrototoxin	43	—	—	—	—	1	1	—	1
Shellfish toxin	4	—	—	—	—	—	—	—	—
Other chemical	—	—	5	—	—	—	—	—	—
Total chemical	102	—	6	—	—	2	1	4	1
Parasitic									
<i>Anisakis</i>	—	—	—	—	—	—	—	—	—
<i>Cryptosporidium parvum</i>	—	—	—	—	—	1	1	—	—
<i>Cyclospora cayentanensis</i>	—	—	—	—	—	5	3	—	—
<i>Giardia intestinalis</i>	—	—	—	—	—	—	—	—	—
<i>Trichinella spiralis</i>	—	—	—	—	—	1	3	—	—
Total parasitic	—	—	—	—	—	7	7	—	—
Viral									
Astrovirus	—	—	—	—	—	—	—	—	—
Hepatitis A	—	—	—	—	1	—	1	—	1
Norovirus	—	—	—	—	1	21	18	3	9
Rotavirus	—	—	—	—	—	—	—	—	—
Total viral	—	—	—	—	2	21	19	3	10
Multiple etiologies									
Multiple etiologies	—	—	—	—	—	2	3	1	—
Confirmed etiology	104	1	6	—	6	212	91	15	131
Unknown etiology	42	1	20	3	7	133	38	10	251
Total 1998–2002	146	2	26	3	13	345	129	25	382

* More than one contributing factor might be reported per outbreak.

† See Appendix A for description of each factor.

TABLE 19. (Continued) Number of reported foodborne-disease outbreaks, by etiology and contributing factors*† — United States, 1998–2002

Etiology	Contamination factors						Outbreaks in which contamination factor reported
	C10	C11	C12	C13	C14	C15	
Bacterial							
<i>Bacillus cereus</i>	1	1	1	3	2	—	10
<i>Brucella</i>	—	—	—	—	—	—	1
<i>Campylobacter</i>	5	2	1	13	3	3	39
<i>Clostridium botulinum</i>	—	—	—	1	—	1	7
<i>Clostridium perfringens</i>	8	2	2	12	6	11	41
<i>Escherichia coli</i>	12	1	6	8	3	3	60
<i>Listeria monocytogenes</i>	—	—	—	1	—	3	6
<i>Salmonella</i>	37	4	64	82	17	23	260
<i>Shigella</i>	12	3	16	5	2	3	25
<i>Staphylococcus aureus</i>	17	5	30	9	5	6	50
<i>Streptococcus</i>	—	—	1	—	—	—	1
<i>Vibrio cholerae</i>	—	—	—	—	—	—	2
<i>Vibrioparahaemolyticus</i>	2	1	1	4	1	—	14
<i>Vibrio, other</i>	—	—	—	—	—	—	—
<i>Yersinia enterocolitica</i>	—	—	—	3	—	—	4
Other bacterial	—	—	—	1	—	—	1
Total bacterial	94	19	122	142	39	53	521
Chemical							
Ciguatera	—	—	—	—	—	—	58
Heavy metals	—	—	—	—	—	—	2
Mushroom toxin	—	—	—	—	—	—	1
Scombrototoxin	—	1	—	1	2	9	55
Shellfish toxin	—	—	—	—	—	—	4
Other chemical	—	—	—	—	1	1	6
Total chemical	—	1	—	1	3	10	126
Parasitic							
Anisakis	—	—	—	—	—	1	1
<i>Cryptosporidium parvum</i>	—	—	—	—	—	—	1
<i>Cyclospora cayentanensis</i>	—	—	—	—	—	1	6
<i>Giardia intestinalis</i>	1	—	2	1	1	—	2
<i>Trichinella spiralis</i>	—	—	—	—	—	—	4
Total parasitic	1	—	2	1	1	2	14
Viral							
Astrovirus	—	—	—	—	—	—	—
Hepatitis A	13	4	16	1	—	3	25
Norovirus	129	30	202	40	7	28	312
Rotavirus	—	—	—	—	—	—	—
Total viral	142	34	218	41	7	31	337
Multiple etiologies	2	1	7	6	2	4	14
Confirmed etiology	239	55	349	191	52	100	1,012
Unknown etiology	526	132	251	477	122	191	1,365
Total 1998–2002	765	187	600	668	174	291	2,377

* More than one contributing factor might be reported per outbreak.

† See Appendix A for description of each factor.

TABLE 19. (Continued) Number of reported foodborne-disease outbreaks, by etiology and contributing factors*† — United States, 1998–2002

Etiology	Proliferation factors							
	P1	P2	P3	P4	P5	P6	P7	P8
Bacterial								
<i>Bacillus cereus</i>	16	1	4	4	—	3	—	—
<i>Brucella</i>	—	—	—	—	—	—	—	—
<i>Campylobacter</i>	6	1	4	—	—	1	—	—
<i>Clostridium botulinum</i>	7	—	—	1	—	—	3	—
<i>Clostridium perfringens</i>	53	50	20	27	—	39	—	—
<i>Escherichia coli</i>	12	2	7	3	—	6	—	—
<i>Listeria monocytogenes</i>	—	—	1	—	—	—	—	—
<i>Salmonella</i>	110	26	53	33	3	28	—	1
<i>Shigella</i>	3	1	5	1	—	—	—	—
<i>Staphylococcus aureus</i>	42	17	13	17	1	15	—	1
<i>Streptococcus</i>	—	—	—	—	—	1	—	—
<i>Vibrio cholerae</i>	—	—	—	—	—	—	—	—
<i>Vibrio parahaemolyticus</i>	5	—	—	2	—	—	—	—
<i>Vibrio, other</i>	—	—	—	—	—	—	—	—
<i>Yersinia enterocolitica</i>	—	—	—	—	—	—	—	—
Other bacterial	1	—	—	—	—	—	—	—
Total bacterial	255	98	107	88	4	93	3	2
Chemical								
Ciguatoxin	—	—	—	—	—	—	—	—
Heavy metals	—	—	—	—	—	—	—	—
Mushroom toxin	—	—	—	—	—	—	—	—
Scombrototoxin	21	—	18	1	—	—	—	—
Shellfish toxin	—	—	—	—	—	—	—	—
Other chemical	—	—	—	—	—	—	—	—
Total chemical	21	—	18	1	—	—	—	—
Parasitic								
<i>Anisakis</i>	—	—	—	—	—	—	—	—
<i>Cryptosporidium parvum</i>	—	—	—	—	—	—	—	—
<i>Cyclospora cayentanensis</i>	—	—	—	—	—	—	—	—
<i>Giardia intestinalis</i>	—	—	—	—	—	—	—	—
<i>Trichinella spiralis</i>	—	—	—	—	—	—	—	—
Total parasitic	—	—	—	—	—	—	—	—
Viral								
Astrovirus	—	—	—	—	—	—	—	—
Hepatitis A	—	—	—	—	—	—	—	—
Norovirus	17	4	8	3	—	6	1	—
Rotavirus	—	—	—	—	—	—	—	—
Total viral	17	4	8	3	—	6	1	—
Multiple etiologies	9	4	3	2	—	3	—	—
Confirmed etiology	296	102	136	93	4	100	4	2
Unknown etiology	589	224	432	159	15	219	1	—
Total 1998–2002	894	330	571	254	19	322	5	2

* More than one contributing factor might be reported per outbreak.

† See Appendix A for description of each factor.

TABLE 19. (Continued) Number of reported foodborne-disease outbreaks, by etiology and contributing factors*† — United States, 1998–2002

Etiology	Proliferation factors				Outbreaks in which proliferation factor reported
	P9	P10	P11	P12	
Bacterial					
<i>Bacillus cereus</i>	—	—	—	—	21
<i>Brucella</i>	—	—	—	1	1
<i>Campylobacter</i>	—	—	1	4	14
<i>Clostridium botulinum</i>	—	1	—	1	11
<i>Clostridium perfringens</i>	4	—	—	5	103
<i>Escherichia coli</i>	—	—	—	3	24
<i>Listeria monocytogenes</i>	—	1	—	—	2
<i>Salmonella</i>	4	—	—	25	186
<i>Shigella</i>	—	—	—	3	10
<i>Staphylococcus aureus</i>	3	—	—	1	63
<i>Streptococcus</i>	—	—	—	—	1
<i>Vibrio cholerae</i>	—	—	—	—	—
<i>Vibrioparahaemolyticus</i>	—	—	—	—	6
<i>Vibrio, other</i>	—	—	—	—	—
<i>Yersinia enterocolitica</i>	—	—	—	1	1
Other bacterial	—	—	—	—	1
Total bacterial	11	2	1	44	444
Chemical					
Ciguatera	—	—	—	—	—
Heavy metals	—	—	—	—	—
Mushroom toxin	—	—	—	—	—
Scombrototoxin	—	—	—	3	31
Shellfish toxin	—	—	—	—	—
Other chemical	—	—	—	—	—
Total chemical	—	—	—	3	31
Parasitic					
<i>Anisakis</i>	—	—	—	—	—
<i>Cryptosporidium parvum</i>	—	—	—	—	—
<i>Cyclospora cayentanensis</i>	—	—	—	—	—
<i>Giardia intestinalis</i>	—	—	—	—	—
<i>Trichinella spiralis</i>	—	—	—	—	—
Total parasitic	—	—	—	—	—
Viral					
Astrovirus	—	—	—	—	—
Hepatitis A	—	—	—	1	1
Norovirus	—	—	—	1	28
Rotavirus	—	—	—	—	—
Total viral	—	—	—	2	29
Multiple etiologies	1	—	—	—	13
Confirmed etiology	11	2	1	50	511
Unknown etiology	41	—	4	40	1,133
Total 1998–2002	53	2	5	90	1,657

* More than one contributing factor might be reported per outbreak.

† See Appendix A for description of each factor.

TABLE 19. (Continued) Number of reported foodborne-disease outbreaks by etiology, and contributing factors*† — United States, 1998–2002

Etiology	Survival factors					Outbreaks in which survival factor reported	Outbreaks in which any contributing factor reported	Total
	S1	S2	S3	S4	S5			
Bacterial								
<i>Bacillus cereus</i>	—	3	—	—	1	4	24	37
<i>Brucella</i>	—	—	—	—	1	1	1	1
<i>Campylobacter</i>	15	—	—	1	6	21	39	61
<i>Clostridium botulinum</i>	5	1	2	—	4	10	12	12
<i>Clostridium perfringens</i>	33	41	—	5	4	61	102	130
<i>Escherichia coli</i>	20	2	—	3	4	27	68	140
<i>Listeria monocytogenes</i>	1	—	—	—	—	1	6	11
<i>Salmonella</i>	104	23	1	5	21	139	326	585
<i>Shigella</i>	—	—	—	—	4	4	27	67
<i>Staphylococcus aureus</i>	8	14	—	2	8	28	73	101
<i>Streptococcus</i>	—	—	—	—	—	—	1	1
<i>Vibrio cholerae</i>	—	—	—	—	—	—	2	3
<i>Vibrioparahaemolyticus</i>	3	—	—	—	1	4	15	25
<i>Vibrio, other</i>	—	—	—	—	—	—	—	1
<i>Yersinia enterocolitica</i>	—	—	—	—	1	1	4	8
Other bacterial	—	—	—	—	—	—	1	1
Total bacterial	189	84	3	16	55	301	695	1,184
Chemical								
Ciguatera	—	—	—	—	—	—	58	84
Heavy metals	—	—	—	—	—	—	2	2
Mushroom toxin	—	—	—	—	—	—	1	2
Scombrototoxin	1	—	—	—	3	4	74	118
Shellfish toxin	—	—	—	—	—	—	4	5
Other chemical	—	—	—	—	—	—	6	10
Total chemical	1	—	—	—	3	4	145	221
Parasitic								
<i>Anisakis</i>	—	—	—	—	—	—	1	1
<i>Cryptosporidium parvum</i>	—	—	—	—	—	—	1	4
<i>Cyclospora cayentanensis</i>	—	—	—	—	—	—	6	9
<i>Giardia intestinalis</i>	—	—	—	—	—	—	2	3
<i>Trichinella spiralis</i>	3	—	—	—	—	3	5	6
Total parasitic	3	—	—	—	—	3	15	23
Viral								
Astrovirus	—	—	—	—	—	—	—	1
Hepatitis A	—	—	—	—	—	—	25	50
Norovirus	5	4	—	—	10	18	319	657
Rotavirus	—	—	—	—	—	—	—	1
Total viral	5	4	—	—	10	18	344	709
Multiple etiologies	4	1	—	1	1	6	20	30
Confirmed etiology	198	88	3	16	69	327	1,238	2,167
Unknown etiology	171	161	2	16	74	369	1,834	4,480
Total 1998–2002	373	250	5	33	144	702	3,072	6,647

* More than one contributing factor might be reported per outbreak.

† See Appendix A for description of each factor.

[Return to top.](#)

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of pages found at these sites. URL addresses listed in *MMWR* were current as of the date of publication.

All *MMWR* HTML versions of articles are electronic conversions from typeset documents. This conversion might result in character translation or format errors in the HTML version. Users are referred to the electronic PDF version (<http://www.cdc.gov/mmwr>) and/or the original *MMWR* paper copy for printable versions of official text, figures, and tables. An original paper copy of this issue can be obtained from the Superintendent of Documents, U.S. Government Printing Office (GPO), Washington, DC 20402-9371; telephone: (202) 512-1800. Contact GPO for current prices.

** Questions or messages regarding errors in formatting should be addressed to mmwrq@cdc.gov.



Page last reviewed: 10/26/2006

Page last updated:

Content source: [Centers for Disease Control and Prevention](#)

[Home](#) [A-Z Index](#) [Policies](#) [Using this Site](#) [Link to Us](#) [Contact CDC](#)

Centers for Disease Control and Prevention 1600 Clifton Rd. Atlanta, GA 30333, USA
800-CDC-INFO (800-232-4636) TTY: (888) 232-6348 - [Contact CDC-INFO](#)

