

UC Davis

UC Davis Previously Published Works

Title

Survey of glucosinolate variation in leaves of Brassica rapa crops

Permalink

<https://escholarship.org/uc/item/9fq1w32x>

Journal

Genetic Resources and Crop Evolution: An International Journal, 57(7)

ISSN

1573-5109

Authors

Yang, Bo

Quiros, Carlos F.

Publication Date

2010-10-01

DOI

10.1007/s10722-010-9549-5

Peer reviewed

Survey of glucosinolate variation in leaves of *Brassica rapa* crops

Bo Yang · Carlos F. Quiros

Received: 15 October 2009 / Accepted: 10 March 2010 / Published online: 31 March 2010
© The Author(s) 2010. This article is published with open access at Springerlink.com

Abstract The breakdown products of glucosinolates (gsl) are biologically active secondary metabolites involved in plant defense and human nutrition. We identified and quantified 14 different gsl present in the young leaves of 82 different varieties of *Brassica rapa*, including the following crops: Chinese cabbage, broccolo, Pak choi and other leafy vegetables, turnip, sarson and rapeseed. We did not find crop specific gsl, but their quantity varied extensively among varieties and crops, except that the Chinese cabbage accessions tended to have similar gsl profile and amount. Gluc-onapin, glucobrassicinapin (aliphatic), neoglucobrassicin, glucobrassicin (indolic), and gluconaturtin (aromatic) are the predominant gsl in most of the varieties surveyed. We also found two gsl not commonly found in *B. rapa*, 2-methyl-2-propenyl and *n*-butyl. Their identities were confirmed by HPLC-MS. Most of the Chinese cabbages contain lower amount of aliphatic than indolic gsl, whereas broccolo, turnip and rapeseed all have much higher aliphatic gsl content than indolic gsl content. The predominant aliphatic gsl in most of the varieties contain 4-carbon side-chains. The lack of significant correlation

observed between the conversion of 3- to 4-carbon side-chain gsl and the conversion of 4- to 5-carbon side-chain gsl suggests that these two elongation cycles are probably under the control of two independent genes in *B. rapa*. The absence of glucoraphanin in all accessions indicates that only functional *Brgsl-Alk* alleles are present in *B. rapa*.

Keywords *Brassica rapa* · Brassicaceae · Glucosinolates · Secondary metabolites

Introduction

Glucosinolates (gsl) have been of considerable interest to plant biologists for many years. These nitrogen- and sulfur-containing plant secondary metabolites and their breakdown products are responsible for the distinctive flavor and taste of the plants and crops that contain them. Certain types of gsl derivatives, such as the isothiocyanate sulforaphane, have been found to possess biological activity as cancer-prevention agents in mammals, and others as plant defense compounds against insect and diseases (Mithen 2001).

In *Arabidopsis thaliana* (L.) Heynh., a *Brassica* relative, it was reported that gsl accumulation varies significantly among different organs and developmental stages, with regards to both composition and concentration. The highest gsl concentrations were found in young leaves and reproductive organs such

Electronic supplementary material The online version of this article (doi:10.1007/s10722-010-9549-5) contains supplementary material, which is available to authorized users.

B. Yang · C. F. Quiros (✉)
Department of Plant Sciences, University of California,
Davis, CA 95616, USA
e-mail: cfquiros@ucdavis.edu

as seeds, siliques, flowers and developing inflorescences (Brown et al. 2003). Exceptionally high concentration of gsl was reported in the sulfur-rich S-cells of the flower stalk (Koroleva et al. 2000). This was the first report of gsl accumulation localized to a specific cell type found just outside the phloem. Roots, stems and fully expanded leaves contain intermediate concentrations of gsl. The lowest concentrations were found in senescing rosette leaves (Brown et al. 2003, Petersen et al. 2002).

With the completion of the *A. thaliana* genome sequence, gsl research progressed rapidly in the past decade (Halkier and Gershenzon 2006). The availability of abundant functional genomic resources along with the large natural gsl variation among ecotypes in *A. thaliana* has greatly facilitated this research. Much of the core biosynthetic pathway of these compounds, their breakdown pathway and their regulation factors in *A. thaliana* has been elucidated. However, this knowledge is just starting to being transferred to various gsl-containing vegetable crops included in human diet. A few major genes involved in the aliphatic, methionine-derived gsl biosynthesis in *B. oleracea* have been genetically characterized and cloned (Li and Quiros 2002; Li and Quiros 2003; Gao et al. 2004). Gsl content have been surveyed in several crops such as broccoli, broccoli sprouts, cauliflower and oilseed rape among others (Barillari et al. 2005; Windsor et al. 2005; Font et al. 2005; Tian et al. 2005; Skutlarek et al. 2004; Branca et al. 2002).

The species *B. rapa* L. em. Metzg. includes a great array of crops commonly used in human diet, particularly in Asian cuisine, such as Chinese cabbage, pak-choi, among many other leafy vegetables, as well as turnip and rapeseed. Limited evaluations of gsl have been performed in different tissues of a few *B. rapa* crops with the objective of determining their potential biological activity (Siemens et al. 2002; Cipollini et al. 2003; Kang et al. 2006; Bellosta et al. 2007). Padilla et al. (2007) determined the gsl diversity of *B. rapa* varieties restricted to turnip greens from northwestern Spain. In the present study, we analyzed gsl profiles from young leaves of 82 different *B. rapa* accessions representing all major crops and constituting a diverse sample of the genetic, geographical and environmental range of this species. We found extensive variation in both the composition and total concentration of gsl in the accessions of this species.

Materials and methods

Plant material

Eighty two varieties (Table 1), representing most major crops of *B. rapa*, were sampled in the present study. All accessions come from the *Brassica* germplasm working collection at the Department of Plant Sciences, UC Davis. These include the following subspecies: *B. rapa* ssp. *pekinensis* (Lour.) Hanelt (Chinese cabbage), *B. rapa* ssp. *chinensis* (L.) Hanelt (pak choi), *B. rapa* ssp. *nipposinica* (L. H. Bailey) Hanelt (syn. *B. ruvo* Bailey) (broccoleto), *B. rapa* ssp. *rapifera* Metzg. (turnip), *B. rapa* ssp. *perviridis* L. H. Bailey (Japanese/Oriental tender green), *B. rapa* ssp. *oleifera* (DC.) Metzg. (rapeseed), *B. rapa* ssp. *dichotoma* (Roxb.) Hanelt (toria) and *B. rapa* ssp. *trilocularis* (Roxb.) Hanelt (Yellow sarson). Seeds from each accession were germinated in 12 × 20 × 1.5 inch trays (135 wells in each tray) and transferred into 6.5 × 6.5 × 7 inch pots at 3 weeks. The trays and pots were filled with “Super Soil” potting mix (Rod McLellan Co., San Mateo, CA) and the plants were grown in the greenhouse. The natural photoperiod was extended to 16 h by metal halide light bulbs (Agrosun 400 W, Philips). Temperature was maintained at 24°C day; 18°C night. The leaf tissue was harvested from each plant 6 weeks after germination. 200 mg of leaf material per sample was used for the gsl extraction. All samples for each variety were analyzed in triplicate in independent experiments.

Glucosinolate extraction and purification

The purification technique used was modified from the basic Sephadex/sulfatase *Arabidopsis* protocol previously described by Kliebenstein et al. (2001a, b). Each 200 mg fresh leaf tissue sample was frozen in liquid nitrogen and ground into fine powder in a 1.5 µl eppendorf tube with a mini pestle. 500 µl of 90% methanol was added to the fine powder, followed by incubation at 80°C for 10 min. After incubating at room temperature for 1 h, the tissue and protein were precipitated by centrifugation for 10 min at 3,500 rpm and the supernatant used for anion-exchange chromatography. Each well of a 96 well filter plate (ISC BioExpress) was loaded with Sephadex DEAE A-25 using the multiscreen filter

Table 1 List of *Brassica rapa* accession sampled in this study

Acc	Crop variety	Origin
1	Rapid cycling 121	Wisconsin, USA
2	Chinese cabbage b, Beijing Fanxinbai	Beijing, China
3	Chinese cabbage, Lubao F1 Hybrid	Beijing, China
4	Chinese cabbage, Zhongfeng F1 Hybrid	Beijing, China
5	Chinese cabbage, 1063, land race	Beijing, China
6	Chinese cabbage, 282 Michihli F-3X	California, USA
7	Chinese cabbage, 50	Beijing, China
8	Chinese cabbage, 536, land race	Beijing, China
9	Chinese cabbage, 582, land race	Beijing, China
10	Chinese cabbage, 76	Beijing, China
11	Chinese cabbage, Bianzao-26	Henan, China
12	Chinese cabbage, CC20-1	Henan, China
13	Chinese cabbage, CC20-2	Henan, China
14	Chinese cabbage, CC20-3	Henan, China
15	Chinese cabbage, CC20-4	Henan, China
16	Chinese cabbage, CC35-1	Henan, China
17	Chinese cabbage, CC35-2	Henan, China
18	Chinese cabbage, CC35-4	Henan, China
19	Chinese cabbage, CC35-5	Henan, China
20	Chinese cabbage, CC35-6	Henan, China
21	Chinese cabbage, China Pride #33 F1	Japan
22	Chinese cabbage, Early Jade Pagoda F1	Japan
23	Chinese cabbage, Early Top F1	Japan
24	Chinese cabbage, F1 #32	Japan
25	Chinese cabbage, Guang 90 E 16	Henan, China
26	Chinese cabbage, Hsia Shang (35 days)	Taiwan
27	Chinese cabbage, Kwan-Hoo Choi	Korea
28	Chinese cabbage, Market Pride	Japan
29	Chinese cabbage, Matsushima	Maine, USA
30	Chinese cabbage, Michihili Jade Pagoda F1	Japan
31	Ch. cabbage landrace	Taiwan
32	Chinese cabbage, Oriental King F1	Japan
33	Chinese cabbage, Ri29-3	Henan, China
34	Chinese cabbage, Spring Triumph #27 F1	Japan
35	Chinese cabbage, Tip top F1	Japan
36	Chinese cabbage, Tropical Delight F1	Japan
37	Chinese cabbage, Tropical Pride (F1)	Japan
38	Chinese cabbage, Winter Giant F1	Japan
39	Chinese cabbage, Winter Knight #17	Japan
40	Chinese cabbage, Wong Bok	Japan
41	Broccoletto, Sessantina A Cima Grande	Wellesbourne, UK
42	Broccoletto, Cima Di Rapa	Wellesbourne, UK
43	Japanese green, Shirona	Wellesbourne, UK
44	Oriental green, Koyona (Mizuna)	Maine, USA
45	Pak Choi, Lei-Choi	California, USA

Table 1 continued

Acc	Crop variety	Origin
46	Pak Choi, Japanese Greens, Taisai	Wellesbourne, UK
47	Pak Choi, Chinese Cabbage	Wellesbourne, UK
48	Turnip Destro	Japan
49	Turnip Kamo Kabu	Japan
50	Turnip, Kanamachi Kokabu	Japan
51	Turnip, Kenshin Kabu	Japan
52	Turnip, Leaf Mustard	Japan
53	Turnip, Market Express (F1)	Japan
54	Turnip unnamed	Wellesbourne, UK
55	Turnip, Navet Rave De Treignac	Wellesbourne, UK
56	Turnip, Ndzawana	Japan
57	Turnip, Togari Kabu	Japan
58	Turnip, Zaruishi	Japan
59	Turnip Rapa Palla di Neve	Wellesbourne, UK
60	Turnip Royal Crown #1 (F1)	Japan
61	Turnip Royal Crown #2 (F1)	Japan
62	Turnip Tokyo Market Second Early	Japan
63	Turnip Tokyo Top (F1)	Japan
64	Turnip White Knight (F1)	Japan
65	Turnip Yorii Spring	Japan
66	Wild rapeseed 770010	Japan
67	Wild rapeseed 770016	Japan
68	Wild rapeseed 770024	Japan
69	Wild rapeseed 770072	Japan
70	Wild rapeseed 771003	Turkey
71	Wild rapeseed 771042	Sweden
72	Wild rapeseed 771048	Afghanistan
73	Wild rapeseed 771059	Iran
74	Wild rapeseed 771072	Thailand
75	Wild rapeseed 771073	USSR
76	Wild rapeseed 771091	Montana, USA
77	Wild rapeseed 770140	Pakistan
78	Wild rapeseed 771006	India
79	Wild rapeseed 771148	USSR
80	Toria 770278K940	Pakistan
81	Yellow Sarson 770407K899	Pakistan
82	Yellow Sarson R2L	Wisconsin, USA

plate column loader (Millipore). 300 μ l of water was added to each well and allowed to equilibrate for 1 h. The filter plate was placed on top of a deep well 96 well plate. After water was spun out to the plate by centrifugation for 2 min at 1,000 rpm, 200 μ l of the supernatant was added to the 96-well Sephadex columns and the liquid spun into the plate by centrifuging for 3 min at 1,200 rpm at room

temperature. All the liquid in the plate was discarded. Then the Sephadex columns were washed one time with 175 μ l of 90% methanol, and two times with 150 μ l of water. To desulfate gsl in the column, 10 μ l of sulfatase solution and 70 μ l of water were added to each column and the plate placed in the dark for overnight sulfatase incubation at room temperature. Desulfo-gsl were eluted by centrifuging for 3 min at

1,200 rpm at room temperature placing a shallow-well 2 ml 96-well plate (Agilent Technologies) in the bottom of the filter column plate.

HPLC analysis

Twenty microliters of the gsl extract was run on a 5- μ m column (Lichrospher 100 RP-18 end capped, 250 mm \times 4.6 mm, Alltech) on an Agilent 1100 series High Performance Liquid Chromatography (HPLC). Compounds were detected by UV at 229 nm and separated utilizing the following program with a water-acetonitrile gradient at a constant 1.00 ml min⁻¹ flow rate. The program included a 5-min gradient from 1.5 to 7.0% (v/v) acetonitrile, a 5 min gradient from 7.0 to 25.0% (v/v) acetonitrile, a 4 min gradient from 25.0 to 80.0% (v/v) acetonitrile, 3 min at 80.0% (v/v) acetonitrile, a 3 min gradient from 80.0 to 35.0% (v/v) acetonitrile, and a final 3.5 min gradient from 35.0 to 1.5% (v/v) acetonitrile.

Glucosinolate identification and quantification

Identification of most of the HPLC peaks was based on a comparison of retention time and UV absorption spectrum as determined on a diode-array detector with those of purified standards. To quantify the

amount of gsl we used the standard methods reported by Brown et al. (2003). Briefly, we ran the standard sinigrin (Sigma-Aldrich) for each independent experiment to calibrate the instrument. We produced a standard curve using the sinigrin standard at a series of concentrations and the slope was computed by performing a regression analysis. The quantification of each individual gsl present in the sample was calculated from HPLC peak areas using the sinigrin standard as a reference. Then, the data were corrected using published UV response factors for different types of gsl (ISO-9167, 1992), and the response factors for indolic gsl were originated from Fiebig and Arens (1992). Quantities results are given as μ mol g FW⁻¹ tissue.

The value shown for each variety is based on the mean of the three extractions for that variety. The list of compound identities based on HPLC is shown in Table 2. The analysis of variance *F*-test and student's *t*-test were used for sample means comparisons among and between different crops or different gsl (Steel et al. 1997).

HPLC-MS/MS assay

The identity of three of the gsl (No. 5, No. 6 and No. 8) was further confirmed by a Finnigan LTQ 2D

Table 2 List of glucosinolates found in 82 varieties of *Brassica rapa*

Ret. time	Abbr.	Common name	Chemical name	Type
7	4OHB	Progoitrin	2-Hydroxy-3-butenyl glucosinolate	Aliphatic glucosinolate
7.5	3PREY	Sinigrin	2-Propenyl glucosinolate	Aliphatic glucosinolate
8.6	5OHP	Gluconapoleiferin	2-Hydroxy-4-pentenyl glucosinolate	Aliphatic glucosinolate
8.8	5MSO	Glucoalyssin	5-Methylsulphinylpentyl glucosinolate	Aliphatic glucosinolate
9	4CMP		2-Methyl-2-propenyl glucosinolate	Aliphatic glucosinolate
10.5	4BTEY	Gluconapin	3-Butenyl glucosinolate	Aliphatic glucosinolate
11.5	4OHI3M	4-Hydroxyglucobrassicin	4-Hydroxy-3-indolymethyl glucosinolate	Indolic glucosinolate
11.7	4BTY	Glucocochlearin	<i>n</i> -Butyl glucosinolate	Aliphatic glucosinolate
12.5	5PTEY	Glucobrassicinapin	4-Pentenyl glucosinolate	Aliphatic glucosinolate
13	4MTB	Glucoerucin	4-Methylthiobutyl glucosinolate	Aliphatic glucosinolate
13.3	I3M	Glucobrassicin	3-Indolymethyl glucosinolate	Indolic glucosinolate
13.8	GNST	Gluconasturtiin	2-Phenylethyl glucosinolate	Aromatic glucosinolate
14.2	4MOI3M	4-Methoxyglucobrassicin	4-Methoxy-3-indolymethyl glucosinolate	Indolic glucosinolate
14.5	NMOI3M	Neoglucobrassicin	<i>N</i> -methoxy-3-indolymethyl glucosinolate	Indolic glucosinolate

The retention time of the peak when it shows in the 20-min HPLC program described in experimental; The abbreviation name of the compound used in supplementary Table 1

linear Ion Trap Mass Spectrometer system (Thermo Electron Corporation). The LC-MS/MS system consisted of a Finnigan Surveyor LC pump which has a 4 μm (Synergi Hydro RP 80, 150 mm \times 3 mm) column, a Finnigan Surveyor LC autosampler, a Finnigan Surveyor PDA detector, and a Finnigan LTQ mass spectrometer.

HPLC separation was performed at a constant flow rate of 0.4 ml min⁻¹ with a gradient from 0 to 100% of solvent B in 30 min before column re-equilibration. Solvent A was 13 mM ammonium acetate (pH 5.5) and solvent B was methanol. A linear-gradient mobile phase from 100% (v/v) of solvent A to 15% (v/v) of solvent B in 3 min, to 55% of solvent B (v/v) in 4 min, to 100% (v/v) of solvent B in 13 min, 2 min at 100% (v/v) of solvent B, and returned to 100% (v/v) of solvent A in 8 min was used for elution.

Both positive and negative ion tandem mass spectrometry (MS/MS) were conducted to detect gsl with selected reaction monitoring (SRM). The scan event cycle used a full scan mass spectrum with a range of 250.00 to 900.00 mass/charge (m/z) and two corresponding data-dependent MS/MS events. The most intense ions detected during full scan MS triggered data dependent scanning. Data dependent scanning was performed without the use of a parent ion list. The microscan count for full and MS/MS scan events were set to unity and a repeat count for dynamic exclusion was set to three for verification. MS/MS activation parameters used an isolation width of 2.0 m/z, minimum signal threshold of 500.0 counts, default charge state of 2, normalized collision energy of 20.0, an activation Q of 0.250, and an activation time of 30 ms.

Data interpretation

The data from HPLC-MS/MS were processed using *Qual Browser* of *XCalibur* (Thermo Electron, San Jose, CA), which provides accurate mass thresholds to filter data. In addition, chemical formula calculator was used to provide chemical formula and saturation values for product ions of gsl. Comparisons between theoretical and experimental product ion spectra further aided in the identification of the gsl structures. The predictive structure was drawn using *ACDLABS* 8.0. This determination was done at the metabolomics facility of the UCD Genome Center by Dr. Vladimir Tolstikov.

Results and discussion

Natural variation of glucosinolate composition

To survey natural variation in the gsl of *B. rapa*, we identified and quantified 14 different gsl present in the young leaves of the different varieties sampled (Fig. 1; Supplementary Tables 1, 2). We found a total of nine aliphatic gsl, four indolic gsl and one aromatic gsl. The predominant aliphatic gsl in most varieties contain four-carbon (4-C) side-chains, except for a rapeseed accession from USSR (Acc #79), which has predominantly sinigrin (No. 2), a three-carbon (3-C) side chain gsl. The composition of indolic gsl was fairly consistent in most of the varieties except a rapeseed accession from Sweden (Acc #71), which lacked this type of gsl. 2-phenylethyl gsl, also named gluconaturtiin (No. 12), was the only aromatic gsl found in these varieties. All the varieties except a rapeseed accession from Pakistan (Acc #77) have this aromatic gsl (Supplementary Table 1).

The identity of three of the compounds, with retention times at 9.8, 10.5 and 11.7 min, was in doubt, so they were further analyzed by HPLC-MS. Turnip 'Kenshin Kabu' (Acc #51), Turnip 'Kamo Kabu' (Acc #49) and Turnip 'Royal Crown' #1 (Acc #60) were used for this determination because they have high content of these compounds (Supplementary Table 1). All three compounds are 4-C aliphatic gsl and their identities could be ascertained based on retention time, UV spectrum and mass spectrum. The results indicate that they are 2-methyl-2-propenyl gsl (RT = 9.8), 3-butenyl gsl (RT = 10.5) and *n*-butyl gsl (RT = 11.7). 2-Methyl-2-propenyl gsl (No. 5) and *n*-butyl gsl (No. 8) are not commonly found in *Brassica* species (Kiddle et al. 2001). The genetic analyses for these compounds have not yet been performed in any species because of their low amounts and their rarity. In this survey, turnip 'Royal Crown #1' (Acc #62) was found to have a high amount of 2-methyl-2-propenyl gsl (No. 5); whereas turnip 'Tokyo Top' (Acc #65) has a high amount of *n*-butyl gsl (No. 8). These two varieties could be used to determine the genetic basis for the synthesis of these two compounds. The other compound, 3-butenyl gsl (No. 6), was commonly found in *B. rapa* in previous reports (Kang et al. 2006; Bellosta et al. 2007; Padilla et al. 2007).




Fig. 1 Relative content of aliphatic, indolic and aromatic gsl accumulation in young leaves of 82 varieties of *B. rapa*. The variety names are listed in Table 1. The empty region of each bar depicts the average content of total aromatic gsl, the filled

region depicts the average content of total indolic gsl the diagonal line region depicts the average content of total aliphatic gsl of each variety

The identities of the main gsl compounds in *B. rapa* reported in this survey (gluconapin, No. 6; glucobrassicin, No. 11; and gluconasturtiin, No. 12) are consistent with those reported in previous studies (Rosa et al. 1997; Ciska et al. 2000; Skutlarek et al. 2004; Kang et al. 2006; Padilla et al. 2007). Kang et al. (2006) studied in 23 accessions of Chinese cabbage the factors affecting the phenotypic variation of gsl content in two different environments. They detected and quantified only four major gsl: glucobrassicin (No. 11), gluconasturtiin (No. 12), neoglucobrassicin (No. 14) and progoitrin (No. 1). Our results included the quantification of each individual gsl, even those with low concentration for 82 accessions representing most crops in this species. For all these accessions, we found that the major aliphatic gsl

present was gluconapin (No. 6). However, similar to the findings of Kang et al. (2006), progoitrin (No. 1) was in higher amount than gluconapin (No. 6). in the Chinese cabbage accessions we sampled. Padilla et al. (2007) found that gluconapin (No. 6) and glucobrassicinapin (No. 9) were the most abundant aliphatic gsl in turnip greens varieties from northwestern Spain. They found that indolic and aromatic gsl concentrations were low and there were fairly constant in these varieties, which is consistent with our results. In any case, these differences are not surprising considering that age of the plants, organ sampled and growing conditions will affect gsl concentration. Our samples were done at 6 weeks in leaves of plants grown at the greenhouse. It is expected that changes in gsl concentration but not profile will occur in older plants

and those grown in the field, specially for indolic gsl that are the most affected by environment (Kang et al. 2006).

Natural variation of glucosinolate concentration

Gsl quantity varied extensively among varieties and crops (Fig. 1; Supplementary Table 1). The global mean of total aliphatic gsl was $0.697 \mu\text{mol g FW}^{-1}$ leaf tissue and the global mean of total indolic and aromatic gsl were 0.213 and $0.035 \mu\text{mol g FW}^{-1}$ leaf tissue, respectively. Turnip ‘Tokyo Top’ (Acc #63) had the highest content of aliphatic gsl, $6.823 \mu\text{mol g FW}^{-1}$ leaf tissue. In contrast, Chinese cabbage ‘Matsushima’ (Acc #29) had only $0.011 \mu\text{mol g FW}^{-1}$ and Chinese Cabbage Acc #31, an unknown variety from Taiwan, lacked aliphatic gsl (Fig. 1). The differences of indolic gsl quantities among accessions are not as dramatic as that observed for of aliphatic gsl. The highest content of indolic gsl was $0.820 \mu\text{mol g FW}^{-1}$ leaf tissue from Chinese cabbage ‘Wong Bok’ (Acc #40); the lowest one was $0.004 \mu\text{mol g FW}^{-1}$ leaf tissue from an oilseed Sarson accession from Pakistan (Acc #81). For aromatic gsl, the highest amount was $0.202 \mu\text{mol g FW}^{-1}$ leaf tissue, also from Chinese cabbage ‘Wong Bok’ (Acc #40); the lowest one was $0.001 \mu\text{mol g FW}^{-1}$ leaf tissue from oilseed sarson (Acc #81) (Fig. 1).

Aliphatic gsl with 3-C, 4-C and 5-C side-chains were found in the accessions sampled, with variable concentrations for each class (Supplementary Table 1). Only fourteen of the accessions analyzed have 3-C side-chain gsl, with an average content of $0.014 \mu\text{mol g FW}^{-1}$ leaf tissue. The crop types of these accessions included Chinese cabbage, Japanese greens, Pak choi, turnip, and rapeseed. Rapeseed accession Acc #79 had the highest content of 3-C side chain gsl ($0.833 \mu\text{mol g FW}^{-1}$ leaf tissue) but very low content of 4-C side chain gsl ($0.017 \mu\text{mol g FW}^{-1}$ leaf tissue). The global mean of 4-C side-chain gsl in all accessions was $0.470 \mu\text{mol g FW}^{-1}$ leaf tissue, the predominant class of aliphatic gsl in the accessions sampled. The highest amount observed was $5.219 \mu\text{mol g FW}^{-1}$ leaf tissue for a doubled haploid ‘rapid cycling’ line from Wisconsin (Acc #1) whereas Acc #31, a Chinese cabbage variety, lacked completely 4-C side chain gsl. The 5-C side-chain gsl quantities are lower than those for the 4-C side-chain gsl in most of the varieties, with the global mean of $0.213 \mu\text{mol g FW}^{-1}$ leaf tissue, except for some

varieties of Chinese cabbage (Acc# 40), turnip (Acc #53 and 54), and rapeseed (Acc #68 and 79). Acc #63, turnip ‘Tokyo Top’, had the highest content of 5-C side-chain gsl, $3.324 \mu\text{mol g FW}^{-1}$ leaf tissue, while its 4-C side-chain gsl content ($3.499 \mu\text{mol g FW}^{-1}$ leaf tissue) was also relatively high compared to other accessions.

3-Butenyl gsl (gluconapin, No. 6) and 4-pentenyl gsl (glucobrassicinapin, No. 9) were the predominant aliphatic gsl in most of the *B. rapa* varieties. The means of these two compounds are 0.406 and $0.205 \mu\text{mol g FW}^{-1}$ leaf tissue, respectively, constituting ~58 and 29% of total aliphatic gsl in leaves. *N*-methoxy-3-indolymethyl gsl (neoglucobrassicin, No. 14) and 3-indolymethyl gsl (glucobrassicin, No. 11) were the predominant indolic gsl in most varieties. The means of these two compounds are 0.120 and $0.085 \mu\text{mol g FW}^{-1}$ leaf tissue, respectively, which constitute ~56 and 40% of total indolic gsl.

Crop specificity

The main crops surveyed in this study include Chinese cabbage, broccoleto, Pak choi and other leafy vegetables, turnip, sarson and rapeseed (Table 1; Fig. 1). We did not find crop-specific gsl, however, after testing crops which included more than ten varieties, such as Chinese cabbage, turnip and rapeseed, we found significant differences in the mean for total gsl ($P \ll 0.01$). Turnip has higher gsl content than Chinese cabbage and rapeseed, but the difference between the last two crops was not significant. The mean total aliphatic gsl in Chinese cabbage, turnip and rapeseed was significantly different ($P \ll 0.01$). Total aliphatic gsl of Chinese cabbage was significantly lower than that of rapeseed and turnip ($P = 0.05$). However, the total aliphatic gsl of rapeseed and turnip was not significantly different. The methylsufinylalkyl/alkenyl (MSO/ALK) means were also significantly different in these three crops ($P < 0.05$), but the difference was only significant between that of Chinese cabbage and turnip ($P < 0.05$).

In general, aliphatic gsl content in Chinese cabbage is significantly lower than indolic gsl content ($P < 0.01$). Pak Choi does not have a high amount of aliphatic gsl either, although its aliphatic gsl content is higher than its indolic gsl content (Fig. 1; Supplementary Table 1). Broccoleto, turnip and rapeseed all have significantly higher aliphatic gsl content than indolic

gsl content ($P \ll 0.01$), and the main aliphatic gsl for broccoleto and turnip are both 4-C and 5-C side-chain gsl. Some of the rapeseed varieties tend to have high content of 3-C side-chain gsl. Compared to other crops, Chinese cabbage and turnip have significantly higher content of aromatic gsl ($P < 0.01$).

Crop domestication

Although we did not find crop specific gsl, there were significant differences in total and aliphatic gsl content. These might be related to the domestication of some of these crops considering that gsl derivatives influence their taste and aroma. Chinese cabbage originated in China over 6,000 years ago and was introduced to Korea in the fifteenth century and into Japan in the early twentieth century. It has been suggested that Chinese cabbage may have resulted from hybridization between pak-choi and turnip (Li 1982; Song et al. 1988). The exact place where turnip was domesticated is unknown. It either originated from “primitive types” in central Asia or derived from two independent domestication events in Europe and Asia (McGrath and Quiros 1992). Rapeseed was distributed from northern Europe to China and Korea as early as 2,000 years ago, with the primary center of diversity in the Himalayan region (Sovero 1993). Thus geographically separate domestication of these crops might explain in part the differences observed in their gsl content.

Glucosinolate biosynthetic pathway controlling loci

Our understanding on the genetic control of gsl composition is fairly extensive thanks to the model plant *A. thaliana* (Halkier and Gershenzon 2006). Most of the loci controlling the biosynthetic pathway for aliphatic gsl in *A. thaliana* and *B. oleracea* have been described previously (Kroymann et al. 2001; Kliebenstein et al. 2001a, b; Li and Quiros 2002, 2003; Gao et al. 2004; Textor et al. 2004; Field et al. 2004). Three partially redundant *MAM* (methylthioalkyl malate synthase) family genes (*MAM1*, *MAM2* and *MAM-L*) in the *GS-Elong* locus control the variation in side chain length of aliphatic gsl in *A. thaliana*. Functional analysis demonstrated that *MAM1* catalyze the first two elongation cycles for the synthesis of 3-C and 4-C gsl. *MAM-L* carries out the subsequent

elongation cycles forming 5 to 8-C long chain gsl. *MAM2* is assumed to control the 3-C gsl synthesis, but this activity has not been functionally demonstrated (Textor et al. 2004; Field et al. 2004). In *B. oleracea*, *Bogsl-Elong* and *Bogsl-Pro* are suggested to control the elongation cycle (Li and Quiros 2002; Gao et al. 2004). *B. rapa* is a rich source of 5-C aliphatic gsl. This leads to the hypothesis that *B. rapa* also has orthologs for all these three genes in the *MAM* family.

The effect of three loci on side chain modification of aliphatic gsl in *B. rapa* could be ascertained for the 82 varieties tested. *Brgsl-OX* controls the conversion of methylthioalkyl to methylsulfinylalkyl gsl. Most varieties carry out this conversion efficiently and typically contain at least as much methylsulfinylalkyl as methylthioalkyl gsl in the leaves. However, Acc #4, 13, 37, 40, 46, 58, 64 and 76 all have significantly higher ($P < 0.01$) concentrations of methylthioalkyl than methylsulfinylalkyl gsl (Supplementary Table 2), indicating that they are impaired in this conversion and presumably contain a different *Brgsl-OX* allele than the other varieties. *Brgsl-Alk* is responsible for the conversion of methylsulfinylalkyl to alkenyl gsl. Most varieties carry out this conversion also very efficiently and typically contain at least twice as much alkenyl as methylsulfinylalkyl gsl in the leaves except for Acc #14 and 22 ($P < 0.01$) (Supplementary Table 2). *Brgsl-OH* controls the production of hydroxyalkenyl gsl (D. Kliebenstein, personal communication). Most varieties carry out this conversion inefficiently and typically contain at most half as much hydroxyalkenyl as alkenyl gsl in the leaves. However, Acc #2, 14, 22, 28, 30, 38, 55 and 76 all have significantly higher concentrations of hydroxyalkenyl than alkenyl gsl ($P < 0.01$) (Supplementary Table 2), suggesting that they may contain a more efficient *Brgsl-OH* allele than the other varieties.

Sulforaphane, the isothiocyanate derived from 4-methylsulphanylbutyl gsl (glucoraphanin) commonly found in broccoli, is of great interest because of its role of inducing Phase II enzymes exerting blocking effects on carcinogens (Mithen 2001). However, contrary to broccoli, none of the accessions in the survey have glucoraphanin, which indicates that only functional *Brgsl-Alk* alleles are present in *B. rapa*. Conventional plant breeding, including the synthesis of artificial *B. napus* by hybridizing *B. rapa* and *B. oleracea*, could be used to replace functional *Brgsl-Alk* alleles with their non-functional homologs

from *B. oleracea* (broccoli). Alternatively, other approaches to develop a variety of *B. rapa* containing this desirable compound is to produce *Brgsl-Alk* knockout lines to efficiently accumulate glucoraphanin in the side-chain modification pathway, or use gene silencing methods such as RNAi to accomplish the same objective. The genetic approaches described above could also be applied to manipulate content of other gsl. For instance, disrupting the functional *Brgsl-OH* allele in *B. rapa* could abolish the accumulation of the detrimental compound progoitrin (No. 1). The study of natural gsl variation in *B. rapa* provides valuable information for answering questions about the biosynthesis, evolution and function of these interesting natural products in this species.

Conclusions

The main contributions of this study are: (1) Determination of gsl profiles and quantification of these compounds in representatives of most crops of in *B. rapa*. (2) Identification of new compounds not reported before for *B. rapa*. These include 2-methyl-2-propenyl (No. 5) and *n*-butyl gsl (No. 8), present in some of the varieties. (3) Although there are no crop specific gsl, there are significant differences in total amounts for these compounds. For example the leafy Chinese cabbages contain higher indolic gsl content than the rest of the *B. rapa* crops, whereas broccoleto and two oilseed vegetables, turnip and rapeseed, all have much higher aliphatic gsl content than the rest of the crops.

Acknowledgments The research was supported by USDA-IFAFS grant #00-52100-9683 “Development of Genomic Tools and Resources for *Brassica*”, and by the National Research Initiative of the USDA Cooperative State Research, Education and Extension Service, grant number #200x-35301-15886 “Cloning and characterization of the major glucosinolate genes involved in the aliphatic glucosinolate pathway of *Brassica* crops”). We acknowledge the assistance from Mr. Vincent D’Antonio raising the plants, to Dr. Dan Qiu for gsl analysis and to Mr. J. Erron Haggard for proofreading the manuscript. We also thank Dr. Daniel Kliebenstein and Dr. Vladimir Tolstikov for aid in compound identification.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Barillari J, Iori R, Rollin P, Hennion F (2005) Glucosinolates in the subantarctic crucifer kerguelen cabbage (*Pringlea antiscorbutica*). *J Nat Prod* 68:234–236
- Bellosta N, Sorensen JC, Sorensen H (2007) Profiling glucosinolates in vegetative and reproductive tissues of four *Brassica* species of the U-triangle for their biofumigation potential. *J Sci Food Agric* 87:1586–1594
- Branca F, Li G, Goyal S, Quiros CF (2002) Survey of aliphatic glucosinolates in Sicilian wild and cultivated Brassicaceae. *Phytochemistry* 59:717–724
- Brown PD, Tokuhisa JG, Reichelt M, Gershenzon J (2003) Variation of glucosinolate accumulation among different organs and developmental stages of *Arabidopsis thaliana*. *Phytochemistry* 62:471–481
- Cipollini DF, Busch JW, Stowe KA, Simms EL, Bergelson J (2003) Genetic variation and relationships of constitutive and herbivore-induced glucosinolates, trypsin inhibitors, and herbivore resistance in *Brassica rapa*. *J Chem Ecol* 29:285–302
- Ciska E, Martyniak-Przybyszewska B, Kozłowska H (2000) Content of glucosinolates in cruciferous vegetables grown at the same site for two years under different climatic conditions. *J Agric Food Chem* 48:2862–2867
- Fiebig HJ, Arens M (1992) Glucosinolates (HPLC method)—survey by a working party of the Dgf, 128th report—German standard methods for investigation of fats, fatty products, tensides and related materials, 98th report—analysis of fat raw-materials. 12. Fett Wissenschaft Technologie—Fat Science Technology, vol 94, pp 199–203
- Field B, Cardon G, Traka M, Botterman J, Vancanneyt G, Mithen R (2004) Glucosinolate and amino acid biosynthesis in *Arabidopsis*. *Plant Physiol* 135:828–839
- Font R, Rio-Celestino MD, Cartea E, Haro-Bailon AD (2005) Quantification of glucosinolates in leaves of leaf rape (*Brassica napus* ssp. *pabularia*) by near-infrared spectroscopy. *Phytochemistry* 66:175–185
- Gao M, Li G, Yang B, McCombie BWR, Quiros CF (2004) Comparative analysis of a *Brassica* BAC clone containing several major aliphatic glucosinolate genes with its corresponding *Arabidopsis* sequence. *Genome* 47:666–679
- Halkier BA, Gershenzon J (2006) Biology and biochemistry of glucosinolates. *Ann Rev Plant Biol* 57:303–333
- ISO 9167-1 (1992) Rapeseeds-determination of glucosinolates content. Part 1. Method using high performance liquid chromatography. International Organization for Standardization. Geneva, Switzerland
- Kang JY, Ibrahim KE, Kim DH, Kang WJ, Juvik JA (2006) Genetic and environmental variation of glucosinolate content in Chinese cabbage. *HortScience* 41:1382–1385
- Kiddle G, Bennett RN, Botting NP, Davidson NE, Robertson AAB, Wallsgrave RM (2001) High-performance liquid chromatographic separation of natural and synthetic desulphoglucosinolates and their chemical validation by UV, NMR and chemical ionisation-MS methods. *Phytochem Anal* 12:226–242
- Kliebenstein DJ, Kroymann J, Brown P, Figuth A, Pedersen D, Gershenzon J, Mitchell-Olds T (2001a) Genetic control of

- natural variation in *Arabidopsis* glucosinolate accumulation. *Plant Physiol* 126:811–825
- Kliebenstein DJ, Lambrix VM, Reichelt M, Gershenzon J, Mitchell-Olds T (2001b) Gene duplication in the diversification of secondary metabolism: tandem 2-oxoglutarate-dependent dioxygenases control glucosinolate biosynthesis in *Arabidopsis*. *Plant Cell* 13:681–693
- Koroleva OA, Davies A, Deeken R, Thorpe MR, Tomos AD, Hedrich R (2000) Identification of a new glucosinolate-rich cell type in *Arabidopsis* flower stalk. *Plant Physiol* 124:599–608
- Kroymann J, Textor S, Tokuhisa JG, Falk KL, Bartram S, Gershenzon J, Mitchell-Olds T (2001) A gene controlling variation in *Arabidopsis* glucosinolate composition is part of the methionine chain elongation pathway. *Plant Physiol* 127:1077–1088
- Li CW (1982) The origin, evolution, taxonomy and hybridization of Chinese cabbage. In: Talekar NS, Griggs TD (eds) Chinese cabbage, Proceedings of first international 1 AVRDC symposium 1981, Taiwan, pp 1–10
- Li G, Quiros CF (2002) Genetic analysis, expression and molecular characterization of Bogsl-ELONG, a major gene involved in the aliphatic glucosinolate pathway of *Brassica* species. *Genetics* 162:1937–1943
- Li G, Quiros CF (2003) In planta side-chain glucosinolate modification in *Arabidopsis* by introduction of dioxygenase *Brassica* homolog Bogsl-ALK. *Theor Appl Genet* 106:1116–1121
- McGrath JM, Quiros CF (1992) Genetic diversity at isozyme and RFLP loci in *Brassica campestris* as related to crop type and geographical origin. *Theor Appl Genet* 83:783–790
- Mithen R (2001) Glucosinolates-biochemistry, genetics and biological activity. *Plant Growth Reg* 34:91–103
- Padilla G, Cartea ME, Velasco P, Haro AD, Ordas A (2007) Variation of glucosinolates in vegetable crops of *Brassica rapa*. *Phytochemistry* 68:536–545
- Petersen BL, Chen S, Hansen CH, Olsen CE, Halkier BA (2002) Composition and content of glucosinolates in developing *Arabidopsis thaliana*. *Planta* 214:562–571
- Rosa EAS, Heancy RK, Fenwick GR, Portas CAM (1997) Glucosinolates in crop plants. *Hort Rev* 19:99–215
- Siemens DH, Garner SH, Mitchell-Olds T, Callaway RM (2002) Cost of defense in the context of plant competition: *Brassica rapa* may grow and defend. *Ecology* 83:505–517
- Skutlarek D, Farber H, Lippert F, Ulbrich A, Wawrzum A, Buning-Pfaue H (2004) Determination of glucosinolate profiles in Chinese vegetables by precursor ion scan and multiple reaction monitoring scan mode (LC-MS/MS). *Eur Food Res Technol* 219:643–649
- Song KM, Osborn TC, Williams PH (1988) *Brassica* taxonomy based on nuclear restriction fragment length polymorphisms (RFLPs). 2. Preliminary analysis of subspecies within *B. rapa* (syn. *campestris*) and *B. oleracea*. *Theor Appl Genet* 76:593–600
- Sovero M (1993) Rapeseed, a new oilseed crop for the United States. In: Janick J, Simon JE (eds) New crops. Wiley, New York, pp 302–307
- Steel RGD, Torrie JH, Dickey DA (1997) Principles and procedures of statistics, a biometrical approach, 3rd edn. McGraw-Hill series in probability and statistics. McGraw-Hill, Boston
- Textor S, Bartram S, Kroymann J, Falk KL, Hick A, Pickett JA, Gershenzon J (2004) Biosynthesis of methionine-derived glucosinolates in *Arabidopsis thaliana*: recombinant expression and characterization of methylthioalkylmalate synthase, the condensing enzyme of the chain-elongation cycle. *Planta* 218:1026–1035
- Tian Q, Rossetto RA, Schwartz SJ (2005) Quantitative determination of intact glucosinolates in broccoli, broccoli sprouts, Brussels sprouts, and cauliflower by high-performance liquid chromatography-electrospray ionization-tandem mass spectrometry. *Anal Biochem* 343:93–99
- Windsor AJ, Reichelt M, Figuth A, Svatos A, Kroymann J, Kleibenstein DJ, Gershenzon J, Mitchell-Olds T (2005) Geographic and evolutionary diversification of glucosinolates among near relatives of *Arabidopsis thaliana* (Brassicaceae). *Phytochemistry* 66:1321–1333