

Symbiont 'bleaching' in planktic foraminifera during the Middle Eocene Climatic Optimum

Edgar, K. M.; Bohaty, S. M.; Gibbs, S. J.; Sexton, P. F.; Norris, R. D.; Wilson, P. A.

DOI:
[10.1130/G33388.1](https://doi.org/10.1130/G33388.1)

License:
None: All rights reserved

Document Version
Peer reviewed version

Citation for published version (Harvard):
Edgar, KM, Bohaty, SM, Gibbs, SJ, Sexton, PF, Norris, RD & Wilson, PA 2013, 'Symbiont 'bleaching' in planktic foraminifera during the Middle Eocene Climatic Optimum', *Geology*, vol. 41, no. 1, pp. 15-18.
<https://doi.org/10.1130/G33388.1>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1 **Symbiont ‘bleaching’ in planktic foraminifera during the Middle Eocene**
2 **Climatic Optimum**

3

4 K.M. Edgar^{a,b,c*}, S.M. Bohaty^a, S.J. Gibbs^a, P.F. Sexton^{c,d}, R.D. Norris^c and P.A.
5 Wilson^a

6 ^aOcean and Earth Science, National Oceanography Centre Southampton, University
7 of Southampton, SO14 3ZH, UK.

8 ^bSchool of Earth and Ocean Sciences, Cardiff University, CF10 3AT, UK.

9 ^cScripps Institution of Oceanography, University of California, San Diego, CA 92093,
10 USA.

11 ^dnow at Centre for Earth, Planetary, Space & Astronomical Research, The Open
12 University, MK7 6AA, UK.

13

14 *Corresponding author. Tel.: +44- (0)2920 874573; Fax: +44 (0)2920 874326

15 E-mail address: EdgarK1@cf.ac.uk

16

17 **ABSTRACT**

18 Many genera of modern planktic foraminifera are adapted to nutrient-poor
19 (oligotrophic) surface waters by hosting photosynthetic symbionts, but it is unknown
20 how they will respond to future changes in surface-ocean temperature and acidity.
21 Here we show that ca. 40 Ma, some fossil photosymbiont-bearing planktic
22 foraminifera were temporarily ‘bleached’ of their symbionts coincident with transient
23 global warming during the Middle Eocene Climatic Optimum (MECO). At Ocean
24 Drilling Program (ODP) Sites 748 and 1051 (Southern Ocean and mid-latitude North
25 Atlantic, respectively), the typically positive relationship between the size of

26 photosymbiont-bearing planktic foraminifer tests and their carbon isotope ratios
27 ($\delta^{13}\text{C}$) was temporarily reduced for \sim 100 k.y. during the peak of the MECO. At the
28 same time, the typically photosymbiont-bearing planktic foraminifera *Acarinina*
29 suffered transient reductions in test size and relative abundance, indicating ecological
30 stress. The coincidence of minimum $\delta^{18}\text{O}$ values and reduction in test size– $\delta^{13}\text{C}$
31 gradients suggests a link between increased sea surface temperatures and bleaching
32 during the MECO, although changes in pH and nutrient availability may also have
33 played a role. Our findings show that host-photosymbiont interactions are not constant
34 through geological time, with implications for both the evolution of trophic strategies
35 in marine plankton and the reliability of geochemical proxy records generated from
36 symbiont-bearing planktic foraminifera.

37

38 Keywords: Middle Eocene Climatic Optimum, bleaching, photosymbionts, planktic
39 foraminifera, Ocean Drilling Program

40

41 INTRODUCTION

42 Photosymbiotic algae play a critical role in the nutrition, reproduction, calcification,
43 growth, and longevity of their planktic foraminiferal hosts. Any changes in
44 photosymbiont activity will therefore have a direct impact on the ‘success’ of the host
45 taxon (Bé et al., 1982; Bijma et al., 1990; Caron et al., 1982; Hemleben et al., 1989).
46 Symbionts modify the chemistry of a foraminifer’s calcifying microenvironment,
47 which impacts the elemental and isotopic ratios of test calcite and imparts
48 characteristic geochemical signatures that are preserved in the sedimentary record
49 (e.g., D’Hondt et al., 1994; Elderfield et al., 2002; Hönisch et al., 2003). There is
50 evidence from the geological record that photosymbionts hosted by planktic

51 foraminifera can be lost or their activity inhibited (i.e. ‘bleaching’). For example, in
52 the late middle Eocene, the gradual breakdown of the host–symbiont relationship over
53 2 m.y. is implicated in the extinction of the large acarininids and morozovelloidids
54 (Wade, 2004; Wade et al., 2008). Furthermore, a rapid increase in surface ocean
55 temperatures during the Paleocene-Eocene Thermal Maximum (PETM) appears to
56 have caused the short-term (<40 k.y.) loss of symbionts from the surface-dwelling
57 planktic foraminifera *Morozovella* and *Acarinina* (Norris, 2007). Yet there is
58 considerable uncertainty regarding how common this loss of symbionts is in the
59 geologic record and, consequently, the mechanism(s) responsible.

60 The Middle Eocene Climatic Optimum (MECO) was a transient global
61 warming event at ca. 40 Ma that interrupted the long-term Eocene cooling trend (Fig.
62 1; Bohaty and Zachos, 2003; Sexton et al., 2006a; Bohaty et al., 2009; Edgar et al.,
63 2010). It lasted for ~500 to 800 k.y. and was marked by gradual ocean warming of ~3
64 to 6 °C, with peak warmth lasting <100 k.y. (Bohaty et al., 2009; Bijl et al., 2010).

65 Here we use the established relationship between test size and $\delta^{13}\text{C}$ in fossil
66 planktic foraminifera as a proxy for photosymbiont activity (Berger et al., 1978;
67 D’Hondt and Zachos, 1993; D’Hondt et al., 1994; Norris, 1996; Pearson et al., 1993;
68 Spero and Lea, 1993). Symbiotic algae preferentially utilize isotopically light carbon
69 (^{12}C) during photosynthesis, leaving dissolved inorganic carbon (DIC) in the
70 foraminifer calcifying microenvironment relatively enriched in ^{13}C . As foraminifera
71 grow and host additional symbionts (or support higher photosymbiont activity), a
72 characteristic increase in $\delta^{13}\text{C}$ with increasing test size occurs (Spero and DeNiro,
73 1987). Using this relationship, we investigate whether host–symbiont interactions
74 were affected by the geologically abrupt environmental changes that accompanied the
75 MECO.

76

77 **MATERIALS AND METHODS**

78 Planktic foraminifera were analyzed from Ocean Drilling Program (ODP) Site 1051
79 (Blake Nose Plateau, subtropical North Atlantic Ocean, 30°03'N 76°21'W) and ODP
80 Site 748 (Kerguelen Plateau, Indian sector of the Southern Ocean, 58°26'S 78°58'E).
81 Middle Eocene paleo-water depths at these sites were ~700–2000 m (Bohaty et al.,
82 2009; Shipboard Scientific Party, 1998; Shipboard Scientific Party, 2004). Planktic
83 foraminifera at both sites are characterized by ‘frosty’ preservation (sensu Sexton et
84 al., 2006b) and show some evidence of recrystallization but are free of infilling. Age
85 models follow Edgar et al. (2010).

86 Planktic foraminiferal $\delta^{13}\text{C}$ data were generated using monospecific separates
87 of the known photosymbiont-bearing genera *Acarinina* (*A. praetopilensis* and *A.*
88 *topilensis* at Site 1051 and *Acarinina primitiva* at Site 748 and *Morozovelloides*
89 *crassatus* (e.g., Pearson et al., 1993; Norris, 1996; Wade et al., 2008). Specimens of
90 the inferred symbiotic genus *Globigerinatheka* and the asymbiotic genus *Subbotina*
91 were also analyzed. All samples were picked from restricted size fractions between
92 150 and 450 μm . Samples were cleaned by ultrasonication, and between 5 and 30
93 individuals (depending on availability) were analysed from each size fraction. Stable
94 isotope values were determined using Europa GEO 20-20 (University of
95 Southampton, UK) and VG Prism (University of California - Santa Cruz, USA) mass
96 spectrometers equipped with automated carbonate preparation devices. Stable isotope
97 results are reported relative to the Vienna PeeDee Belemnite (VPDB) standard with
98 an external analytical precision of $\pm 0.05\text{\textperthousand}$. Relative abundance data were generated
99 from sample splits of the >300 μm size fraction on ~400 individuals.

100

101 **RESULTS**

102 Pre- and post-MECO assemblages of *Acarinina*, *Globigerinatheka*, and
103 *Morozovelloides* show a distinct increase in $\delta^{13}\text{C}$ values with increasing test size (Fig.
104 2A and C), consistent with a surface habitat and hosting dinoflagellate
105 photosymbiosis akin to modern taxa (Pearson et al., 1993; Norris, 1996; Sexton et al.,
106 2006c; Tables DR1-DR4 in the GSA Data Repository). To our knowledge, these are
107 the first published *Globigerinatheka* test size- $\delta^{13}\text{C}$ data, and confirm the long-held
108 view that this group was symbiotic. *Acarinina* and *Morozovelloides* specimens
109 display the highest absolute $\delta^{13}\text{C}$ values in each of the samples, with
110 *Globigerinatheka* offset to slightly lower $\delta^{13}\text{C}$ values (Fig. 2). However, *Acarinina*
111 and *Globigerinatheka* test size- $\delta^{13}\text{C}$ gradients are higher at Site 1051 than at Site 748,
112 which is likely a function of either reduced light conditions and/or temperatures at
113 higher latitudes, lower Symbiont density, or different symbionts (Table DR3). In
114 contrast, the subbotinids exhibit no size-related increase in $\delta^{13}\text{C}$ values at either of the
115 sites investigated and have lower $\delta^{13}\text{C}$ values than other analyzed taxa, consistent
116 with an asymbiotic ecology and thermocline habitat (Pearson et al., 1993; Norris,
117 1996; Sexton et al., 2006c). During the peak of the MECO at ca. 40 Ma, the positive
118 test size- $\delta^{13}\text{C}$ trend in *Acarinina* is temporarily reduced at both study sites, and in
119 *Globigerinatheka* at Site 748 only, resulting in test size- $\delta^{13}\text{C}$ gradients more similar
120 to the asymbiotic genus *Subbotina*. In contrast, *Morozovelloides*, a thermophilic genus
121 confined to (sub)tropical areas and present only at Site 1051, shows no significant
122 gradient reduction during the MECO, but a low gradient prior to the event.

123 *Acarinina* are the dominant surface-dwelling taxa at Site 1051 during the pre-
124 and ‘initial’ MECO (Fig. 3A). They subsequently decrease in relative abundance,
125 reaching lowest abundance during the peak warming interval of the event coincident

126 with their smallest maximum test size (Fig. 3A; Table DR5) and lowest test size- $\delta^{13}\text{C}$
127 gradients (Fig. 2B). In contrast, *Morozovelloides* and *Globigerinatheka* generally
128 increase in relative abundance and maximum test size (Figs. 3B and C) during the
129 event, with a decrease or little change in abundance or test size following the MECO.

130

131 DISCUSSION

132 Mechanisms for a reduction of test size- $\delta^{13}\text{C}$ gradients

133 A reduction and/or loss of the test size- $\delta^{13}\text{C}$ gradients in some photosymbiont-bearing
134 foraminifera during the peak of the MECO may have resulted from (1) gametogenic
135 or ontogenetic overprinting of the symbiont $\delta^{13}\text{C}$ signal, (2) a switch in the type of
136 symbiont hosted, (3) an increase in the average habitat depth during later stages of
137 ontogeny, and/or 4) a loss or inhibition of photosymbionts.

138 First, we do not consider increased inclusion of metabolic CO_2 during late
139 ontogeny, or enhanced calcite precipitation during gametogenesis, as viable
140 explanations for the reduced $\delta^{13}\text{C}$ -size trends observed in the MECO at ODP Sites
141 1051 and 748. Modern culture and \S experiments do not provide support for either
142 hypothesis because metabolic activity is highest in juvenile specimens ($<100\ \mu\text{m}$) and
143 decreases during later growth stages (Berger et al., 1978). There is also little evidence
144 for depth migration of acarininids during late ontogeny (e.g., D'Hondt et al., 1994;
145 Norris, 1996) and the addition of gametogenic calcite, even in heavily calcified
146 globigerinathekids, is insufficient to remove any existing test size- $\delta^{13}\text{C}$ trend (Fig. 2).

147 A second possibility is that the primary algal symbiont groups in planktic
148 foraminifera changed during the MECO, affecting test size- $\delta^{13}\text{C}$ relationships (e.g.,
149 dinoflagellates versus chrysophytes; Bornemann and Norris 2007). Modern
150 foraminifera such as *Globigerinella siphonifera* that host chrysophyte symbionts have

151 a much lower $\delta^{13}\text{C}$ -size gradient than those hosting dinoflagellates, e.g.,
152 *Globigerinoides ruber* (Hemleben et al., 1989). Thus, if acarininids switched from
153 hosting dinoflagellates to chrysophytes during the MECO, their test size- $\delta^{13}\text{C}$
154 relationship might be indistinguishable from asymbiotic taxa, even though they were
155 still symbiont bearing. While modern data are sparse, there is no evidence to suggest
156 that individual taxa switch their symbiont type during their life cycle or between
157 succeeding generations (e.g., Hemleben, 1989, Gast and Caron, 1996), although
158 modern foraminifera are flexible with regards to the genetic subgroups of
159 dinoflagellate that they host (Shaked and de Vargas, 2006). However, if taxa
160 remained symbiotic we might not expect any coincident change in species test size or
161 relative abundance.

162 Third, coincident with environmental change during the MECO, mixed-layer-
163 dwelling foraminifera may have temporarily occupied a deeper position in the water
164 column during late stages of ontogeny. A deeper habitat would also directly inhibit
165 symbiont activity via a reduction in irradiance levels (Spero and DeNiro, 1987; Spero
166 and Lea, 1993; Spero et al., 1997). Thus, foraminifera may have either passively or
167 actively lost their symbionts and migrated to deeper waters to predate on the more
168 abundant algae in the deep chlorophyll maximum. This scenario is analogous to
169 events proposed for the PETM, when tropical ‘excursion’ taxa *M. allisonensis* and *A.*
170 *sibaiensis* are thought to have occupied a deeper ecological niche, more similar to
171 *Subbotina* (Kelly et al., 1996) and yield low $\delta^{13}\text{C}$ -size gradients consistent with
172 asymbiotic or chrysophyte-bearing planktic foraminifera (Kelly et al., 1998;
173 Bornemann and Norris, 2007). Although changes in calcification depth of planktic
174 foraminifera are not unprecedented on long (geological) time scales (e.g., Coxall et
175 al., 2007), available $\delta^{18}\text{O}$ data for the MECO are ambiguous in this regard (Fig. DR1).

176 However, $\delta^{13}\text{C}$ data indicate maintenance of the offset between mixed-layer and
177 thermocline taxa throughout the MECO, suggesting continued separation of depth
178 habitats between taxa.

179 Fourth, a decrease in symbiont activity and/or symbiont concentration could
180 explain the absence of a positive test size– $\delta^{13}\text{C}$ trend in typically symbiont-hosting
181 foraminifera. Laboratory experiments show that the artificial removal of
182 dinoflagellate symbionts from modern foraminifera species (simulating ‘bleaching’) is
183 accompanied by decreases in test size (Bé et al., 1982; Caron et al., 1982),
184 presumably owing to the ecological stress imposed by symbiont eradication.
185 Similarly, the loss of symbionts from *Morozovelloides* in the late middle Eocene is
186 coincident with a decrease in maximum test size (Wade and Olsson, 2009). Hence, the
187 disappearance during the MECO of the normally positive test size– $\delta^{13}\text{C}$ trend in
188 *Acarinina* and the associated pronounced decreases in their size and abundance is
189 consistent with loss of their photosymbionts (Fig. 3C).

190

191 **Bleaching mechanisms**

192 Studies of modern marine taxa in stressed environments may provide some
193 insight into the foraminiferal response to the MECO. However, direct analogy to
194 bleaching events observed in modern coral and benthic foraminifera in the natural
195 environment, and simulated in laboratory cultures is limited, owing to (1) different
196 habitats (planktic versus benthic), (2) the likelihood that culture experiments are not
197 directly representative of the natural environment, and (3) the different relative time
198 scales (annual versus millennial) and number of generations involved. Furthermore,
199 planktic foraminifera cannot be readily observed *in situ*; thus, we do not know if there

200 have been detrimental losses of photosymbionts in response to modern environmental
201 change.

202 If symbiosis is obligate in acarininids (as implied by analogy to modern taxa),
203 bleaching is most likely not a direct stress response given the timescales of
204 environmental change during the MECO. But perhaps cumulatively, environmental
205 changes may have crossed a threshold beyond which foraminifer or their symbionts
206 were unable to successfully operate, triggering the breakdown of the symbiotic
207 relationship. The variable response of the three genera investigated here highlights
208 differential relative sensitivities to the same environmental changes occurring during
209 the MECO. The acarininids were the most sensitive genus to environmental changes,
210 perhaps implying that they were living close to their environmental limits.

211 It is compelling that reduced test size- $\delta^{13}\text{C}$ gradients at both study sites (Fig. 3)
212 occur within the short-lived interval of peak warmth (Figs 1 and 2); yet surface waters
213 also experienced increased nutrient availability (Luciani et al., 2010; Witkowski et al.,
214 2012) and an inferred pH reduction across the MECO (Bijl et al., 2010). However, it's
215 unclear how changes in the trophic state relate to warming; the responses of marine
216 organisms to ΔpH are variable (Hofmann et al., 2010) and culture experiments
217 assessing the impact of carbonate chemistry on the $\delta^{13}\text{C}$ of planktic foraminiferal
218 calcite appear to show little impact on the host-symbiont relationship (Spero et al.,
219 1997). Consequently, while nutrient and pH changes may have exacerbated
220 environmental stress, the temperature increase across the MECO was most likely the
221 primary factor leading to the inhibition of photosymbiosis in *Acarinina* on a global
222 scale. Regardless of the environmental control on foraminiferal bleaching during the
223 MECO, all affected taxa were able to live and maintain populations, implying that, at
224 least on geological short timescales, symbiosis is not essential to their survival. There

225 are several modern mixed-layer taxa that do not harbour symbionts, e.g., *Globigerina*
226 *bulloides* (Hemleben et al., 1989), indicating that symbionts are not essential for
227 survival in the mixed layer.

228 If symbiosis is not essential for foraminifer survival, the exclusion of
229 photosymbionts may represent an adaptive response to changing environmental
230 and/or biotic pressures. Indeed, bleaching has been suggested to be an adaptive
231 mechanism in corals allowing them to be recolonized by new types of algae better
232 suited to short-lived conditions of environmental stress (Brown, 1997). Moreover,
233 symbiont loss in *Acarinina* may have been passive: an indirect consequence of
234 migration to a slightly deeper (i.e. aphotic) depth habitat during the MECO. Yet
235 regardless of whether the loss or inhibition of symbionts was an adaptive or passive
236 mechanism, it came at a cost, highlighted by the fact that *Acarinina* declined in size
237 and abundance across the MECO compared with the other major surface-dwelling
238 planktic foraminiferal groups (Fig. 3). The rapid recovery of test size- $\delta^{13}\text{C}$ gradients
239 (and abundance and test size) to pre-event values following the MECO indicates that,
240 once environmental conditions became more favourable for these planktic
241 foraminifers or their symbionts, the photosymbiotic relationship was re-established at
242 pre-event levels.

243

244 **ACKNOWLEDGEMENTS**

245 This work used samples provided by the Ocean Drilling Program. We thank M.
246 Bolshaw and D. Spanner for help with laboratory work and O. Friedrich, T. Aze, B.
247 Wade and two anonymous reviewers for their insightful comments. Financial support
248 for this research was provided by WUN and NERC fellowships to Edgar, a Royal
249 Society University Research Fellowship to Gibbs, a European Commission Marie

250 Curie Outgoing International Fellowship to Sexton and a NERC small grant to Gibbs
251 and Wilson.

252

253 **FIGURE CAPTIONS**

254 Figure 1. Benthic foraminiferal stable isotope records across the Middle Eocene
255 Climatic Optimum (MECO) from Ocean Drilling Program Sites 1051 (Edgar et al.,
256 2010), 738 and 748 (Bohaty and Zachos, 2003; Bohaty et al., 2009). Isotope
257 stratigraphies at Sites 738 and 748 are aligned to Site 1051. Subdivisions indicate
258 different climatic phases of the MECO.

259

260 Figure 2. Trends in $\delta^{13}\text{C}$ versus mean sieve size related trends in planktic foraminifera
261 for Ocean Drilling Program (ODP) Sites 1051 and 748 across the Middle Eocene
262 Climatic Optimum (MECO). Acar. = *Acarinina* spp (solid diamonds); Glob. =
263 *Globigerinatheka* spp (open circles); Moro. = *Morozovelloides crassatus* (solid
264 triangles) and Subb. = *Subbotina* spp (solid squares). Different coloured symbols from
265 different samples.

266

267 Figure 3. Relative abundance changes in >300 μm -sieve size fraction (lines) and
268 changes in maximum test size diameter (solid symbols) of the dominant surface-
269 dwelling planktic foraminifera across the Middle Eocene Climatic Optimum (MECO)
270 at Ocean Drilling Program Site 1051. Mean diameter of the 20 (where possible)
271 largest specimens in each group is shown and plotted with 1σ .

272

273 **REFERENCES**

- 274 Bé, A.W.H., Spero, H.J. and Anderson, O.R., 1982, Effect of symbiont elimination
275 and reinfection on the life processes of the planktonic foraminifera *Globigerinoides*
276 *sacculifer*: Marine Biology v. 70, p. 73-86.
- 277
- 278 Berger, W.H., Killingley, J.S. and Vincent, E., 1978, Stable isotopes in deep-sea
279 carbonates: Box core ERDC-92, west equatorial Pacific: Oceanologica Acta, v. 1, p.
280 203-216.
- 281
- 282 Bijma, J., Faber, W.W. and Hemleben, C., 1990, Temperature and salinity limits for
283 growth and survival of some planktonic foraminifers in laboratory cultures: Journal of
284 Foraminiferal Research, v. 20, p. 95-116.
- 285
- 286 Bijl, P.K., Houben, A.J.P., Schouten, S., Bohaty, S.M., Sluijs, A., Reichart, G-J.,
287 Sinninghe Damsté, J.S. and Brinkhuis, H., 2010, Transient middle Eocene
288 atmospheric CO₂ and temperature variations: Science v. 330, p. 819-821.
- 289
- 290 Bohaty, S.M. and Zachos, J.C., 2003, Significant Southern Ocean warming event in
291 the late middle Eocene: Geology v. 31, p.1017-1020.
- 292
- 293 Bohaty, S.M., Zachos, J.C., Florindo, F. and Delaney, M.L., 2009, Coupled
294 greenhouse warming and deep-sea acidification in the middle Eocene:
295 Paleoceanography v. 24, doi: 10.1029/2008PA001676.
- 296

- 297 Bornemann, A. and Norris, R.D., 2007. Size-related stable isotope changes in Late
298 Cretaceous planktic foraminifera: Implications for paleoecology and photosymbiosis:
299 Marine Micropaleontology v. 65, p. 32-42.
- 300
- 301 Brown, B.E., 1997, Coral bleaching: causes and consequences: Coral Reefs, v. 16,
302 p.129-138.
- 303
- 304 Caron, D.A., Bé, A.W.H. and Anderson, O.R., 1982, Effects of variations in light
305 intensity on life processes of the planktonic foraminifer *Globigerinoides sacculifer* in
306 laboratory culture: Journal of the Marine Biological Association UK, v. 62, p. 435-
307 451.
- 308
- 309 Coxall, H.K., Wilson, P.A., Pearson, P.N., and Sexton, P.F., 2007, Iterative evolution
310 of digitate planktonic foraminifera: Paleobiology, v. 33, p. 495-516,
311 doi:10.1666/06034.1.
- 312
- 313 D'Hondt, S. and Zachos, J.C., 1993, On stable isotopic variation and earliest
314 Paleocene planktonic foraminifera: Paleoceanography v. 8, p. 527-547.
- 315
- 316 D'Hondt, S., Zachos, J.C. and Schultz, G., 1994, Stable isotopic signals and
317 photosymbiosis in Late Paleocene planktic foraminifera: Paleobiology, v. 20, p. 391-
318 406.
- 319

- 320 Douglas, R.G. and Savin, S.M., 1978, Oxygen isotopic evidence for the depth
321 stratification of Tertiary and Cretaceous planktic foraminifera: Marine
322 Micropaleontology v. 3, p. 175-196.
- 323
- 324 Edgar, K.M., Wilson, P.A., Sexton, P.F., Gibbs, S.J., Roberts, A.P. and Norris, R.D.,
325 2010, New biostratigraphic, magnetostratigraphic and isotopic insights in the Middle
326 Eocene Climatic optimum in low latitudes: Palaeogeography, Palaeoclimatology,
327 Palaeoecology, v. 297, p. 670-682.
- 328
- 329 Elderfield, H., Vautravers, M., and Cooper, M., 2002. The relationship between
330 Mg.Ca, Sr/Ca, $\delta^{18}\text{O}$, and $\delta^{13}\text{C}$ of species of planktonic foraminifera: Geochemistry,
331 Geophysics, Geosystems, v. 3, doi:10.1029/2001GC00194.
- 332
- 333 Gast, R.J. and Caron, D.A., 1996. Molecular phylogeny of symbiotic dinoflagellates
334 from planktonic foraminifera and radiolaria: Molecular Biology and Evolution v. 13,
335 p. 1192-1197.
- 336
- 337 Hemleben, C., Spindler, M. and Anderson, O.R., 1989, Modern planktonic
338 foraminifera: Springer-Verlag (New York), 363 p.
- 339
- 340 Hofmann, G.E., Barry, J.P., Edmunds, P.J., Gates, R.D., Hutchins, D.A., Klinger, T.,
341 and Sewell, M.A., 2010, The effects of ocean acidification on calcifying organisms in
342 marine ecosystems: An organism –to-ecosystem perspective: Annual Review of
343 Ecology, Evolution and Systematics, V. 41, p. 127-147.

- 344 Hönnisch, B., Bijma, J., Russell, A.D., Spero, H.J., Palmer, M.R., Zeebe, R.E. and
345 Eisenhauer, A., 2003, The influence of symbiont photosynthesis on the boron isotopic
346 composition of foraminifera shells: *Marine Micropaleontology* v. 49, p. 87-96.
- 347
- 348 Kelly, D.C., Bralower, T.J., Zachos, J.C., Premoli Silva, I. and Thomas, E., 1996,
349 Rapid diversification of planktonic foraminifera in the tropical Pacific (ODP Site 865)
350 during the late Paleocene thermal maximum: *Geology* v.24, p. 423-426.
- 351
- 352 Kelly, D.C., Bralower, T.J. and Zachos, J.C., 1998, Evolutionary consequences of the
353 latest Paleocene thermal maximum for tropical planktonic foraminifera:
354 *Palaeogeography, Palaeoclimatology, Palaeoecology* v.141, p. 139-161.
- 355
- 356 Luciani, V., Giusberti, L., Agnini, C., Fornaciari, E., Rio, D., Spofforth, D.J.A. and
357 Pälike, H., 2010, Ecological and evolutionary response of Tethyan planktonic
358 foraminifera to the middle Eocene climatic optimum (MECO) form the Alano section
359 (NE Italy): *Palaeogeography, Palaeoclimatology, Palaeoecology* v. 292, p. 82-95.
- 360
- 361 Norris, R.D., 1996, Symbiosis as an Evolutionary Innovation in the Radiation of
362 Paleocene Planktic Foraminifera; *Paleobiology*, v. 22, p. 461-480.
- 363
- 364 Norris, R.D., 1998, Recognition and evolutionary significance of photosymbiosis in
365 molluscs, corals, and foraminifera; in Manger, W.L., and Meeks, K.L., eds., *Isotope*
366 *paleobiology and paleoecology: Paleontological Society Papers*, v. 4,p. 68-100.
- 367

- 368 Norris, R.D., 2007, Bleaching of symbiotic foraminifera during extreme global
369 warming at the Paleocene-Eocene boundary; EOS Trans, AGU 88(52), Fall Meeting
370 Suppl., Abstract OS14A-07.
- 371
- 372 Pearson, P.N., Shackleton, N.J. and Hall, M., 1993, Stable isotope paleoceanography of
373 middle Eocene planktonic foraminifera and multispecies isotope stratigraphy: Journal
374 of Foraminiferal Research v. 23, p. 123-140.
- 375
- 376 Premoli-Silva, I. and Boersma, A., 1988, Atlantic Eocene planktonic foraminiferal
377 historical biogeography and paleohydrographic indices: Palaeogeography,
378 Palaeoclimatology, Palaeoecology v. 67, p. 315-356.
- 379
- 380 Quillévéré, F., Norris, R.D., Moussa, I., Berggren, W.A., 2001, Role of
381 photosymbiosis and biogeography in the diversification of early Paleogene
382 Acarininids (planktonic foraminifera): Paleobiology v. 27, p. 311-326.
- 383
- 384 Sexton, P.F., Wilson, P.A. and Norris, R.D., 2006a, Testing the Cenozoic multisite
385 composite $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ curves: new monospecific Eocene records from a single
386 locality, Demerara Rise (Ocean Drilling Program Leg 207): Paleoceanography v. 21,
387 PA2019, doi:10.1029/2005PA001253.
- 388
- 389 Sexton, P.F., Wilson, P.A. and Pearson, P.N., 2006b, Microstructural and
390 geochemical perspectives on planktic foraminiferal preservation: ‘Glassy’ versus
391 ‘Frosty’: Geochemistry, Geophysics, Geosystems, v. 7, doi:10.1029/2006GC001291.
- 392

393 Sexton, P.F., Wilson, P.A. and Pearson, P.N., 2006c, Palaeoecology of late middle
394 Eocene planktic foraminifera and evolutionary implications: Marine
395 Micropaleontology, v. 60, p. 1-16.

396

397 Shaked, Y. and de Vargas, C., 2006, Pelagic photosymbiosis: rDNA assessment of
398 diversity and evolution of dinoflagellate symbionts and planktonic foraminiferal
399 hosts: Marine Ecology Progress Series v. 325, p. 59-71.

400

401 Shipboard Scientific Party, 1998, Site 1051, *in* Norris, R.D., Kroon, D. and Klaus, A.
402 et al., Proceedings of the Ocean Drilling Program, Volume 171B: Washington, D.C.,
403 U.S. Government Printing Office, p. 171-239.

404

405 Shipboard Scientific Party, 2004, Site 1260, *in* Erbacher, J., Mosher, D.C. and
406 Malone, M.J. et al., Proceedings of the Ocean Drilling Program, Volume 207:
407 Washington, D.C., U.S. Government Printing Office,
408 doi:10.2973/odp.proc.ir.207.108.2004.

409

410 Spero, H.J. and DeNiro, M.J., 1987. The influence of photosynthesis on the $\delta^{18}\text{O}$ and
411 $\delta^{13}\text{C}$ values of planktonic foraminiferal shell calcite: Symbiosis, v. 4, p. 213-228.

412


413 Spero, H.J. and Lea, D.W., 1993, Intraspecific stable isotope variability in the planktic
414 foraminifera *Globigerinoides sacculifer*: Results from laboratory experiments: Marine
415 Micropaleontology v. 22, p. 221-234.


416


- 417 Spero, H.J., Bijma, J., Lea, D.W. and Bemis, B.E., 1997, Effect of seawater carbonate
418 concentration on foraminiferal carbon and oxygen isotopes: Nature v. 390, p. 497-
419 500.
- 420
- 421 Wade, B.S., 2004, Planktonic foraminiferal biostratigraphy and mechanisms in the
422 extinction of *Morozovella* in the late middle Eocene: Marine Micropaleontology v.
423 51, p23-38.
- 424
- 425 Wade, B.S., Al-Sabouni, N., Hemleben, C. and Kroon, D., 2007, Symbiont bleaching
426 in fossil planktonic foraminifera: Evolutionary Ecology v. 22, p. 253-265.
- 427
- 428 Wade, B.S. and Olsson, R.K., 2009, Investigation of pre-extinction dwarfing in
429 Cenozoic planktonic foraminifera; Palaeogeography, Palaeoclimatology,
430 Palaeoecology, v. 284, p. 39-46.
- 431
- 432 Witkowski, J., Bohaty, S.M., McCartney, K., and Harwood,D.M., 2012, Enhanced
433 siliceous plankton productivity in response to middle Eocene warming at Southern
434 Ocean ODP Sites 748 and 749: Palaeogeography, Palaeoclimatology, Palaeoecology,
435 v. 326-328, p. 78-94.
- 436
- 437 Zachos, J.C., Dickens, G.R. and Zeebe, R.E., 2008, An early Cenozoic perspective on
438 greenhouse warming and carbon-cycle dynamics: Nature v. 451, p. 279-283.
- 439
- 440 ¹GSA Data Repository item 2013002, stable isotope, test size data, and SEM images;
441 Figure DR1-DR4; and Tables DR1-5, is available online at

442 www.geosociety.org/pubs/ft2009.htm, or on request from editing@geosociety.org or

443 Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.









Data Repository Material: Symbiont ‘bleaching’ in planktic foraminifera during the Middle Eocene Climatic Optimum by Edgar et al.


Data Repository Figure 1. $\delta^{18}\text{O}$ versus mean sieve size trends in planktic foraminifera at ODP Sites 1051 and 748 across the MECO. Abbreviations and symbols are: Acar. = *Acarinina* spp (solid diamonds); Glob. = *Globigerinatheka* spp (open circles); Moro. = *Morozovelloides crassatus* (solid triangles) and Subb. = *Subbotina* spp (solid squares). Different coloured symbols of the same shape are from different samples. For clarity, individual samples are plotted in Figs DR2 and DR3. $\delta^{18}\text{O}$ -size related trends are shown for four different timeslices (rather than three as in Fig. 2) to distinguish between pre- and initial-MECO conditions. At Site 748, there is a clear offset in absolute $\delta^{18}\text{O}$ values between inferred surface (*Acarinina* and *Globigerinatheka*) and thermocline dwelling taxa (*Subbotina*) across the MECO providing little support for *Acarinina* or *Globigerinatheka* occupying a deeper position in the water column during the MECO. At Site 1051, it is more difficult to assess any changes in the relative depth ordering of taxa across the MECO. Perhaps, in part, because of the multiple species of *Globigerinatheka* and *Subbotina* combined for isotope analysis but also because of potential diagenetic alteration of $\delta^{18}\text{O}$ isotope values at this site. [Site 748 planktic foraminifer are likely less susceptible to diagenetic alteration because of weaker vertical thermal water column gradients at higher latitudes]. We note that $\delta^{13}\text{C}$ values are more resiliant to diagenetic alteration than $\delta^{18}\text{O}$ values and there is little difference in values reported between ‘glassy’ and ‘frosty’ planktic foraminifer (e.g., Pearson et al., 2001; Sexton et al., 2006). The overall reduced $\delta^{18}\text{O}$ offsets between taxa and lower test size- $\delta^{18}\text{O}$ gradients at Site 748 than at Site 1051 are most easily explained by the presence of a less thermally stratified water column at high latitude. Note the different y-axis values between ODP Sites 1051 and 748.




Data Repository Figure 2. $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ -mean sieve size related trends in planktic foraminifera in individual samples at ODP Site 1051 across the MECO. Different coloured symbols represent different genera of planktic foraminifera: *Acarinina topilensis* = solid red diamonds; *Morozovelloides crassatus* = solid purple triangles; *Globigerinatheka* spp = open green circles and *Subbotina* spp = solid blue squares. The globigerinathekids typically show a negative sieve size- $\delta^{18}\text{O}$ gradient reflecting as they sink through the water column towards the end of their life cycle and precipitate a thick clacite crust (gametogenic calcite) from water with more positive $\delta^{18}\text{O}$ values than the surface waters that they originally precipitated their test from.



Data Repository Figure 3. $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ -mean sieve size related trends in planktic foraminifera in individual samples at ODP Site 748 across the MECO. Different coloured symbols represent different planktic foraminifera: *Acarinina primitiva* = solid diamonds; *Globigerinatheka index* = open circles and *Subbotina* spp = solid squares.





Data Repository Figure 4 - Scanning electron microscope images from ODP Sites 1051 and 748 illustrating species concepts adopted in this study. Planktic foraminifera are ‘frosty’ not ‘glassy’ indicating some diagenetic alteration but are free of infilling. Scale bars are 100 µm in a-h and 10 µm in i. a. *Acarinina primitiva*, Sample 748B 19H-2, 127-129 cm. b and c. *Globigerinatheka index*, Sample 748B 19H-2, 127-129 cm. d. *Morozovelloides crassatus*, Sample 1051B 8H-6, 5-7 cm. e and f. *Acarinina topilensis*, Sample 1051B 8H-6, 5-7 cm. g and h. *Acarinina praetopilensis*, Sample 1051B 8H-6, 5-7 cm. i. close up of *A. topilensis* wall texture, Sample 1051B 8H-6, 5-7 cm. *A. topilensis* sensu stricto (e) is restricted to large size fractions, thus in this study we have adopted a broad species concept for *A. topilensis* that includes less elaborate morphological forms that fall within *A. praetopilensis*.




TABLE DR1. PLANKTIC FORAMINIFERAL SIZE FRACTION- $\delta^{13}\text{C}$ AND $\delta^{18}\text{O}$ DATA FROM ODP SITE 1051

Sample	Depth (mbsf)	CK95* Age (Ma)	Taxon †	Sieve size fraction (microns)	$\delta^{13}\text{C}$ VPDB	$\delta^{18}\text{O}$ VPDB
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	300-355	3.704	-1.104
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	250-300	3.370	-1.046
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	212-250	2.865	-0.946
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	180-212	2.891	-0.919
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Morozovelloides crassatus</i>	250-300	3.261	-1.379
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Morozovelloides crassatus</i>	212-250	2.915	-1.030
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Morozovelloides crassatus</i>	180-212	2.794	-1.130
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Morozovelloides crassatus</i>	150-180	2.532	-1.082
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	355-400	3.043	-1.452
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	300-355	2.731	-1.451
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	250-300	2.324	-1.435
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	212-250	2.002	-1.604
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	180-212	1.489	-1.807
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Subbotina</i> spp	300-355	1.455	-0.333
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Subbotina</i> spp	250-300	1.243	-0.241
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Subbotina</i> spp	212-250	1.205	-0.273
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Subbotina</i> spp	180-212	1.515	-0.271
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Acarinina topilensis</i>	250-300	2.573	-0.907
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Acarinina topilensis</i>	212-250	2.528	-0.882
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Acarinina topilensis</i>	180-212	2.459	-0.708
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	300-355	2.948	-1.197
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	250-300	2.704	-0.775
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	212-250	2.590	-0.702
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	180-212	2.374	-0.668
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	150-180	2.080	-0.858
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	355-400	2.130	-0.959
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	300-355	2.004	-1.786
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	250-300	2.299	-0.959
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	212-250	1.911	-1.380
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	180-212	1.335	-1.407
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Subbotina</i> spp	250-300	1.248	-0.388
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Subbotina</i> spp	212-250	1.263	-1.163
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Subbotina</i> spp	180-212	1.080	-0.923
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Acarinina topilensis</i>	300-355	3.276	-0.697
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Acarinina topilensis</i>	250-300	2.922	-0.740
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Acarinina topilensis</i>	212-250	2.936	-0.010
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Acarinina topilensis</i>	180-212	1.821	-1.944
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Morozovelloides crassatus</i>	400-450	3.458	-0.713
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Morozovelloides crassatus</i>	355-400	3.095	-1.152
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Morozovelloides crassatus</i>	250-300	2.904	-0.865
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Morozovelloides crassatus</i>	212-250	2.062	-1.914
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Morozovelloides crassatus</i>	180-212	2.404	-0.803
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Morozovelloides crassatus</i>	150-180	1.797	-1.086
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Globigerinatheka</i> spp	400-450	2.543	-1.251
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Globigerinatheka</i> spp	355-400	2.604	-0.853
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Globigerinatheka</i> spp	300-355	2.398	-0.844
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Globigerinatheka</i> spp	250-300	2.187	-1.041
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Globigerinatheka</i> spp	212-250	1.847	-1.238
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Globigerinatheka</i> spp	180-212	1.392	-1.283
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Subbotina</i> spp	355-400	1.670	-0.273
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Subbotina</i> spp	300-355	1.499	-0.385
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Subbotina</i> spp	250-300	1.490	-0.170
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Subbotina</i> spp	212-250	1.909	0.308
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Subbotina</i> spp	180-212	1.480	-0.165

1051B-11H-4, 45-47 cm	92.25	40.4	<i>Acarinina topilensis</i>	355-400	3.821	-1.621
1051B-11H-4, 45-47 cm	92.25	40.4	<i>Acarinina topilensis</i>	300-355	3.332	-1.512
1051B-11H-4, 45-47 cm	92.25	40.4	<i>Acarinina topilensis</i>	250-300	3.384	-1.093
1051B-11H-4, 45-47 cm	92.25	40.4	<i>Acarinina topilensis</i>	212-250	2.808	-1.187
1051B-11H-4, 45-47 cm	92.25	40.4	<i>Acarinina topilensis</i>	180-212	2.831	-0.805
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	300-355	3.294	-1.066
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	250-300	3.104	-0.410
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	212-250	2.730	-1.234
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	180-212	2.607	-0.891
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	150-180	2.158	-0.880
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	355-400	2.762	-0.958
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	300-355	2.554	-0.880
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	250-300	2.151	-1.160
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	212-250	1.905	-1.312
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	180-212	1.562	-1.473
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Subbotina</i> spp	300-355	1.535	-0.669
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Subbotina</i> spp	250-300	1.407	-0.693
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Subbotina</i> spp	212-250	1.562	-0.373
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Subbotina</i> spp	180-212	1.117	-0.352
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Acarinina topilensis</i>	355-400	3.299	-0.472
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Acarinina topilensis</i>	300-355	3.299	-0.340
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Acarinina topilensis</i>	250-300	2.994	-0.200
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Acarinina topilensis</i>	212-250	2.675	-0.511
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Acarinina topilensis</i>	180-212	2.152	-0.521
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Morozovelloides crassatus</i>	300-355	2.880	-0.521
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Morozovelloides crassatus</i>	250-300	2.562	-0.899
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Morozovelloides crassatus</i>	212-250	2.504	-0.728
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Morozovelloides crassatus</i>	180-212	2.349	-0.453
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Morozovelloides crassatus</i>	150-180	2.174	-0.430
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Globigerinatheka</i> spp	300-355	2.754	-0.859
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Globigerinatheka</i> spp	250-300	2.200	-0.671
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Globigerinatheka</i> spp	212-250	2.079	-0.225
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Globigerinatheka</i> spp	180-212	1.218	-1.182
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Subbotina</i> spp	400-450	1.524	-0.254
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Subbotina</i> spp	355-400	1.393	-0.223
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Subbotina</i> spp	300-355	1.193	-0.091
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Subbotina</i> spp	250-300	1.360	-0.261
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Subbotina</i> spp	212-250	1.328	-0.149
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Subbotina</i> spp	180-212	1.590	-0.088
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Acarinina topilensis</i>	300-355	3.330	-0.682
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Acarinina topilensis</i>	250-300	3.195	-0.512
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Acarinina topilensis</i>	212-250	2.632	-0.682
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Acarinina topilensis</i>	180-212	2.482	-0.706
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Morozovelloides crassatus</i>	300-355	2.580	-0.706
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Morozovelloides crassatus</i>	250-300	2.687	-0.780
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Morozovelloides crassatus</i>	212-250	2.960	-0.504
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Morozovelloides crassatus</i>	180-212	2.865	-0.329
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Morozovelloides crassatus</i>	150-180	2.475	-0.537
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	300-355	2.427	-0.658
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	250-300	2.405	-0.723
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	212-250	1.759	-0.811
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	180-212	1.442	-1.141
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Subbotina</i> spp	250-300	1.186	-0.050
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Subbotina</i> spp	212-250	1.718	-0.295

*CK95 = Age scale of Cande and Kent (1995).

†A broad species concept was adopted for *A. topilensis* (see Fig. DR4).

TABLE DR2. PLANKTIC FORAMINIFERAL SIZE FRACTION- $\delta^{13}\text{C}$ AND $\delta^{18}\text{O}$ DATA FROM ODP SITE 748

Sample	Depth (mbsf)	CK95* Age (Ma)	Taxon	Sieve size fraction (microns)	$\delta^{13}\text{C}$ VPDB	Average $\delta^{13}\text{C}$ VPDB	$\delta^{18}\text{O}$ VPDB	Average $\delta^{18}\text{O}$ VPDB
748B-18H-7, 23-25 cm	161.33	39.35	<i>Acarinina primitiva</i>	300-355	3.01		-0.12	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Acarinina primitiva</i>	250-300	3.19		-0.02	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Acarinina primitiva</i>	212-250	2.84		0.01	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Acarinina primitiva</i>	180-212	2.55		0.00	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Acarinina primitiva</i>	150-180	2.39		0.07	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Globigerinatheka index</i>	>425	3.18		0.16	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Globigerinatheka index</i>	355-425	2.98		0.44	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Globigerinatheka index</i>	300-355	2.91		0.42	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Globigerinatheka index</i>	250-300	2.77		0.40	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Globigerinatheka index</i>	212-250	2.66		0.22	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Globigerinatheka index</i>	180-212	2.55		0.28	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Globigerinatheka index</i>	150-180	2.33		0.22	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Subbotina spp.</i>	355-425	1.88		0.77	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Subbotina spp.</i>	300-355	1.87		0.75	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Subbotina spp.</i>	250-300	1.88		0.79	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Subbotina spp.</i>	212-250	1.82		0.81	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Subbotina spp.</i>	180-212	1.81		0.66	
748B-18H-7, 23-25 cm	161.33	39.35	<i>Subbotina spp.</i>	150-180	1.74		0.54	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	250-300	2.79		0.07	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	250-300	2.86	2.91	0.10	0.04
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	250-300	3.07		-0.05	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	212-250	2.81		-0.06	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	212-250	2.97	2.83	-0.19	-0.15
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	212-250	2.88		-0.24	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	212-250	2.66		-0.10	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	180-212	2.47		0.19	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	180-212	2.46		0.07	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	180-212	2.52	2.55	-0.14	0.00
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	180-212	2.65		-0.01	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	180-212	2.63		-0.11	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	150-180	2.21		-0.03	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	150-180	2.33	2.37	-0.07	-0.06
748B-18H-7, 32-34 cm	161.42	39.36	<i>Acarinina primitiva</i>	150-180	2.58		-0.08	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	355-425	2.74		0.38	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	355-425	2.97	2.85	0.25	0.31
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	355-425	2.85		0.30	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	300-355	2.73		0.31	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	300-355	2.77	2.79	0.39	0.33
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	300-355	2.94		0.29	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	300-355	2.70		0.34	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	250-300	2.73		0.36	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	250-300	2.73	2.69	0.19	0.30
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	250-300	2.61		0.35	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	212-250	2.34		0.40	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	212-250	2.54	2.50	0.23	0.27
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	212-250	2.63		0.20	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	180-212	2.42	2.44	0.17	0.11
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	180-212	2.47		0.12	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	180-212	2.44		0.04	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	150-180	2.08		0.30	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	150-180	2.25	2.24	0.13	0.15
748B-18H-7, 32-34 cm	161.42	39.36	<i>Globigerinatheka index</i>	150-180	2.41		0.01	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Subbotina spp.</i>	250-300	1.67		0.63	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Subbotina spp.</i>	250-300	1.69	1.70	0.69	0.67
748B-18H-7, 32-34 cm	161.42	39.36	<i>Subbotina spp.</i>	250-300	1.75		0.67	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Subbotina spp.</i>	212-250	1.69		0.65	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Subbotina spp.</i>	212-250	1.69	1.69	0.67	0.66
748B-18H-7, 32-34 cm	161.42	39.36	<i>Subbotina spp.</i>	212-250	1.70		0.65	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Subbotina spp.</i>	180-212	1.47	1.59	0.64	0.58

748B-18H-7, 32-34 cm	161.42	39.36	<i>Subbotina spp.</i>	180-212	1.67	0.54	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Subbotina spp.</i>	180-212	1.69	0.50	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Subbotina spp.</i>	180-212	1.53	0.65	
748B-18H-7, 32-34 cm	161.42	39.36	<i>Subbotina spp.</i>	150-180	1.54	1.57	0.39
748B-18H-7, 32-34 cm	161.42	39.36	<i>Subbotina spp.</i>	150-180	1.59	0.46	0.42
748B-19H-2, 127-129 cm	164.37	40.04	<i>Acarinina primitiva</i>	300-355	2.10	-0.26	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Acarinina primitiva</i>	250-300	2.25	-0.38	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Acarinina primitiva</i>	212-250	1.95	-0.41	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Acarinina primitiva</i>	180-212	1.94	-0.52	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Acarinina primitiva</i>	150-180	1.86	-0.65	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Globigerinatheka index</i>	355-425	1.98	-0.27	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Globigerinatheka index</i>	355-400	1.99	-0.38	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Globigerinatheka index</i>	300-355	1.94	2.06	-0.38
748B-19H-2, 127-129 cm	164.37	40.04	<i>Globigerinatheka index</i>	300-355	2.17	-0.32	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Globigerinatheka index</i>	250-300	1.78	-0.32	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Globigerinatheka index</i>	212-250	1.92	-0.34	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Globigerinatheka index</i>	180-212	1.85	-0.23	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Globigerinatheka index</i>	150-180	1.40	-0.43	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Subbotina spp.</i>	300-355	1.00	-0.21	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Subbotina spp.</i>	250-300	1.10	-0.32	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Subbotina spp.</i>	212-250	1.28	-0.16	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Subbotina spp.</i>	180-212	1.20	-0.18	
748B-19H-2, 127-129 cm	164.37	40.04	<i>Subbotina spp.</i>	150-180	1.25	-0.24	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Acarinina primitiva</i>	300-355	2.07	-0.40	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Acarinina primitiva</i>	250-300	1.92	-0.42	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Acarinina primitiva</i>	212-250	1.81	-0.59	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Acarinina primitiva</i>	180-212	1.90	-0.47	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Acarinina primitiva</i>	150-180	1.80	-0.38	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Globigerinatheka index</i>	>400	2.24	-0.33	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Globigerinatheka index</i>	355-400	2.02	-0.40	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Globigerinatheka index</i>	300-355	1.83	-0.36	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Globigerinatheka index</i>	250-300	1.79	-0.35	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Globigerinatheka index</i>	212-250	1.76	-0.30	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Globigerinatheka index</i>	180-212	1.87	-0.40	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Globigerinatheka index</i>	150-180	1.76	-0.19	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Globigerinatheka index</i>	125-150	1.49	-0.20	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Subbotina spp.</i>	300-355	1.12	-0.14	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Subbotina spp.</i>	250-300	1.22	-0.03	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Subbotina spp.</i>	212-250	1.18	0.12	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Subbotina spp.</i>	180-212	1.31	0.00	
748B-19H-3, 31-33 cm	164.91	40.08	<i>Subbotina spp.</i>	150-180	1.27	-0.13	
748B-19H-3, 103-105 cm	165.63	40.13	<i>Acarinina primitiva</i>	300-355	2.11	-0.14	
748B-19H-3, 103-105 cm	165.63	40.13	<i>Acarinina primitiva</i>	250-300	1.86	-0.15	
748B-19H-3, 103-105 cm	165.63	40.13	<i>Acarinina primitiva</i>	212-250	1.93	-0.18	
748B-19H-3, 103-105 cm	165.63	40.13	<i>Acarinina primitiva</i>	180-212	2.10	-0.34	
748B-19H-3, 103-105 cm	165.63	40.13	<i>Acarinina primitiva</i>	150-180	2.21	-0.63	
748B-19H-3, 103-105 cm	165.63	40.13	<i>Acarinina primitiva</i>	125-150	2.08	-0.50	
748B-19H-4, 103-105 cm	167.13	40.24	<i>Acarinina primitiva</i>	300-355	2.27	0.24	
748B-19H-4, 103-105 cm	167.13	40.24	<i>Acarinina primitiva</i>	250-300	1.89	0.04	
748B-19H-4, 103-105 cm	167.13	40.24	<i>Acarinina primitiva</i>	180-212	1.78	0.10	
748B-19H-4, 103-105 cm	167.13	40.24	<i>Acarinina primitiva</i>	150-180	1.84	0.23	
748B-19H-4, 103-105 cm	167.13	40.24	<i>Acarinina primitiva</i>	125-150	1.99	-0.16	
748B-19H-5, 71-73 cm	168.31	40.30	<i>Acarinina primitiva</i>	300-355	2.04	0.12	
748B-19H-5, 71-73 cm	168.31	40.30	<i>Acarinina primitiva</i>	250-300	1.94	0.11	
748B-19H-5, 71-73 cm	168.31	40.30	<i>Acarinina primitiva</i>	212-250	1.94	0.07	
748B-19H-5, 71-73 cm	168.31	40.30	<i>Acarinina primitiva</i>	180-212	1.90	0.00	
748B-19H-5, 71-73 cm	168.31	40.30	<i>Acarinina primitiva</i>	150-180	1.75	0.14	
748B-19H-6, 55-57 cm	169.65	40.38	<i>Acarinina primitiva</i>	300-355	2.06	0.03	
748B-19H-6, 55-57 cm	169.65	40.38	<i>Acarinina primitiva</i>	250-300	2.03	0.11	
748B-19H-6, 55-57 cm	169.65	40.38	<i>Acarinina primitiva</i>	212-250	1.79	0.08	

748B-19H-6, 55-57 cm	169.65	40.38	<i>Acarinina primitiva</i>	180-212	1.95	-0.29		
748B-19H-6, 55-57 cm	169.65	40.38	<i>Acarinina primitiva</i>	150-180	1.62	0.09		
748B-19H-6, 55-57 cm	169.65	40.38	<i>Acarinina primitiva</i>	125-150	1.72	0.19		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Acarinina primitiva</i>	300-355	1.93	0.42		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Acarinina primitiva</i>	250-300	1.85	0.22		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Acarinina primitiva</i>	212-250	1.63	0.25		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Acarinina primitiva</i>	180-212	1.46	0.29		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Acarinina primitiva</i>	150-180	1.37	-0.14		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Globigerinatheka index</i>	355-400	2.52	0.12		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Globigerinatheka index</i>	300-355	2.46	-0.06		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Globigerinatheka index</i>	250-300	2.23	0.05		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Globigerinatheka index</i>	212-250	1.90	0.15		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Globigerinatheka index</i>	180-212	1.80	0.16		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Globigerinatheka index</i>	150-180	1.38	0.23		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Subbotina spp.</i>	300-355	1.15	0.53		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Subbotina spp.</i>	250-300	1.25	0.40		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Subbotina spp.</i>	212-250	1.16	0.39		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Subbotina spp.</i>	180-212	1.20	0.50		
748B-20H-1, 16-18 cm	171.26	41.41	<i>Subbotina spp.</i>	150-180	0.95	0.39		
748B-20H-3, 76-78 cm	174.86	41.79	<i>Acarinina primitiva</i>	300-355	1.57	-0.32		
748B-20H-3, 76-78 cm	174.86	41.79	<i>Acarinina primitiva</i>	250-300	1.25	-0.32		
748B-20H-3, 76-78 cm	174.86	41.79	<i>Acarinina primitiva</i>	212-250	0.92	1.01	-0.27	-0.24
748B-20H-3, 76-78 cm	174.86	41.79	<i>Acarinina primitiva</i>	212-250	1.11		-0.21	
748B-20H-3, 76-78 cm	174.86	41.79	<i>Acarinina primitiva</i>	180-212	0.85	0.89	-0.26	-0.30
748B-20H-3, 76-78 cm	174.86	41.79	<i>Acarinina primitiva</i>	180-212	0.92		-0.33	
748B-20H-3, 76-78 cm	174.86	41.79	<i>Acarinina primitiva</i>	150-180	0.79		-0.44	
748B-20H-3, 76-78 cm	174.86	41.79	<i>Globigerinatheka index</i>	355-400	1.60		-0.38	
748B-20H-3, 76-78 cm	174.86	41.79	<i>Globigerinatheka index</i>	300-355	1.39		-0.36	
748B-20H-3, 76-78 cm	174.86	41.79	<i>Globigerinatheka index</i>	250-300	1.37		-0.35	
748B-20H-3, 76-78 cm	174.86	41.79	<i>Globigerinatheka index</i>	212-250	1.17		-0.28	
748B-20H-3, 76-78 cm	174.86	41.79	<i>Globigerinatheka index</i>	180-212	1.14		-0.31	
748B-20H-3, 76-78 cm	174.86	41.79	<i>Globigerinatheka index</i>	150-180	0.81		-0.31	
748B-20H-3, 76-78 cm	174.86	41.79	<i>Subbotina spp.</i>	300-355	0.33		-0.01	
748B-20H-3, 76-78 cm	174.86	41.79	<i>Subbotina spp.</i>	250-300	0.56		-0.03	
748B-20H-3, 76-78 cm	174.86	41.79	<i>Subbotina spp.</i>	212-250	0.50		-0.08	
748B-20H-3, 76-78 cm	174.86	41.79	<i>Subbotina spp.</i>	180-212	0.54		-0.05	
748B-20H-3, 76-78 cm	174.86	41.79	<i>Subbotina spp.</i>	150-180	0.68		0.02	

*CK95 = Age scale of Cande and Kent (1995).

TABLE DR3. CHANGE IN THE GRADIENT OF CARBON ISOTOPES WITH SIZE ($\delta^{13}\text{C}/100$ MICRONS) OF PLANKTIC FORAMINIFERA

Sample	Depth (mbsf)	CK95* Age (Ma)	CK95 Age from Bohaty et al., 2009	<i>Acarinina</i> spp† ($\Delta\delta^{13}\text{C}/100$ microns)	<i>Morozovelloides</i> <i>crassatus</i> ($\Delta\delta^{13}\text{C}/100$ microns)	<i>Globigerinatheka</i> spp§ ($\Delta\delta^{13}\text{C}/100$ microns)	<i>Subbotina</i> spp ($\Delta\delta^{13}\text{C}/100$ microns)	Interval
1051A-8H-6, 5-7 cm	69.14	39.90	—	0.68	0.63	0.83	0.00	Post-MECO
1051B-9H-5, 5-7 cm	77.35	40.05	—	0.14	0.50	0.34	0.20	Peak MECO
1051B-9H-6, 75-77 cm	79.47	40.16	—	0.95	0.75	0.49	0.01	Initial' MECO
1051B-11H-4, 45-47 cm	92.25	40.40	—	0.55	0.67	0.66	0.23	Initial' MECO
1051A-12H-3, 131.5-133 cm	105.1	40.78	—	0.62	0.40	1.06	0.00	Initial' MECO
1051B-13H-5, 63-66 cm	111.4	40.87	—	0.70	0.00	0.80	—	Pre-MECO
748B-18H-7, 23-25 cm	161.33	39.40	39.35	0.44	—	0.27	0.03	Post-MECO
748B-18H-7, 32-34 cm	161.42	39.41	39.36	0.51	—	0.29	0.15	Post-MECO
748B-19H-2, 127-129 cm	164.37	40.09	40.04	0.19	—	0.19	-0.15	Peak MECO
748B-19H-3, 31-33 cm	164.91	40.13	40.08	0.14	—	0.18	-0.14	Peak MECO
748B-19H-3, 103-105 cm	165.63	40.18	40.13	0.07	—	—	—	Peak MECO
748B-19H-4, 103-105 cm	167.13	40.29	40.24	0.15	—	—	—	Initial' MECO?
748B-19H-5, 71-73 cm	168.31	40.35	40.30	0.15	—	—	—	Initial' MECO?
748B-19H-6, 55-57 cm	169.65	40.43	40.38	0.21	—	—	—	Initial' MECO?
748B-20H-1, 16-18 cm	171.26	41.46	41.41	0.37	—	0.52	-0.03	Pre-MECO
748B-20H-3, 76-78 cm	174.86	41.84	41.79	0.47	—	0.32	-0.16	Pre-MECO

Note: All $\delta^{13}\text{C}/100$ micron values are calculated independently for each taxon in every sample using linear regression and multiplying the resulting gradient by 100.

*CK95 = Age scale of Cande and Kent (1995).

†*Acarinina topilensis* and *Acarinina primitiva* analysed at ODP Sites 1051 and 748, respectively.

§*Globigerinatheka* index analysed at ODP Site 748.

TABLE DR4. CHANGE IN THE TEST SIZE- $\delta^{13}\text{C}$ GRADIENT ($\delta^{13}\text{C}/100$ MICRONS) OF MODERN AND PALEOGENE PLANKTIC FORAMINIFERA

Sample	Species	Age (Ma)	$\Delta\delta^{13}\text{C}/100$ microns	Data source
384, 11H-1, 128-136 cm	<i>Morozovella angulata</i>	Paleocene	0.55	Norris, 1996
384, 10H-CC	<i>Morozovella conicotruncata</i>	Paleocene	0.54	Norris, 1996
384, 6H-1, 30-32 cm	<i>Morozovella velascoensis</i>	Paleocene	0.74	Norris, 1996
384, 3H-4, 60-62 cm	<i>Morozovella acutaspira</i>	Paleocene	0.54	Norris, 1996
384, 6H-1, 30-32 cm	<i>Acarinina mckanni</i>	Paleocene	0.88	Norris, 1996
758A, 28-4, 24-26 cm	<i>Morozovella subbotinae</i>	Paleocene	0.46	D'Hondt et al., 1994
758A, 28-4, 24-26 cm	<i>Morozovella velascoensis</i>	Paleocene	0.35	D'Hondt et al., 1994
758A, 28-4, 24-26 cm	<i>Acarinina nitida</i>	Paleocene	0.62	D'Hondt et al., 1994
BOFS 31K	<i>Globigerinoides ruber</i> (white)	Modern	0.31	Elderfield et al., 2002
BOFS 31K	<i>Globigerinoides ruber</i> (pink)	Modern	-0.13	Elderfield et al., 2002
BOFS 31K	<i>Globigerinoides sacculifer</i>	Modern	0.67	Elderfield et al., 2002
BOFS 31K	<i>Orbulina universa</i>	Modern	0.54	Elderfield et al., 2002
61BC, 0-1 cm	<i>Globigerinoides ruber</i> (pink)	Modern	0.48	Bornemann and Norris, 2007
61BC, 0-1 cm	<i>Globigerinoides sacculifer</i>	Modern	0.41	Bornemann and Norris, 2007
KNR110, 1-3 cm	<i>Globigerinoides sacculifer</i>	Modern	0.42	Bornemann and Norris, 2007

Note: All $\delta^{13}\text{C}/100$ micron values are calculated independently for each taxon in every sample using linear regression and multiplying the resulting gradient by 100. For consistency and to avoid kinetic effects on $\delta^{13}\text{C}$ values at small test sizes we only use sieve size fractions from >180 microns. Based on test size- $\delta^{13}\text{C}$ gradients alone, the Paleogene muricates and globigerinethicks were acquiring and using symbionts in a similar manner and as effectively as modern cancellate spinose forms. However, we note that as highlighted in this study, test size- $\delta^{13}\text{C}$ gradients are likely to vary spatially and through time.

TABLE DR5. MAXIMUM TEST DIAMETER OF PLANKTIC FORAMINIFERA AT ODP SITE 1051

Sample	Depth (mbsf)	CK95* Age (Ma)	Taxon†	Maximum test diameter (microns)
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	580
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	537
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	512
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	509
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	506
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	506
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	504
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	486
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	481
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	478
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	468
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	465
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	465
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	459
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	458
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	449
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	431
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	426
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Acarinina topilensis</i>	402
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Morozovelloides crassatus</i>	498
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Morozovelloides crassatus</i>	489
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Morozovelloides crassatus</i>	468
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Morozovelloides crassatus</i>	444
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	411
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	403
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	403
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	402
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	401
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	397
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	396
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	395
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	395
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	393
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	393
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	391
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	389
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	388
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	386
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	386
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	384
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	383
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	372
1051A-8H-6, 5-7 cm	69.14	39.90	<i>Globigerinatheka</i> spp	359
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Acarinina praetopilensis-topilensis</i>	411
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Acarinina praetopilensis-topilensis</i>	374

1051B-9H-5, 5-7 cm	77.35	40.05	<i>Acarinina praetopilensis-topilensis</i>	306
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Acarinina praetopilensis-topilensis</i>	282
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	561
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	553
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	545
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	521
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	511
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	505
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	504
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	504
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	501
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	496
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	489
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	480
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	479
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	479
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	477
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	476
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	460
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	458
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Morozovelloides crassatus</i>	433
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	667
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	603
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	574
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	569
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	563
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	542
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	541
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	541
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	539
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	535
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	532
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	530
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	509
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	476
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	453
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	448
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	437
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	426
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	426
1051B-9H-5, 5-7 cm	77.35	40.05	<i>Globigerinatheka</i> spp	418
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Acarinina topilensis</i>	541
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Acarinina topilensis</i>	503
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Acarinina topilensis</i>	497
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Acarinina topilensis</i>	489
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Acarinina topilensis</i>	473
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Acarinina topilensis</i>	472
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Acarinina topilensis</i>	463
1051B-9H-6, 75-77 cm	79.47	40.16	<i>Acarinina topilensis</i>	462

1051B-9H-6, 75-77 cm	79.47	40.16	<i>Globigerinatheka</i> spp	455
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	588
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	542
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	538
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	528
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	506
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	506
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	495
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	494
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	480
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	471
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	461
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	460
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	456
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	438
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Acarinina topilensis</i>	431
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	529
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	526
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	519
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	519
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	506
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	503
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	501
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	494
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	487
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Morozovelloides crassatus</i>	478
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	495
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	412
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	411
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	406
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	404
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	403
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	396
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	395
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	393
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	391
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	389
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	388
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	388
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	384
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	384
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	378
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	378
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	378
1051B-11H-4, 45-47 cm	92.25	40.40	<i>Globigerinatheka</i> spp	377
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Acarinina topilensis</i>	617
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Acarinina topilensis</i>	614
1051A-12H-3, 131.5-133 cm	105.12	40.78	<i>Acarinina topilensis</i>	596

1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	384
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	381
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	380
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	380
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	379
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	379
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	378
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	367
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	363
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	362
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	361
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	360
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	359
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	355
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	353
1051B-13H-5, 63-66 cm	111.43	40.87	<i>Globigerinatheka</i> spp	352

*CK95 = Age scale of Cande and Kent (1995).

†A broad species concept was adopted for *A. topilensis* (see Fig. DR4).

References

- Bornemann, A. and Norris, R.D., 2007. Size-related stable isotope changes in Late Cretaceous planktic foraminifera: Implications for paleoecology and photosymbiosis: *Marine Micropaleontology* v. 65, p. 32-42.
- Cande, S.C. and Kent, D.V., 1995. Revised calibration of the geomagnetic polarity timescale for the Late Cretaceous and Cenozoic: *Journal of Geophysical Research*, v. 100, p. 6093-6095.
- D'Hondt, S., Zachos, J.C. and Schultz, G., 1994, Stable isotopic signals and photosymbiosis in Late Paleocene planktic foraminifera: *Paleobiology*, v. 20, p. 391-406.
- Elderfield, H., Vautravers, M., Cooper, M., 2002. The relationship between Mg/Ca, Sr/Ca, $\delta^{18}\text{O}$, and $\delta^{13}\text{C}$ of species of planktonic foraminifera. *Geochemistry, Geophysics, Geosystems*, v. 3, doi:10.1029/2001GC000194.
- Norris, R.D., 1996, Symbiosis as an Evolutionary Innovation in the Radiation of Paleocene Planktic Foraminifera; *Paleobiology*, v. 22, p. 461-480.
- Pearson et al., 2001, Warm tropical seas surface temperatures in the Late Cretaceous and Eocene epochs; *Nature*, v. 413, p. 481-488.
- Sexton, P.F., Wilson, P.A. and Pearson, P.N., 2006, Microstructural and geochemical perspectives on planktic foraminiferal preservation: 'Glassy' versus 'Frosty', *Geochemistry, Geophysics, Geosystems*, 7, Q12P19, doi:10.1029/2006GC001291.