

# Synergistic Effect of Sulindac and Simvastatin on Apoptosis in Lung Cancer A549 Cells through AKT-Dependent Downregulation of Survivin

Young-Suk Kim, PhD  
Chang-Hwan Seol, MD  
Jae-Wan Jung, MD  
Su-Jin Oh, MD  
Ki-Eun Hwang, MD  
Hwi-Jung Kim, MD  
Eun-Taik Jeong, PhD  
Hak-Ryul Kim, MD, PhD

Department of Internal Medicine,  
Institute of Wonkwang Medical Science,  
Wonkwang University School of Medicine,  
Iksan, Korea

Correspondence: Hak-Ryul Kim, MD, PhD  
Department of Internal Medicine,  
Wonkwang University School of Medicine,  
460 Iksan-daero, Iksan 570-974, Korea  
Tel: 82-63-859-2583  
Fax: 82-63-855-2025  
E-mail: kshryj@wonkwang.ac.kr

Received October 7, 2013  
Accepted December 27, 2013  
Published online October 27, 2014

\*Young-Suk Kim and Chang-Hwan Seol  
contributed equally to this work.

## Purpose

Non-steroidal anti-inflammatory drugs (NSAIDs) and statins are potential chemopreventive or chemotherapeutic agents. The mechanism underlying the deregulation of survivin by NSAIDs and statins in human non-small cell lung cancer cells has not been elucidated. In this study, we investigated the synergistic interaction of sulindac and simvastatin in lung cancer A549 cells.

## Materials and Methods

Cell viability was measured by an MTT assay, while the expression of apoptotic markers, AKT, and survivin in response to sulindac and simvastatin was examined by Western blotting. DNA fragmentation by apoptosis was analyzed by flow cytometry in A549 cells. Reactive oxygen species (ROS) generation was measured by flow cytometry using H<sub>2</sub>DCFDA and MitoSOX Red, and the effects of pretreatment with N-acetylcysteine were tested. The effects of AKT on survivin expression in sulindac- and simvastatin-treated cells were assessed. Survivin was knocked down or overexpressed to determine its role in apoptosis induced by sulindac and simvastatin.

## Results

Sulindac and simvastatin synergistically augmented apoptotic activity and intracellular ROS production in A549 cells. Inhibition of AKT by siRNA or LY294002 inhibited survivin, while AKT overexpression markedly increased survivin expression, even in the presence of sulindac and simvastatin. Moreover, survivin siRNA enhanced sulindac- and simvastatin-induced apoptosis. In contrast, survivin upregulation protected against sulindac- and simvastatin-induced apoptosis.

## Conclusion

Combined treatment with sulindac and simvastatin augmented their apoptotic potential in lung cancer cells through AKT signaling-dependent downregulation of survivin. These results indicate that sulindac and simvastatin may be clinically promising therapies for the prevention of lung cancer.

## Key words

Sulindac, Simvastatin, Apoptosis, Lung neoplasms,  
Oncogene protein AKT, Survivin

## Introduction

Many strategies shown to be effective in the chemoprevention of cancer or the precancerous stage change the signaling

pathways targeted by therapies currently being used in the lung; however, no drug effective as a lung cancer chemotherapy agent has been identified to date [1]. Accordingly, prevention of lung cancer may be a better approach, and the use of multiple agents that target different molecules

involved in carcinogenesis holds great promise [2,3]. Several studies have shown that low-dose administration of a combination of cancer preventive agents with different modes of action may produce synergistic effects on efficacy and minimize possible side effects associated with high-dose administration.

Non-steroidal anti-inflammatory drugs (NSAIDs) have potential chemotherapeutic efficacy against carcinogenesis that occurs via inhibition of cyclooxygenase (COX) enzyme [4], which catalyzes the production of prostaglandins from arachidonic acid and therefore plays an important role in inflammation and cancer [5]. There is accumulating evidence of the cancer preventive effects of NSAIDs. However, the relatively high doses required for the observed chemopreventive effect in human trials may discourage the long-term use of NSAIDs for lung cancer prevention owing to the possibility of increased risk of serious gastrointestinal and cardiovascular side effects [6,7]. An alternative approach is to combine NSAIDs with other chemopreventive agents.

Statins, or 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors, are a class of drugs that inhibit the rate-limiting step of the mevalonate pathway [8]. In addition to their use in the treatment of lipid disorders, statins have been investigated for their anticarcinogenic effects in several models, including carcinomas of the colon and rectum, prostate, breast, lung, and skin [9,10]. The mechanism of apoptosis induction is well-known, and statin-induced suppression of angiogenesis via vascular endothelial growth factor and of tumor invasiveness and metastatic potential through interaction with adhesion molecules have shown anticancer effects [11]. The prevention of cardiovascular side effects of these statins, while in theory, as a safe drug is already widely used in clinical therapy with NSAIDs as interesting combination. In recent studies, NSAIDs and statins showed synergistic effects in combination with other therapeutics, such as peroxisome proliferator-activated receptor  $\gamma$  ligands, inhibitors of the epidermal growth factor receptor family, TRAIL receptor ligands, cisplatin, and doxorubicin [12]. We previously suggested that sulindac and simvastatin-induced reactive oxygen species (ROS) generation in A549 lung cancer cells causes ROS accumulation in mitochondria, triggering the release of apoptogenic molecules from the mitochondria to the cytosol, leading to caspase activation and cell death [13]. However, studies of the mechanism by which NSAIDs and statins combine to induce apoptosis and inhibit the growth of lung cancer cells are insufficient. Furthermore, the mechanism underlying deregulation of survivin by NSAIDs and statins in human non-small cell lung cancer cells has not been elucidated. In this study, we demonstrated the synergistic interaction of sulindac and simvastatin and elucidated the mechanisms underlying this synergy.

## Materials and Methods

### 1. Reagents

RPMI 1640, fetal bovine serum (FBS), and antibiotics were purchased from Gibco-BRL (Grand Island, NY). Sulindac, simvastatin, propidium iodide (PI), 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide (MTT), bichinchonic acid, dimethyl sulfoxide (DMSO), N-acetylcysteine (NAC), and LY294002 were purchased from Sigma-Aldrich (St. Louis, MO). The following primary antibodies were used caspase-3, -8, -9, poly(ADP-ribose) polymerase (PARP; Santa Cruz Biotechnology, Santa Cruz, CA), serine/threonine protein kinase (AKT), phospho-AKT, survivin, X-linked inhibitor of apoptosis protein (XIAP), and glyceraldehyde 3-phosphate dehydrogenase (GAPDH; Cell Signaling Technology, Beverly, MA). Anti-rabbit IgG-conjugated horseradish peroxidase (HRP) antibodies and an enhanced chemiluminescent (ECL) kit were purchased from Amersham Pharmacia Biotech (Buckinghamshire, UK).

### 2. Cell culture


Human lung cancer A549 cells were purchased from the Korean Cell Line Bank (Seoul, Korea). Cells were maintained in RPMI-1640 supplemented with 10% FBS, 100 U/mL penicillin, and 100  $\mu$ g/mL streptomycin in a humidified atmosphere of 5% CO<sub>2</sub> at 37°C.

### 3. Cell viability assay

The effects of sulindac, simvastatin, and sulindac plus simvastatin treatment on cell viability were determined by MTT assay. Briefly, A549 cells were seeded at a density of  $5 \times 10^3$  cells per well in 96-well plates in triplicate. After 24 hours, sulindac, simvastatin, and sulindac plus simvastatin were added at various concentrations, and the cells were incubated for 48 hours. To determine the cell viability, MTT was added to the cell suspension for 4 hours. After 3 washes with phosphate buffered saline (PBS; pH, 7.4) the insoluble formazan product was dissolved in DMSO. The optical density (OD) of each well was then measured using a microplate reader (Titertek Multiskan, Flow Laboratories, North Ryde, Australia) at 590 nm. The OD derived from formazan production in control cells was defined as 100% viability, and all other measurements were expressed as a percentage of the control cell value.

### 4. Detection of apoptosis

The degree of apoptosis was determined by measuring the



**Fig. 1.** Effect of combined treatment with sulindac and simvastatin on the growth of A549 cells. (A) Cells were treated with different concentrations of sulindac and simvastatin for 48 hours, after which viability was determined by an MTT assay. Survival relative to that of control cells is presented. (B) Cells were incubated with 300  $\mu$ M sulindac and/or 5  $\mu$ M simvastatin for 48 hours, after which apoptosis was quantified by DNA content analysis. The percentage of hypodiploid DNA content is indicated for each test condition. Bars represent the mean  $\pm$  standard deviation of three independent experiments. Sim, simvastatin; Sul, sulindac. \* $p < 0.05$  compared to control.

percentage of cells showing hypodiploid DNA content upon flow cytometric analysis following staining of the cells with PI using the method described by Crissman and Steinkamp [14]. Cell cycle analysis was conducted using a FACScan equipped with the CellQuest software (Becton Dickinson, San Jose, CA).

## 5. Determination of ROS

To measure intracellular ROS, A549 cells were incubated in the dark with 10  $\mu$ mol/L 5-(and-6)-carboxy-2',7'-dichlorodihydrofluorescein diacetate (carboxy- $H_2$ DCFDA; Molecular Probes, Eugene, OR) for 30 minutes. Cells were

then washed, scraped gently, resuspended in PBS, and kept on ice for immediate analysis by FACScan flow cytometry using an argon laser (488 nm) for excitation. Green fluorescence due to intracellularly trapped DCF was collected on the FL1 channel on a log scale. Data were acquired and analyzed with the CellQuest program (Becton Dickinson).

To measure mitochondria-derived ROS, the mitochondria-targeted,  $O_2^-$  sensitive, hydroethidine analog probe MitoSOX (M36008, Invitrogen Life Technologies, Carlsbad, CA) was used to determine relative  $O_2^-$  levels. Briefly, cells were incubated with 5  $\mu$ M MitoSOX for 10 minutes in RPMI-1640, washed twice with PBS, and then analyzed by FACScan flow cytometry.

## 6. Immunoblot analysis

Cells were harvested and lysed in protein extraction solution (PRO-PREP; Intron Biotechnology, Seoul, Korea) for 20 minutes on ice. Next, cell lysates were centrifuged at 15,000 rpm for 20 minutes at 4°C, after which the supernatant was mixed with a one-fifth volume of 5× sodium dodecyl sulfate (SDS) sample buffer, boiled for 5 minutes, and then separated by 10% SDS–polyacrylamide gel electrophoresis. Following electrophoresis, proteins were transferred to a membrane that had been blocked with TBS-T (25 mM Tris [pH, 7.6], 138 mM NaCl, and 0.05% Tween-20) containing 5% skim milk and incubated with primary antibodies at dilutions of 1:1,000–1:5,000. After a series of washes, the membranes were further incubated with secondary antibody (at 1:2,000–1:10,000) conjugated with HRP. The immunoreactive signal was detected using an ECL detection system (Amersham Bioscience, Buckinghamshire, UK).

## 7. Small interference RNA transfection

Transcriptional expression was specifically suppressed by the introduction of a 21-nucleotide duplex siRNA that targeted the nucleotides of the AKT or survivin mRNA coding sequence [15]. Briefly, cells ( $10^5$  cells per well) were plated in 6-well plates and transiently transfected with 2  $\mu$ g per well of AKT or survivin siRNA (Cell Signaling Technology) mixed with X-tremeGENE siRNA transfection reagent (Roche Applied Science, Penzberg, Germany) according to the manufacturer's directions. Silencer Negative Scramble siRNA (Roche Applied Science) was introduced into the cells as a negative control using the same protocol.

## 8. Plasmid preparation overexpressing AKT or survivin

Human survivin cDNA was obtained by reverse transcription–PCR of RNA derived from A549 cells with the following primers based on GenBank accession No. U75285 (forward,

5'-GGGAATTCATGGGTGCCCCGACGTTGCC-3'; reverse 5'-CTCTCGAGTCAATCCATGGCAGCCAGCT-3'). PCR products were digested with *Eco*RI and *Xho*I and ligated into pcDNA3.1myc [16]. The myr-AKT1 cDNA was subcloned into the *Eco*RI and *Xho*I site of the pcDNA3.1myr-myc. For transient transfection, A549 cells were transfected with 2  $\mu$ g Myr-AKT (catalogue No. 9008, Addgene, Cambridge, MA) or control vector using Lipofectamine 2000.

## 9. Statistical analysis

Each experiment was performed at least three times, and all values are presented as the mean $\pm$ SD of triplicate experiments. A Student's t test was used to determine the statistical significance of the results. A  $p < 0.05$  was considered statistically significant.


# Results

## 1. Sulindac and simvastatin synergistically augment the apoptotic activity of A549 cells

A549 cells were treated with sulindac and simvastatin, and their viability was measured by MTT assay. As shown in Fig. 1A, the combination of sulindac and simvastatin showed increased dose-dependent growth-inhibitory effects on A549 cells relative to treatment with sulindac or simvastatin alone. We next examined whether sulindac and simvastatin induced apoptosis in A549 cells. DNA fragmentation by apoptosis was considered the percentage of hypodiploid DNA content in A549 cells stained with PI measured by flow cytometry (Fig. 1B). The combination of sulindac and simvastatin caused a significantly greater increase in the accumulation of hypodiploid DNA content than either therapy alone. Taken together, these data indicate that combined treatment with sulindac and simvastatin triggers the apoptotic signaling pathway in A549 cells.

## 2. Combination of sulindac and simvastatin enhances intracellular ROS production

We next investigated the upstream regulatory mechanisms leading to the induction of apoptosis by combined treatment with sulindac and simvastatin. Intracellular ROS generation was assessed by flow cytometry using the total ROS marker  $H_2$ DCFDA and the mitochondrial superoxide marker MitoSOX Red. Elevated ROS levels were detected 48 hours after combined treatment with sulindac and simvastatin (Fig. 2, blue bars), with marked ROS generation being observed




**Fig. 2.** Contribution of reactive oxygen species (ROS) generation to the anticancer activity of combination therapy in A549 cells. Cells were treated with 300  $\mu$ M sulindac and/or 5  $\mu$ M simvastatin for 48 hours, then loaded with H<sub>2</sub>DCFDA (blue bars) and MitoSOX Red (green bars). Fluorescence measurements were carried out using a FACSCalibur flow cytometer. The data represent the mean  $\pm$  standard deviation of three independent experiments. Sim, simvastatin; Sul, sulindac. \* $p < 0.05$  compared to the control.

in cells treated with both sulindac and simvastatin relative to cells treated with either compound alone. Flow cytometry measurement using MitoSOX revealed that intracellular O<sub>2</sub><sup>-</sup> levels increased significantly, which correlated well with the onset of total ROS production (Fig. 2, green bars). FACS analysis demonstrated an increase in the fluorescence intensity of MitoSOX Red in A549 cells. Specifically, the M<sub>2</sub> levels were 9% and 6%, respectively, after 10 minutes of exposure to 300  $\mu$ M sulindac or 5  $\mu$ M simvastatin, but the M<sub>2</sub> level was 18% in cells treated with both drugs.


### 3. Pretreatment with NAC prevents apoptosis and downregulates survivin by sulindac and simvastatin

To identify the molecular mechanism underlying apoptosis induced by combined treatment with sulindac and simvastatin, we examined the expression of AKT and the inhibitor of apoptosis protein (IAP) family members survivin and XIAP by immunoblot analysis in A549 cells treated with sulindac and/or simvastatin for 24 hours. As shown in Fig. 3A, treatment of A549 cells with sulindac and simvastatin resulted in significantly lower survivin levels relative to treatment with either drug alone, but had no effect on the expression of XIAP. In addition, combined treatment with sulindac and simvastatin decreased phosphorylated AKT expression. To determine whether elevated ROS participates



**Fig. 3.** Effect of *N*-acetylcysteine (NAC) on sulindac- and simvastatin-induced apoptosis and survivin expression. (A) Cells were treated with 300  $\mu$ M sulindac and/or 5  $\mu$ M simvastatin for 48 hours. (B) Cells were pretreated with or without the free radical scavenger NAC. The expression of phosphorylated AKT, AKT, survivin, X-linked inhibitor of apoptosis protein (XIAP), cleaved poly(ADP-ribose) polymerase (PARP), cleaved caspase-3, -8, -9, and glyceraldehyde 3-phosphate dehydrogenase (GAPDH) was measured in the cell lysates. Data are representative of two individual experiments. Sim, simvastatin; Sul, sulindac.






**Fig. 4.** Effect of AKT on survivin expression in sulindac- and simvastatin-treated cells. (A, B) Effect of AKT inhibition on survivin expression. Cells were pretreated with the phosphoinositide-3 kinase inhibitor LY294002 or transfected with AKT siRNA and then further incubated with sulindac and/or simvastatin for 48 hours. Scrambled RNA containing the same number of each nucleotide found in AKT siRNA was used as the transfection control. (C) Effect of AKT activation on survivin expression. Cells were transiently transfected with constructs expressing constitutively active AKT and then incubated with sulindac and simvastatin for 48 hours. The expression of phosphorylated AKT, AKT, cleaved poly(ADP-ribose) polymerase (PARP), cleaved caspase-3, -8, -9, and survivin was measured in the cell lysates. Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) was used as a loading control. Data are representative of two individual experiments.

in apoptosis induced by the drug combination, cells were incubated with a free radical scavenger, NAC, before treatment. Western blot analysis of A549 cell lysates (Fig. 3B) showed that the combination of sulindac and simvastatin enhanced expression of the 85-kDa form of PARP and expression of caspase-3, -8, and -9, while pretreatment with NAC blocked this effect. Together, these findings indicate that ROS generation plays a primary role in apoptosis induced by sulindac and simvastatin.

#### 4. Downregulation of survivin by sulindac and simvastatin is associated with suppression of the AKT signaling pathway


Because survivin downregulation involves the AKT signaling pathway, we tested the effects of the phosphatidylinositol 3-kinase (PI3K) inhibitor, LY294002, on A549 cells. To accomplish this, cells were pre-treated with or without LY294002 and then treated with sulindac and simvastatin,



**Fig. 5.** Downregulation of survivin with siRNA augments apoptosis induced by sulindac and simvastatin. (A) Cells were transfected with siRNA specific for survivin, and survivin protein expression was assessed by immunoblot analysis 24 hours after transfection. Scrambled RNA containing the same number of each nucleotide found in survivin siRNA was used as the transfection control. The expression of survivin, cleaved poly(ADP-ribose) polymerase (PARP), and cleaved caspase-9 was measured in the cell lysates. (B) Apoptosis was evaluated by green fluorescent protein-annexin V+propidium iodide. Bars represent the mean±standard deviation of three independent experiments. \**p* < 0.05 for survivin-siRNA and combined treatment with sulindac and simvastatin versus either scrambled siRNA and combined treatment or combined treatment alone. Sul, sulindac; Sim, simvastatin; siSCR, scrambled siRNA; siSUR, survivin-siRNA; GAPDH, glyceraldehyde 3-phosphate dehydrogenase.

after which the cell lysates were assessed by Western blotting. As shown in Fig. 4A, pre-treatment with LY294002 decreased the activation of AKT, while treatment of cells with the combination of sulindac and simvastatin markedly decreased the levels of phospho-AKT. This effect was associated with significantly increased apoptosis, as

evidenced by the cleavage of PARP and caspase-3, -8, and -9. The immunoreactive band of survivin was markedly decreased in cells treated with LY294002, as well as in cells treated with the combination of sulindac and simvastatin. We next knocked down AKT using siRNA. As shown in Fig. 4B, the introduction of AKT siRNA abrogated AKT protein



**Fig. 6.** Effect of survivin overexpression on apoptosis induced by sulindac and simvastatin in A549 cells. (A) Cells were transfected with a survivin gene expression vector, pcDNA3-myc-survivin, and survivin protein expression was assessed by immunoblot analysis 48 hours after transfection. The expression of survivin, cleaved poly(ADP-ribose) polymerase (PARP), and cleaved caspase-3, -8, and -9 was measured in the cell lysates. (B) Apoptosis was evaluated by green fluorescent protein-annexin V+propidium iodide. The data represent the mean±standard deviation of three independent experiments. GAPDH, glyceraldehyde 3-phosphate dehydrogenase; Sul, sulindac; Sim, simvastatin; SUR, survivin. \* $p < 0.05$  for the indicated comparisons.

expression 48 hours after transfection. Combined treatment with AKT siRNA, sulindac, and simvastatin resulted in greater cell death than combined treatment with scrambled siRNA, sulindac, and simvastatin. Next, we tested the effect

of AKT activation by expressing a constitutively active form of AKT. As shown in Fig. 4C, treatment of A549 cells overexpressing AKT-myc with sulindac and simvastatin for 48 hours resulted in increased AKT and survivin relative to the



control group. AKT overexpression markedly increased the p-AKT level, even in the presence of sulindac and simvastatin. The expression of survivin was also markedly elevated when compared with combination treatment. Collectively, these data suggest that the AKT pathway regulates survivin expression.

### 5. Downregulation of survivin augments apoptosis by sulindac and simvastatin

To determine the role of survivin in sulindac- and simvastatin-induced apoptosis in lung cancer, we knocked down survivin with siRNA. Introduction of survivin siRNA attenuated survivin protein expression 48 hours after transfection. No reduction in survivin protein was observed in cells transfected with the scrambled siRNA, which contained the same number of each nucleotide found in survivin siRNA. Cell death was higher in cells transfected with survivin siRNA than in those transfected with control siRNA. Moreover, the combination of sulindac, simvastatin, and survivin siRNA resulted in greater cell death than the combination of sulindac, simvastatin, and control siRNA (Fig. 5A). The number of annexin V-positive cells also increased in the survivin siRNA transfectants, and the combination of sulindac, simvastatin, and survivin siRNA increased the percentage of annexin V-positive cells to 32.4% (Fig. 5B). Taken together, these data indicate that the induction of apoptosis by sulindac and simvastatin is due, at least in part, to the downregulation of survivin.

### 6. Upregulation of survivin attenuates apoptosis induced by sulindac and simvastatin

We next determined whether survivin upregulation played a role in apoptosis induced by sulindac and simvastatin. To accomplish this, we introduced a survivin gene expression vector, pcDNA3-myc-survivin, into A549 cells. The survivin gene transfectants expressed myc-tagged survivin protection in addition to endogenous survivin protection. The expression of cleaved PARP, caspase-3, -8, and -9 (Fig. 6A) was suppressed in survivin-overexpressing cells. Moreover, annexin V staining revealed that the number of apoptotic cells in control vector transfectants treated with sulindac and simvastatin for 48 hours was 34.2% higher than in untreated A549 cells. In contrast, the number of apoptotic cells in survivin gene transfectants treated with sulindac and simvastatin decreased by 19.5% (Fig. 6B). Together, these data indicate that the reduction of apoptosis by sulindac and simvastatin was due, at least in part, to the upregulation of survivin.

## Discussion

In the present study, we showed that sulindac and simvastatin synergistically inhibit growth and induce apoptosis of A549 lung cancer cells, but that these effects were not observed when either drug was administered alone. The synergistic effect of sulindac and simvastatin on apoptosis has the potential to increase chemoprevention and treatment efficacy in lung cancer, which could enable a reduction in the dose of sulindac to minimize possible detrimental side effects. Additionally, simvastatin could be beneficial to cardiovascular health [10].

Apoptosis is one of the most vital pathways through which chemopreventive agents inhibit the overall growth of cancer cells; thus, it is important to determine whether the inhibition of cell proliferation and induction of apoptosis by chemopreventive agents are associated with the downregulation of antiapoptotic genes, such as survivin, which plays a critical role in cancer cell progression [17].

Many chemotherapeutic strategies have been designed to significantly increase cellular ROS levels with the goal of inducing irreparable tumor cell damage and death. Intracellular oxidative status is important for simvastatin sensitivity, and sulindac increases ROS levels more efficiently than selective COX-2 inhibitors [18]. A novel fluoroprobe, MitoSOX, was recently introduced for the selective detection of ROS in the mitochondria of live cells, and a quantitative method for the detection of mitochondrial ROS generation by flow cytometry was established [19]. Here, we adopted a staining technique using H<sub>2</sub>DCFDA and MitoSOX to concurrently detect intracellular and mitochondrial ROS generation in sulindac- and simvastatin-treated A549 cells. This simple method enables convenient observation of quantitative changes in ROS, both inside and outside the mitochondria. For novel therapeutic strategies it will be necessary to fine-tune intracellular ROS signaling to enable effective therapeutic gains.

Survivin, an IAP, regulates mitosis and inhibits apoptosis, primarily by suppressing the activity of caspase-3 [20]. Disruption of survivin-microtubule interactions results in loss of survivin anti-apoptotic function, increased caspase-3 activity, and cleavage of PARP. In addition, inhibition of the function of survivin induced apoptosis and increases in the responsiveness to chemotherapy or radiation therapy have been recognized as targets during the treatment of cancer [21]. There is evidence that the AKT-mediated survival pathway inhibits apoptosis by stimulating survivin synthesis in various cancer cell lines [22,23]. Therefore, we used siRNA-mediated knockdown of AKT or the PI3K inhibitor

LY294002 in combination with sulindac and simvastatin treatment to determine whether the mechanism of enhanced apoptosis by combined treatment involves inhibition of the AKT pathway in lung cancer cells. It is known that the survivin signaling pathway to the insulin-like growth factor/mammalian target of rapamycin signaling pathway occurs via rapid changes in the mRNA transcription step [24]. Ras oncogene-related proteins or signal instruments and the activity of three transcription factors of the Wnt signaling pathway have been reported [25]. For future therapy, the upper signaling pathways that directly control survivin activity should be investigated.

## Conclusion

Our study elucidated the involvement of AKT signaling-dependent downregulation of survivin in the anti-cancer effects of sulindac and simvastatin against lung cancer. These

data provide a theoretical basis for future clinical use of sulindac and simvastatin in lung cancer chemoprevention and therapy and important information that will be useful for mechanistic and biomarker studies in clinical trials.

## Conflicts of Interest

Conflict of interest relevant to this article was not reported.

## Acknowledgments

This work was supported by a grant from the Korean Association for the Study of Lung Cancer Research Fund in 2011.

## References

1. Omenn GS. Chemoprevention of lung cancer is proving difficult and frustrating, requiring new approaches. *J Natl Cancer Inst.* 2000;92:959-60.
2. Narayanan BA, Narayanan NK, Pittman B, Reddy BS. Regression of mouse prostatic intraepithelial neoplasia by nonsteroidal anti-inflammatory drugs in the transgenic adenocarcinoma mouse prostate model. *Clin Cancer Res.* 2004;10:7727-37.
3. Ulrich CM, Bigler J, Potter JD. Non-steroidal anti-inflammatory drugs for cancer prevention: promise, perils and pharmacogenetics. *Nat Rev Cancer.* 2006;6:130-40.
4. Albayrak A, Polat B, Cadirci E, Hacimuftuoglu A, Halici Z, Gulapoglu M, et al. Gastric anti-ulcerative and anti-inflammatory activity of metyrosine in rats. *Pharmacol Rep.* 2010;62:113-9.
5. Krysan K, Reckamp KL, Sharma S, Dubinett SM. The potential and rationale for COX-2 inhibitors in lung cancer. *Anticancer Agents Med Chem.* 2006;6:209-20.
6. Solomon SD, McMurray JJ, Pfeffer MA, Wittes J, Fowler R, Finn P, et al. Cardiovascular risk associated with celecoxib in a clinical trial for colorectal adenoma prevention. *N Engl J Med.* 2005;352:1071-80.
7. Psaty BM, Potter JD. Risks and benefits of celecoxib to prevent recurrent adenomas. *N Engl J Med.* 2006;355:950-2.
8. Goldstein JL, Brown MS. Regulation of the mevalonate pathway. *Nature.* 1990;343:425-30.
9. Chan KK, Oza AM, Siu LL. The statins as anticancer agents. *Clin Cancer Res.* 2003;9:10-9.
10. Demierre MF, Higgins PD, Gruber SB, Hawk E, Lippman SM. Statins and cancer prevention. *Nat Rev Cancer.* 2005;5:930-42.
11. Cafforio P, Dammacco F, Gernone A, Silvestris F. Statins activate the mitochondrial pathway of apoptosis in human lymphoblasts and myeloma cells. *Carcinogenesis.* 2005;26:883-91.
12. Jalving M, Koornstra JJ, De Jong S, De Vries EG, Kleibeuker JH. Review article: the potential of combinational regimen with non-steroidal anti-inflammatory drugs in the chemoprevention of colorectal cancer. *Aliment Pharmacol Ther.* 2005;21:321-39.
13. Hwang KE, Park C, Kwon SJ, Kim YS, Park DS, Lee MK, et al. Synergistic induction of apoptosis by sulindac and simvastatin in A549 human lung cancer cells via reactive oxygen species-dependent mitochondrial dysfunction. *Int J Oncol.* 2013;43:262-70.
14. Crissman HA, Steinkamp JA. Cell cycle-related changes in chromatin structure detected by flow cytometry using multiple DNA fluorochromes. *Eur J Histochem.* 1993;37:129-38.
15. Elbashir SM, Harborth J, Lendeckel W, Yalcin A, Weber K, Tuschl T. Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature.* 2001;411:494-8.
16. Tamm I, Wang Y, Sausville E, Scudiero DA, Vigna N, Oltersdorf T, et al. IAP-family protein survivin inhibits caspase activity and apoptosis induced by Fas (CD95), Bax,

- caspace, and anticancer drugs. *Cancer Res.* 1998;58:5315-20.
17. Krajewska M, Krajewski S, Banares S, Huang X, Turner B, Bubendorf L, et al. Elevated expression of inhibitor of apoptosis proteins in prostate cancer. *Clin Cancer Res.* 2003;9:4914-25.
  18. Minami T, Adachi M, Kawamura R, Zhang Y, Shinomura Y, Imai K. Sulindac enhances the proteasome inhibitor bortezomib-mediated oxidative stress and anticancer activity. *Clin Cancer Res.* 2005;11:5248-56.
  19. Mukhopadhyay P, Rajesh M, Yoshihiro K, Hasko G, Pacher P. Simple quantitative detection of mitochondrial superoxide production in live cells. *Biochem Biophys Res Commun.* 2007;358:203-8.
  20. Li F, Ambrosini G, Chu EY, Plescia J, Tognin S, Marchisio PC, et al. Control of apoptosis and mitotic spindle checkpoint by survivin. *Nature.* 1998;396:580-4.
  21. Kaneko R, Tsuji N, Asanuma K, Tanabe H, Kobayashi D, Watanabe N. Survivin down-regulation plays a crucial role in 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor-induced apoptosis in cancer. *J Biol Chem.* 2007;282:19273-81.
  22. Ohashi H, Takagi H, Oh H, Suzuma K, Suzuma I, Miyamoto N, et al. Phosphatidylinositol 3-kinase/Akt regulates angiotensin II-induced inhibition of apoptosis in microvascular endothelial cells by governing survivin expression and suppression of caspase-3 activity. *Circ Res.* 2004;94:785-93.
  23. Fornaro M, Plescia J, Chheang S, Tallini G, Zhu YM, King M, et al. Fibronectin protects prostate cancer cells from tumor necrosis factor-alpha-induced apoptosis via the AKT/survivin pathway. *J Biol Chem.* 2003;278:50402-11.
  24. Vaira V, Lee CW, Goel HL, Bosari S, Languino LR, Altieri DC. Regulation of survivin expression by IGF-1/mTOR signaling. *Oncogene.* 2007;26:2678-84.
  25. You L, He B, Xu Z, Uematsu K, Mazieres J, Mikami I, et al. Inhibition of Wnt-2-mediated signaling induces programmed cell death in non-small-cell lung cancer cells. *Oncogene.* 2004;23:6170-4.