



## Synovitis detected on magnetic resonance imaging and its relation to pain and cartilage loss in knee osteoarthritis

Catherine L Hill, David J Hunter, Jingbo Niu, Margaret Clancy, Ali Guermazi, Harry Genant, Daniel Gale, Andrew Grainger, Philip Conaghan and David T Felson

*Ann Rheum Dis* 2007;66;1599-1603; originally published online 9 May 2007;  
doi:10.1136/ard.2006.067470

---

Updated information and services can be found at:

<http://ard.bmj.com/cgi/content/full/66/12/1599>

---

*These include:*

### References

This article cites 23 articles, 5 of which can be accessed free at:

<http://ard.bmj.com/cgi/content/full/66/12/1599#BIBL>

1 online articles that cite this article can be accessed at:

<http://ard.bmj.com/cgi/content/full/66/12/1599#otherarticles>

### Rapid responses

You can respond to this article at:

<http://ard.bmj.com/cgi/eletter-submit/66/12/1599>

### Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

---

### Notes

---

To order reprints of this article go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to *Annals of the Rheumatic Diseases* go to:

<http://journals.bmj.com/subscriptions/>

## EXTENDED REPORT

## Synovitis detected on magnetic resonance imaging and its relation to pain and cartilage loss in knee osteoarthritis

C L Hill, D J Hunter, J Niu, M Clancy, A Guermazi, H Genant, D Gale, A Grainger, P Conaghan, D T Felson

*Ann Rheum Dis* 2007;66:1599–1603. doi: 10.1136/ard.2006.067470

**Objective:** To examine the relationship between longitudinal fluctuations in synovitis with change in pain and cartilage in knee osteoarthritis.

**Methods:** Study subjects were patients 45 years of age and older with symptomatic knee osteoarthritis from the Boston Osteoarthritis of the Knee Study. Baseline and follow-up assessments at 15 and 30 months included knee magnetic resonance imaging (MRI), BMI and pain assessment (VAS) over the last week. Synovitis was scored at 3 locations (infrapatellar fat pad, suprapatellar and intercondylar regions) using a semiquantitative scale (0–3) at all 3 time points on MRI. Scores at each site were added to give a summary synovitis score (0–9).

**Results:** We assessed 270 subjects whose mean (SD) age was 66.7 (9.2) years, BMI 31.5 (5.7) kg/m<sup>2</sup>; 42% were female. There was no correlation of baseline synovitis with baseline pain score ( $r=0.09$ ,  $p=0.17$ ). The change in summary synovitis score was correlated with the change in pain ( $r=0.21$ ,  $p=0.0003$ ). An increase of one unit in summary synovitis score resulted in a 3.15-mm increase in VAS pain score (0–100 scale). Effusion change was not associated with pain change. Of the 3 locations for synovitis, changes in the infrapatellar fat pad were most strongly related to pain change. Despite cartilage loss occurring in over 50% of knees, synovitis was not associated with cartilage loss in either tibiofemoral or patellofemoral compartment.

**Conclusions:** Change in synovitis was correlated with change in knee pain, but not loss of cartilage. Treatment of pain in knee osteoarthritis (OA) needs to consider treatment of synovitis.

See end of article for authors' affiliations

Correspondence to:  
D T Felson, Boston University School of Medicine, A203, 80 E. Concord Street, Boston, MA 02118, USA; [dfelson@bu.edu](mailto:dfelson@bu.edu)

Accepted 19 April 2007  
Published Online First 9 May 2007

The cause of pain in knee osteoarthritis remains elusive as the primary site of pathology in OA, cartilage, has no pain fibres.<sup>1</sup> Many other structures around the knee such as the periosteum, subchondral bone, the fat pad, capsule and, inconsistently, the synovium have been shown to contain nociceptive fibres.<sup>1</sup> In addition, inflammation itself appears to play a role in increasing input from peripheral nociceptors.<sup>2</sup> Biopsies of patients with both early and late knee OA have shown low-grade chronic synovitis with production of pro-inflammatory cytokines.<sup>3,4</sup>

Magnetic resonance imaging (MRI) allows evaluation of multiple structures within the knee, including synovium, cartilage, menisci, bone marrow lesions and effusion. In cross-sectional studies of MRI in knee osteoarthritis (OA), bone-marrow lesions, periarticular lesions, knee effusions and synovitis have been shown to be more often present in persons with knee pain than in persons with a comparable amount of radiographic knee osteoarthritis but without pain.<sup>5–8</sup> Fernandez-Madrid *et al* demonstrated that synovial thickening seen on non-contrast enhanced MRI in the infrapatellar region of knees with OA showed low-grade synovial inflammation on biopsy. This feature was present in 73% of knees with early OA.<sup>9</sup>

We have previously shown that this synovial thickening is present in persons with knee pain and OA much more often than in persons with OA but without pain.<sup>7</sup> Among those with pain, the presence of MRI synovial thickening identified those with more severe pain. While this evidence suggests that synovial thickening may affect pain, these data are cross-sectional, making it impossible to evaluate the temporal relation of pain with synovial thickening. More persuasive evidence would emanate from a longitudinal study in which fluctuations in synovial thickening could be tied to fluctuations in the severity of knee pain. Herein, we provide such evidence.

Our aims were to study the association between baseline and longitudinal changes in MRI-detected synovitis and changes in knee pain, and also to study the association between baseline and longitudinal changes in synovitis and cartilage loss in patients with symptomatic knee OA.

## PATIENTS AND METHODS

## Study population

Patients were recruited to participate in a natural history study of symptomatic knee OA, the Boston Osteoarthritis of the Knee Study. The recruitment for this study has been described in detail elsewhere.<sup>10</sup> Briefly, patients were recruited from two prospective studies, one in men and one in women, of quality of life among veterans; from clinics at Boston Medical Center in Boston, Massachusetts; and from advertisements in local newspapers. Potential participants were asked two questions: "Do you have pain, aching, or stiffness in one or both knees on most days?" and "Has a doctor ever told you that you have knee arthritis?" For patients who answered yes to both questions, we conducted a follow-up interview in which we asked about other types of arthritis that could cause knee symptoms. If no other forms of arthritis were identified, then the individual was eligible for recruitment. A series of knee radiographs (PA, lateral and skyline) were obtained for each patient to determine whether radiographic OA was present. If patients had a definite osteophyte on any view in the symptomatic knee, they were eligible for the study. Because they had frequent knee symptoms and radiographic OA, all patients met American College of Rheumatology criteria for symptomatic knee OA.<sup>11</sup>

**Abbreviations:** MRI, magnetic resonance imaging; OA, osteoarthritis; VAS, visual analogue scale; WOMBS, whole-organ magnetic resonance imaging score

The study included a baseline and follow-up examinations at 15 and 30 months. At baseline, patients who did not have contraindications to MRIs had an MRI of the more symptomatic knee. MRIs of the same knee were also performed at follow-up visits. At each visit, pain in the imaged knee over the past week was assessed using a visual analogue scale (VAS, 0–100), and subjects were weighed with shoes off on a balance-beam scale. The Institutional Review Boards of Boston University Medical Center and the Veterans Administration Boston Health Care System approved the study.

### MRI measurements

All MRIs were performed with a Signa 1.5T system (General Electric, Milwaukee, WI) using a phased-array knee coil. A positioning device was used to ensure uniformity among patients. Coronal, sagittal and axial images were obtained. Fat-suppressed spin-echo proton density and T2-weighted images (repetition time, 2200 ms; echo time, 20/80 ms) with a slice thickness of 3 mm, a 1-mm interslice gap, one excitation, a field of view of 11–12 cm, and a matrix of 256×128 pixels were obtained.

Cartilage morphology was assessed by a musculoskeletal radiologist (AG) using a semiquantitative, multi-feature scoring method (whole-organ magnetic resonance imaging score, WOMBS) for whole-organ evaluation of the knee that is applicable to conventional MRI techniques.<sup>12</sup> Intraclass correlation coefficients of agreement among the readers for cartilage readings ranged from 0.72 to 0.97.

Tibiofemoral cartilage on MRI was scored on all 5 plates (central and posterior femur; anterior, central and posterior tibia) in each of the medial and lateral tibiofemoral joints. The anterior femur was not included in this analysis, as this is part of the patellofemoral joint. Patellofemoral cartilage was scored on 4 plates (medial and lateral patella, and medial and lateral anterior femur). These were read using the fat-suppressed T2-weighted FSE images on a 7-point scale: 0 = normal thickness and signal; 1 = normal thickness but increased signal on T2-weighted images; 2 = partial-thickness focal defect <1 cm in greatest width; 3 = multiple areas of partial-thickness (Grade 2) defects intermixed with areas of normal thickness, or a Grade 2 defect wider than 1 cm but <75% of the region; 4 = diffuse ( $\geq 75\%$  of the region) partial-thickness loss; 5 = multiple areas of full-thickness loss wider than 1 cm but <75% of the region; 6 = diffuse ( $\geq 75\%$  of the region) full-thickness loss.

In WOMBS, grade 1 does not represent a morphological abnormality but rather represents a change in signal in cartilage of otherwise normal morphology. Grades 2 and 3 represent similar types of abnormality of the cartilage, focal defects without overall thinning. Scores of 1 and 2 were exceedingly unusual. Therefore, to create a consistent and logical scale for evaluation of cartilage morphological change, we collapsed the WOMBS cartilage score to a 0–4 scale, where the original WOMBS scores of 0 and 1 were collapsed to 0, the original scores of 2 and 3 were collapsed to 1, and the original scores of 4, 5 and 6 were considered 2, 3 and 4, respectively, in the new scale.<sup>13</sup> The intraobserver agreement for reading of cartilage morphology ranged from 0.65 to 0.78 (kappa). We defined a lesion as occurring in either the medial or lateral tibiofemoral compartment if it was present in the femur or tibia of that compartment. While we conducted analyses using this collapsed WOMBS cartilage scale, analyses using the original scale yielded the same results.

On the baseline and follow-up MRIs, effusion was scored 0–3 based on volume. Bone-marrow lesions were scored only on the baseline MRIs using the WOMBS scale also in which lesions are

scored according to their size (0–3) within quadrants of the femur and tibia.

### Synovial reading

Synovial thickening on MRI using sagittal T2-weighted and proton-density sequences was scored separately at 3 locations (infrapatellar fat pad, suprapatellar and intercondylar regions) using a semiquantitative scale (0–3) at all 3 time points (fig 1). Given the confirmation that these MRI findings connote inflammation in the synovium, we shall label these findings as synovitis.

One reader (CLH) read synovitis on MRI's. For each subject, MRIs were blinded to subject's identity and read paired and in sequential order. The intraobserver agreement (kappa) for



**Figure 1** (A) T2-weighted MR image, sagittal view, with soft tissue density and surrounding synovitis in intercondylar region. (B) T2-weighted MR image, sagittal view, with synovitis (grade 2) in infrapatellar region.

infrapatellar fat pad synovitis score was 0.63, intercondylar 0.49 and suprapatellar 0.20.

### Validation of non-gadolinium synovitis scoring

To validate non-gadolinium scoring, 20 subjects with knee OA at University of Leeds underwent MRI using a sagittal T2 weighted fat suppressed sequence and a gadolinium enhanced T1 weighted fat-suppressed sequence. A trained reader scored the infrapatellar fat pad for synovial thickening on a 0–3 scale using just the T2 sequence without reference to other sequences (as above). The same reader then scored the infrapatellar synovitis using the same scoring system using just the postgadolinium sagittal sequence again without reference to other sequences. The films were blinded and presented to the reader in random order for the 2 reads, which took place 1 week apart. Of the 20 knees, 13 showed contrast (gadolinium) enhanced and non-contrast enhanced scores that were identical (ranging from 0 to 3). As expected, in 6 knees, the non-gadolinium images underestimated the amount of synovial thickening seen on the contrast enhanced image. Only one knee showed an over-reading of synovial thickening on the non-contrast image (score of 2 vs 1).

### Analysis

Synovitis scores at each location were added to give a summary synovitis score at each time point (0–9). Changes in synovitis score were calculated at each time point. In addition, an analysis was carried out for synovitis scores at each individual site.

To examine whether differences in VAS pain score can be explained by differences in synovitis both cross-sectionally and longitudinally, we applied the generalised estimating equation to test this hypothesis with the following statistical model:

$$Y_{it} = \beta_0 + \beta_1 X_{i0} + \beta_2 (X_{it} - X_{i0})$$

where  $Y_{it}$  is the pain score assessed at baseline, at 15 months and at 30 months.  $X_{i0}$  is the synovitis assessed at baseline, and  $X_{it}$  is the corresponding measure of synovitis assessed at time  $t$ , that is, baseline, 15 months and 30 months, respectively. The coefficient  $\beta_1$  measures the cross-sectional association of the synovitis at baseline and VAS pain score, and  $\beta_2$  measures the effect of changes in synovitis on changes of VAS pain score.

The interpretation of the estimate from the model is the expected change in pain over time (from baseline to follow-up

including both 15 and 30 months) per unit change in synovitis score in the corresponding follow-up period for a given subject.

To examine whether cartilage loss can be explained by baseline synovitis, we used cartilage loss in each compartment (medial tibiofemoral, lateral tibiofemoral, and patellofemoral) in 30-month follow-up for analyses unless unavailable, in which case cartilage loss in 15 months was used. Cartilage loss took whole number values from 0 (no loss) to 4 (maximum loss) and was analysed as ordered categories using the proportional odds logistic regression model. A generalised estimating equations correction was applied to the regression model to account for the association in the cartilage loss outcome between regions within a joint. For change in pain and cartilage loss, analyses were adjusted for baseline cartilage scores, age, sex, BMI, effusion score (0–3) and bone-marrow lesion score (using WORMS), and change in both bone-marrow lesion score and effusion score. For cartilage loss, we adjusted additionally for baseline WORMS based meniscal score.

A similar method was used to examine whether cartilage loss can be explained by synovitis change. The change in synovitis in a 30-month follow-up was used for analyses unless unavailable, in which case the change of synovitis in 15 month follow-up was used.

### RESULTS

We assessed 270 subjects (158 male, 112 female) with at least one follow-up MRI. Two hundred and thirty-three subjects were followed for 30 months, and 37 subjects were followed for 15 months. Demographic details are recorded in table 1. The mean age of the subjects was 66.7 years, BMI 31.5 kg/m<sup>2</sup> and VAS pain score 44.2 mm. Most knees had evidence of synovitis at one of the 3 sites at baseline, and about 40% had a change in knee synovitis over time (table 1). The presence of synovitis at baseline was significantly correlated with the baseline Kellgren–Lawrence radiological grade. The Spearman correlation was 0.44, 0.32, 0.27 and 0.39 for the Kellgren–Lawrence grade and summary synovitis, synovitis in infrapatellar fat pad, intercondylar and suprapatellar regions, respectively.

There was no correlation of baseline synovitis with baseline pain score (Pearson correlation coefficient  $r = 0.09$ ,  $p = 0.17$ ); an increase in synovitis score at baseline was correlated with a slightly higher baseline VAS pain score (adjusted estimate per increase in one unit of synovitis 0.72, 95% CI; –1.15, 2.59). However, there was a correlation of change of pain with change of synovitis score ( $r = 0.21$ ,  $p = 0.0003$ ; adjusted estimate of 3.15 VAS score change (on 0–100 scale) per unit change in

**Table 1** Characteristics of subjects

Total	270 subjects
Gender	158 (58.5%) males, 112 females
Age (mean)	66.7 (9.2) years
BMI (mean)	31.5 (5.7) kg/m <sup>2</sup>
Kellgren–Lawrence grade* (median, range)	3 (0 to 4)
Baseline pain† (mean, SD) (0–100)	44.2 (25.2) mm
Change in pain (mean, SD) from baseline to follow-up	–1.9 (25.5) mm
Baseline synovitis score 0–9	
Baseline infrapatellar fat pad score, 0/1/2/3 (%)	21.5/40.0/33.9/4.6
Baseline intercondylar notch score, 0/1/2/3 (%)	31.5/52.3/15.0/1.2
Baseline suprapatellar notch score, 0/1/2/3 (%)	24.5/41.2/22.2/12.1
Change in synovitis score	
Change in infrapatellar fat pad score, decrease/no change/increase (%)	15.2/63.1/21.7
Change in intercondylar notch score, decrease/no change/increase (%)	14.3/66.8/18.9
Change in suprapatellar notch score, decrease/no change/increase (%)	20.4/59.2/20.4
Baseline cartilage score in 14 plates (mean, SD)	16.1 (9.2)
Baseline summary bone-marrow lesion (mean, SD)	3.2 (3.1)
Baseline effusion score (mean, SD)	0.9 (0.8)

\*Subjects with ‘‘Knee pain/OA’’, who had Kellgren–Lawrence grade 0 due to normal PA views, were defined as having radiographic OA due to definite osteophytes in patello-femoral joint; †pain was measured on a 100-mm visual analogue scale.

**Table 2** Change in pain as a function of change in synovitis at individual sites\*

	Baseline synovitis and baseline VAS pain			Change of synovitis and change of VAS pain		
	Estimate	95% CI	p value	Estimate	95% CI	p value
Infrapatellar fat pad	2.31	-1.54, 6.16	0.24	4.89	0.42, 9.36	0.03
Intercondylar	2.26	-2.92, 7.45	0.39	5.74	0.34, 11.14	0.04
Suprapatellar	-0.07	-3.97, 3.82	0.97	3.35	-0.34, 7.05	0.08

\*Analyses adjusted for age, sex, BMI, cartilage score at baseline, effusion score (0-3), bone-marrow lesion score (WORMS), change in bone-marrow lesion score and change in effusion score.

**Table 3** Risk of cartilage loss by synovitis (adjusted odds ratio, 95% CI)\*

	Odds ratio for loss of cartilage in medial tibiofemoral compartment	Odds ratio for loss of cartilage in lateral tibiofemoral compartment	Odds ratio for loss of cartilage in patellofemoral compartment
Baseline summary synovitis score			
Summary synovitis score (0-2)	Referent	Referent	Referent
Summary synovitis score (3-4)	2.9 (1.3, 6.5)	0.9 (0.4, 2.2)	1.3 (0.5, 3.3)
Summary synovitis score (5-9)	1.5 (0.6, 4.1)	1.0 (0.3, 2.9)	0.9 (0.2, 3.9)
Change in synovitis score			
Improvement in synovitis score	Referent	Referent	Referent
Stable synovitis score	0.7 (0.3, 1.6)	0.6 (0.2, 1.7)	1.9 (0.6, 5.7)
Worsening synovitis score	0.6 (0.2, 1.3)	1.1 (0.5, 2.9)	2.5 (0.8, 7.5)

\*Analyses adjusted for age, sex, BMI, cartilage score at baseline, effusion score (0-3), bone-marrow lesion score (WORMS), change in bone-marrow lesion score and change in effusion score.

synovitis score 95% CI; 1.04, 5.26.  $p = 0.003$ ). However, the change in effusion score over time was not associated with pain change (adjusted estimate 1.19, 95% CI; -8.08, 10.46.  $p = 0.80$ ). Similar results were seen in both males and females.

Of the 3 locations for synovitis examined separately, none of the baseline synovitis scores was associated with baseline pain (table 2). However, for each site, the change in synovial score was associated with pain change. This relation was strongest for the change in synovitis in infrapatellar and intercondylar fat pads in which there was a 5.7-mm and 4.9-mm increase in pain, respectively, per unit increase in synovitis. The relation between change in synovitis at the suprapatellar region and change in pain was not significant.

Despite cartilage loss occurring in over 50% of knees from baseline to follow-up, baseline synovitis score was not associated with cartilage loss in the patellofemoral compartment and only in those with higher grades of synovitis in the tibiofemoral compartment (table 3). The change in synovitis score was not associated with cartilage loss in medial or lateral tibiofemoral compartments, but there was a trend towards increased cartilage loss in the patellofemoral compartment in those with worsening synovitis scores (table 3).

## DISCUSSION

Our study demonstrates that the change in synovitis over time was correlated modestly with changes in knee pain, but not at all with loss of cartilage. The correlation was direct—an increase in synovitis was associated with worsening pain and a decrease with less severe pain. This effect on pain was independent of changes in knee-joint effusion. We have demonstrated for the first time that longitudinal changes in synovitis reflect changes in pain in knee OA. We failed to find any association of synovial inflammation with cartilage loss.

Our findings regarding synovitis and cartilage loss are in keeping with previous cross-sectional findings of no association between arthroscopic synovitis or degree of synovial thickening on MRI and cartilage loss on MRI in a cross-sectional study.<sup>14</sup> We found little association between the change in synovitis score and cartilage loss, suggesting that the two processes, synovitis and cartilage loss, are, for the most part, independent.

Synovial thickening in the infrapatellar region, detected on non-contrast MRI imaging, like that used in this paper, has been shown to represent low-grade synovial inflammation when biopsied in knee OA.<sup>9</sup> The synovial scoring we used was based on a previous study that demonstrated a good correlation between the MRI semiquantitative scores and MRI synovial volume measurements, suggesting that semiquantitative synovitis scores are valid and enable feasible evaluation of the synovium in OA cohorts.<sup>15</sup> Gadolinium-enhanced MRI remains the gold standard for the assessment of synovitis on MRI. Although, we did not use gadolinium in our study, we did validate non-gadolinium scoring and found that non-contrast images can provide a valid measure of synovitis, as demonstrated by gold-standard gadolinium-enhanced MR images in knee OA.

Intra-articular sources of pain in knee OA such as the synovium have been suggested by other studies which have demonstrated the analgesic properties of intra-articular instillation of local anaesthetic,<sup>16</sup> low-dose morphine<sup>17</sup> and steroid.<sup>18-20</sup> Another recent study suggested that intra-articular anakinra (interleukin-1 receptor antagonist) given to 7 patients with knee OA resulted in an improvement in pain, which paralleled improvements in MRI synovial scores.<sup>21</sup> Also, NSAIDs appear to have a superior effect over acetaminophen for treatment of knee OA in several studies, and this could be related to their anti-inflammatory properties.<sup>22-23</sup> These studies, along with our observation that changes in pain are associated with changes in synovitis, suggest that treatment of synovitis may be a key element in the management of pain in knee OA. A limitation of our study was that we did not collect sufficiently detailed information on the use of analgesics and NSAIDs to allow this to be utilised in the analysis.

The relation we found of synovitis change with pain change was significant but only modest. This weak correlation ( $r = .21$ ) may be because there are other causes of change in pain intensity in persons with knee osteoarthritis but may also be due to the crude scale for synovitis change we used and our inability to score synovitis in all sites in which it occurs within the knee. Further, we had difficulty identifying suprapatellar synovitis on non-contrast images, and that accounts for the

poor reliability of readings at this site. It may also explain why we found stronger relations of pain change with synovitis at other knee sites. Even so, our findings suggest that synovitis is likely to contribute to the fluctuations in pain experienced by persons with symptomatic knee OA and needs confirmation.

In conclusion, in this longitudinal MRI study, we found that a change in synovitis score was associated with changes in pain score. We did not demonstrate any association between synovitis or its change and cartilage loss. Our findings have clear-cut implications for the treatment of painful osteoarthritis of the knee, as they suggest that treatments targeted to synovitis may improve pain.

## ACKNOWLEDGEMENTS

We are indebted to the study participants for generously giving their time.

## Authors' affiliations

**Catherine L Hill**, Rheumatology Unit, The Queen Elizabeth Hospital, Woodville, Australia

**David J Hunter, Jingbo Niu, Margaret Clancy, David T Felson**, Boston University Clinical Epidemiology Research and Training Unit, Boston, MA, USA

**Ali Guermazi, Harry Genant**, Synarc, Inc, San Francisco, CA, USA and University of California San Francisco, San Francisco, CA, USA

**Daniel Gale, David T Felson**, VA Boston Healthcare System, Boston, MA, USA

**Andrew Grainger, Philip Conaghan**, Departments of Radiology and Rheumatology, Leeds Teaching Hospitals Trust & Leeds Primary Care Trust, Leeds, UK

Supported by NIH AR47785.

Competing interests: None declared.

## REFERENCES

- Dye SF**, Vaupel GL, Dye CC. Conscious neurosensory mapping of the internal structures of the human knee without intraarticular anesthesia. *Am J Sports Med* 1998;**26**:1-4.
- Schaible HG**, Grubb BD. Afferent and spinal mechanisms of joint pain. *Pain* 1993;**55**:5-54.
- Smith MD**, Wetherall M, Darby T, Esterman A, Slavotinek J, Roberts-Thomson P, *et al*. A randomized placebo-controlled trial of arthroscopic lavage versus lavage plus intra-articular corticosteroids in the management of symptomatic osteoarthritis of the knee. *Rheumatology (Oxford)* 2003;**42**:1477-85.
- Benito MJ**, Veale DJ, FitzGerald O, van den Berg WB, Bresnihan B. Synovial tissue inflammation in early and late osteoarthritis. *Ann Rheum Dis* 2005;**64**:1263-7.
- Felson DT**, Chaisson CE, Hill CL, Totterman SM, Gale ME, Skinner KM, *et al*. The association of bone marrow lesions with pain in knee osteoarthritis. *Ann Intern Med* 2001;**134**:541-9.
- Sowers MF**, Hayes C, Jamadar D, Capul D, Lachance L, Jannausch M, Welch G. Magnetic resonance-detected subchondral bone marrow and cartilage defect characteristics associated with pain and x-ray-defined knee osteoarthritis. *Osteoarthritis Cartilage* 2003;**11**:387-93.
- Hill CL**, Gale DG, Chaisson CE, Kazis L, Totterman S, Gale ME, Felson DT. Knee effusions, popliteal cysts and synovial thickening: Association with knee pain in those with and without osteoarthritis. *J Rheumatol* 2001;**28**:1330-7.
- Hill CL**, Gale D, Chaisson CE, Skinner K, Kazis L, Gale ME, Felson DT. Periarticular lesions detected on magnetic resonance imaging: prevalence in knees with and without symptoms. *Arthritis Rheum* 2003;**48**:2836-44.
- Fernandez-Madrid F**, Karvonen RL, Teitge RA, Miller PR, An T, Negendank WG. Synovial thickening detected by MR imaging in osteoarthritis of the knee confirmed by biopsy as synovitis. *Mag Res Imaging* 1995;**13**:177-83.
- Felson DT**, McLaughlin S, Goggins J, LaValley MP, Gale ME, Totterman S, *et al*. Bone marrow edema and its relation to progression of knee osteoarthritis. *Ann Intern Med* 2003;**139**:330-6.
- Altman R**, Asch, Bloch D, Bole G, Borenstein D, Brandt K, *et al*. The American College of Rheumatology criteria for the classification and reporting of osteoarthritis of the knee. *Arthritis Rheum* 1986;**29**:1039-49.
- Peterfy CG**, Guermazi A, Zaim S, Tirman PF, Miaux Y, White D, *et al*. Whole-Organ Magnetic Resonance Imaging Score (WORMS) of the knee in osteoarthritis. *Osteoarthritis Cartilage* 2004;**12**:177-90.
- Amin S**, LaValley MP, Guermazi A, Grigoryan M, Hunter DJ, Clancy M, *et al*. The relationship between cartilage loss on magnetic resonance imaging and radiographic progression in men and women with knee osteoarthritis. *Arthritis Rheum* 2005;**52**:3152-9.
- Loeulle D**, Chary-Valckenaere I, Champigneulle J, Rat AC, Toussaint F, Pinzano-Watrin A, *et al*. Macroscopic and microscopic features of synovial membrane inflammation in the osteoarthritic knee. *Arthritis Rheum* 2005;**52**:3492-501.
- Rhodes LA**, Grainger AJ, Keenan A, Thomas C, Emery P, Conaghan PG. The validation of simple scoring methods for evaluating compartment-specific synovitis detected by MRI in knee osteoarthritis. *Rheumatology (Oxford)* 2005;**44**:1569-73.
- Creamer P**, Hunt M, Dieppe P. Pain mechanisms in osteoarthritis of the knee: effect of intraarticular anesthetic. *J Rheumatol* 1996;**23**:1031-6.
- Likar R**, Schafer M, Paulak F, Sittl R, Pipam W, Schalk H, *et al*. Intraarticular morphine analgesia in chronic pain patients with osteoarthritis. *Anesth Analg* 1997 Jun;**84**:1313-7.
- Ostergaard M**, Stoltenberg M, Gideon P, Sorensen K, Hendriksen O, Lorenzen I. Changes in synovial membrane and joint effusion volumes after intraarticular methylprednisolone. Quantitative assessment of inflammatory and destructive changes in arthritis by MRI. *J Rheumatol* 1996;**23**:1151-61.
- Dieppe PA**, Sathapatayavongs B, Jones HE, Bacon PA, Ring EF. Intra-articular steroids in osteoarthritis. *Rheumatol Rehabil* 1980;**19**:212-7.
- Raynauld JP**, Buckland-Wright C, Ward R, Choquette D, Haraoui B, Martel-Pelletier J, *et al*. Safety and efficacy of long-term intraarticular steroid injections in osteoarthritis of the knee: a randomized, double-blind, placebo-controlled trial. *Arthritis Rheum* 2003;**48**:370-7.
- Loeulle D**, Chary-Valckenaere I, Goebel C, Rat AC, Blum A, Kiefer P, *et al*. MRI evaluation of the synovial membrane after a single intraarticular injection of anakinra in 7 patients with osteoarthritis of the knee. *Arthritis Rheum* 2005;**9**:S70.
- Pincus T**, Koch GG, Sokka T, Lefkowitz J, Wolfe F, Jordan JM, *et al*. A randomized, double-blind, crossover clinical trial of diclofenac plus misoprostol versus acetaminophen in patients with osteoarthritis of the hip or knee. *Arthritis Rheum* 2001;**44**:1477-80.
- Altman RD**, IAP Study Group. Ibuprofen, acetaminophen and placebo in osteoarthritis of the knee: a six-day double-blind study [abstract]. *Arthritis Rheum* 1999;**42**(Suppl 9):S403.