

2017-12-01

Synthesis and in vitro antimycobacterial activity of novel n-arylpiperazines containing an ethane-1,2-diyl connecting chain

Tomás Gonec
Veterinární a farmaceutická univerzita Brno

Ivan Malík
Comenius University

Jozef Csöllei
Veterinární a farmaceutická univerzita Brno

Josef Jampílek
Comenius University

Jirina Stolaríková
Zdravotního ústavu se sídlem v Ostrave

See next page for additional authors

Follow this and additional works at: <https://sword.cit.ie/dptbiosciart>

 Part of the [Biology Commons](#)

Recommended Citation

Goněc, T.; Malík, I.; Csöllei, J.; Jampílek, J.; Stolaríková, J.; Solovič, I.; Mikuš, P.; Keltošová, S.; Kollár, P.; O'Mahony, J.; Coffey, A. Synthesis and In Vitro Antimycobacterial Activity of Novel N-Arylpiperazines Containing an Ethane-1,2-diyl Connecting Chain. *Molecules* 2017, 22, 2100.



This Article is brought to you for free and open access by the Biological Sciences at SWORD - South West Open Research Deposit. It has been accepted for inclusion in Department of Biological Sciences Publications by an authorized administrator of SWORD - South West Open Research Deposit. For more information, please contact sword@cit.ie.

Authors

Tomás Gonec, Ivan Malík, Jozef Csöllei, Josef Jampílek, Jirina Stolaríková, Ivan Solovic, Peter Miku, Stanislava Keltoová, Peter Kollár, Jim O'Mahony, and Aidan Coffey

Article

Synthesis and *In Vitro* Antimycobacterial Activity of Novel *N*-Arylpiperazines Containing an Ethane-1,2-diyl Connecting Chain

Tomáš Goněč ¹, Ivan Malík ^{2,*}, Jozef Csöllei ¹, Josef Jampílek ² , Jiřina Stolaříková ³, Ivan Solovič ^{4,5}, Peter Mikuš ⁶, Stanislava Keltošová ⁷, Peter Kollár ⁷, Jim O'Mahony ⁸ and Aidan Coffey ⁸ 

¹ Department of Chemical Drugs, Faculty of Pharmacy, University of Veterinary and Pharmaceutical Sciences in Brno, Palackého 1946/1, Brno CZ-612 42, Czech Republic; gonect@vfu.cz (T.G.); csolleij@vfu.cz (J.Cs.)

² Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Comenius University in Bratislava, Odbojárov 10, Bratislava SK-832 32, Slovak Republic; malikivan001@gmail.com (I.M.); josef.jampilek@gmail.com (J.J.)

³ Laboratory for Mycobacterial Diagnostics and Tuberculosis, Regional Institute of Public Health, Partyzánské náměstí 7, Ostrava CZ-702 00, Czech Republic; Jirina.Stolarikova@zu.cz (J.S.)

⁴ Clinic for Tuberculosis and Lung Diseases, National Institute for Tuberculosis, Lung Diseases and Thoracic Surgery, Vyšné Hágy, Vysoké Tatry SK-059 84, Slovak Republic; solovic@hagy.sk (I.S.)

⁵ Department of Public Health, Faculty of Health, Catholic University in Ružomberok, Hrabovská cesta 1A, Ružomberok SK-034 01, Slovak Republic; ivan.solovic@ku.sk (I.S.)

⁶ Department of Pharmaceutical Analysis and Nuclear Pharmacy, Faculty of Pharmacy, Comenius University in Bratislava, Odbojárov 10, Bratislava SK-832 32, Slovak Republic; mikus@fpharm.uniba.sk (P.M.)

⁷ Department of Human Pharmacology and Toxicology, University of Veterinary and Pharmaceutical Sciences in Brno, Palackého 1946/1, Brno CZ-612 42, Czech Republic; kollarp@vfu.cz (P.K.); keltošovas@vfu.cz (S.K.)

⁸ Department of Biological Sciences, Cork Institute of Technology, Bishopstown, Cork T12 P928, Ireland; jim.omahony@cit.ie (J.O.M.); aidan.coffey@cit.ie (A.C.)

* Correspondence: malikivan001@gmail.com; Tel.: +421-2-501-117-227

Received: 30 October 2017; Accepted: 27 November 2017; Published: 30 November 2017

Abstract: Novel 1-(2-{3-/4-[(alkoxycarbonyl)amino]phenyl}-2-hydroxyethyl)-4-(2-fluorophenyl)-piperazin-1-ium chlorides (alkoxy = methoxy to butoxy; **8a–h**) have been designed and synthesized through multistep reactions as a part of on-going research programme focused on finding new antimycobacterials. Lipophilic properties of these compounds were estimated by RP-HPLC using methanol/water mobile phases with a various volume fraction of the organic modifier. The log k_w values, which were extrapolated from intercepts of a linear relationship between the logarithm of a retention factor k ($\log k$) and volume fraction of a mobile phase modifier (φ_M), varied from 2.113 (compound **8e**) to 2.930 (**8h**) and indicated relatively high lipophilicity of these salts. Electronic properties of the molecules **8a–h** were investigated by evaluation of their UV/Vis spectra. In a next phase of the research, the compounds **8a–h** were *in vitro* screened against *M. tuberculosis* CNCTC My 331/88 (identical with H₃₇R_v and ATCC 2794), *M. kansasii* CNCTC My 235/80 (identical with ATCC 12478), a *M. kansasii* 6 509/96 clinical isolate, *M. avium* CNCTC My 330/80 (identical with ATCC 25291) and *M. avium intracellulare* ATCC 13950, respectively, as well as against *M. kansasii* CIT11/06, *M. avium* subsp. *paratuberculosis* CIT03 and *M. avium hominissuis* CIT10/08 clinical isolates using isoniazid, ethambutol, ofloxacin, ciprofloxacin or pyrazinamide as reference drugs. The tested compounds **8a–h** were found to be the most promising against *M. tuberculosis*; a MIC = 8 μ M was observed for the most effective 1-(2-{4-[(butoxycarbonyl)amino]phenyl}-2-hydroxyethyl)-4-(2-fluorophenyl)piperazin-1-ium chloride (**8h**). In addition, all of them showed low (insignificant) *in vitro* toxicity against a human monocytic

leukemia THP-1 cell line, as observed LD_{50} values $> 30 \mu\text{M}$ indicated. The structure–antimycobacterial activity relationships of the analyzed **8a–h** series are also discussed.

Keywords: *N*-arylpiperazines; arylaminoethanols; lipophilicity; electronic properties; *Mycobacterium tuberculosis* H₃₇R_v

1. Introduction

An *N*-arylpiperazine privileged scaffold [1] has been found in the chemical structure of many effective antimycobacterial agents [2–8]. Some of these compounds have been characterized by a very typical structural arrangement [5–8], i.e., the *N*-aryl- or variously substituted *N*-phenylpiperazine moiety, connecting hydrocarbon chain and terminal heterocyclic fragment. Efforts to combine the *N*-arylpiperazine and “ethambutol-like” structural frameworks were supported by comprehensive structure–activity relationships (SAR) studies of homopiperazin-1,4-diyl-containing derivatives (e.g., a molecule **SQ775**; Figure 1a), ethambutol (EMB; Figure 1b) and the diamines structurally based on EMB [9–12]. Among the synthesized compounds, *N*-geranyl-*N'*-(2-adamantan-1-yl)ethane-1,2-diamine (Figure 1c) showed promising efficiency [12]. The outlined systematic research led to *N*-ArPA molecules (Figure 1d), in which structure the distance between two nitrogens, presence of β -aminoalcohol motifs and short connecting chains were crucial for their *in vitro* antimycobacterial activity [13]. The derivatives *N*-ArPA were very effective against *Mycobacterium tuberculosis* CNCTC My 331/88 (identical with *M. tuberculosis* H₃₇R_v) and multi-drug resistant (MDR) *M. tuberculosis* 43 strain, which showed resistance to rifampicin (RIF) and isoniazid (INH).




Figure 1. Chemical structures of: (a) compound **SQ775**; (b) ethambutol (EMB); (c) *N*-geranyl-*N'*-(2-adamantan-1-yl)ethane-1,2-diamine; and (d) chiral *N*-arylpiperazine-based aminoalcohols (*N*-ArPA), which showed a notable *in vitro* efficiency against *M. tuberculosis* CNCTC My 331/88 [9–13].

It was also concluded that removal or significant alteration of basicity of either amino group led to loss of potency [9,13]. In addition, the presence of the $R = 2-/4\text{-F}$ substituent and OH group with the (*R*)-configuration at the carbon of a connecting chain (Figure 1d) resulted in higher *in vitro* antimycobacterial efficiency than in a case of EMB. On the other hand, the remaining *R* substituents (H, Cl; Figure 1d) caused decrease in activity [13].

Regarding the design of original antimycobacterials, a carbamate (NHCOO) functionality is structurally related to hybrid amide-ester features and, in general, displays very good chemical and

proteolytic stabilities [14]. The carbamate functionality imposes a degree of conformational restriction due to the delocalization of non-bonded electrons on nitrogen into the carboxyl moiety. In addition, this functionality participates in hydrogen bonding through the carboxyl group and the backbone NH [15]. Therefore, a substitution on the *O*- and *N*-termini of the carbamate offers opportunities to modulate biological properties and improve stability and pharmacokinetic features [14,15].

The idea to introduce a lipophilic 3-/4-alkoxy or 3-/4-alkoxycarbonylamino moiety (alkoxy = methoxy to butoxy) into a chemical structure of newly designed molecules was based on previous studies focused on the synthesis and *in vitro* biological evaluation of the *N*-arylpiperazines and phenylcarbamic acid derivatives [16–21]. Their antimycobacterial activity increased with elongation of this chain until a maximum in efficiency was reached [19–21]. Further increase in its length led to the decrease in potency. The observed dependence was approximated by a parabolic function and described as a cut-off effect. That phenomenon was comprehensively reviewed and rationalized in number of mechanistic ways by Hansch and Clayton [22] as well as Balgavý and Devínský [23] more than two decades later.

Regarding the conclusions published in papers [19–21], it might be expected that eventual incorporation of the alkoxycarbonylamino group into a 2-position of a phenyl ring would cause the decrease in antimycobacterial activity.

It was also believed that the presence of the linear alkoxy side chain would be very favorable in terms of interactions with target structures located in biomembranes of mycobacterial strains, especially *M. tuberculosis* H₃₇R_v. Branching or substitution of the alkoxy with the alkyl group is reported to have caused decrease in efficiency [19–21,24].

It was found that a suitable modification of the aromatic ring attached to a piperazin-1,4-diyl framework might not result in loss of activity. The derivatives containing a pyrimidin-2-yl fragment were slightly more efficient than the ones with a pyridin-2-yl moiety (a series MM; Figure 2). The molecules MM [18] were able to effectively *in vitro* fight *M. tuberculosis* My 331/88, *M. kansasii* 6 509/96, *M. tuberculosis* 7375/1998, a strain resistant to INH, RIF, rifabutine (RFB) and streptomycin (STM), respectively, as well as *M. tuberculosis* Prague 1, an extremely-resistant strain to INH, RIF, RFB, STM, EMB, ofloxacin (OFLX), gentamicin (GTM) and amikacin (AK), respectively [18].




Figure 2. *N*-Arylpiperazine derivatives containing a 2-hydroxyethane-1,2-diyl connecting chain (a series MM), which were *in vitro* screened against some mycobacterial strains [18].

Inspired and encouraged by given importance of the 3-/4-alkoxycarbonylamino, β -aminoalcohol and 4-(2-fluorophenyl)piperazin-1-yl fragments in a chemical structure of effective antimycobacterials, the present study was focused on the synthesis of original racemic compounds in order to find out if they would be efficient against some of following mycobacterial strains *in vitro*, namely, *M. tuberculosis* CNCTC My 331/88 (identical with H₃₇R_v and ATCC 2794), *M. kansasii* CNCTC My 235/80 (identical with ATCC 12478), a *M. kansasii* 6 509/96 clinical isolate, *M. avium* CNCTC My 330/80 (identical with ATCC 25291) as well as *M. avium intracellulare* ATCC 13950, *M. kansasii* CIT11/06, *M. avium* subsp. *paratuberculosis* CIT03 and *M. avium hominissuis* CIT10/08 clinical isolates, respectively.

Despite a fact that currently designed molecules contained a stereogenic centre (Table 1), the synthesis of racemates has been regarded as a reasonable strategy in conceptual development of original antimycobacterial agents to verify the relevancy of proposed structural frameworks [16,18,25–27].

Table 1. Chemical structure of the compounds **8a–h**, their lipophilicity indices R_M (RP-TLC) and $\log k$ (RP-HPLC) as well as retention times t_r (RP-HPLC) estimated in the mobile phases consisted of a various volume ratio (v/v) of a methanol (MeOH) organic modifier and water.

Comp.	$^1 R_M$	Mobile phase MeOH/water (v/v)							
		60:40		70:30		80:20		85:15	
		t_r (min)	$\log k$	t_r (min)	$\log k$	t_r (min)	$\log k$	t_r (min)	$\log k$
8a	−0.55	6.593	0.612	3.587	0.248	2.493	−0.034	2.200	−0.156
8b	−0.35	7.707	0.695	4.893	0.444	2.907	0.095	2.433	−0.056
8c	−0.16	8.933	0.771	5.907	0.552	3.193	0.166	2.620	0.010
8d	0.01	10.021	0.829	7.820	0.702	3.586	0.245	2.830	0.074
8e	−0.02	5.307	0.491	3.180	0.163	2.360	−0.085	2.120	−0.196
8f	0.19	7.153	0.656	3.953	0.312	2.620	0.010	2.275	−0.121
8g	0.39	8.280	0.732	5.320	0.492	3.033	0.128	2.533	−0.020
8h	0.63	11.035	0.881	7.287	0.665	3.543	0.240	2.814	0.069

$^1 R_M$, Lipophilicity index (RP-TLC). Silica gel plates (stationary phases) were impregnated by 1% silicone oil in heptane.


2. Results and Discussion

2.1. Chemistry


2.1.1. Synthesis and Spectral Characteristics

Designer *N*-arylpiperazines were synthesized *via* multistep reactions, exploring the impact of their lipohydrophilic and electronic properties on the *in vitro* activity against selected mycobacterial strains. In addition, the *in vitro* toxicity profile of the final molecules against a human monocytic leukemia THP-1 cell line was inspected.

Following the outlined objectives, the compounds were synthesized according to Schemes 1 and 2 as follows. Initially, 3-aminoacetophenone (**1a**) and 4-aminoacetophenone (**1b**) were employed as convenient starting compounds. They were treated with alkyl chloroformates **2a–d** (alkyl = methyl to butyl) at presence of pyridine in acetone to afford colourless alkyl (3-/4-acetylphenyl)-carbamates **3a–h** [28] in the yields that varied from 89% to 99% (Scheme 1).



Scheme 1. Synthesis of the alkyl [3-/4-(bromoacetyl)phenyl]carbamates **4a–h** (alkyl = methyl to butyl), Reagents and conditions: (i) ClCOOR' ($R' =$ methyl to butyl; **2a–d**), pyridine; (ii) Br_2 , chloroform.



4a, 6a, 7a, 8a: R=3-NHCOOCH₃, 4b, 6b, 7b, 8b: R=3-NHCOOC₂H₅, 4c, 6c, 7c, 8c: R=3-NHCOOC₃H₇,
 4d, 6d, 7d, 8d: R=3-NHCOOC₄H₉, 4e, 6e, 7e, 8e: R=4-NHCOOCH₃, 4f, 6f, 7f, 8f: R=4-NHCOOC₂H₅,
 4g, 6g, 7g, 8g: R=4-NHCOOC₃H₇, 4h, 6h, 7h, 8h: R=4-NHCOOC₄H₉

Scheme 2. Synthesis of the final 1-(2-{3-/4-[(alkoxycarbonyl)amino]phenyl}-2-hydroxyethyl)-4-(2-fluorophenyl)piperazin-1-ium chlorides **8a–h** (alkoxy = methoxy to butoxy), *Reagents and conditions:* (i) TEA, THF; (ii) NaBH₄, methanol; (iii) a saturated solution of hydrogen chloride in diethyl ether.

When designing the chemical structure of target molecules, 2-aminoacetophenone was not considered a suitable starting structure due to a possible undesired cyclization of resulting intermediates [29,30]. In addition, a weak *in vitro* antimycobacterial activity of final products would be probably observed [19–21].

The molecules **3a–h** underwent α -bromination of an acetyl group because of the dropwise addition bromine in chloroform. This reaction procedure took place at a sufficiently high rate in chloroform at room temperature with constant stirring [31]. Resulting alkyl {3-/4-(bromoacetyl)phenyl}carbamates **4a–h** (alkyl = methyl to butyl; Scheme 1) were achieved with 75% to 93% yields.

A substitution of bromine by 1-(2-fluorophenyl)piperazine **5** [32] in the presence of triethylamine (TEA) in anhydrous tetrahydrofuran (THF) led to colourless alkyl {3-/4-[(4-(2-fluorophenyl)piperazin-1-yl)acetyl]phenyl}carbamates **6a–h** (Scheme 2). The intermediates **6a–h** were prepared in moderate to good yields that ranged from 75% to 96%. Next, they were transformed into alkyl {3-/4-[2-(4-(2-fluorophenyl)piperazin-1-yl)-1-hydroxyethyl]phenyl}carbamates **7a–h** by nucleophilic addition of hydride anions (Scheme 2) using simple and convenient reduction with sodium borohydride [32]. The molecules **7a–h** were synthesized in 80% to 94% yields.

Detailed spectral characteristics (¹H-NMR, ¹³C-NMR, HR-MS or ESI-MS) of the thirty-two prepared intermediates **3a–h**, **4–h**, **6a–h** and **7a–h**, are given in the Materials and Methods section of a current paper.

Addition of a saturated solution of hydrogen chloride in diethyl ether into a particular solution of the compounds **7a–h** in chloroform led to the desired 1-(2-{3-/4-[(alkoxycarbonyl)amino]phenyl}-2-hydroxyethyl)-4-(2-fluorophenyl)piperazin-1-ium chlorides **8a–h**. The crude products **8a–h** were purified by recrystallization from acetone providing 84% to 96% yields (Table 1, Scheme 2).

Purity of the final salts **8a–h** was verified by thin-layer chromatography (TLC) using petroleum ether/diethyl amine eluant (10:3, *v/v*) as a mobile phase. Spots were observed under iodine vapours/UV light at a wavelength (λ) of 254 nm. The position and length of an 3-/4-alkoxycarbonylamino fragment influenced values of a retardation factor (R_f), as expected. The 3-positional isomers **8a–d** showed slightly higher R_f values ($R_f = 0.40$ – 0.73) compared to those of

the 4-substituted ones **8e–h** (0.31–0.56). The R_f values have been provided in detail in the Materials and Methods section of a current paper.

The newly synthesized target substances **8a–h** were fully characterized by their IR, $^1\text{H-NMR}$, $^{13}\text{C-NMR}$ and ESI-MS spectral values, which were in full accordance with proposed structures. Analyzing the IR spectra of **8a–h**, bands typical for stretching vibrations ν ($\text{C}=\text{O}$) were observed in the region from 1730 cm^{-1} to 1718 cm^{-1} . Identity of aromatic rings was confirmed by presence of ν ($\text{C}=\text{C}$) at around 1600 cm^{-1} . The recorded IR spectra also afforded vibrations in the range from 1552 cm^{-1} to 1543 cm^{-1} due to δ (N-H). The bands between 1234 cm^{-1} and 1221 cm^{-1} were related to asymmetric stretching of a C-O-C fragment. The in-plane deformation vibrations (δ_{ip}) at around 1020 cm^{-1} and out-of-plane deformation vibrations (δ_{oop}) at around 850 cm^{-1} of a $=\text{C-H}$ group were also observed.

In the $^1\text{H-NMR}$, signals of particular protons were verified on basis of their chemical shift (δ), multiplicities and coupling constants in $\text{DMSO-}d_6$. Regarding the 3-alkoxycarbonylamino substituent-containing molecules **8a–d**, a proton signal of a carbamoyloxy group was detected in the δ interval from 9.60 ppm to 9.57 ppm. A shift of this chain to a 4-position (compounds **8e–h**) led to slightly higher δ values recognized from 9.71 ppm to 9.67 ppm. The δ chemical shift between 154.90 ppm and 153.25 ppm (doublet) was assigned to the carbon atom of a C-F bond in the $^{13}\text{C-NMR}$ spectra of prepared salts **8a–h**. The carbon of a carbamoyloxy group was identified in the δ range from 153.91 ppm to 152.91 ppm.

Elemental analyses of the synthesized derivatives **8a–h** indicated that addition of a saturated solution of hydrogen chloride in diethyl ether caused a protonation of only one nitrogen of a piperazin-1,4-diyl fragment. This was due to a positive mesomeric effect of the nitrogen atom towards an aromatic ring. This conclusion was also evidenced by mass spectral values of these compounds, for which particular $[\text{M} + \text{H}]^+$ molecular peaks were observed. Current elemental analyses results (% C, H, N) were within $\pm 0.40\%$ of theoretical values for the proposed monohydrochlorides.

2.1.2. Lipohydrophilic Properties

Lipophilicity has been the physicochemical parameter continually attracting prime interest in QSAR and SAR studies as a predominant descriptor of pharmacodynamic, pharmacokinetic and toxic aspects of the antimycobacterial drugs [33–36].

A partition coefficient P (or its logarithm) between water or a phosphate buffer and octan-1-ol has been used as a preferential experimental expression of lipophilic properties of a compound. However, the $\log P$ parameter is losing that role as the method of a choice due to some methodological drawbacks and limitations, which were extensively described and explained in [37]. Chromatographic methods have been therefore developed and used successfully to estimate the lipophilicity of organic compounds [37].

Lipophilicity indices R_M and k ($\log k$) of the compounds **8a–h** were estimated by the reversed-phase thin-layer chromatography (RP-TLC) and reversed-phase high-performance liquid chromatography (RP-HPLC). The reason for more detailed chromatographic characterization of the salts **8a–h** was their better solubility in polar media (mobile phases) compared to free bases **7a–h**.

In addition, previous *in vitro* antimycobacterial assays [18] employed structurally very similar compounds **MM** as hydrochlorides (Figure 2). This approach was very beneficial due to improvement in their solubility in tested media compared to free bases.

Based on previous experience, it would be more precise to evaluate the lipohydrophilic properties and antimycobacterial activities of **8a–h** and compare the observed values to those related to the **MM** series.

Calculations of the R_M and $\log k$ parameters were detailed in the Materials and Methods section of a current paper.

In the RP-TLC, silica gel plates impregnated by a hand with a variously concentrated silicone oil in heptane (a strong hydrophobic agent) were used as a non-polar stationary phase [37]. Optimal

differences in R_M values within both homological groups **8a–d** and **8e–h** were observed if 1% silicone oil in heptane was chosen (Tables 1 and S1 in Supplementary Materials).

The calculated R_M values for the **8a–d** series varied from -0.55 to 0.01 , the molecules **8e–h** showed higher R_M s from -0.02 to 0.63 (Table 1). These R_M parameters were considered at least useful as a “quick and rough” estimation of lipophilicity.

Octadecyl-functionalized silica gel was used as a stationary phase in the RP-HPLC evaluation of **8a–h**. A gradient of two solvents at different volume ratios modulated retention properties of a stationary phase [38]. Liquid binary mixtures of methanol (MeOH) with water were employed as mobile phases in a present isocratic RP-HPLC method. The MeOH organic modifier was preferred because of making a reversed-phase chromatographic system closer to the octan-1-ol/water partitioning one in terms of sensitivity to H -bond donor properties of investigated compounds [39,40]. The modifier was applied in different volume concentrations that varied from 60% to 85% (v/v).

The isocratic separation was possible and in addition, reasonable retention of the analyzed compounds **8a–h** was observed in all mobile phases. The estimated k parameters were found in an acceptable interval from 0.5 to 20 [39] and were listed in Table S2 (Supplementary Materials).

Increase in a volume concentration of MeOH led to shortening of t_r and $\log k$ values for all molecules **8a–h** (Table 1).

The 3-alkoxycarbonylamino substituent-containing derivatives **8a–d** showed higher t_r and $\log k$ parameters than their 4-positional isomers **8e–h** (Table 1), albeit excluding compounds **8d** and **8h**. Elongation of an R substituent led to the increase in t_r and $\log k$ values within both groups **8a–d** and **8e–h** (Table 1). Lower $\log k$ parameters of a molecule **8e** compared to those of a derivative **8a** (Table S2 in Supplementary Materials) would be a result of “linearity” of the 4-substituted molecule as well as interactions between the mobile phase and methoxy moiety of **8e**. Hydrogen atoms of this group were more acidic due to movement of electrons as a consequence of a different electronegativity of carbon and oxygen so the ability of the compound **8e** to form hydrogen bonds with a particular MeOH-containing mobile phase would be enhanced.

The highest $\log k$ values were observed for 1-(2-[3-[(butoxycarbonyl)amino]phenyl]-2-hydroxyethyl)-4-(2-fluorophenyl)piperazin-1-ium chloride (**8d**) and its positional isomer, 1-(2-[4-[(butoxycarbonyl)amino]phenyl]-2-hydroxyethyl)-4-(2-fluorophenyl)piperazin-1-ium chloride (**8h**; Table 1).

Extrapolation of the estimated $\log k$ parameters to elution with 100% water, i.e., the calculation of $\log k_w$ values, has become a widely accepted approach. The $\log k_w$ descriptor has been considered more efficient predictor of a biological activity than the $\log k$ itself because it reduced influence of an organic mobile phase modifier what was necessary to obtain measurable elution [41–43]. The $\log k_w$ values were extrapolated from intercepts of a linear relationship between the $\log k$ and volume fraction of a mobile phase modifier (φ_M) using a Snyder-Soczewiński relationship [42–44]. The linear relationship was justified by a correlation coefficient (R) > 0.9900 and adjusted coefficient of determination ($Adj. R^2$) > 0.9700 for all fitting models, excluding the one connected with the compound **8d** (Table 2). Anyway, values of the calculated statistical descriptors related to **8d** were also satisfactory ($R = 0.9730$, $Adj. R^2 = 0.9201$; Table 2).

The extrapolated $\log k_w$ values of the analyzed compounds **8a–h** (Table 2) were in accordance with their elution order and hydrophobicity and ranged from 2.113 (**8e**) to 2.930 (**8h**). Higher $\log k_w$ values were observed for the derivatives **8a–c** compared to **8e–g**. Butoxycarbonylamino substituent-containing compounds **8d** and **8h** were found to be the most lipophilic, as proven by their $\log k_w$ of 2.796 (**8d**) and 2.930 (**8h**), respectively (Table 2).

The slope S of a regression line used to obtain $\log k_w$ encoded notable information regarding a specific hydrophobic surface area and could serve as indicative measure of uniformity of a retention mechanism. If uniformity was observed, a convenient model between the slope(s) and intercept(s) was anticipated [45]. The currently calculated S parameters varied from 2.7386 (**8e**) to 3.3441 (**8h**; Table 2). The slope S was related to a specific hydrophobic surface of a compound and could be used as alternative measure of its lipophilicity [46].

Table 2. Extrapolated $\log k_w$ values (RP-HPLC) of the analyzed molecules **8a–h** and statistical descriptors (RSS , R , $Adj. R^2$, $RMSE$, NoR , F and $Prob > F$), which characterized a linear relationship between the $\log k$ and φ_M values for a particular compound. The φ_M parameter was a volume fraction of MeOH in the isocratic elution RP-HPLC.

Comp.	$\log k_w$	¹ S	² RSS	³ R	⁴ $Adj. R^2$	⁵ $RMSE$	⁶ NoR	⁷ F	⁸ $Prob > F$
8a	2.430	3.0678	0.0023	0.9967	0.9902	0.0337	0.0477	305.70	0.0033 **
8b	2.546	3.0529	0.0017	0.9975	0.9925	0.0294	0.0415	398.96	0.0025 **
8c	2.679	3.1244	0.0051	0.9930	0.9791	0.0504	0.0713	141.72	0.0070 **
8d	2.796	3.1641	0.0208	0.9730	0.9201	0.1019	0.1441	35.58	0.0270 *
8e	2.113	2.7386	0.0019	0.9965	0.9896	0.0311	0.0440	285.13	0.0035 **
8f	2.512	3.1156	0.0009	0.9988	0.9964	0.0208	0.0294	826.63	0.0012 **
8g	2.600	3.0739	0.0027	0.9962	0.9885	0.0367	0.0519	259.09	0.0038 **
8h	2.930	3.3441	0.0081	0.9903	0.9710	0.0637	0.0901	101.50	0.0097 **

¹ S , Slope; ² RSS , residual sum of squares; ³ R , correlation coefficient; ⁴ $Adj. R^2$, adjusted coefficient of determination; ⁵ $RMSE$, root mean squared error (standard deviation); ⁶ NoR , norm of residuals; ⁷ F , Fisher's significance ratio (Fisher's F -test); ⁸ $Prob > F$, probability of obtaining the F Ratio (significance of a whole model). Indication of a significance level of the F Ratio was as follows: * (one star), significant; ** (two stars), very significant.

Statistically extremely significant relationship between the $\log k_w$ and S values was described by Equation (1). The model was characterized by the $Prob > F$ parameter, which was in the range from 0 to <0.0010 :

$$S = 0.6457 (\pm 0.0882) \times \log k_w + 1.4219 (\pm 0.2208) \quad (1)$$

$$RSS = 0.0199, R = 0.9483, Adj. R^2 = 0.8826, RMSE = 0.0575, NoR = 0.1409, F = 53.61, Prob > F = 0.0003, n = 8$$

Based on the calculated statistical descriptors provided above, the uniformity of a retention mechanism of the studied derivatives **8a–h** was proven and suitability of selected mobile phases was confirmed for lipophilicity evaluation.

2.1.3. Electronic Properties

Electronic properties of the inspected compounds **8a–h** (Table 3) were characterized by logarithms of molar absorption coefficients ($\log \epsilon$) of their methanolic solutions ($c = 3.0 \times 10^{-5}$ M) investigated in the UV/Vis region of the spectrum.

Table 3. Wavelengths of the observed absorption maxima (λ_1 , $\lambda_{2(\text{Ch-T})}$ and λ_3) and logarithms of the molar absorption coefficients ($\log \epsilon$) of compounds' methanolic solutions ($c = 3.0 \times 10^{-5}$ M), which were investigated in the UV/Vis region of a spectrum.

Comp.	λ_1 (nm)	$\log \epsilon_1$	$\lambda_{2(\text{Ch-T})}$ (nm)	¹ $\log \epsilon_{2(\text{Ch-T})}$	λ_3 (nm)	$\log \epsilon_3$
8a	210	4.30	238	4.30	276	3.45
8b	210	4.31	238	4.33	276	3.40
8c	210	4.30	238	4.37	276	3.42
8d	210	4.31	238	4.32	276	3.49
8e	210	4.61	240	4.67	274	3.67
8f	208	4.59	240	4.60	274	3.60
8g	210	4.47	240	4.54	274	3.52
8h	208	4.34	240	4.42	274	3.42

¹ $\log \epsilon_{2(\text{Ch-T})}$, Logarithms of molar absorption coefficients observed at the charge-transfer absorption maximum $\lambda_{2(\text{Ch-T})} = 238\text{--}240$ nm.

The solutions showed three absorption maxima in a near ultraviolet (quartz) region of the electromagnetic spectrum between 200 and 400 nm [47], e.g., $\lambda_1 = 208\text{--}210$ nm, $\lambda_{2(\text{Ch-T})} = 238\text{--}240$ nm and $\lambda_3 = 274\text{--}276$ nm, respectively (Table 3).

The $\log \epsilon_{2(\text{Ch-T})}$ parameters of the compounds **8a–d** observed at a charge-transfer absorption maximum $\lambda_{2(\text{Ch-T})}$ were found in a narrow interval from 4.30 (**8a**) to 4.37 (**8c**). The methanolic solutions

of **8e–h** were characterized by higher $\log \varepsilon_{2(\text{Ch-T})}$ values than the ones of **8a–d** and varied from 4.42 (**8h**) to 4.67 (**8e**; Table 3). In addition, elongation of the 4-side chain led to lower $\log \varepsilon$ values related to all observed absorption maxima (Table 3).

2.2. Biological Assays

2.2.1. In Vitro Antimycobacterial Activity and Structure–Activity Relationships

The compounds **8a–h** were initially tested *in vitro* against *M. tuberculosis* CNCTC My 331/88 (identical with H₃₇R_v and ATCC 2794), *M. avium* CNCTC My 330/80 (identical with ATCC 25291), *M. avium intracellulare* ATCC 13950 and *M. kansasii* CNCTC My 235/80 (identical with ATCC 12478), respectively, as well as against *M. kansasii* 6 509/96, *M. kansasii* CIT11/06, *M. avium* subsp. *paratuberculosis* CIT03 and *M. avium hominissuis* CIT10/08 clinical isolates by methods described earlier [17,48–50].

The MIC was defined as the lowest concentration of a particular compound, which (i) inhibited growth of *M. tuberculosis* CNCTC My 331/88, *M. avium* CNCTC My 330/80, *M. kansasii* CNCTC My 235/80 or *M. kansasii* 6 509/96 [48]; (ii) prevented a visual colour change from blue to pink when testing susceptibility of the *M. kansasii* CIT11/06, *M. avium* subsp. *paratuberculosis* CIT03, *M. avium hominissuis* CIT10/08 or *M. avium intracellulare* ATCC 13950 strain. The MIC for given mycobacteria was defined as 90% or greater reduction of their growth (*IC*₉₀) compared to a control [17,49,50].

The efficiency of newly synthesized molecules **8a–h** was compared to the activity of reference drugs, i.e., isonicotinic acid hydrazide (isoniazid, INH), ethambutol (EMB), ofloxacin (OFLX), ciprofloxacin (CPX) or pyrazinamide (PZA) under same experimental conditions.

Next sections of the current research were focused specifically on the most susceptible strains, i.e., *M. tuberculosis* CNCTC My 331/88, *M. kansasii* CNCTC My 235/80 and *M. kansasii* 6 509/96, respectively. The *in vitro* activities (MIC values) of the most promising *N*-arylpiperazines were highlighted by a bold font style in gray (Table 4).

Table 4. The *in vitro* activity (the MIC values were expressed in the μM units) of currently screened compounds **8a–h** and reference drugs isoniazid (INH), ethambutol (EMB) and ofloxacin (OFLX) against *M. tuberculosis* My 331/88 (*M. tuberculosis* H₃₇R_v; MT My 331/88), *M. kansasii* My 235/80 (MK My 235/80), *M. kansasii* 6 509/96 (MK 6 509/96) and *M. avium* My 330/88 (MA My 330/88), respectively.

Comp.	MIC (μM)									
	MT My 331/88		MK My 235/80			MK 6 509/96			MA My 330/88	
	¹ 14 d	² 21 d	³ 7 d	14 d	21 d	7 d	14 d	21 d	14 d	21 d
8a	250	250	125	500	1000	125	500	500	500	500
8b	125	125	62.5	250	250	62.5	250	250	250	250
8c	62.5	62.5	62.5	125	125	32	125	125	125	250
8d	32	32	32	62.5	62.5	16	32	62.5	62.5	62.5
8e	125	125	125	500	500	125	500	500	250	500
8f	32	62.5	125	>250	>250	62.5	>125	>125	>250	>250
8g	16	16	125	>250	>250	125	>250	>250	>250	>250
8h	8	8	62.5	>125	>125	125	>250	>250	>250	250
INH	0.5	0.5	>250	>250	>250	4	8	8	>250	>250
EMB	1	2	1	2	2	1	2	2	16	16
OFLX	1	2	0.5	1	1	0.5	0.5	1	32	62.5

¹ 14 d, 14-Day cultivation; ² 21 d, 21-day cultivation; ³ 7 d, 7-day cultivation. The *in vitro* activities (MIC values) of the most promising *N*-arylpiperazines were highlighted by a bold font style in gray.

The position of a side chain notably affected the activity of tested derivatives **8a–h** against *M. tuberculosis* CNCTC My 331/88 (Table 4). After 14- and 21-day cultivation (14-d/21-d), the 4-positional isomers were more active, with the MIC values ranging from 8 μM (**8h**) to 125 μM (**8e**), than the 3-positional ones, which possessed the MICs from 32 μM (**8d**) to 250 μM (**8a**).

Among all the *in vitro* screened molecules, INH standard was found to be the most active with the MIC = 0.5 μ M (14-d/21-d).

Introduction of a 4-(pyrimidin-2-yl)piperazin-1-yl fragment instead of the 4-(2-fluorophenyl)piperazin-1-yl one led to the derivatives MM, which showed a comparable efficiency [18] to the compounds **8h** and **8d**, especially if they contained $R = C_3H_7/C_4H_9$ (Figure 2). Similarly, presence of an 3-alkoxyphenylcarbamoyloxy moiety (alkoxy = methoxy to butoxy) and elongation of a connecting chain resulted in the molecules **IM** (Figure 3) with a comparable *in vitro* activity [51] to **8a–h**.




Figure 3. *N*-Arylpiperazines containing a 2-hydroxypropane-1,3-diyl connecting chain (a series **IM**), which were *in vitro* screened against *M. tuberculosis* H₃₇R_v [51].

Isosteric replacement of the carbamoyloxy with a carboxy group in a structure of the compounds **IM** and introduction of an 4-alkoxycarbonylamino side chain (alkoxy = methoxy to butoxy) at the aromatic ring resulted in decreased *in vitro* efficiency of such modified racemic derivatives against *M. tuberculosis* H₃₇R_v [16].

The compounds **8a–d** were more efficient against *M. kansasii* My 235/80 and *M. kansasii* 6 509/96 than the substances **8e–h**. The most active compound against both mycobacteria was **8d** with the MIC = 16 μ M and 62.5 μ M, respectively, depending on a particular strain and also on the number of days of incubation. Increase in length of the side chain resulted in lower MIC values of **8a–d** against both tested *M. kansasii* strains. The observed MIC values were, however, higher compared to the ones related to EMB with the MIC = 1 μ M and 2 μ M (14-d/21-d), or OFLX, which showed the MIC = 0.5 μ M and 1 μ M, respectively (14-d/21-d; Table 4).

The *in vitro* activity of screened compounds **8a–h** against a non-tuberculous INH-resistant *M. avium* CNCTC My 330/80 was apparently dependent on the position of the alkoxy carbonylamino chain *R*. Its presence in the 3-position (**8a–d**) led to the MIC values varying from 62.5 μ M (**8d**) to 500 μ M (**8a**; 14-d/21-d). However, a potential of 4-substituent-containing derivatives (**8e–h**) to fight given mycobacterium was insufficient (MIC > 250 μ M; Table 4).

The activity of the most active substance **8d** (MIC = 62.5 μ M; 14-d/21-d) against *M. avium* CNCTC My 330/80 was comparable to the effectiveness of OFLX (MIC = 32 μ M and 62.5 μ M, respectively; 14-d/21-d). The EMB reference drug was slightly more active (MIC = 16 μ M; 14-d/21-d) than **8d**. Elongation of an 3-*R* side chain led to more potent compounds (Table 4).

The molecules **8a–h** were *in vitro* practically inactive against *M. kansasii* CIT11/06, *M. avium* subsp. *paratuberculosis* CIT03, *M. avium* *intracellulare* ATCC 13950 and *M. avium* *hominissuis* CIT10/08, respectively (MIC \geq 295 μ M; Table 5). The CPX standard was the most effective among all investigated compounds (MIC > 91 μ M and 181 μ M, respectively; Table 5).

The lipophilicity has been considered one of the most important factors, which critically influenced a compound's activity in penetrating mycobacterial cell walls [48,52,53]. To explore this statement in detail, relationships between the log k_w (independent variable) and activity values (dependent variable) of the compounds **8a–h** were inspected. For purposes of the current SAR study, observed MIC values were transformed into log (1/MIC [M]) units.

The INH, EMB and OFLX standard drugs were not included in the investigated models because of being structurally different and, in addition, different modes of their action have been proposed [54–60].

Table 5. The *in vitro* activity (the MIC values were expressed in the μM units) of the inspected compounds **8a–h** and reference drugs isoniazid (INH), ciprofloxacin (CPX) and pyrazinamide (PZA) against *M. kansasii* CIT11/06 (MK CIT11/06), *M. avium* subsp. *paratuberculosis* CIT03 (MAP CIT03), *M. avium intracellulare* ATCC 13950 (MAI ATCC 13950) and *M. avium hominissuis* CIT10/08 (MAH CIT10/08), respectively.

Comp.	MIC (μM)			
	MK	MAP	MAI	MAH
	CIT11/06	CIT03	ATCC 13950	CIT10/08
8a	>610	>610	>610	>610
8b	295	>590	>590	>590
8c	>571	>571	>571	>571
8d	>553	> 53	>553	>553
8e	>610	>610	>610	>610
8f	295	>590	>590	>590
8g	>571	>571	>71	>571
8h	>553	>553	>553	>553
INH	>1823	>1823	>1823	>1823
CPX	>91	181	181	181
PZA	>2031	>2031	>2031	487

Linear regression analyses were carried out using the Origin Pro 9.0.0 software (OriginLab Corporation, Northampton, MA, USA). More details about particular statistical parameters and significance levels were provided in the Materials and Methods section of a current paper.

Regarding the **8a–h** set, analyses related to *M. tuberculosis* My 331/88 (*M. tuberculosis* H₃₇R_v; 14-d/21-d) were expressed by Equations (2) and (3):

MT My 331/88 (14-d), **8a–h**:

$$\log (1/\text{MIC} [\text{M}]) = 1.4383 (\pm 0.5893) \times \log k_w + 0.6071 (\pm 1.5239) \quad (2)$$

RSS = 0.8867, $R = 0.7059$, Adj. $R^2 = 0.4146$, RMSE = 0.3844, NoR = 0.9417, $F = 5.95$, $\text{Prob} > F = 0.0504$, $n = 8$

MT My 331/88 (21-d), **8a–h**:

$$\log (1/\text{MIC} [\text{M}]) = 1.4819 (\pm 0.5598) \times \log k_w + 0.4586 (\pm 1.4476) \quad (3)$$

RSS = 0.8002, $R = 0.7340$, Adj. $R^2 = 0.4619$, RMSE = 0.3652, NoR = 0.8945, $F = 7.01$, $\text{Prob} > F = 0.0382$, $n = 8$

Based on given statistical parameters, only Equation (3) described a statistically significant model, for which the $\text{Prob} > F$ value was found the interval from 0.0100 to <0.0500.

If the analysis was separately applied to the groups **8a–d** and **8e–h**, much more convenient values of statistical descriptors were calculated for **8a–d** (14-d/21-d) as follows: RSS = 0.0003, $R = 0.9997$, Adj. $R^2 = 0.9991$, RMSE = 0.0119, NoR = 0.0168, $F = 3143.04$, $\text{Prob} > F = 0.0003$, $n = 4$. The statistically extremely significant models (Table S3 in Supplementary Materials) were characterized by Equations (S1) and (S3).

On the other hand, the relationships connected with the **8e–h** series were statistically insignificant, as proven by Equation (S2) and (S4), respectively (Table S3).

Focusing on the *M. kansasii* My 235/80 strain after 7-day cultivation (7-d), only the derivatives **8b–d** showed interesting MIC values of 32 μM or 62.5 μM (Table 4). However, the linear regression model involving the $\log k_w$ and $\log (1/\text{MIC} [\text{M}])$ values related to the **8a–d** set was statistically insignificant, as expected (Equation (S5) in Table S3).

Differing effectacy of the compounds **8a–d** against *M. tuberculosis* My 331/88 (*M. tuberculosis* H₃₇R_v) and *M. kansasii* My 235/80 (Table 4) was probably a consequence of diverse composition of the bacterial membrane of those strains [61,62].

Lipophilicity could play a crucial role in a mechanism of action of the compounds **8a–d** against the *M. kansasii* 6 509/96 clinical isolate (7-d; Table 4), as provided by Equation (4).

MK 6 509/96 (7-d), **8a–d**:

$$\log (1/MIC [M]) = 2.4098 (\pm 0.0576) \times \log k_w - 1.9466 (\pm 0.1507) \quad (4)$$

$RSS = 0.0005$, $R = 0.9994$, $Adj. R^2 = 0.9982$, $RMSE = 0.0158$, $NoR = 0.0224$, $F = 1750.57$, $Prob > F = 0.0005$, $n = 4$

Despite of very convenient values of the statistical descriptors, a main limitation of the approach was a low number of the compounds ($n = 4$) involved in this analysis.

The $\log \epsilon_{2(\text{Ch-T})}$ values (independent variable), which were observed at the charge-transfer absorption maximum $\lambda_{2(\text{Ch-T})}$, were taken into a special consideration (Table 3), because they could be the most sensitive to the differences in position and electronic properties of a particular alkoxy-carbonylamino substituent [47].

The relationship between the $\log \epsilon_{2(\text{Ch-T})}$ parameters and $\log (1/MIC [M])$ values connected with the *in vitro* screening of **8a–h** against *M. tuberculosis* My 331/88 (*M. tuberculosis* H₃₇R_v; 14-d) provided a bilinear course. Based on this fitting, maximal efficiency of the tested compounds could be observed if their $\log \epsilon_{2(\text{Ch-T})}$ values were approximately 4.43 (Figure 4).




Figure 4. Bilinear relationship between the $\log \epsilon_{2(\text{Ch-T})}$ and $\log (1/MIC [M])$ parameters of the investigated compounds **8a–h**. The dependent variable values were taken from the *in vitro* screening of these derivatives against *M. tuberculosis* CNCTC My 331/88 (*M. tuberculosis* H₃₇R_v; 14-d).

Regarding the *M. kansasii* My 235/80 and *M. kansasii* 6 509/96 strains, no significant relationships between the *in vitro* activity of the compounds **8a–h** and their electronic features were observed.

2.2.2. In Vitro Cytotoxicity Screening

Cytotoxicity of the compounds **8a–h** was inspected as the LD_{50} value, i.e., a lethal dose to 50% of a cell population, which was derived from survival rate curves. The highest dose of all tested derivatives in a medium (30 μM) did not lead to a significant lethal effect on a human monocytic leukemia THP-1 cell line.

The tested molecules showed low (insignificant) toxicity of $LD_{50} > 30 \mu\text{M}$ against given cell line. Moreover, the compounds **8a** and **8e** increased proliferation of the THP-1 cells in 24 h when compared

to a control. Relative survival rate (in percentages) of the THP-1 cells for all tested derivatives was found to be over 80%, excluding the molecule **8d**, where it was 79% at the highest tested concentration of 30 μM (Figure S1 in Supplementary Materials). Only the compounds with the $IC_{50} < 10 \mu\text{M}$ could be considered antiproliferative (cytotoxic) agents [63], and the highest tested concentration used for the current toxicity assay was 3-fold this value.

As the LD_{50} values of oxaliplatin and camptothecin standard drugs assessed in this cell line were considerably lower ($1.7 \pm 0.64 \mu\text{M}$ and $0.16 \pm 0.07 \mu\text{M}$, respectively), the discussed compounds **8a–h** were deemed non-toxic agents suitable for further design and development of novel antimycobacterials.

3. Materials and Methods

3.1. General Information

All reagents used for syntheses were commercially available from Alpha Aesar (Lancashire, UK), Fluka Chemie (Buchs, Switzerland), Lachema (Brno, Czech Republic), LachNer (Neratovice, Czech Republic), Lancaster (Ward Hill, MA, USA), Merck (Darmstadt, Germany) or Sigma-Aldrich (Dorset, UK) in sufficient quality and were used without additional purification. Solvents were dried and freshly distilled before use.

Thin-layer chromatography (TLC) Kieselgel 60 F₂₅₄ plates (Merck) visualized by UV irradiation ($\lambda = 254 \text{ nm}$) were used to monitor reactions and purity of synthesized compounds.

Melting point (Mp or mp) values of prepared intermediates and final compounds, respectively, were determined on the Kofler hot plate apparatus HMK (Franz Kustner Nacht GK, Dresden, Germany) and left uncorrected. The mps of some intermediates were already published, i.e., **3a**: 102–104 °C [64]; **3b**: 111–112 °C [64] and 113–114 °C [65], respectively; **3d**: 53–55 °C [64]; **3e**: 168 °C [64] and 160–162 °C [66], respectively; **3f**: 158 °C [67], 157–158 °C [68,69] and 160–161 °C [64], respectively; **3h**: 87–88.5 °C [64]; **4a**: 99–103 °C [70]; **4b**: 108–110 °C [71]; **4d**: 80–86 °C [70]; **4e**: 200–201 °C [72].

The R_f values of prepared salts **8a–h** were obtained by the TLC on 10-cm aluminium sheets pre-coated with silica gel 60 F₂₅₄ (0.25 mm thickness; Merck) in glass developing chambers using petroleum ether/diethylamine (10:3, *v/v*) eluant as a mobile phase. Spots were located under iodine vapours/UV light at $\lambda = 254 \text{ nm}$.

The ¹H- and ¹³C-NMR spectral analyses were carried out on the FT-NMR spectrometer (Varian Co., Palo Alto, CA, USA) operating at 300 MHz (¹H-NMR) and 75 MHz (¹³C-NMR), respectively, in dried DMSO-*d*₆ using tetramethylsilane (TMS; Sigma-Aldrich, Darmstadt, Germany) as an internal standard.

Chemical shifts were reported in a δ scale in parts *per* million (ppm), coupling constants *J* were given in Hertz (Hz) and spin multiplicities were expressed as follows: s (singlet), d (doublet), dd (double doublet), t (triplet), q (quartet), m (multiplet). A complete assignment of ¹H- and ¹³C-NMR resonances was based on an interpretation of standard NMR values.

The FT-IR (IR) spectra were obtained by the ATR technique on the FT-IR spectrophotometer Impact 410 (Thermo Fisher Scientific, West Palm Beach, FL, USA). The absorption frequencies $\tilde{\nu}_{\text{max}}$ were reported in reciprocal centimeters (cm^{-1}) in a recorded range from 4000 cm^{-1} to 400 cm^{-1} .

The mass spectra (HR-MS) of prepared intermediates **3a–h**, **4a–h** and **6a–h** (Schemes 1 and 2) were measured using the high-performance liquid chromatograph Dionex UltiMate® 3000 (Thermo Fisher Scientific) coupled with the LTQ Orbitrap XL™ Hybrid Ion Trap-Orbitrap Fourier Transform Mass Spectrometer (Thermo Fischer Scientific) with injection into HESI-II in a positive mode.

The liquid chromatography mass spectra of the compounds **7a–h** and **8a–h** (Scheme 2, Table 1) were carried out on the Agilent 1100 LC/MSD Trap (Agilent Technologies, Santa Clara, CA, USA) in a positive mode using electrospray ionization at atmospheric pressure. The particular compound was dissolved in methanol ($c = 1 \text{ mg/mL}$) and the solution was passed through the XDB SOX 2.1 mm column (Agilent Technologies) with a 1.8 μm particle size at the pressure of 400 bar. The UV detection was performed at $\lambda = 254 \text{ nm}$. Nebulization gas (N_2) flow was 8 L/min, pressure was 40 psi. The MS electrospray operated at capillary voltage of 3.5 kV and temperature was set to 350 °C (ESI-MS). Fragments were described

as a relationship between atomic mass units and charge (m/z), a recorded interval was from 50 m/z to 1000 m/z .

The elemental analysis (% C, H, N) of the compounds **8a–h** was carried out by the Perkin-Elmer 2400 Series-II Elemental Analyzer (Perkin-Elmer, Waltham, MA, USA) and all the derivatives were within $\pm 0.40\%$ of calculation.

The chromatographic HPLC-Diode Array Detection apparatus for the determination of capacity (retention) factor k ($\log k$) values was the LC Agilent Infinity System (Agilent Technologies, Santa Clara, CA, USA) equipped with a Infinity 1260 gradient pump with a degasser, a 1260 HiPals automatic injector, a column thermostat 1290, a photo-diode array detector Infinity 1290 (all the equipments were obtained by Agilent Technologies) and personal computer with the Agilent ChemStation software (Agilent Technologies) for the registration of values and calibration procedures. The chromatographic column Eclipse plus RP C₁₈, 150 \times 4.6 mm i.d., a 5 μ m particle size (Agilent Technologies), was used and thermostated at 35 °C.

The analyses were performed at pressure ranged from 7 MPa to 15 MPa. The detection wavelength was set to 260 nm, injection sample volume was 5 μ L with a flow rate of 1.0 mL/min in all RP-HPLC analyses.

LC-MS Methanol (J. T. Baker Chemicals Co., Phillipsburg, NJ, USA) and HPLC-quality water (Sigma-Aldrich, Darmstadt, Germany) were used for a preparation of mobile phases. Water was firstly deionized and purified by the Millipore Simplicity 185 Ultrapure water purification system (Millipore, Billerica, MA, USA).

The UV/Vis spectra of methanolic solutions of the analyzed compounds **8a–h** ($c = 3.0 \times 10^{-5}$ M) were estimated on the 8452A Diode Array spectrophotometer HP-8452A (Hewlett Packard, Palo Alto, CA, USA). Methanol for UV-spectroscopy (Merck) was used for the preparation of these solutions.

Results of the UV/Vis analyses were collected and stored digitally using the ChemStation controller software (Agilent Technologies, Waldbronn, Germany). The HP-8452A apparatus measured a complete range of a spectrum from 190 nm to 820 nm.

3.2. Synthesis of Compounds

3.2.1. General Procedure For the Preparation of Alkyl (3-/4-Acetylphenyl)carbamates (**3a–h**)

Into a stirred solution of 3-aminoacetophenone **1a** (CAS Registry Number 99-03-6; 5.00 g, 37 mmol) or 4-aminoacetophenone **1b** (CAS Registry Number 99-92-3; 5.00 g, 37 mmol) and pyridine (3.0 mL, 37 mmol) in 20 mL of acetone, a solution of methyl chloroformate **2a** (CAS Registry Number 79-22-1; 3.5 mL, 37 mmol), ethyl chloroformate **2b** (CAS Registry Number 541-41-3; 4.0 mL, 37 mmol), propyl chloroformate **2c** (CAS Registry Number 109-61-5; 4.5 mL, 37 mmol) or butyl chloroformate **2d** (CAS Registry Number 592-34-7; 5.0 mL, 37 mmol) in 5 mL of acetone, was added dropwise. The particular mixture was heated to reflux for 3 h [28]. When the reaction was completed (TLC control), the solvents were removed in vacuo, crude solid products **3a–h** were washed with distilled water and recrystallized from absolute ethanol. Full characterization data for the compounds **3a–h** (Scheme 1), isolated as colourless solids, are given below.

Methyl (3-acetylphenyl)carbamate (3a); CAS Registry Number 87743-55-3). Yield 6.80 g (95%); M_r 193.19; Mp 103–104 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 9.87 (s, 1H, NHCOO), 8.06 (s, 1H, Ar-H), 7.71 (d, 1H, Ar-H, $J = 7.7$ Hz), 7.61 (d, 1H, Ar-H, $J = 7.3$ Hz), 7.44 (t, 1H, Ar-H, $J = 8.2$ Hz), 3.69 (s, 3H, COOCH₃), 2.55 (s, 3H, COCH₃); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 197.47, 153.92, 139.49, 137.37, 129.01, 122.56, 122.36, 117.26, 51.53, 26.51. HR-MS: for C₁₀H₁₁O₃N [M – H]⁺ calculated 192.06552 m/z , found 192.06728 m/z .

Ethyl (3-acetylphenyl)carbamate (3b); CAS Registry Number 39569-24-9). Yield 7.50 g (98%); M_r 207.19; Mp 112–113 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 9.83 (s, 1H, NHCOO), 8.07 (s, 1H, Ar-H), 7.68 (d, 1H, Ar-H, $J = 8.1$ Hz), 7.60 (d, 1H, Ar-H, $J = 7.7$ Hz), 7.42 (t, 1H, Ar-H, $J = 7.8$ Hz), 4.14 (q, 2H, CH₂CH₃,

$J = 7.1$ Hz), 2.53 (s, 3H, COCH₃), 1.24 (t, 3H, CH₂CH₃, $J = 7.1$ Hz); ¹³C-NMR (DMSO-*d*₆) δ_C (ppm): 197.53, 153.50, 139.60, 137.35, 129.01, 122.60, 122.33, 117.25, 60.21, 26.57, 14.37. HR-MS: for C₁₁H₁₃O₃N [M – H]⁺ calculated 206.08117 m/z , found 206.08297 m/z .

Propyl (3-acetylphenyl)carbamate (3c). Yield 7.30 g (89%); M_r 221.19; Mp 101–103 °C; ¹H-NMR (DMSO-*d*₆) δ_H (ppm): 9.85 (s, 1H, NHCOO), 8.09 (s, 1H, Ar–H), 7.70 (d, 1H, Ar–H, $J = 8.1$ Hz), 7.61 (d, 1H, Ar–H, $J = 8.1$ Hz), 7.44 (t, 1H, Ar–H, $J = 7.9$ Hz), 4.06 (t, 2H, CH₂CH₂CH₃, $J = 6.6$ Hz), 2.53 (s, 3H, COCH₃), 1.72–1.56 (m, 2H, CH₂CH₂CH₃), 0.94 (t, 3H, CH₂CH₂CH₃, $J = 7.5$ Hz); ¹³C-NMR (DMSO-*d*₆) δ_C (ppm): 197.45, 153.59, 139.60, 137.36, 128.96, 122.57, 122.27, 117.28, 65.73, 26.51, 21.76, 10.06. HR-MS: for C₁₂H₁₅O₃N [M – H]⁺ calculated 220.09682 m/z , found 220.09854 m/z .

Butyl (3-acetylphenyl)carbamate (3d); CAS Registry Number 72531-03-4). Yield 7.90 g (91%); M_r 235.19; Mp 58–59 °C; ¹H-NMR (DMSO-*d*₆) δ_H (ppm): 9.83 (s, 1H, NHCOO), 8.07 (s, 1H, Ar–H), 7.68 (d, 1H, Ar–H, $J = 7.7$ Hz), 7.60 (d, 1H, Ar–H, $J = 7.7$ Hz), 7.42 (t, 1H, Ar–H, $J = 7.9$ Hz), 4.09 (t, 2H, CH₂CH₂CH₂CH₃, $J = 6.6$ Hz), 2.54 (s, 3H, COCH₃), 1.64–1.54 (m, 2H, CH₂CH₂CH₂CH₃), 1.46–1.33 (m, 2H, CH₂CH₂CH₂CH₃), 0.92 (t, 3H, CH₂CH₂CH₂CH₃, $J = 6.2$ Hz); ¹³C-NMR (DMSO-*d*₆) δ_C (ppm): 197.53, 153.60, 139.60, 137.35, 129.01, 122.59, 122.30, 117.28, 63.94, 30.46, 26.56, 18.49, 13.45. HR-MS: for C₁₃H₁₇O₃N [M – H]⁺ calculated 234.11247 m/z , found 234.11425 m/z .

Methyl (4-acetylphenyl)carbamate (3e); CAS Registry Number 60677-43-2). Yield 7.05 g (99%); M_r 193.19; Mp 168–170 °C; ¹H-NMR (DMSO-*d*₆) δ_H (ppm): 10.09 (s, 1H, NHCOO), 7.91 (d, 2H, Ar–H, $J = 8.8$ Hz), 7.59 (d, 2H, Ar–H, $J = 8.8$ Hz), 3.70 (s, 3H, COOCH₃), 2.52 (s, 3H, COCH₃); ¹³C-NMR (DMSO-*d*₆) δ_C (ppm): 196.28, 153.72, 143.60, 130.99, 129.43, 117.20, 51.79, 26.24. HR-MS: for C₁₀H₁₁O₃N [M – H]⁺ calculated 192.06552 m/z , found 192.06728 m/z .

Ethyl (4-acetylphenyl)carbamate (3f); CAS Registry Number 5520-79-6). Yield 7.60 g (99%); M_r 207.19; Mp 161–163 °C; ¹H-NMR (DMSO-*d*₆) δ_H (ppm): 10.08 (s, 1H, NHCOO), 7.91 (d, 2H, Ar–H, $J = 8.8$ Hz), 7.59 (d, 2H, Ar–H, $J = 8.8$ Hz), 4.13 (q, 2H, CH₂CH₃, $J = 7.1$ Hz), 2.52 (s, 3H, COCH₃), 1.24 (t, 3H, CH₂CH₃, $J = 7.1$ Hz); ¹³C-NMR (DMSO-*d*₆) δ_C (ppm): 196.28, 153.38, 143.73, 130.94, 129.42, 117.19, 60.44, 26.23, 14.39. HR-MS: for C₁₁H₁₃O₃N [M – H]⁺ calculated 206.08117 m/z , found 206.08295 m/z .

Propyl (4-acetylphenyl)carbamate (3g). Yield 8.05 g (98%); M_r 221.19; Mp 125–126 °C; ¹H-NMR (DMSO-*d*₆) δ_H (ppm): 10.07 (s, 1H, NHCOO), 7.91 (d, 2H, Ar–H, $J = 8.8$ Hz), 7.60 (d, 2H, Ar–H, $J = 8.8$ Hz), 4.07 (t, 2H, CH₂CH₂CH₃, $J = 6.6$ Hz), 2.51 (s, 3H, COCH₃), 1.75–1.57 (m, 2H, CH₂CH₂CH₃), 0.94 (t, 3H, CH₂CH₂CH₃, $J = 7.3$ Hz); ¹³C-NMR (DMSO-*d*₆) δ_C (ppm): 196.28, 153.37, 143.71, 130.93, 129.42, 117.20, 65.96, 26.22, 21.73, 10.11. HR-MS: for C₁₂H₁₅O₃N [M – H]⁺ calculated 220.09682 m/z , found 220.09852 m/z .

Butyl (4-acetylphenyl)carbamate (3h); CAS Registry Number 72531-04-5). Yield 8.23 g (95%); M_r 235.19; Mp 89–91 °C; ¹H-NMR (DMSO-*d*₆) δ_H (ppm): 10.05 (s, 1H, NHCOO), 7.91 (d, 2H, Ar–H, $J = 8.8$ Hz), 7.60 (d, 2H, Ar–H, $J = 8.8$ Hz), 4.11 (t, 2H, CH₂CH₂CH₂CH₃, $J = 6.6$ Hz), 2.51 (s, 3H, COCH₃), 1.69–1.55 (m, 2H, CH₂CH₂CH₂CH₃), 1.48–1.30 (m, 2H, CH₂CH₂CH₂CH₃), 0.92 (t, 3H, CH₂CH₂CH₂CH₃, $J = 7.3$ Hz); ¹³C-NMR (DMSO-*d*₆) δ_C (ppm): 196.27, 153.36, 143.69, 130.92, 129.40, 117.17, 64.14, 30.40, 26.22, 18.47, 13.45. HR-MS: for C₁₃H₁₇O₃N [M – H]⁺ calculated 234.11247 m/z , found 234.11425 m/z .

3.2.2. General Procedure For the Preparation of Alkyl [3-/4-(Bromoacetyl)phenyl]carbamates (4a–h)

Into a stirred solution of a particular alkyl (3-/4-acetylphenyl)carbamate, i.e., **3a**, **3e** (6.96 g, 36 mmol), **3b**, **3f** (7.46 g, 36 mmol), **3c**, **3g** (7.97 g, 36 mmol), **3d** or **3h** (8.47 g, 36 mmol), in 80 mL of chloroform, a solution of bromine (1.9 mL, 36 mmol) in 10 mL of chloroform was added dropwise and stirred for 3 h at laboratory temperature. When the reaction was completed (TLC control), the solvent was removed *in vacuo* giving a crude solid product [31]. Intermediates **4a–h** (Scheme 1) were recrystallized from propan-2-ol. Full characterization parameters for the compounds **4a–h**, isolated as colourless solids, are given below.

Methyl [3-(bromoacetyl)phenyl]carbamate (4a). Yield 7.40 g (75%); M_r 272.09; Mp 99–103 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 9.92 (s, 1H, NHCO), 8.08 (s, 1H, Ar-H), 7.78–7.66 (m, 2H, Ar-H), 7.47 (t, 1H, Ar-H, $J = 8.1$ Hz), 4.89 (s, 2H, COCH₂Br), 3.68 (s, 3H, COOCH₃); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 191.39, 153.92, 139.70, 134.51, 129.17, 123.28, 122.98, 117.62, 51.64, 33.61. HR-MS: for C₁₀H₁₀O₃BrN [M – H]⁺ calculated 269.97603 m/z , found 269.97781 m/z .

Ethyl [3-(bromoacetyl)phenyl]carbamate (4b; CAS Registry Number 88541-97-3). Yield 9.30 g (90%); M_r 286.09; Mp 105–108 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 9.89 (s, 1H, NHCO), 8.10 (s, 1H, Ar-H), 7.78–7.66 (m, 2H, Ar-H), 7.46 (t, 1H, Ar-H, $J = 7.7$ Hz), 4.88 (s, 2H, COCH₂Br), 4.14 (q, 2H, CH₂CH₃, $J = 7.3$ Hz), 1.25 (t, 3H, CH₂CH₃, $J = 7.3$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 191.43, 153.48, 139.67, 134.51, 129.17, 123.29, 122.95, 117.60, 60.27, 33.72, 14.36. HR-MS: for C₁₁H₁₂O₃BrN [M – H]⁺ calculated 283.99168 m/z , found 283.99340 m/z .

Propyl [3-(bromoacetyl)phenyl]carbamate (4c). Yield 9.60 g (89%); M_r 300.09; Mp 104–107 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 9.90 (s, 1H, NHCO), 8.10 (s, 1H, Ar-H), 7.78–7.65 (m, 2H, Ar-H), 7.46 (t, 1H, Ar-H, $J = 7.9$ Hz), 4.88 (s, 2H, COCH₂Br), 4.05 (t, 2H, CH₂CH₂CH₃, $J = 6.6$ Hz), 1.73–1.55 (m, 2H, CH₂CH₂CH₃), 0.93 (t, 3H, CH₂CH₂CH₃, $J = 7.3$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 191.43, 153.59, 139.81, 134.51, 129.17, 123.30, 122.93, 117.61, 65.81, 33.70, 21.76, 10.13. HR-MS: for C₁₂H₁₄O₃BrN [M – H]⁺ calculated 298.00733 m/z , found 298.00913 m/z .

Butyl [3-(bromoacetyl)phenyl]carbamate (4d). Yield 9.60 g (85%); M_r 314.19; Mp 82–85 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 9.89 (s, 1H, NHCO), 8.10 (s, 1H, Ar-H), 7.77–7.65 (m, 2H, Ar-H), 7.46 (t, 1H, Ar-H, $J = 7.9$ Hz), 4.88 (s, 2H, COCH₂Br), 4.10 (t, 2H, CH₂CH₂CH₂CH₃, $J = 6.6$ Hz), 1.64–1.54 (m, 2H, CH₂CH₂CH₂CH₃), 1.47–1.34 (m, 2H, CH₂CH₂CH₂CH₃), 0.92 (t, 3H, CH₂CH₂CH₂CH₃, $J = 7.3$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 191.41, 153.57, 139.79, 134.50, 129.16, 123.27, 122.91, 117.61, 63.99, 33.64, 30.43, 18.46, 13.43. HR-MS: for C₁₃H₁₆O₃BrN [M – H]⁺ calculated 312.02298 m/z , found 312.02474 m/z .

Methyl [4-(bromoacetyl)phenyl]carbamate (4e; CAS Registry Number 942316-98-5). Yield 7.71 g (79%); M_r 272.09; Mp 200–202 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 10.17 (s, 1H, NHCOO), 7.97 (d, 2H, Ar-H, $J = 8.8$ Hz), 7.61 (d, 2H, Ar-H, $J = 8.8$ Hz), 4.85 (s, 2H, COCH₂Br), 3.71 (s, 3H, COOCH₃); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 190.15, 153.71, 144.30, 130.17, 127.90, 117.32, 51.90, 33.46. HR-MS: for C₁₀H₁₀O₃BrN [M – H]⁺ calculated 269.97603 m/z , found 269.97781 m/z .

Ethyl [4-(bromoacetyl)phenyl]carbamate (4f). Yield 9.63 g (93%); M_r 286.09; Mp 174–176 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 10.14 (s, 1H, NHCOO), 7.96 (d, 2H, Ar-H, $J = 8.8$ Hz), 7.62 (d, 2H, Ar-H, $J = 8.8$ Hz), 4.85 (s, 2H, COCH₂Br), 4.17 (q, 2H, CH₂CH₃, $J = 6.7$ Hz), 1.27 (t, 3H, CH₂CH₃, $J = 7.3$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 190.12, 153.22, 144.38, 130.14, 127.81, 117.29, 60.56, 33.46, 14.31. HR-MS: for C₁₁H₁₂O₃BrN [M – H]⁺ calculated 283.99168 m/z , found 283.9938 m/z .

Propyl [4-(bromoacetyl)phenyl]carbamate (4g). Yield 9.26 g (86%); M_r 300.09; Mp 154–155 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 10.14 (s, 1H, NHCOO), 7.96 (d, 2H, Ar-H, $J = 8.8$ Hz), 7.62 (d, 2H, Ar-H, $J = 8.8$ Hz), 4.84 (s, 2H, COCH₂Br), 4.08 (t, 2H, CH₂CH₂CH₃, $J = 6.6$ Hz), 1.75–1.57 (m, 2H, CH₂CH₂CH₃), 0.94 (t, 3H, CH₂CH₂CH₃, $J = 7.3$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 190.15, 153.36, 144.41, 130.16, 127.82, 117.32, 66.07, 33.48, 21.73, 10.14. HR-MS: for C₁₂H₁₄O₃BrN [M – H]⁺ calculated 298.00733 m/z , found 298.00911 m/z .

Butyl [4-(bromoacetyl)phenyl]carbamate (4h). Yield 9.64 g (85%); M_r 314.19; Mp 152–154 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 10.13 (s, 1H, NHCOO), 7.96 (d, 2H, Ar-H, $J = 8.8$ Hz), 7.62 (d, 2H, Ar-H, $J = 8.8$ Hz), 4.84 (s, 2H, COCH₂Br), 4.12 (t, 2H, CH₂CH₂CH₂CH₃, $J = 6.6$ Hz), 1.70–1.56 (m, 2H, CH₂CH₂CH₂CH₃), 1.48–1.30 (m, 2H, CH₂CH₂CH₂CH₃), 0.92 (t, 3H, CH₂CH₂CH₂CH₃, $J = 7.3$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 190.12, 153.31, 144.38, 130.13, 127.79, 117.28, 64.23, 33.43, 30.38, 18.47, 13.45. HR-MS: for C₁₃H₁₆O₃BrN [M – H]⁺ 312.02298 m/z , found 312.02474 m/z .

3.2.3. General Procedure For the Preparation of Alkyl {3-/4-[(4-(2-Fluorophenyl)piperazin-1-yl)-acetyl]phenyl}carbamates (6a–h)

A solution of a particular alkyl [3-/4-(bromoacetyl)phenyl]carbamate, i.e., **4a**, **4e** (1.50 g, 5.5 mmol), **4b**, **4f** (1.57 g, 5.5 mmol), **4c**, **4g** (1.65 g, 5.5 mmol), **4d** or **4h** (1.73 g, 5.5 mmol) in 30 mL of anhydrous tetrahydrofuran (THF) was added dropwise into a stirred solution of 1-(2-fluorophenyl)piperazine **5** (CAS Registry Number 111-15-0; 1.00 g, 5.5 mmol) and triethylamine (TEA; 0.8 mL, 5.5 mmol) in 20 mL of anhydrous THF [32]. The particular mixture was stirred for 3 h at laboratory temperature. When the reaction was completed (TLC control), the solvents were removed *in vacuo* and remaining solid was treated with 100 mL of distilled water and 100 mL of chloroform. The organic layer was washed with distilled water, dried over anhydrous sodium sulphate and evaporated *in vacuo* giving crude solid products. Prepared intermediates **6a–h** (Scheme 2) were recrystallized from acetone. Full characterization parameters for the compounds **6a–h**, isolated as colourless solids, are provided below.

Methyl {3-[(4-(2-fluorophenyl)piperazin-1-yl)acetyl]phenyl}carbamate (6a). Yield 1.80 g (86%); M_r 371.39; Mp 103–105 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 9.86 (s, 1H, NHCO), 8.09 (s, 1H, Ar-H), 7.75–7.60 (m, 2H, Ar-H), 7.40 (t, 1H, Ar-H, $J = 7.9$ Hz), 7.07–6.98 (m, 4H, Ar-H), 3.87 (s, 2H, COCH₂N), 3.68 (s, 3H, COOCH₃), 3.10–3.00 (m, 4H, 3,5-piperazine), 2.75–2.60 (m, 4H, 2,6-piperazine); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 196.47, 154.86 (d, $J = 242.8$ Hz), 153.92, 139.76 (d, $J = 8.4$ Hz), 139.45, 136.44, 128.89, 124.66 (d, $J = 3.0$ Hz), 122.63, 122.19, 122.09 (d, $J = 7.6$ Hz), 119.15 (d, $J = 2.3$ Hz), 117.35, 115.37 (d, $J = 20.5$ Hz), 63.55, 52.56, 51.58, 50.00 (d, $J = 3.1$ Hz). HR-MS: for C₂₀H₂₂O₃FN₃ [M – H]⁺ calculated 370.15615 m/z , found 370.15791 m/z .

Ethyl {3-[(4-(2-fluorophenyl)piperazin-1-yl)acetyl]phenyl}carbamate (6b). Yield 1.64 g (75%); M_r 385.44; Mp 127–129 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 9.84 (s, 1H, NHCO), 8.11 (s, 1H, Ar-H), 7.75–7.60 (m, 2H, Ar-H), 7.42 (t, 1H, Ar-H, $J = 7.9$ Hz), 7.10–6.94 (m, 4H, Ar-H), 4.13 (q, 2H, CH₂CH₃, $J = 7.1$ Hz), 3.87 (s, 2H, COCH₂N), 3.10–3.00 (m, 4H, 3,5-piperazine), 2.75–2.60 (m, 4H, 2,6-piperazine), 1.24 (t, 3H, CH₂CH₃, $J = 7.0$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 196.47, 154.89 (d, $J = 242.8$ Hz), 153.50, 139.77 (d, $J = 8.4$ Hz), 139.55, 136.41, 128.92, 124.68 (d, $J = 3.0$ Hz), 122.66, 122.15, 122.15 (d, $J = 7.6$ Hz), 119.17 (d, $J = 2.3$ Hz), 117.32, 115.78 (d, $J = 20.5$ Hz), 63.55, 60.21, 52.58, 49.99 (d, $J = 3.1$ Hz), 14.37. HR-MS: for C₂₁H₂₄O₃FN₃ [M – H]⁺ calculated 384.17180 m/z , found 384.17360 m/z .

Propyl {3-[(4-(2-fluorophenyl)piperazin-1-yl)acetyl]phenyl}carbamate (6c). Yield 1.78 g (80%); M_r 399.47; Mp 124–126 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 9.84 (s, 1H, NHCO), 8.11 (s, 1H, Ar-H), 7.75–7.60 (m, 2H, Ar-H), 7.42 (t, 1H, Ar-H, $J = 7.9$ Hz), 7.10–6.94 (m, 4H, Ar-H), 4.06 (t, 2H, CH₂CH₂CH₃, $J = 6.6$ Hz), 3.87 (s, 2H, COCH₂N), 3.10–3.00 (m, 4H, 3,5-piperazine), 2.70–2.55 (m, 4H, 2,6-piperazine), 1.74–1.56 (m, 2H, CH₂CH₂CH₃), 0.94 (t, 3H, CH₂CH₂CH₃, $J = 7.3$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 196.47, 154.89 (d, $J = 242.8$ Hz), 153.62, 139.79 (d, $J = 8.4$ Hz), 139.58, 136.41, 128.95, 124.74 (d, $J = 3.0$ Hz), 122.66, 122.15 (d, $J = 7.6$ Hz), 122.16, 119.17 (d, $J = 2.3$ Hz), 117.32, 115.80 (d, $J = 20.5$ Hz), 65.77, 63.56, 52.61, 49.98 (d, $J = 3.1$ Hz), 21.79, 10.14. HR-MS: for C₂₂H₂₆O₃FN₃ [M – H]⁺ calculated 398.18745 m/z , found 398.18917 m/z .

Butyl {3-[(4-(2-fluorophenyl)piperazin-1-yl)acetyl]phenyl}carbamate (6d). Yield 1.85 g (81%); M_r 413.50; Mp 112–114 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 9.84 (s, 1H, NHCO), 8.11 (s, 1H, Ar-H), 7.75–7.60 (m, 2H, Ar-H), 7.42 (t, 1H, Ar-H, $J = 7.9$ Hz), 7.11–6.94 (m, 4H, Ar-H), 4.10 (t, 2H, CH₂CH₂CH₂CH₃, $J = 6.2$ Hz), 3.87 (s, 2H, COCH₂N), 3.15–3.00 (m, 4H, 3,5-piperazine), 2.80–2.60 (m, 4H, 2,6-piperazine), 1.68–1.55 (m, 2H, CH₂CH₂CH₂CH₃), 1.48–1.33 (m, 2H, CH₂CH₂CH₂CH₃), 0.91 (t, 3H, CH₂CH₂CH₂CH₃, $J = 6.8$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 196.47, 154.89 (d, $J = 242.8$ Hz), 153.59, 139.77 (d, $J = 8.4$ Hz), 139.55, 136.43, 128.87, 124.65 (d, $J = 3.0$ Hz), 122.63, 122.10, 122.10 (d, $J = 7.6$ Hz), 119.13 (d, $J = 2.3$ Hz), 117.35, 115.75 (d, $J = 20.5$ Hz), 63.93, 63.56, 52.58, 49.98 (d, $J = 3.1$ Hz), 30.46, 18.47, 13.42. HR-MS: for C₂₃H₂₈O₃FN₃ [M – H]⁺ calculated 412.20310 m/z , found 412.20488 m/z .

Methyl 4-[(4-(2-fluorophenyl)piperazin-1-yl)acetyl]phenyl]carbamate (6e). Yield 1.71 g (81%); M_r 371.39; Mp 178–180 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 10.09 (s, 1H, NHCO), 7.97 (d, 2H, Ar-H, $J = 8.4$ Hz), 7.59 (d, 2H, Ar-H, $J = 8.4$ Hz), 7.17–6.91 (m, 4H, Ar-H), 3.84 (s, 2H, COCH₂N), 3.70 (s, 3H, COOCH₃), 3.05–2.95 (m, 4H, 3,5-piperazine), 2.75–2.60 (m, 4H, 2,6-piperazine); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 195.13, 154.85 (d, $J = 242.8$ Hz), 153.66, 143.63, 139.70 (d, $J = 8.4$ Hz), 129.92, 129.39, 124.62 (d, $J = 3.0$ Hz), 122.10 (d, $J = 7.6$ Hz), 119.14 (d, $J = 2.3$ Hz), 117.14, 115.74 (d, $J = 20.5$ Hz), 63.32, 52.56, 51.73, 49.89 (d, $J = 3.1$ Hz). HR-MS: for C₂₀H₂₂O₃FN₃ [M – H]⁺ calculated 370.15615 m/z , found 370.15787 m/z .

Ethyl 4-[(4-(2-fluorophenyl)piperazin-1-yl)acetyl]phenyl]carbamate (6f). Yield 1.82 g (86%); M_r 385.44; Mp 172–174 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 10.05 (s, 1H, NHCO), 7.96 (d, 2H, Ar-H, $J = 8.4$ Hz), 7.59 (d, 2H, Ar-H, $J = 8.4$ Hz), 7.17–6.91 (m, 4H, Ar-H), 4.13 (q, 2H, CH₂CH₃, $J = 7.0$ Hz), 3.82 (s, 2H, COCH₂N), 3.05–2.95 (m, 4H, 3,5-piperazine), 2.75–2.60 (m, 4H, 2,6-piperazine), 1.25 (t, 3H, CH₂CH₃, $J = 7.0$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 195.27, 154.85 (d, $J = 242.8$ Hz), 153.21, 143.69, 139.73 (d, $J = 8.4$ Hz), 129.92, 129.36, 124.62 (d, $J = 3.0$ Hz), 122.13 (d, $J = 7.6$ Hz), 119.14 (d, $J = 2.3$ Hz), 117.13, 115.76 (d, $J = 20.5$ Hz), 63.38, 60.39, 52.58, 49.94 (d, $J = 3.1$ Hz), 14.25. HR-MS: for C₂₁H₂₄O₃FN₃ [M – H]⁺ calculated 384.17180 m/z , found 384.17360 m/z .

Propyl 4-[(4-(2-fluorophenyl)piperazin-1-yl)acetyl]phenyl]carbamate (6g). Yield 1.71 g (75%); M_r 399.47; Mp 140–142 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 10.06 (s, 1H, NHCO), 7.96 (d, 2H, Ar-H, $J = 8.4$ Hz), 7.59 (d, 2H, Ar-H, $J = 8.4$ Hz), 7.17–6.91 (m, 4H, Ar-H), 4.06 (t, 2H, CH₂CH₂CH₃, $J = 6.6$ Hz), 3.83 (s, 2H, COCH₂N), 3.05–2.95 (m, 4H, 3,5-piperazine), 2.75–2.60 (m, 4H, 2,6-piperazine), 1.75–1.57 (m, 2H, CH₂CH₂CH₃), 0.93 (t, 3H, CH₂CH₂CH₃, $J = 7.3$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 195.28, 154.87 (d, $J = 242.8$ Hz), 153.33, 143.72, 139.75 (d, $J = 8.4$ Hz), 129.92, 129.37, 124.64 (d, $J = 3.0$ Hz), 122.08 (d, $J = 7.6$ Hz), 119.16 (d, $J = 2.3$ Hz), 117.14, 115.74 (d, $J = 20.5$ Hz), 65.93, 63.40, 52.59, 49.98 (d, $J = 3.1$ Hz), 21.69, 10.05. HR-MS: for C₂₂H₂₆O₃FN₃ [M – H]⁺ calculated 398.18745 m/z , found 398.18917 m/z .

Butyl 4-[(4-(2-fluorophenyl)piperazin-1-yl)acetyl]phenyl]carbamate (6h). Yield 2.23 g (96%); M_r 413.50; Mp 154–156 °C; $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 10.06 (s, 1H, NHCO), 7.96 (d, 2H, Ar-H, $J = 8.4$ Hz), 7.59 (d, 2H, Ar-H, $J = 8.4$ Hz), 7.16–6.91 (m, 4H, Ar-H), 4.10 (t, 2H, CH₂CH₂CH₂CH₃, $J = 6.6$ Hz), 3.82 (s, 2H, COCH₂N), 3.05–2.95 (m, 4H, 3,5-piperazine), 2.75–2.60 (m, 4H, 2,6-piperazine), 1.69–1.55 (m, 2H, CH₂CH₂CH₂CH₃), 1.46–1.30 (m, 2H, CH₂CH₂CH₂CH₃), 0.91 (t, 3H, CH₂CH₂CH₂CH₃, $J = 7.3$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_C (ppm): 195.33, 154.90 (d, $J = 242.8$ Hz), 153.36, 143.74, 139.78 (d, $J = 8.4$ Hz), 129.95, 129.40, 124.67 (d, $J = 3.0$ Hz), 122.13 (d, $J = 7.6$ Hz), 119.17 (d, $J = 2.3$ Hz), 117.17, 115.77 (d, $J = 20.5$ Hz), 64.16, 63.42, 52.61, 50.00 (d, $J = 3.1$ Hz), 30.39, 18.46, 13.40. HR-MS: for C₂₃H₂₈O₃FN₃ [M – H]⁺ calculated 412.20310 m/z , found 412.20490 m/z .

3.2.4. General Procedure For the Preparation of Alkyl {3-/4-[2-(4-(2-Fluorophenyl)piperazin-1-yl)-1-hydroxyethyl]phenyl}carbamates (7a–h)

The synthesized alkyl {3-/4-[2-(4-(2-fluorophenyl)piperazin-1-yl)acetyl]phenyl}carbamates, i.e., **6a**, **6e** (1.49 g, 4.0 mmol), **6b**, **6f** (1.54 g, 4.0 mmol), **6c**, **6g** (1.60 g, 4.0 mmol), **6d** or **6h** (1.45 g, 4.0 mmol), were dissolved in hot methanol (50 mL) and solid NaBH₄ (0.30 g, 8.0 mmol) was added in small portions [32]. The mixtures were refluxed for 1 h. When the reaction was completed (TLC control), the solvent was evaporated in vacuo, residua were treated with 100 mL of distilled water and 100 mL of chloroform. The organic layer was washed with distilled water, dried over anhydrous sodium sulphate and evaporated *in vacuo* to give crude solid products **7a–h**, which were crystallized from acetone. Full characterization parameters for the compounds **7a–h** (Scheme 2), isolated as colourless solids, are given below.

Methyl 3-[2-(4-(2-fluorophenyl)piperazin-1-yl)-1-hydroxyethyl]phenyl]carbamate (7a). Yield 1.20 g (83%); M_r 373.41; Mp 103–105 °C; IR (ATR, cm⁻¹): 3382 (ν NH), 2951 (ν_{as} CH₂), 2816 (ν_s CH₂), 1726 (ν C = O), 1553 (δ NH), 1497 (ν CN), 1229 (ν_{as} COC), 1079 (ν_s CO), 1020 (δ_{ip} = C–H), 850 (δ_{oop} = C–H); $^1\text{H-NMR}$ (DMSO- d_6) δ_H (ppm): 9.61 (s, 1H, NHCO), 7.48 (s, 1H, Ar–H), 7.35 (d, 1H, Ar–H, $J = 8.1$ Hz), 7.22 (t,

1H, Ar-H, $J = 8.1$ Hz), 7.16–6.90 (m, 5H, Ar-H), 5.06 (d, 1H, CHOH , $J = 3.7$ Hz), 4.73–4.65 (m, 1H, OH), 3.66 (s, 3H, COOCH_3), 3.10–2.95 (m, 4H, 3,5-piperazine), 2.75–2.60 (m, 4H, 2,6-piperazine), 2.55–2.38 (m, 2H, $\text{CH(OH)CH}_2\text{N}$); $^{13}\text{C-NMR}$ ($\text{DMSO-}d_6$) δ_{C} (ppm): 154.83 (d, $J = 242.8$ Hz), 153.88, 145.20, 139.80 (d, $J = 8.4$ Hz), 138.73, 128.05, 124.61 (d, $J = 3.0$ Hz), 121.97 (d, $J = 7.6$ Hz), 120.08, 119.02 (d, $J = 2.3$ Hz), 116.76, 116.00, 115.72 (d, $J = 20.5$ Hz), 69.76, 66.10, 53.02, 51.34, 50.00 (d, $J = 3.1$ Hz); ESI-MS: for $\text{C}_{20}\text{H}_{24}\text{O}_3\text{FN}_3$ $[\text{M} + \text{H}]^+$ calculated 373.42138 m/z , found 373.42095 m/z .

Ethyl {3-[2-(4-(2-fluorophenyl)piperazin-1-yl)-1-hydroxyethyl]phenyl}carbamate (**7b**). Yield 1.30 g (82%); M_{r} 387.46; Mp 118–120 °C; IR (ATR, cm^{-1}): 3463 (ν NH), 2937 (ν_{as} CH_2), 2833 (ν_{s} CH_2), 1703 (ν C = O), 1548 (δ NH), 1498 (ν CN), 1240 (ν_{as} COC), 1082 (ν_{s} CO), 1018 (δ_{ip} = C–H), 852 (δ_{oop} = C–H); $^1\text{H-NMR}$ ($\text{DMSO-}d_6$) δ_{H} (ppm): 9.57 (s, 1H, NHCO), 7.50 (s, 1H, Ar–H), 7.34 (d, 1H, Ar–H, $J = 8.1$ Hz), 7.21 (t, 1H, Ar–H, $J = 8.1$ Hz), 7.13–6.91 (m, 5H, Ar–H), 5.05 (d, 1H, CHOH , $J = 3.7$ Hz), 4.73–4.65 (m, 1H, OH), 4.12 (q, 2H, CH_2CH_3 , $J = 7.1$ Hz), 3.10–2.95 (m, 4H, 3,5-piperazine), 2.75–2.60 (m, 4H, 2,6-piperazine), 2.52–2.37 (m, 2H, $\text{CH(OH)CH}_2\text{N}$), 1.24 (t, 3H, CH_2CH_3 , $J = 7.0$ Hz); $^{13}\text{C-NMR}$ ($\text{DMSO-}d_6$) δ_{C} (ppm): 154.83 (d, $J = 242.8$ Hz), 153.44, 145.18, 139.82 (d, $J = 8.4$ Hz), 138.82, 128.04, 124.64 (d, $J = 3.0$ Hz), 121.98 (d, $J = 7.6$ Hz), 120.01, 119.02 (d, $J = 2.3$ Hz), 116.78, 115.99, 115.76 (d, $J = 20.5$ Hz), 69.79, 66.11, 59.86, 53.03, 50.01 (d, $J = 3.1$ Hz), 14.39; ESI-MS: for $\text{C}_{21}\text{H}_{26}\text{O}_3\text{FN}_3$ $[\text{M} + \text{H}]^+$ calculated 387.44796 m/z , found 387.44805 m/z .

Propyl {3-[2-(4-(2-fluorophenyl)piperazin-1-yl)-1-hydroxyethyl]phenyl}carbamate (**7c**). Yield 1.30 g (80%); M_{r} 401.49; Mp 103–105 °C; IR (ATR, cm^{-1}): 3262 (ν NH), 2942 (ν_{as} CH_2), 2829 (ν_{s} CH_2), 1726 (ν C = O), 1545 (δ NH), 1497 (ν CN), 1226 (ν_{as} COC), 1081 (ν_{s} CO), 1025 (δ_{ip} = C–H), 848 (δ_{oop} = C–H); $^1\text{H-NMR}$ ($\text{DMSO-}d_6$) δ_{H} (ppm): 9.58 (s, 1H, NHCO), 7.51 (s, 1H, Ar–H), 7.34 (d, 1H, Ar–H, $J = 8.1$ Hz), 7.21 (t, 1H, Ar–H, $J = 8.1$ Hz), 7.13–6.91 (m, 5H, Ar–H), 5.05 (d, 1H, CHOH , $J = 3.7$ Hz), 4.73–4.65 (m, 1H, OH), 4.03 (t, 2H, $\text{CH}_2\text{CH}_2\text{CH}_3$, $J = 6.6$ Hz), 3.07–2.93 (m, 4H, 3,5-piperazine), 2.73–2.59 (m, 4H, 2,6-piperazine), 2.55–2.38 (m, 2H, $\text{CH(OH)CH}_2\text{N}$), 1.73–1.55 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_3$), 0.93 (t, 3H, $\text{CH}_2\text{CH}_2\text{CH}_3$, $J = 7.3$ Hz); $^{13}\text{C-NMR}$ ($\text{DMSO-}d_6$) δ_{C} (ppm): 154.83 (d, $J = 242.8$ Hz), 153.57, 145.23, 139.85 (d, $J = 8.4$ Hz), 138.87, 128.08, 124.67 (d, $J = 3.0$ Hz), 122.03 (d, $J = 7.6$ Hz), 120.04, 119.06 (d, $J = 2.3$ Hz), 116.81, 116.02, 115.77 (d, $J = 20.5$ Hz), 69.80, 66.16, 65.46, 53.07, 50.05 (d, $J = 3.1$ Hz), 21.82, 10.13; ESI-MS: for $\text{C}_{22}\text{H}_{28}\text{O}_3\text{FN}_3$ $[\text{M} + \text{H}]^+$ calculated 401.47454 m/z , found 401.47421 m/z .

Butyl {3-[2-(4-(2-fluorophenyl)piperazin-1-yl)-1-hydroxyethyl]phenyl}carbamate (**7d**). Yield 1.50 g (90%); M_{r} 415.52; Mp 107–110 °C; IR (ATR, cm^{-1}): 3294 (ν NH), 2940 (ν_{as} CH_2), 2830 (ν_{s} CH_2), 1706 (ν C = O), 1541 (δ NH), 1497 (ν CN), 1228 (ν_{as} COC), 1082 (ν_{s} CO), 1018 (δ_{ip} = C–H), 845 (δ_{oop} = C–H); $^1\text{H-NMR}$ ($\text{DMSO-}d_6$) δ_{H} (ppm): 9.57 (s, 1H, NHCO), 7.51 (s, 1H, Ar–H), 7.33 (d, 1H, Ar–H, $J = 8.1$ Hz), 7.20 (t, 1H, Ar–H, $J = 8.1$ Hz), 7.13–6.91 (m, 5H, Ar–H), 5.04 (d, 1H, CHOH , $J = 3.7$ Hz), 4.73–4.65 (m, 1H, OH), 4.07 (t, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $J = 6.6$ Hz), 3.10–2.95 (m, 4H, 3,5-piperazine), 2.75–2.60 (m, 4H, 2,6-piperazine), 2.53–2.37 (m, 2H, $\text{CH(OH)CH}_2\text{N}$), 1.67–1.53 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.47–1.29 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 0.92 (t, 3H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $J = 7.3$ Hz); $^{13}\text{C-NMR}$ ($\text{DMSO-}d_6$) δ_{C} (ppm): 154.85 (d, $J = 242.8$ Hz), 153.56, 145.21, 139.83 (d, $J = 8.4$ Hz), 138.85, 128.07, 124.65 (d, $J = 3.0$ Hz), 122.01 (d, $J = 7.6$ Hz), 120.02, 119.03 (d, $J = 2.3$ Hz), 116.78, 115.97, 115.77 (d, $J = 20.5$ Hz), 69.80, 66.14, 63.64, 53.07, 50.04 (d, $J = 3.1$ Hz), 30.52, 18.49, 13.45; ESI-MS: for $\text{C}_{23}\text{H}_{30}\text{O}_3\text{FN}_3$ $[\text{M} + \text{H}]^+$ calculated 415.50112 m/z , found 415.50207 m/z .

Methyl {4-[2-(4-(2-fluorophenyl)piperazin-1-yl)-1-hydroxyethyl]phenyl}carbamate (**7e**). Yield 1.42 g (94%); M_{r} 373.41; Mp 143–146 °C; IR (ATR, cm^{-1}): 3340 (ν NH), 2975 (ν_{as} CH_2), 2808 (ν_{s} CH_2), 1702 (ν C = O), 1552 (δ NH), 1502 (ν CN), 1234 (ν_{as} COC), 1063 (ν_{s} CO), 1021 (δ_{ip} = C–H), 855 (δ_{oop} = C–H); $^1\text{H-NMR}$ ($\text{DMSO-}d_6$) δ_{H} (ppm): 9.59 (s, 1H, NHCO), 7.38 (d, 2H, Ar–H, $J = 8.4$ Hz), 7.25 (d, 2H, Ar–H, $J = 8.4$ Hz), 7.16–6.90 (m, 4H, Ar–H), 4.96 (d, 1H, CHOH , $J = 3.7$ Hz), 4.75–4.63 (m, 1H, OH), 3.64 (s, 3H, COOCH_3), 3.12–2.90 (m, 4H, 3,5-piperazine), 2.75–2.55 (m, 4H, 2,6-piperazine), 2.55–2.35 (m, 2H, $\text{CH(OH)CH}_2\text{N}$); $^{13}\text{C-NMR}$ ($\text{DMSO-}d_6$) δ_{C} (ppm): 154.85 (d, $J = 242.8$ Hz), 153.92, 139.82 (d, $J = 8.4$ Hz), 138.55, 137.69, 126.26, 124.64 (d, $J = 3.0$ Hz), 121.98 (d, $J = 7.6$ Hz), 119.05 (d, $J = 2.3$ Hz), 117.87, 115.74 (d, $J = 20.5$ Hz),

69.36, 66.04, 53.03, 51.37, 50.05 (d, $J = 3.1$ Hz); ESI-MS: for $C_{20}H_{24}O_3FN_3$ $[M + H]^+$ calculated 373.42138 m/z , found 373.42142 m/z .

Ethyl {4-[2-(4-(2-fluorophenyl)piperazin-1-yl)-1-hydroxyethyl]phenyl}carbamate (**7f**). Yield 1.40 g (88%); M_r 387.46; Mp 142–145 °C; IR (ATR, cm^{-1}): 3340 (ν NH), 2977 (ν_{as} CH_2), 2808 (ν_s CH_2), 1711 (ν C = O), 1549 (δ NH), 1503 (ν CN), 1234 (ν_{as} COC), 1060 (ν_s CO), 1018 (δ_{ip} = C–H), 860 (δ_{oop} = C–H); 1H -NMR (DMSO- d_6) δ_H (ppm): 9.54 (s, 1H, NHCO), 7.38 (d, 2H, Ar–H, $J = 8.4$ Hz), 7.22 (d, 2H, Ar–H, $J = 8.4$ Hz), 7.15–6.91 (m, 4H, Ar–H), 4.96 (d, 1H, $\underline{C}HOH$, $J = 3.7$ Hz), 4.71–4.61 (m, 1H, OH), 4.08 (q, 2H, $\underline{C}H_2CH_3$, $J = 7.0$ Hz), 3.06–2.91 (m, 4H, 3,5-piperazine), 2.71–2.54 (m, 4H, 2,6-piperazine), 2.52–2.34 (m, 2H, $\underline{C}H(OH)\underline{C}H_2N$), 1.21 (t, 3H, $\underline{C}H_2\underline{C}H_3$, $J = 7.0$ Hz); ^{13}C -NMR (DMSO- d_6) δ_C (ppm): 154.85 (d, $J = 242.8$ Hz), 153.47, 139.82 (d, $J = 8.4$ Hz), 138.46, 137.76, 126.23, 124.63 (d, $J = 3.0$ Hz), 121.97 (d, $J = 7.6$ Hz), 119.03 (d, $J = 2.3$ Hz), 117.82, 115.73 (d, $J = 20.5$ Hz), 69.36, 66.05, 59.88, 53.03, 50.07 (d, $J = 3.1$ Hz), 14.39; ESI-MS: for $C_{21}H_{26}O_3FN_3$ $[M + H]^+$ calculated 387.44796 m/z , found 387.44822 m/z .

Propyl {4-[2-(4-(2-fluorophenyl)piperazin-1-yl)-1-hydroxyethyl]phenyl}carbamate (**7g**). Yield 1.53 g (92%); M_r 401.49; Mp 149–152 °C; IR (ATR, cm^{-1}): 3339 (ν NH), 2966 (ν_{as} CH_2), 2808 (ν_s CH_2), 1698 (ν C = O), 1550 (δ NH), 1505 (ν CN), 1235 (ν_{as} COC), 1073 (ν_s CO), 1015 (δ_{ip} = C–H), 857 (δ_{oop} = C–H); 1H -NMR (DMSO- d_6) δ_H (ppm): 9.56 (s, 1H, NHCO), 7.39 (d, 2H, Ar–H, $J = 8.4$ Hz), 7.24 (d, 2H, Ar–H, $J = 8.4$ Hz), 7.16–6.89 (m, 4H, Ar–H), 4.98 (d, 1H, $\underline{C}HOH$, $J = 3.7$ Hz), 4.73–4.61 (m, 1H, OH), 4.01 (t, 2H, $\underline{C}H_2CH_2CH_3$, $J = 6.6$ Hz), 3.08–2.91 (m, 4H, 3,5-piperazine), 2.68–2.54 (m, 4H, 2,6-piperazine), 2.54–2.33 (m, 2H, $\underline{C}H(OH)\underline{C}H_2N$), 1.71–1.52 (m, 2H, $\underline{C}H_2CH_2CH_3$), 0.92 (t, 3H, $\underline{C}H_2CH_2CH_3$, $J = 7.3$ Hz); ^{13}C -NMR (DMSO- d_6) δ_C (ppm): 154.86 (d, $J = 242.8$ Hz), 153.60, 139.82 (d, $J = 8.4$ Hz), 138.46, 137.79, 126.24, 124.64 (d, $J = 3.0$ Hz), 122.00 (d, $J = 7.6$ Hz), 119.05 (d, $J = 2.3$ Hz), 117.84, 115.75 (d, $J = 20.5$ Hz), 69.38, 66.05, 65.48, 53.05, 50.06 (d, $J = 3.1$ Hz), 21.79, 10.11; ESI-MS: for $C_{22}H_{28}O_3FN_3$ $[M + H]^+$ calculated 401.47454 m/z , found 401.47510 m/z .

Butyl {4-[2-(4-(2-fluorophenyl)piperazin-1-yl)-1-hydroxyethyl]phenyl}carbamate (**7h**). Yield 1.62 g (94%); M_r 415.52; Mp 140–143 °C; IR (ATR, cm^{-1}): 3336 (ν NH), 2954 (ν_{as} CH_2), 2808 (ν_s CH_2), 1698 (ν C = O), 1549 (δ NH), 1527 (ν CN), 1228 (ν_{as} COC), 1076 (ν_s CO), 1018 (δ_{ip} = C–H), 852 (δ_{oop} = C–H); 1H -NMR (DMSO- d_6) δ_H (ppm): 9.56 (s, 1H, NHCO), 7.40 (d, 2H, Ar–H, $J = 8.4$ Hz), 7.25 (d, 2H, Ar–H, $J = 8.4$ Hz), 7.16–6.89 (m, 4H, Ar–H), 4.99 (d, 1H, $\underline{C}HOH$, $J = 3.7$ Hz), 4.72–4.59 (m, 1H, OH), 4.06 (t, 2H, $\underline{C}H_2CH_2CH_2CH_3$, $J = 6.6$ Hz), 3.09–2.94 (m, 4H, 3,5-piperazine), 2.64–2.55 (m, 4H, 2,6-piperazine), 2.55–2.36 (m, 2H, $\underline{C}H(OH)\underline{C}H_2N$), 1.65–1.52 (m, 2H, $\underline{C}H_2CH_2CH_2CH_3$), 1.46–1.28 (m, 2H, $\underline{C}H_2CH_2CH_2CH_3$), 0.91 (t, 3H, $\underline{C}H_2CH_2CH_2CH_3$, $J = 7.3$ Hz); ^{13}C -NMR (DMSO- d_6) δ_C (ppm): 154.87 (d, $J = 242.8$ Hz), 153.62, 139.84 (d, $J = 8.4$ Hz), 138.47, 137.81, 126.26, 124.66 (d, $J = 3.0$ Hz), 122.02 (d, $J = 7.6$ Hz), 119.07 (d, $J = 2.3$ Hz), 117.86, 115.76 (d, $J = 20.5$ Hz), 69.38, 66.07, 63.67, 53.05, 50.06 (d, $J = 3.1$ Hz), 30.52, 18.49, 13.43; ESI-MS: for $C_{23}H_{30}O_3FN_3$ $[M + H]^+$ calculated 415.50112 m/z , found 415.50154 m/z .

3.2.5. General Procedure For the Preparation of 1-(2-(3-/4-[(Alkoxy carbonyl)amino]phenyl)-2-hydroxyethyl)-4-(2-fluorophenyl)piperazin-1-ium Chlorides (**8a–h**)

The solution of a particular alkyl {3-/4-[2-(4-(2-fluorophenyl)piperazin-1-yl)-1-hydroxyethyl]phenyl}carbamate, i.e. **7a**, **7e** (0.71 g, 1.9 mmol), **7b**, **7f** (0.74 g, 1.9 mmol), **7c**, **7g** (0.76 g, 1.9 mmol), **7d** or **7h** (0.79 g, 1.9 mmol), in 40 mL of chloroform was treated with a saturated solution of hydrogen chloride in diethyl ether and stirred for 5 h at laboratory temperature. The solvents were removed *in vacuo* and solid crude products **8a–h** were crystallized from acetone. Full characterization parameters for the target compounds **8a–h** (Scheme 2), isolated as colourless solids, are provided below.

1-(2-Hydroxy-2-(3-[(methoxycarbonyl)amino]phenyl)ethyl)-4-(2-fluorophenyl)piperazin-1-ium chloride (8a). Yield 0.75 g (88%); M_r 409.89; Mp 139–141 °C; R_f 0.40; IR (ATR, cm^{-1}): 3266 (ν NH), 2952 (ν_{as} CH_2), 2815 (ν_s CH_2), 1718 (ν C = O), 1612 (ν C = C), 1552 (δ NH), 1491 (ν CN), 1233 (ν_{as} COC), 1081 (ν_s CO), 1020

($\delta_{ip} = C-H$), 848 ($\delta_{oop} = C-H$); 1H -NMR (DMSO- d_6) δ_H (ppm): 10.71 (m, 1H, NH⁺), 9.60 (s, 1H, NHCO), 7.51 (s, 1H, Ar-H), 7.44 (t, 1H, $J = 8.0$ Hz), 7.35 (d, 1H, Ar-H, $J = 8.0$ Hz), 7.25 (t, 1H, Ar-H, $J = 8.0$ Hz), 7.30–7.15 (m, 5H, Ar-H), 5.25 (dd, 1H, CHOH, $J = 4.1$ Hz, $J = 9.0$ Hz), 4.70–4.65 (m, 1H, OH), 3.65 (s, 3H, COOCH₃), 3.65–3.55 (m, 4H, 3,5-piperazine), 4.10–3.25 (m, 6H, CH(OH)CH₂N + 2,6-piperazine); ^{13}C -NMR (DMSO- d_6) δ_C (ppm): 153.25 (d, $J = 249.1$ Hz), 152.91, 144.68, 139.37 (d, $J = 8.3$ Hz), 137.21, 128.16, 124.35 (d, $J = 3.1$ Hz), 121.42 (d, $J = 7.6$ Hz), 120.15, 118.46 (d, $J = 2.2$ Hz), 116.24, 116.07, 115.11 (d, $J = 20.4$ Hz), 69.55, 66.14, 53.34, 51.26, 50.07 (d, $J = 3.0$ Hz); ESI-MS: for C₂₀H₂₅O₃FN₃ [M + H]⁺ calculated 374.42932 m/z , found 374.42873 m/z . Anal. Calcd. for C₂₀H₂₅O₃ClFN₃ (409.89): C, 58.61%; H, 6.15%; N, 10.25%. Found: C, 58.65%; H, 6.18%; N, 10.15%.

1-(2-{3-[(Ethoxycarbonyl)amino]phenyl}-2-hydroxyethyl)-4-(2-fluorophenyl)piperazin-1-ium chloride (**8b**). Yield 0.80 g (91%); M_r 423.91; Mp 117–119 °C; R_f 0.53; IR (ATR, cm⁻¹): 3425 (ν NH), 2983 (ν_{as} CH₂), 2852 (ν_s CH₂), 1718 (ν C = O), 1610 (ν C = C), 1547 (δ NH), 1496 (ν CN), 1229 (ν_{as} COC), 1068 (ν_s CO), 1018 ($\delta_{ip} = C-H$), 852 ($\delta_{oop} = C-H$); 1H -NMR (DMSO- d_6) δ_H (ppm): 10.73 (m, 1H, NH⁺), 9.57 (s, 1H, NHCO), 7.50 (s, 1H, Ar-H), 7.45 (t, 1H, Ar-H, $J = 8.0$ Hz), 7.35 (d, 1H, Ar-H, $J = 8.1$ Hz), 7.27–7.15 (m, 5H, Ar-H), 5.24 (dd, 1H, CHOH, $J = 4.1$ Hz, $J = 9.0$ Hz), 4.73–4.65 (m, 1H, OH), 4.20 (q, 2H, CH₂CH₃, $J = 7.0$ Hz), 3.60–3.50 (m, 4H, 3,5-piperazine), 4.00–3.25 (m, 6H, CH(OH)CH₂N + 2,6-piperazine), 1.30 (t, 3H, CH₂CH₃, $J = 7.0$ Hz); ^{13}C -NMR (DMSO- d_6) δ_C (ppm): 154.81 (d, $J = 241.4$ Hz), 153.47, 146.78, 139.94 (d, $J = 8.4$ Hz), 138.57, 128.16, 124.02 (d, $J = 3.0$ Hz), 121.64 (d, $J = 7.5$ Hz), 119.78, 119.07 (d, $J = 2.3$ Hz), 116.35, 116.28, 115.01 (d, $J = 20.4$ Hz), 69.74, 65.89, 59.34, 53.07, 50.12 (d, $J = 3.1$ Hz), 14.21; ESI-MS: for C₂₁H₂₇O₃FN₃ [M + H]⁺ calculated 388.45590 m/z , found 388.45615 m/z . Anal. Calcd. for C₂₁H₂₇O₃ClFN₃ (423.91): C, 59.50%; H, 6.42%; N, 9.91%. Found: C, 59.33%; H, 6.31%; N, 10.06%.

1-(2-Hydroxy-2-{3-[(propoxycarbonyl)amino]phenyl}ethyl)-4-(2-fluorophenyl)piperazin-1-ium chloride (**8c**). Yield 0.82 g (91%); M_r 437.94; Mp 99–101 °C; R_f 0.66; IR (ATR, cm⁻¹): 3402 (ν NH), 2964 (ν_{as} CH₂), 2837 (ν_s CH₂), 1724 (ν C = O), 1612 (ν C = C), 1546 (δ NH), 1498 (ν CN), 1224 (ν_{as} COC), 1072 (ν_s CO), 1022 ($\delta_{ip} = C-H$), 850 ($\delta_{oop} = C-H$); 1H -NMR (DMSO- d_6) δ_H (ppm): 10.78 (m, 1H, NH⁺), 9.58 (s, 1H, NHCO), 7.50 (s, 1H, Ar-H), 7.44 (t, 1H, Ar-H, $J = 8.0$ Hz), 7.35 (d, 1H, Ar-H, $J = 8.0$ Hz), 7.20 (t, 1H, Ar-H, $J = 8.1$ Hz), 7.30–7.18 (m, 5H, Ar-H), 5.24 (dd, 1H, CHOH, $J = 4.0$ Hz, $J = 9.0$ Hz), 4.73–4.65 (m, 1H, OH), 4.12 (t, 2H, CH₂CH₂CH₃, $J = 7.0$ Hz), 3.60–3.45 (m, 4H, 3,5-piperazine), 4.00–3.42 (m, 6H, CH(OH)CH₂N + 2,6-piperazine), 1.69–1.57 (m, 2H, CH₂CH₂CH₃), 0.95 (t, 3H, CH₂CH₂CH₃, $J = 7.0$ Hz); ^{13}C -NMR (DMSO- d_6) δ_C (ppm): 154.80 (d, $J = 242.7$ Hz), 153.78, 145.15, 138.19 (d, $J = 8.4$ Hz), 137.45, 128.49, 124.02 (d, $J = 3.0$ Hz), 122.09 (d, $J = 7.7$ Hz), 119.97, 119.03 (d, $J = 2.3$ Hz), 116.76, 116.08, 115.29 (d, $J = 20.5$ Hz), 69.54, 66.17, 65.76, 53.27, 50.03 (d, $J = 3.1$ Hz), 21.81, 10.15; ESI-MS: for C₂₂H₂₉O₃FN₃ [M + H]⁺ calculated 402.48248 m/z , found 402.48197 m/z . Anal. Calcd. for C₂₂H₂₉O₃ClFN₃ (437.94): C, 60.34%; H, 6.67%; N, 9.59%. Found: C, 60.12%; H, 6.81%; N, 9.37%.

1-(2-{3-[(Butoxycarbonyl)amino]phenyl}-2-hydroxyethyl)-4-(2-fluorophenyl)piperazin-1-ium chloride (**8d**). Yield 0.88 g (95%); M_r 451.97; Mp 98–100 °C; R_f 0.73; IR (ATR, cm⁻¹): 3416 (ν NH), 2959 (ν_{as} CH₂), 2837 (ν_s CH₂), 1730 (ν C = O), 1604 (ν C = C), 1543 (δ NH), 1495 (ν CN), 1221 (ν_{as} COC), 1074 (ν_s CO), 1016 ($\delta_{ip} = C-H$), 840 ($\delta_{oop} = C-H$); 1H -NMR (DMSO- d_6) δ_H (ppm): 10.78 (m, 1H, NH⁺), 9.57 (s, 1H, NHCO), 7.50 (s, 1H, Ar-H), 7.44 (t, 1H, Ar-H, $J = 8.1$ Hz), 7.34 (d, 1H, Ar-H, $J = 8.1$ Hz), 7.25–7.18 (m, 5H, Ar-H), 5.25 (dd, 1H, CHOH, $J = 4.0$ Hz, $J = 9.0$ Hz), 4.73–4.65 (m, 1H, OH), 4.18 (t, 2H, CH₂CH₂CH₂CH₃, $J = 7.0$ Hz), 3.60–3.45 (m, 4H, 3,5-piperazine), 4.00–3.40 (m, 6H, CH(OH)CH₂N + 2,6-piperazine), 1.67–1.58 (m, 2H, CH₂CH₂CH₂CH₃, $J = 7.0$), 1.40–1.30 (m, 2H, CH₂CH₂CH₂CH₃), 0.93 (t, 3H, CH₂CH₂CH₂CH₃, $J = 7.0$ Hz); ^{13}C -NMR (DMSO- d_6) δ_C (ppm): 154.90 (d, $J = 241.4$ Hz), 153.13, 145.61, 139.70 (d, $J = 8.5$ Hz), 138.62, 128.17, 124.34 (d, $J = 3.1$ Hz), 122.14 (d, $J = 7.6$ Hz), 120.07, 119.26 (d, $J = 2.3$ Hz), 116.35, 115.82, 115.61 (d, $J = 20.4$ Hz), 69.81, 66.38, 63.94, 53.23, 50.17 (d, $J = 3.1$ Hz), 30.60, 18.26, 13.33; ESI-MS: for C₂₃H₃₁O₃FN₃ [M + H]⁺ calculated 416.50906 m/z , found 416.50915 m/z . Anal. Calcd. for C₂₃H₃₁O₃ClFN₃ (451.97): C, 61.12%; H, 6.91%; N, 9.30%. Found: C, 61.02%; H, 6.85%; N, 9.42%.

1-(2-Hydroxy-2-{4-[(methoxycarbonyl)amino]phenyl}ethyl)-4-(2-fluorophenyl)piperazin-1-ium chloride (**8e**). Yield 0.78 g (84%); M_r 409.89; Mp 218–220 °C; R_f 0.31; IR (ATR, cm^{-1}): 3377 (ν NH), 2963 (ν_{as} CH_2), 2831 (ν_{s} CH_2), 1719 (ν C = O), 1610 (ν C = C), 1550 (δ NH), 1487 (ν CN), 1234 (ν_{as} COC), 1073 (ν_{s} CO), 1020 (δ_{ip} = C–H), 854 (δ_{oop} = C–H); $^1\text{H-NMR}$ (DMSO- d_6) δ_{H} (ppm): 10.42 (s, 1H, NH^+), 9.71 (s, 1H, NHCO), 7.47 (d, 2H, Ar–H, $J = 8.5$ Hz), 7.32 (d, 2H, Ar–H, $J = 8.5$ Hz), 7.23–7.00 (m, 4H, Ar–H), 6.20 (s, 1H, OH), 5.14 (dd, 1H, CH_2OH , $J = 4.8$ Hz, $J = 7.8$ Hz), 3.66 (s, 3H, COOCH_3), 3.75–3.11 (m, 10H, $\text{CH}(\text{OH})\text{CH}_2\text{N}$ + piperazine); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_{C} (ppm): 154.73 (d, $J = 244.6$ Hz), 153.91, 138.62, 138.25 (d, $J = 8.6$ Hz), 135.32, 126.41, 124.94 (d, $J = 3.1$ Hz), 123.27 (d, $J = 7.8$ Hz), 119.51 (d, $J = 2.2$ Hz), 117.93, 116.10 (d, $J = 20.3$ Hz), 66.15, 61.59, 52.32, 51.54, 50.30, 46.82, 46.51; ESI-MS: for $\text{C}_{20}\text{H}_{25}\text{O}_3\text{FN}_3$ $[\text{M} + \text{H}]^+$ calculated 374.42932 m/z , found 374.42951 m/z . Anal. Calcd. for $\text{C}_{20}\text{H}_{25}\text{O}_3\text{ClFN}_3$ (409.89): C, 58.61%; H, 6.15%; N, 10.25%. Found: C, 58.62%; H, 6.31%; N, 10.06%.

1-(2-{4-[(Ethoxycarbonyl)amino]phenyl}-2-hydroxyethyl)-4-(2-fluorophenyl)piperazin-1-ium chloride (**8f**). Yield 0.79 g (91%); M_r 423.91; Mp 208–210 °C; R_f 0.45; IR (ATR, cm^{-1}): 3354 (ν NH), 2980 (ν_{as} CH_2), 2851 (ν_{s} CH_2), 1719 (ν C = O), 1600 (ν C = C), 1552 (δ NH), 1519 (ν CN), 1232 (ν_{as} COC), 1079 (ν_{s} CO), 1016 (δ_{ip} = C–H), 858 (δ_{oop} = C–H); $^1\text{H-NMR}$ (DMSO- d_6) δ_{H} (ppm): 10.46 (m, 1H, NH^+), 9.67 (s, 1H, NHCO), 7.47 (d, 2H, Ar–H, $J = 8.5$ Hz), 7.32 (d, 2H, Ar–H, $J = 8.5$ Hz), 7.25–7.00 (m, 4H, Ar–H), 6.21 (s, 1H, OH), 5.14 (dd, 1H, CH_2OH , $J = 4.8$ Hz, $J = 7.8$ Hz), 4.12 (q, 2H, CH_2CH_3 , $J = 7.0$ Hz), 3.78–3.14 (m, 10H, $\text{CH}(\text{OH})\text{CH}_2\text{N}$ + piperazine), 1.24 (t, 3H, CH_2CH_3 , $J = 7.0$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_{C} (ppm): 154.74 (d, $J = 244.4$ Hz), 153.36, 138.78, 138.21 (d, $J = 8.7$ Hz), 135.20, 126.34, 124.89 (d, $J = 3.3$ Hz), 123.31 (d, $J = 8.0$ Hz), 119.45 (d, $J = 2.2$ Hz), 117.91, 116.11 (d, $J = 20.5$ Hz), 66.33, 61.74, 60.56, 52.34, 50.30, 46.76, 46.64, 14.38; ESI-MS: for $\text{C}_{21}\text{H}_{27}\text{O}_3\text{FN}_3$ $[\text{M} + \text{H}]^+$ calculated 388.45590 m/z , found 388.45541 m/z . Anal. Calcd. for $\text{C}_{21}\text{H}_{27}\text{O}_3\text{ClFN}_3$ (423.91): C, 59.50%; H, 6.42%; N, 9.91%. Found: C, 59.45%; H, 6.39%; N, 10.02%.

1-(2-Hydroxy-2-{4-[(propoxycarbonyl)amino]phenyl}ethyl)-4-(2-fluorophenyl)piperazin-1-ium chloride (**8g**). Yield 0.77 g (86%); M_r 437.94; Mp 212–214 °C; R_f 0.50; IR (ATR, cm^{-1}): 3392 (ν NH), 2973 (ν_{as} CH_2), 2847 (ν_{s} CH_2), 1720 (ν C = O), 1605 (ν C = C), 1543 (δ NH), 1501 (ν CN), 1231 (ν_{as} COC), 1079 (ν_{s} CO), 1012 (δ_{ip} = C–H), 855 (δ_{oop} = C–H); $^1\text{H-NMR}$ (DMSO- d_6) δ_{H} (ppm): 10.56 (m, 1H, NH^+), 9.68 (s, 1H, NHCO), 7.48 (d, 2H, Ar–H, $J = 8.5$ Hz), 7.33 (d, 2H, Ar–H, $J = 8.5$ Hz), 7.23–7.00 (m, 4H, Ar–H), 6.20 (s, 1H, OH), 5.15 (dd, 1H, CH_2OH , $J = 4.8$ Hz, $J = 7.9$ Hz), 4.03 (t, 2H, $\text{CH}_2\text{CH}_2\text{CH}_3$, $J = 6.8$ Hz), 3.79–3.16 (m, 10H, $\text{CH}(\text{OH})\text{CH}_2\text{N}$ + piperazine), 1.63 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_3$, $J = 7.1$ Hz), 0.93 (t, 3H, $\text{CH}_2\text{CH}_2\text{CH}_3$, $J = 7.3$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_{C} (ppm): 154.74 (d, $J = 244.7$ Hz), 153.45, 138.78, 138.21 (d, $J = 8.7$ Hz), 135.32, 126.42, 124.89 (d, $J = 3.3$ Hz), 123.31 (d, $J = 7.9$ Hz), 119.45 (d, $J = 2.2$ Hz), 117.91, 116.10 (d, $J = 20.2$ Hz), 66.29, 65.56, 61.77, 52.23, 50.42, 46.80, 46.64, 21.78, 10.21; ESI-MS: for $\text{C}_{22}\text{H}_{29}\text{O}_3\text{FN}_3$ $[\text{M} + \text{H}]^+$ calculated 402.48248 m/z , found 402.48217 m/z . Anal. Calcd. for $\text{C}_{22}\text{H}_{29}\text{O}_3\text{ClFN}_3$ (437.94): C, 60.34%; H, 6.67%; N, 9.59%. Found: C, 60.21%; H, 6.75%; N, 9.43%.

1-(2-{4-[(Butoxycarbonyl)amino]phenyl}-2-hydroxyethyl)-4-(2-fluorophenyl)piperazin-1-ium chloride (**8h**). Yield 0.89 g (96%); M_r 451.97; Mp 219–220 °C; R_f 0.56; IR (ATR, cm^{-1}): 3361 (ν NH), 2957 (ν_{as} CH_2), 2825 (ν_{s} CH_2), 1722 (ν C = O), 1610 (ν C = C), 1546 (δ NH), 1499 (ν CN), 1232 (ν_{as} COC), 1075 (ν_{s} CO), 1020 (δ_{ip} = C–H), 852 (δ_{oop} = C–H); $^1\text{H-NMR}$ (DMSO- d_6) δ_{H} (ppm): 10.46 (m, 1H, NH^+), 9.67 (s, 1H, NHCO), 7.48 (d, 2H, Ar–H, $J = 8.5$ Hz), 7.33 (d, 2H, Ar–H, $J = 8.5$ Hz), 7.24–7.01 (m, 4H, Ar–H), 6.22 (s, 1H, OH), 5.14 (dd, 1H, CH_2OH , $J = 4.8$ Hz, $J = 7.8$ Hz), 4.07 (t, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $J = 6.6$ Hz), 3.78–3.16 (m, 10H, $\text{CH}(\text{OH})\text{CH}_2\text{N}$ + piperazine), 1.62 (q, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $J = 7.0$ Hz), 1.38 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $J = 7.4$ Hz), 0.92 (t, 3H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $J = 7.3$ Hz); $^{13}\text{C-NMR}$ (DMSO- d_6) δ_{C} (ppm): 154.74 (d, $J = 244.4$ Hz), 153.52, 138.81, 138.23 (d, $J = 8.6$ Hz), 135.35, 126.41, 124.89 (d, $J = 3.1$ Hz), 123.33 (d, $J = 7.3$ Hz), 119.50 (d, $J = 2.2$ Hz), 117.92, 116.10 (d, $J = 20.2$ Hz), 66.29, 63.78, 61.74, 52.34, 50.32, 46.82, 46.61, 30.52, 18.48, 13.47; ESI-MS: for $\text{C}_{23}\text{H}_{31}\text{O}_3\text{FN}_3$ $[\text{M} + \text{H}]^+$ calculated 416.50906 m/z , found 416.50934 m/z . Anal. Calcd. for $\text{C}_{23}\text{H}_{31}\text{O}_3\text{ClFN}_3$ (451.97): C, 61.12%; H, 6.91%; N, 9.30%. Found: C, 61.05%; H, 6.90%; N, 9.37%.

3.3. Lipophilicity Parameter Determination

Lipohydrophilic properties of the final compounds **8a–h** were characterized by the R_M (RP-TLC), $\log k$ and $\log k_w$ (RP-HPLC) parameters, respectively.

3.3.1. Reversed-Phase Thin-Layer Chromatography (RP-TLC)

The R_M values were calculated according to Equation (5) from the R_f parameters observed in a S_M mobile phase: hydrochloric acid ($c = 1.0$ M)/acetone (4:1, v/v):

$$R_M = \log (1/R_f - 1) \quad (5)$$

For the experiments, aluminium sheets pre-coated with silica gel 60 F₂₅₄ (0.25 mm thickness; Merck) were impregnated by a variously concentrated silicone oil in heptane, which ranged from 1% to 5%. The plates were separately spotted with 2 μ L of methanolic solutions of each compound ($c = 1$ mg/mL), starting points were 1 cm from a bottom edge of these plates. The chromatographic plates were developed in glass developing chambers saturated for 30 min by the S_M mobile phase. Development was carried out upon 15 cm from a starting line by an ascending technique [73,74]. After being developed, the plates were dried at room temperature. Detection of zones was performed under iodine vapours/UV light at $\lambda = 254$ nm. Each experiment was run in triplicate at 21 °C, the R_M values of analytes were calculated separately for each run.

Optimal differences in R_M values within both homological groups **8a–d** and **8e–h** were observed if 1% silicone oil in heptane was chosen. The R_f s were found in a range from 0.19 (**8h**) to 0.78 (**8a**; Table 1). All the calculated average R_M values were summarized in Table S1 (Supplementary Materials).

3.3.2. Reversed-Phase High-Performance Liquid Chromatography (RP-HPLC)

Methanol (MeOH)/water mobile phases with a various volume ratio of the organic modifier and water (60:40, 70:30, 80:20 and 85:15, respectively; v/v) were chosen. A concentration of all analyzed compounds **8a–h** was $c = 0.25$ mg/mL. An aqueous solution of potassium nitrate ($c = 0.1$ M) was used for the determination of dead time $t_0 = 1.295$ min. The capacity (retention) factor k values (Table 1) were calculated by Equation (6) as follows:

$$k = (t_r - t_0)/t_0 \quad (6)$$

where the t_r and t_0 parameters were the retention times of a solute (t_r) and unretained compound (potassium nitrate; t_0), respectively. The observed retention (t_r) and dead (t_0) times were means of three independent determinations [73,74].

The $\log k_w$ values, i.e., the logarithms of extrapolated capacity (retention) factors for 100% water in the isocratic RP-HPLC, were determined from intercepts of linear plots between the $\log k$ and φ_M (a volume fraction of an organic modifier in the isocratic elution RP-HPLC) according to Equation (7):

$$\log k = \log k_w - S \times \varphi_M \quad (7)$$

where the S parameter represented the slope of a regression curve, which was related to the solvent strength of a pure organic solvent [46,75].

3.4. Electronic Properties Determination

The $\log \epsilon$ values characterizing methanolic solutions of the analyzed compounds **8a–h** ($c = 3.0 \times 10^{-5}$ M) were estimated at $\lambda_1 = 208$ –210 nm, $\lambda_{2(\text{Ch-T})} = 238$ –240 nm and $\lambda_3 = 274$ –276 nm (Table 3), respectively, in a near ultraviolet (quartz) region of the electromagnetic spectrum between 200 and

400 nm [47]. The log ϵ values for the observed absorption maxima were calculated according to the Lambert-Beer's law discussed in [47], for example, and expressed by Equation (8):

$$A = \epsilon \times c \times l \quad (8)$$

where the A parameter represented the absorbance of a solution, the descriptor ϵ was a molar absorption coefficient (in the L/mol/cm units) and l was the path length (in the cm units).

3.5. Biological Assays

3.5.1. In Vitro Antimycobacterial Evaluation

Mycobacterial strains. The *in vitro* activity of compounds **8a–h** was inspected against *Mycobacterium tuberculosis* CNCTC My 331/88 (identical with H₃₇R_v and ATCC 2794, respectively; dilution of the strain was 10^{−3} M), *M. kansasii* CNCTC My 235/80 (identical with ATCC 12478; 10^{−4} M), the *M. kansasii* 6 509/96 clinical isolate (10^{−4} M) and *M. avium* CNCTC My 330/80 (identical with ATCC 25291; 10^{−5} M), respectively, in the Laboratory for Mycobacterial Diagnosis and Tuberculosis (Institute of Public Health in Ostrava, Czech Republic). These strains were purchased from the National Reference Laboratory – Czech National Collection of Type Cultures (CNCTC; The National Institute of Public Health, Prague, Czech Republic), excluding *M. kansasii* 6 509/96, which was clinically isolated because the INH-resistant *M. kansasii* strains have not been found in Czech Republic or Slovak Republic yet.

M. avium intracellulare ATCC 13950 as well as the *M. kansasii* CIT11/06, *M. avium* subsp. *paratuberculosis* CIT03 and *M. avium hominissuis* CIT10/08 clinical isolates, respectively, were obtained from the Department of Biological Sciences (Cork Institute of Technology, Bishopstown, Cork, Ireland).

Standard drugs. The isoniazid (INH), ethambutol (EMB), ofloxacin (OFLX), ciprofloxacin (CPX) and pyrazinamide (PZA) reference drugs were purchased from Sigma-Aldrich (Darmstadt, Germany), showing the purity of analytical grade.

Determination of a minimum inhibitory concentration (MIC) against *M. tuberculosis* CNCTC My 331/88, *M. kansasii* CNCTC My 235/80, *M. kansasii* 6 509/96 and *M. avium* CNCTC My 330/80. Efficiency of the compounds **8a–h** and standard drugs against given mycobacteria were determined in the Šula semisynthetic medium (Sevac, Prague, Czech Republic) by a dilution-micromethod [48,76].

In a brief, each mycobacterial strain was simultaneously inoculated into Petri plates containing a Löwenstein-Jensen medium for sterility control and growth of inoculum. All inspected compounds were added to the medium as solutions in dimethyl sulfoxide (DMSO; Sigma-Aldrich, Irvine, UK). In the assays, following concentrations of the solutions were used: 1000, 500, 250, 125, 62.5, 32, 16, 8, 4, 2, 1, 0.5, 0.25 and 0.125 μ M, respectively. The inoculated plates kept in microtone bags were incubated at 37 °C. Particular reading was carried out on a stand with a bottom magnifying mirror, macroscopically, with the use of a magnifying glass. The growth in the plates was evaluated after 7, 14 and 21 days (*M. kansasii* CNCTC My 235/80, *M. kansasii* 6 509/96) and after 14 and 21 days (*M. tuberculosis* CNCTC My 331/88, *M. avium* CNCTC My 330/80), respectively [48,76].

The value of a minimum inhibitory concentration (MIC) was the lowest concentration (on the above concentration scale) of a tested compound, which inhibited growth of the mycobacteria [48,76]. The evaluation was repeated three times and the MIC values, reported in Table 4 in the μ M units, were the same.

Determination of a minimum inhibitory concentration (MIC) against *M. avium intracellulare* ATCC 13950, *M. kansasii* CIT11/06, *M. avium* subsp. *paratuberculosis* CIT03 and *M. avium hominissuis* CIT10/08. Those mycobacterial strains were grown in a Middlebrook broth (MB), supplemented with Oleic-Albumin-Dextrose-Catalase Supplement (OADC; Becton Dickinson, Oxford, UK). Identification of isolates was performed using biochemical and molecular protocols.

At log phase growth, a culture sample (10 mL) was centrifuged at 15,000 rpm/20 min using a bench top centrifuge Model CR 4-12 (Jouan Inc., London, UK). Following removal of a supernatant, the pellet was washed in a fresh Middlebrook 7H9GC broth (Difco, Detroit, MI, USA), and re-suspended

in a fresh supplemented MB (10 mL). Turbidity was adjusted to match the McFarland standard No. 1 containing 3×10^8 Colony Forming Units (CFU) with the MB broth. Further 1:20 dilution of a culture was performed in the broth.

Susceptibility of given mycobacterial strains was investigated in a 96-well plate format. In the experiments, sterile deionised water (300 μ L) was added to all outer-perimeter wells of plates to minimize evaporation of a medium in the test wells during the incubation process. Each tested compound (100 μ L) was incubated with each of mycobacterial species (100 μ L). The dilutions of tested compounds were prepared in triplicate, their final concentrations ranged from 500 μ g/mL to 15 μ g/mL. The screened derivatives were prepared in DMSO (Sigma-Aldrich, London, UK) and dilutions were made in the supplemented MB broth. The plates were sealed with a parafilm and incubated at 37 °C for 7 days. Following the incubation, 10% addition of a water-soluble dye, the alamarBlue reagent (AbD Serotec, Kidlington, UK) was mixed into each well. Absorbance readings at $\lambda = 570$ nm and 600 nm were taken, initially for a background subtraction and after 24 h re-incubation. The subtraction is necessary for strongly coloured compounds, where the colour may interfere with interpretation of any colour change. For non-interfering compounds, a blue colour in a well was interpreted as absence of growth and pink colour was scored as growth [17,49,50].

The MIC value was defined as the lowest concentration of a compound, at which no visible bacterial growth was observed. In other words, the MIC was the lowest concentration that prevented a visual colour change from blue to pink. The MIC for mentioned mycobacteria was defined as 90% or greater growth reduction (IC_{90}) compared to a control. The MIC (IC_{90}) parameter has been routinely and widely used in bacterial assays as a standard detection limit according to the Clinical and Laboratory Standards Institute [49,50].

Clinically used antimycobacterial drugs INH, CPX and PZA, respectively, were applied as standards. The observed MIC values were listed in Table 5 in the μ M units.

3.5.2. *In Vitro* Antiproliferative (Cytotoxicity) Screening

Human monocytic leukemia THP-1 cells were obtained from the European Collection of Cell Cultures (ECACC; Salisbury, UK; Methods of characterization: DNA Fingerprinting (Multilocus probes) and isoenzyme analysis). The cells were routinely cultured in the RPMI 1640 medium (Lonza, Verviers, Belgium) supplemented with 10% fetal bovine serum (Sigma-Aldrich, Darmstadt, Germany), 2% L-glutamine, 1% penicillin and streptomycin (Lonza, Verviers, Belgium) at 37 °C with 5% CO₂. The cells were passaged at approximately 1-week intervals and were routinely tested for absence of mycoplasma by a Hoechst 33258 staining method.

The tested compounds **8a–h** were dissolved in DMSO (Sigma-Aldrich, Darmstadt, Germany) and added in five increasing concentrations to the cell suspension in the culture medium. The maximum concentration of DMSO in the assays never exceeded 0.1%. Subsequently, the cells were incubated for 24 h at 37 °C with 5% CO₂ at various compounds' concentrations varying from 0.37 μ M to 30 μ M in the RPMI 1640 medium.

Cell toxicity was determined using the Cytotoxicity Detection KitPLUS Lactate Dehydrogenase (LDH) assay kit (Roche Diagnostics, Mannheim, Germany) and used according to manufacturer's instructions. For the LDH assays, the cells were seeded into 96-well plates (5×10^4 cells/well in 100 μ L culture medium) in triplicate in the serum-free RPMI 1640 medium, and measurements at $\lambda = 492$ nm (Synergy 2 Multi-Mode Microplate Reader; BioTek, Winooski, VT, USA) were taken 24 h after treatment with the tested compounds [77,78].

Median lethal dose values, LD_{50} , were deduced through the construction of a dose-response curve. All values were evaluated using the GraphPad Prism 5.00 software (GraphPad Software, San Diego, CA, USA).

3.6. Calculations and Statistical Analyses

Regression equations and statistical characteristics were calculated and visualized by the Origin Pro 9.0.0 software (OriginLab Corporation, Northampton, MA, USA).

In a current research, those statistical parameters were calculated: Residual sum of squares (*RSS*), correlation coefficient (*R*), adjusted coefficient of determination (*Adj. R*²), root mean squared error (standard deviation; *RMSE*) and norm of residuals (*NoR*), respectively. The study provided Analysis of Variance (ANOVA) outputs as well, i.e., Fisher's *F*-test (Fisher's significance ratio; *F*) and probability of obtaining the *F* Ratio (*Prob > F*).

The *RSS* descriptor was used to measure amount of variance in a data set that was not explained by the regression model. The *RSS* parameter was a measure of amount of error remaining between the regression function and data set. Relatively smaller *RSS* explained greater amount of data [79,80].

The *R* parameter was based on a method of covariance. The coefficient was used to measure the strength of a relationship between two (continuous) variables. The *Adj. R*² value penalized the *R*² data for addition of regressors, which did not contribute to explanatory power of a model. Indeed, the *Adj. R*² value was never larger than the *R*² one; it would be decreased by the adding of other regressors, and might be even negative for poorly fitting models [81–83].

The *RMSE* was an unambiguous indicator of error for numerical predictions. The parameter provided standard deviation of a model prediction error. Relatively smaller value indicated better model performance [84]. The *NoR* parameter was the square root of *RMSE* and was used as measure for goodness of fit when comparing different fits [85].

The *F* value, as a parameter obtained by the ANOVA, could determine whether the means of three or more groups were different. The ANOVA approach tested an effect of a categorical predictor variable on a continuous dependent variable and used the *F*-test to statistically test equality of means [79].

The *Prob > F* data provided a *p*-value for the test and measured probability of obtaining the *F* Ratio as large as what was observed, given that all parameters except the intercept were zero [81,82].

Indication of significance level of the *F* Ratio by stars was as follows: one star, i.e., statistically significant relationship defined by the *Prob > F* value in the range from 0.0100 to <0.0500; two stars, i.e., statistically very significant model defined by the *Prob > F* parameter in the interval from 0.0010 to <0.0100; and three stars, i.e., statistically extremely significant relationship with the *Prob > F* value in the interval from 0 to <0.0010. Finally, a statistically insignificant model was connected with the *Prob > F* parameter ≥ 0.0500 [82].

4. Conclusions

In summary, original *N*-arylpiperazines **8a–h** were prepared by multistep procedures and characterized by spectral values (¹H-NMR, ¹³C-NMR, IR and ESI-MS) and elemental analyses (% C, H, N). The compounds contained a flexible 3-/4-alkoxycarbonylamino group (alkoxy = methoxy to butoxy), 2-hydroxyethane-1,2-diyl connecting chain and 4-(2-fluorophenyl)piperazin-1-yl moiety (Table 1). These fragments have been separately found in a chemical structure of various compounds with a notable *in vitro* efficiency against some tuberculous strains of mycobacteria.

Lipohydrophilic properties of the molecules **8a–h** were preliminary evaluated by the RP-TLC using silica gel plates impregnated with 1% silicone oil in heptane. Calculated *R*_M values of 3-alkoxycarbonylamino substituent-containing compounds (**8a–d**) ranged from −0.55 (**8a**) to 0.01 (**8d**), the 4-substituted ones (**8e–h**) showed higher *R*_M parameters varying from −0.02 (**8e**) to 0.63 (**8h**).

Linearly extrapolated logarithms of retention factors corresponding to 100% water as a mobile phase, log *k*_w values (RP-HPLC), characterized the lipohydrophilic properties of the molecules **8a–h** (Table 2) more reliably and precisely than any arbitrary selected isocratic log *k* parameters. These log *k*_w values were in accordance with elution order and hydrophobicity of **8a–h** and ranged from 2.113 (**8e**) to 2.930 (**8h**). The derivatives **8a–c** showed higher log *k*_w values (2.430–2.796) than the molecules **8e–g** (2.113–2.600). The compounds containing the longest side chain, i.e., 1-(2-{3-[(butoxycarbonyl)amino]phenyl}-2-hydroxyethyl)-4-(2-fluorophenyl)piperazin-1-ium chloride

(8d) and 1-(2-{4-[(butoxycarbonyl)amino]phenyl}-2-hydroxyethyl)-4-(2-fluorophenyl)-piperazin-1-ium chloride (8h) were found to be the most lipophilic, as proven by their $\log k_w$ of 2.796 (8d) and 2.930 (8h), respectively (Table 2).

Regarding the extrapolation calculations, the slopes S of regression lines varied from 2.7386 (8e) to 3.3441 (8h; Table 2). The S parameter was related to a specific hydrophobic surface of a particular compound and could be used as the alternative measure of its lipophilicity.

Electronic properties of the inspected compounds 8a–h were characterized by logarithms of molar absorption coefficients ($\log \epsilon$) of their methanolic solutions ($c = 3.0 \times 10^{-5}$ M) investigated in the UV/Vis region of a spectrum.

The solutions showed three absorption maxima in a near ultraviolet (quartz) region of the electromagnetic spectrum, e.g., $\lambda_1 = 208\text{--}210$ nm, $\lambda_{2(\text{Ch-T})} = 238\text{--}240$ nm and $\lambda_3 = 274\text{--}276$ nm, respectively (Table 3). The $\log \epsilon_{2(\text{Ch-T})}$ parameters of the compounds 8a–d observed at a charge-transfer absorption maximum $\lambda_{2(\text{Ch-T})}$ were found in a narrow interval from 4.30 (8a) to 4.37 (8c). The methanolic solutions of 8e–h were characterized by higher $\log \epsilon_{2(\text{Ch-T})}$ values than the ones of 8a–d and these parameters ranged from 4.42 (8h) to 4.67 (8e; Table 3). In addition, elongation of a 4-side chain led to lower $\log \epsilon$ values related to all observed absorption maxima (Table 3).

The racemic compounds 8a–h were *in vitro* screened against *Mycobacterium tuberculosis* CNCTC My 331/88 (identical with H₃₇R_v and ATCC 2794), *M. kansasii* CNCTC My 235/80 (identical with ATCC 12478), a *M. kansasii* 6 509/96 clinical isolate, *M. avium* CNCTC My 330/80 (identical with ATCC 25291) and *M. avium intracellulare* ATCC 13950 as well as against *M. kansasii* CIT11/06, *M. avium* subsp. *paratuberculosis* CIT03 and *M. avium hominissuis* CIT10/08 clinical isolates, respectively (Tables 4 and 5). The biological evaluation revealed the most promising potential of the 8a–h set against *M. tuberculosis*, *M. kansasii* My 235/80 and *M. kansasii* 6 509/96, respectively.

A position and length of a side chain notably affected the activity of presently investigated *N*-arylpiperazine derivatives against *M. tuberculosis* CNCTC My 331/88. The 4-positional isomers were more effective, with the MIC values ranging from 8 μM (8h) to 125 μM (8e), than the 3-positional ones, which possessed the MICs from 32 μM (8d) to 250 μM (8a). Among all *in vitro* screened molecules, the INH standard was found to be the most active with the MIC = 0.5 μM (14-d/21-d; Table 4).

The compounds 8a–d were more efficient against *M. kansasii* My 235/80 and *M. kansasii* 6 509/96 than the derivatives 8e–h. The most active molecule against given mycobacteria was 8d with the MIC = 16 μM and 62.5 μM , respectively, depending on a particular strain and also on the number of days of incubation. Increase in length of the side chain resulted in lower MIC values of 8a–d against these mycobacterial strains. The observed MIC parameters were, however, higher compared to the ones related to EMB with the MIC = 1 μM and 2 μM (14-d/21-d), or OFLX, which showed the MIC = 0.5 μM and 1 μM , respectively (14-d/21-d; Table 4).

The activity of 8a–h against a non-tuberculous INH-resistant *M. avium* CNCTC My 330/80 was apparently dependent on the position of an alkoxy-carbonylamino chain. Its presence in the 3-position (8a–d) led to the MIC values varying from 62.5 μM (8d) to 500 μM (8a; 14-d/21-d). However, a potential of the 4-substituent-containing derivatives (8e–h) to fight given *mycobacterium* was insufficient (MIC > 250 μM ; Table 4).

The efficiency of the most active substance 8d (MIC = 62.5 μM ; 14-d/21-d) against *M. avium* CNCTC My 330/80 was comparable to the effectiveness of OFLX (MIC = 32 μM and 62.5 μM , respectively; 14-d/21-d); reference EMB drug was moderately more active (MIC = 16 μM ; 14-d/21-d). Elongation of a 3-side chain led to more promising compounds (Table 4).

Regarding current SAR studies, lipophilic properties represented by extrapolated $\log k_w$ values seemed to be considerably more important for the *in vitro* activity of the 8a–d set against *M. tuberculosis* and *M. kansasii* 6 509/96 compared to the lipophilic features of the compounds 8e–h.

The $\log \epsilon_{2(\text{Ch-T})}$ values (Table 3) observed at the charge-transfer absorption maximum $\lambda_{2(\text{Ch-T})}$ were also taken into a special consideration, because they could be the most sensitive to the differences in a position and electronic properties of the alkoxy-carbonylamino substituent.

A relationship between the $\log \varepsilon_{2(\text{Ch-T})}$ parameters and $\log (1/\text{MIC} [\text{M}])$ values connected with the *in vitro* screening of **8a–h** against *M. tuberculosis* My 331/88 (14-d) provided a bilinear course. Maximal efficiency of these compounds could be observed if their $\log \varepsilon_{2(\text{Ch-T})}$ values were approximately 4.43 (Figure 4).

If attention was paid to the *M. kansasii* My 235/80 and *M. kansasii* 6 509/96 strains, no significant relationships between the *in vitro* activity of the compounds **8a–h** and their electronic features were observed.

Favorable cytotoxicity profiles of the molecules **8a–h** were proved by the LD_{50} values $> 30 \mu\text{M}$, which were estimated on a human monocytic leukemia THP-1 cell line. Moreover, the least lipophilic methoxy group-containing derivatives **8e** ($\log k_w = 2.113$) and **8a** ($\log k_w = 2.430$) increased proliferation of the THP-1 cells in 24 h when compared to a control.

Overall, the results of current *in vitro* biological evaluation and initial SAR investigations of the molecules **8a–h** considered them very promising candidates for further structural optimization, which could lead to even more effective antimycobacterials. Regarding these findings, the authors of the study were inspired and encouraged to synthesize enantiomerically pure compounds and explore their *in vitro* efficiency, especially against *M. tuberculosis* CNCTC My 331/88, *M. kansasii* My 235/80 or *M. kansasii* 6 509/96, in further phases of the research programme

Supplementary Materials: The supplementary materials are available online at www.mdpi.com/link.

Acknowledgments: The authors very gratefully acknowledge a financial support received especially from the Faculty of Pharmacy, Comenius University in Bratislava (Slovak Republic). The research was also supported by the grant projects VEGA 1/0873/15, KEGA 022UK-4/2015 and Science Foundation Ireland Project Ref: 12/R1/2335. Part of the experiments was carried out in the Toxicological and Antidoping Center at the Faculty of Pharmacy, Comenius University in Bratislava (Slovak Republic) and this support was also very acknowledged. The research was also partially supported by Sanofi-Aventis Pharma Slovakia, s.r.o. (Slovak Republic).

Author Contributions: T.G. synthesized and spectrally characterized the compounds; I.M. spectrally characterized the compounds, investigated their lipophilic and electronic properties, analyzed the data related to the *in vitro* antimycobacterial investigation, created the concept and designed the study, investigated, statistically characterized and interpreted the SAR results, wrote and revised the paper; J.Cs. designed the chemical structure of compounds under the study, contributed reagents/materials tools; J.J. and I.S. conceived the *in vitro* antimycobacterial screening of synthesized compounds, analyzed data related to the *in vitro* antimycobacterial evaluation of the molecules; J.S., J.O.M. and A.C. performed the *in vitro* antimycobacterial evaluation of the compounds and interpreted the results, contributed reagents/materials tools; A.C. revised the paper; P.M. investigated spectral and lipophilic properties of the synthesized compounds, contributed reagents/materials tools; S.K. and P.K. performed the *in vitro* antiproliferative (cytotoxic) activity of the synthesized molecules, contributed reagents/materials tools. The authors have approved the final version of manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

The following abbreviations are used in this manuscript:

7-d	7-Day incubation
14-d/21-d	14-/21-Day incubation
φ_M	Volume fraction of a mobile phase modifier (RP-HPLC)
CPX	Ciprofloxacin
<i>Adj. R</i> ²	Adjusted coefficient of determination (statistical analysis)
DMSO	Dimethyl sulfoxide
EMB	Ethambutol
<i>F</i>	Fisher's <i>F</i> -test (Fisher's significance ratio; statistical analysis)
INH	Isoniazid
<i>k</i>	Capacity (retention) factor (RP-HPLC)
$\log k_w$	Lipophilicity index; values extrapolated from intercepts of a linear relationship between the logarithm of retention factor <i>k</i> ($\log k$) and volume fraction of a mobile phase modifier (φ_M ; RP-HPLC)
MDR	Multi-drug resistant

MeOH	Methanol
MA	<i>Mycobacterium avium</i>
MIC	Minimum inhibitory concentration (in the μM units)
MK	<i>Mycobacterium kansasii</i>
Mp/mp	Melting point
MT	<i>Mycobacterium tuberculosis</i>
NoR	Norm of residuals (statistical analysis)
OFLX	Ofloxacin
$Prob > F$	Probability of obtaining the F Ratio (statistical analysis)
PZA	Pyrazinamide
R_M	Lipophilicity index (RP-TLC)
RMSE	Root mean squared error (statistical analysis)
RSS	Residual sum of squares (statistical analysis)
S	Slope (RP-HPLC)
SAR	Structure–activity relationship(s)
t_r	Retention time of a compound (RP-HPLC)

References

- Evans, B.E.; Rittle, K.E.; Bock, M.G.; DiPardo, R.M.; Freidinger, R.M.; Whitter, W.L.; Lundell, G.F.; Veber, D.F.; Anderson, P.S.; Chang, R.S.L.; et al. Methods for drug discovery: Development of potent, selective, orally effective cholecystokinin antagonists. *J. Med. Chem.* **1988**, *31*, 2235–2246. [[CrossRef](#)] [[PubMed](#)]
- Shaquiquzzaman, M.; Verma, G.; Marella, A.; Akhter, M.; Akhtar, W.; Khan, M.F.; Tasneem, S.; Alam, M.M. Piperazine scaffold: A remarkable tool in generation of diverse pharmacological agents. *Eur. J. Med. Chem.* **2015**, *102*, 487–529. [[CrossRef](#)] [[PubMed](#)]
- Bobesh, K.A.; Renuka, J.; Srilakshmi, R.R.; Yellanki, S.; Kulkarni, P.; Yogeewari, P.; Sriram, D. Replacement of cardiotoxic aminopiperidine linker with piperazine moiety reduces cardiotoxicity? *Mycobacterium tuberculosis* novel bacterial topoisomerase inhibitors. *Bioorg. Med. Chem.* **2016**, *24*, 42–52. [[CrossRef](#)] [[PubMed](#)]
- Xu, Z.; Zhang, S.; Feng, L.-S.; Li, X.-N.; Huang, G.-Ch.; Chai, Y.; Lv, Z.-S.; Guo, H.-Y.; Liu, M.-L. Synthesis and *in vitro* antimycobacterial and antibacterial activity of 8-OMe ciprofloxacin-hydrozone/azole hybrids. *Molecules* **2017**, *22*, 1171. [[CrossRef](#)] [[PubMed](#)]
- Kayukova, L.A.; Orazbaeva, M.A.; Bismilda, V.L.; Chingisova, L.T. Synthesis and antituberculosis activity of *O*-aroyl- β -(4-phenylpiperazin-1-yl)propioamidooximes. *Pharm. Chem. J.* **2010**, *44*, 17–20. [[CrossRef](#)]
- Keng Yoon, Y.; Ashraf Ali, M.; Choon, T.S.; Ismail, R.; Chee Wei, A.; Suresh Kumar, R.; Osman, H.; Beevi, F. Antituberculosis: Synthesis and antimycobacterial activity of novel benzimidazole derivatives. *Biomed. Res. Int.* **2013**. [[CrossRef](#)] [[PubMed](#)]
- Sriram, D.; Yogeewari, P.; Senthilkumar, P.; Sangaraju, D.; Nelli, R.; Banerjee, D.; Bhat, P.; Manjashetty, T.H. Synthesis and antimycobacterial evaluation of novel phthalazin-4-ylacetamides against log- and starved phase cultures. *Chem. Biol. Drug. Des.* **2010**, *75*, 381–391. [[CrossRef](#)] [[PubMed](#)]
- Malinka, W.; Świątek, P.; Śliwińska, M.; Szponar, B.; Gamian, A.; Karczmarzyk, Z.; Fruziński, A. Synthesis of novel isothiazolopyridines and their *in vitro* evaluation against *Mycobacterium* and *Propionibacterium acnes*. *Bioorg. Med. Chem.* **2013**, *21*, 5282–5291. [[CrossRef](#)] [[PubMed](#)]
- Bogatcheva, E.; Hanrahan, C.; Nikonenko, B.; Samala, R.; Chen, P.; Gearhart, J.; Barbosa, F.; Einck, L.; Nacy, C.A.; Protopopova, M. Identification of new diamine scaffolds with activity against *Mycobacterium tuberculosis*. *J. Med. Chem.* **2006**, *49*, 3045–3048. [[CrossRef](#)] [[PubMed](#)]
- Shepherd, R.G.; Baughn, C.; Cantrall, M.L.; Goodstein, B.; Thomas, J.P.; Wilkinson, R.G. Structure–activity studies leading to ethambutol, a new type of antituberculous compound. *Ann. N. Y. Acad. Sci.* **1966**, *135*, 686–710. [[CrossRef](#)] [[PubMed](#)]
- Lee, R.E.; Protopopova, M.; Crooks, E.; Slayden, R.A.; Terrot, M.; Barry, C.E., III. Combinatorial lead optimization of [1,2]-diamines based on ethambutol as potential antituberculosis preclinical candidates. *J. Comb. Chem.* **2003**, *5*, 172–187. [[CrossRef](#)] [[PubMed](#)]
- Stavrov, G.; Valcheva, V.; Philipova, I.; Doytchinova, I. Novel camphane-based anti-tuberculosis agents with nanomolar activity. *Eur. J. Med. Chem.* **2013**, *70*, 372–379. [[CrossRef](#)] [[PubMed](#)]

13. Petkova, Z.; Valcheva, V.; Momekov, G.; Petrov, P.; Dimitrov, V.; Doytchinova, I.; Stavrakov, G.; Stoyanova, M. Antimycobacterial activity of chiral aminoalcohols with camphane scaffold. *Eur. J. Med. Chem.* **2014**, *81*, 150–157. [[CrossRef](#)] [[PubMed](#)]
14. Ghosh, A.K.; Brindisi, M. Organic carbamates in drug design and medicinal chemistry. *J. Med. Chem.* **2015**, *58*, 2895–2940. [[CrossRef](#)] [[PubMed](#)]
15. Moraczewski, A.L.; Banaszynski, L.A.; From, A.M.; White, C.E.; Smith, B.D. Using hydrogen bonding to control carbamate C–N rotamer equilibria. *J. Org. Chem.* **1998**, *63*, 7258–7262. [[CrossRef](#)] [[PubMed](#)]
16. Kečkėšová, S.; Sedlářová, E.; Čižmárik, J.; Garaj, V.; Csöllei, J.; Mokřý, P.; Andriamainty, F.; Malík, I.; Kaustová, J. Antimycobacterial activity of novel derivatives of arylcarbonyloxyaminopropanols. *Čes. Slov. Farm.* **2009**, *58*, 203–207.
17. Tengler, J.; Kapustíková, I.; Peško, M.; Govender, R.; Keltošová, S.; Mokřý, P.; Kollár, P.; O'Mahony, J.; Coffey, A.; Král'ová, K.; et al. Synthesis and biological evaluation of 2-hydroxy-3-[(2-aryloxyethyl) amino]propyl-4-[(alkoxycarbonyl)amino]benzoates. *Sci. World J.* **2013**, *2013*. [[CrossRef](#)] [[PubMed](#)]
18. Maruniak, M.; Sedlářová, E.; Csöllei, J.; Kapustíková, I.; Mokřý, P.; Malík, I.; Havranová Sichrovská, L.; Stanzel, L. Study of physicochemical properties and antimycobacterial activity of phenylcarbamic acid derivatives. In *Advances in Pharmaceutical Chemistry*, 1st ed.; Sedlářová, E., Malík, I., Garaj, V., Maruniak, M., Eds.; KO and KA Company: Bratislava, Slovak Republic, 2016; pp. 68–76.
19. Waisser, K.; Dražková, K.; Čižmárik, J.; Kaustová, J. Antimycobacterial activity of basic ethylesters of alkoxy-substituted phenylcarbamic acids. *Folia Microbiol.* **2003**, *48*, 45–50. [[CrossRef](#)]
20. Waisser, K.; Dražková, K.; Čižmárik, J.; Kaustová, J. Antimycobacterial activity of piperidinylpropyl esters of alkoxy-substituted phenylcarbamic acids. *Folia Microbiol.* **2003**, *48*, 585–587. [[CrossRef](#)]
21. Waisser, K.; Dražková, K.; Čižmárik, J.; Kaustová, J. A new group of potential antituberculotics: Hydrochlorides of piperidinylalkyl esters of alkoxy-substituted phenylcarbamic acids. *Folia Microbiol.* **2004**, *49*, 265–268. [[CrossRef](#)]
22. Hansch, C.; Clayton, J.M. Lipophilic character and biological activity of drugs II. The parabolic case. *J. Pharm. Sci.* **1973**, *62*, 1–21. [[CrossRef](#)] [[PubMed](#)]
23. Balgavý, P.; Devínsky, F. Cut-off effects in biological activities of surfactants. *Adv. Colloid Interface Sci.* **1996**, *12*, 23–63. [[CrossRef](#)]
24. Waisser, K.; Dražková, K.; Čižmárik, J.; Kaustová, J. Influence of lipophilicity on the antimycobacterial activity of the hydrochlorides of piperidinyethyl esters of *ortho*-substituted phenylcarbamic acids. *Sci. Pharm.* **2004**, *72*, 43–49. [[CrossRef](#)]
25. Upadhayaya, R.S.; Kulkarni, G.M.; Vasireddy, N.R.; Vandavasi, J.K.; Dixit, S.S.; Sharma, V.; Chattopadhyaya, J. Design, synthesis and biological evaluation of novel triazole, urea and thiourea derivatives of quinoline against *Mycobacterium tuberculosis*. *Bioorg. Med. Chem.* **2009**, *13*, 4681–4692. [[CrossRef](#)] [[PubMed](#)]
26. Upadhayaya, R.S.; Vandavasi, J.K.; Kardile, R.A.; Lahore, S.V.; Dixit, S.S.; Deokar, H.S.; Shinde, P.D.; Sarmah, M.P.; Chattopadhyaya, J. Novel quinoline and naphthalene derivatives as potent antimycobacterial agents. *Eur. J. Med. Chem.* **2010**, *45*, 1854–1867. [[CrossRef](#)] [[PubMed](#)]
27. Parai, M.K.; Panda, G.; Chaturvedi, V.; Manju, Y.K.; Sinha, S. Thiophene containing triarylmethanes as antitubercular agents. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 289–292. [[CrossRef](#)] [[PubMed](#)]
28. Kettmann, V.; Csöllei, J.; Račanská, E.; Švec, P. Synthesis and structure–activity relationships of new β -adrenoreceptor antagonists. Evidence for the electrostatic requirements for β -adrenoreceptor antagonists. *Eur. J. Med. Chem.* **1991**, *26*, 843–851. [[CrossRef](#)]
29. Kiss, A.; Potor, A.; Hell, Z. Heterogeneous catalytic solvent-free synthesis of quinoline derivatives *via* the Friedländer Reaction. *Catal. Lett.* **2008**, *125*, 250–253. [[CrossRef](#)]
30. Broutin, P.-E.; Hilty, P.; Thomas, A.W. An efficient synthesis of *ortho*-*N*-Boc-arylmethyl ketone derivatives. *Tetrahedron Lett.* **2003**, *44*, 6429–6432. [[CrossRef](#)]
31. Kolosov, M.A.; Orlov, V.D. 5-Thiazolyl derivatives of 4-aryl-3,4-dihydropyrimidin-2(1H)-ones. *Chem. Heterocycl. Compd.* **2008**, *44*, 1418–1420. [[CrossRef](#)]
32. Hu, B.; Ellingboe, J.; Han, S.; Largis, E.; Lim, K.; Malamas, M.; Mulvey, R.; Niu, C.; Oliphant, A.; Pelletier, J.; et al. Novel (4-piperidin-1-yl)-phenyl sulfonamides as potent and selective human β_3 agonists. *Bioorg. Med. Chem.* **2001**, *9*, 2045–2059. [[CrossRef](#)]

33. Pan, Y.; Li, P.; Xie, S.; Tao, Y.; Chen, D.; Dai, M.; Hao, H.; Huang, L.; Wang, Y.; Wang, L.; et al. Synthesis, 3D-QSAR analysis and biological evaluation of quinoxaline 1,4-di-N-oxide derivatives as antituberculosis agents. *Bioorg. Med. Chem. Lett.* **2016**, *26*, 4146–4153. [[CrossRef](#)] [[PubMed](#)]
34. Pancholia, S.; Dhameliya, T.M.; Shah, P.; Jadhavar, P.S.; Sridevi, J.P.; Yogeshwari, P.; Sriram, D.; Chakraborti, A.K. Benzo[d]thiazol-2-yl(piperazin-1-yl)methanones as new anti-mycobacterial chemotypes: Design, synthesis, biological evaluation and 3D-QSAR studies. *Eur. J. Med. Chem.* **2016**, *116*, 187–199. [[CrossRef](#)] [[PubMed](#)]
35. Rajkhowa, S.; Deka, R.C. DFT Based QSAR/QSPR models in the development of novel anti-tuberculosis drugs targeting *Mycobacterium tuberculosis*. *Curr. Pharm. Des.* **2014**, *20*, 4455–4473. [[CrossRef](#)] [[PubMed](#)]
36. Joshi, S.D.; More, U.A.; Aminabhavi, T.M.; Badiger, A.M. Two- and three-dimensional QSAR studies on a set of antimycobacterial pyrroles: CoMFA, topomer CoMFA, and HQSAR. *Med. Chem. Res.* **2014**, *23*, 107–126. [[CrossRef](#)]
37. Pliška, V.; Testa, B.; van de Waterbeemd, H. Lipophilicity in drug action and toxicology. In *Methods and Principles of Medicinal Chemistry*; Mannhold, R., Kubinyi, H., Timmerman, H., Eds.; Wiley-VCh Publishers: Weinheim, Germany, 1996; Volume 4, pp. 1–6.
38. Ottaviani, M.F.; Leonardis, I.; Cappiello, A.; Cangiotti, M.; Mazzeo, R.; Truffelli, H.; Palma, P. Structural modifications and adsorption capability of C₁₈-silica/binary solvent interphases studied by EPR and RP-HPLC. *J. Colloid Interface Sci.* **2010**, *352*, 512–519. [[CrossRef](#)] [[PubMed](#)]
39. Snyder, L.R.; Dolan, J.W. Initial experiments in high-performance liquid chromatographic method development I. Use of a starting gradient run. *J. Chromatogr. A.* **1996**, *721*, 3–14. [[CrossRef](#)]
40. Du, Ch.M.; Valko, K.; Bevan, Ch.; Reynolds, D.; Abraham, M.H. Rapid method for estimating octanol–water partition coefficient ($\log P_{\text{oct}}$) from isocratic RP-HPLC and a hydrogen bond acidity term (A). *J. Liquid Chromatogr. Relat. Technol.* **2001**, *24*, 635–649. [[CrossRef](#)]
41. Terada, H. Determination of $\log P_{\text{oct}}$ by high-performance liquid chromatography, and its application in the study of Quantitative Structure–Activity Relationships. *Quant. Struct. Act. Relat.* **1986**, *5*, 81–88. [[CrossRef](#)]
42. Snyder, L.R.; Dolan, J.W.; Grant, J.R. Gradient elution in high-performance liquid chromatography: I. Theoretical basis for reversed-phase systems. *J. Chromatogr. A* **1979**, *165*, 3–30. [[CrossRef](#)]
43. Valkó, K.; Snyder, L.R.; Glajch, J.L. Retention in reversed-phase liquid chromatography as a function of mobile-phase composition. *J. Chromatogr. A* **1993**, *656*, 501–520. [[CrossRef](#)]
44. Soczewiński, E. Mechanistic molecular model of liquid–solid chromatography: Retention–eluent composition relationships. *J. Chromatogr. A* **2002**, *965*, 109–116. [[CrossRef](#)]
45. Vrakas, D.; Panderi, I.; Hadjipavlou-Litina, D.; Tsantili-Kakoulidou, A. Investigation of the relationships between $\log P$ and various chromatographic indices for a series of substituted coumarins. Evaluation of their similarity/dissimilarity using multivariate statistics. *QSAR Comb. Sci.* **2005**, *24*, 254–260. [[CrossRef](#)]
46. Sztanke, K.; Markowski, W.; Świeboda, R.; Polak, B. Lipophilicity of novel antitumour and analgesic active 8-aryl-2,6,7,8-tetrahydroimidazo[2,1-c][1,2,4]triazine-3,4-dione derivatives determined by reversed-phase HPLC and computational methods. *Eur. J. Med. Chem.* **2010**, *45*, 2644–2649. [[CrossRef](#)] [[PubMed](#)]
47. Yadav, L.D.S. Ultraviolet and visible spectroscopy. In *Organic Spectroscopy*; Yadav, L.D.S., Ed.; Springer: Amsterdam, The Netherlands, 2005; pp. 7–51.
48. Ferriz, J.M.; Vávrová, K.; Kunc, F.; Imramovský, A.; Stolaříková, J.; Vavříková, E.; Vinšová, J. Salicylanilide carbamates: Antitubercular agents active against multidrug-resistant *Mycobacterium tuberculosis* strains. *Bioorg. Med. Chem.* **2010**, *18*, 1054–1061. [[CrossRef](#)] [[PubMed](#)]
49. Clinical and Laboratory Standards Institute (CLSI). *Methods for Antimicrobial Susceptibility Testing of Anaerobic Bacteria; Approved Standard*, 8th ed.; CLSI Document M11-A8; CLSI: Wayne, NJ, USA, 2012; pp. 10–56.
50. Clinical and Laboratory Standards Institute (CLSI). *Performance Standards for Antimicrobial Susceptibility Testing*, 24th ed.; Informational Supplement M100-S24; CLSI: Wayne, NJ, USA, 2014; pp. 106–211.
51. Waisser, K.; Doležal, R.; Čižmárik, J.; Malík, I.; Kaustová, J. The potential antitubercotics of the series of 2-hydroxy-3-(4-phenylpiperazin-1-yl)-propylphenylcarbamates. *Folia Pharm. Univ. Carol.* **2007**, *35–36*, 45–48.
52. Doležal, M.; Zitko, J.; Kešetovičová, D.; Kuneš, J.; Svobodová, M. Substituted N-phenylpyrazine-2-carboxamides: Synthesis and antimycobacterial evaluation. *Molecules* **2009**, *14*, 4180–4189. [[CrossRef](#)] [[PubMed](#)]
53. Čižmárik, J.; Waisser, K.; Doležal, R. QSAR Study of antimicrobial activity of esters of substituted phenylcarbamic acid. *Acta Fac. Pharm. Univ. Comen.* **2008**, *55*, 90–95.

54. Timmins, G.S.; Deretic, V. Mechanisms of action of isoniazid. *Mol. Microbiol.* **2006**, *62*, 1220–1227. [[CrossRef](#)] [[PubMed](#)]
55. Forbes, M.; Kuck, N.A.; Peets, E.A. Mode of action of ethambutol. *J. Bacteriol.* **1962**, *84*, 1099–1103. [[PubMed](#)]
56. Jena, L.; Waghmare, P.; Kashikar, S.; Kumar, S.; Harinath, B.C. Computational approach to understanding the mechanism of action of isoniazid, an anti-TB drug. *Int. J. Mycobacteriol.* **2014**, *3*, 276–282. [[CrossRef](#)] [[PubMed](#)]
57. Kuck, N.A.; Peets, E.A.; Forbes, M. Mode of action of ethambutol on *Mycobacterium tuberculosis*, strain H₃₇R_v. *Am. Rev. Respir. Dis.* **1963**, *87*, 905–906. [[PubMed](#)]
58. Mikusová, K.; Slayden, R.A.; Besra, G.S.; Brennan, P.J. Biogenesis of the mycobacterial cell wall and the site of action of ethambutol. *Antimicrob. Agents Chemother.* **1995**, *39*, 2484–2489. [[CrossRef](#)] [[PubMed](#)]
59. Lata, M.; Sharma, D.; Kumar, B.; Deo, N.; Tiwari, P.K.; Bisht, D.; Venkatesan, K. Proteome analysis of ofloxacin and moxifloxacin induced *Mycobacterium tuberculosis* isolates by proteomic approach. *Protein Pept. Lett.* **2015**, *22*, 362–371. [[CrossRef](#)] [[PubMed](#)]
60. Aubry, A.; Pan, X.S.; Fisher, L.M.; Jarlier, V.; Cambau, E. *Mycobacterium tuberculosis* DNA gyrase: Interaction with quinolones and correlation with antimycobacterial drug activity. *Antimicrob. Agents Chemother.* **2004**, *48*, 1281–1288. [[CrossRef](#)] [[PubMed](#)]
61. Brennan, P.J. Structure, function, and biogenesis of the cell wall of *Mycobacterium tuberculosis*. *Tuberculosis* **2003**, *83*, 91–97. [[CrossRef](#)]
62. De Wijs, H.; Jollès, P. Cell walls of three strains of mycobacteria (*Mycobacterium phlei*, *Mycobacterium fortuitum* and *Mycobacterium kansasii*): Preparation, analysis and digestion by lysozymes of different origins. *Biochim. Biophys. Acta* **1964**, *83*, 326–332. [[CrossRef](#)]
63. Suffness, M.; Douros, J. Current status of the NCI plant and animal product program. *J. Nat. Prod.* **1982**, *45*, 1–14. [[CrossRef](#)] [[PubMed](#)]
64. Witek, S.; Bielawski, J.; Bielawska, A. Synthesis of *N*-(formylphenyl)- and *N*-(acetophenyl) derivatives of urea and carbamic acid. *J. Prakt. Chem.* **1979**, *321*, 804–812. [[CrossRef](#)]
65. Takeuchi, H.; Mastubara, E. Electrophilic aromatic *N*-substitution by ethoxycarbonylnitrenium ion generated from ethyl azidoformate in the presence of trifluoroacetic acid. *J. Chem. Soc. Perkin Trans.* **1984**, *1*, 981–985. [[CrossRef](#)]
66. Park, Ch.-H.; Givens, R.S. New photoactivated protecting groups. 6. *p*-Hydroxyphenacyl: A phototrigger for chemical and biochemical probes. *J. Am. Chem. Soc.* **1997**, *119*, 2453–2463. [[CrossRef](#)]
67. Basterfield, S.; Woods, E.L.; Wright, H.N. Studies in urethans. III. The preparation of various substituted urethans. *J. Am. Chem. Soc.* **1926**, *48*, 2371–2375. [[CrossRef](#)]
68. Smith Broadbent, H.; Chu, C.-Y. The carbethoxylation products of *p*-aminoacetophenone and *p*-dimethylaminoacetophenone. *J. Am. Chem. Soc.* **1953**, *75*, 226–227. [[CrossRef](#)]
69. Sigman, E.M.; Autrey, T.; Schuster, G.B. Arylnitrenes with singlet ground states: Photochemistry of acetyl-substituted aroyl and aryloxycarbonyl azides. *J. Am. Chem. Soc.* **1988**, *110*, 4297–4305. [[CrossRef](#)]
70. Vettorazzi, M.; Angelina, E.; Lima, S.; Gonec, T.; Otevrel, J.; Marvanova, P.; Padrtova, T.; Mokry, P.; Bobal, P.; Acosta, L.M.; et al. An integrative study to identify novel scaffolds for sphingosine kinase 1 inhibitors. *Eur. J. Med. Chem.* **2017**, *139*, 461–481. [[CrossRef](#)] [[PubMed](#)]
71. Bietti, G.; Cereda, E.; Donetti, A.; del Soldato, P.; Giachetti, A.; Micheletti, R. Guanidino-heterocycl-phenyl-amidines and Salts Thereof. U.S. Patent No. US4548944 A. Available online: <https://encrypted.google.com/patents/US4548944?cl=un> (accessed on 12 November 2017).
72. Rather, J.B.; Reid, E.E. The identification of acids. IV. Phenacyl esters. *J. Am. Chem. Soc.* **1919**, *41*, 75–83. [[CrossRef](#)]
73. Dross, K.; Rekker, R.F.; de Vries, G.; Mannhold, R. The lipophilic behaviour of organic compounds: 3. The search for interconnections between reversed-phase chromatographic data and log *P*_{f_{oct} values. *Quant. Struct. Act. Relat.* **1999**, *18*, 549–557. [[CrossRef](#)]}
74. Kulig, K.; Malawska, B. Estimation of the lipophilicity of antiarrhythmic and antihypertensive active 1-substituted pyrrolidin-2-one and pyrrolidine derivatives. *Biomed. Chromatogr.* **2003**, *17*, 318–324. [[CrossRef](#)] [[PubMed](#)]
75. Özden, S.; Atabey, D.; Yıldız, S.; Göker, H. Synthesis, potent anti-staphylococcal activity and QSARs of some novel 2-anilinobenzazoles. *Eur. J. Med. Chem.* **2008**, *43*, 1390–1402. [[CrossRef](#)] [[PubMed](#)]

76. Imramovsky, A.; Pesko, M.; Kralova, K.; Vejsova, M.; Stolarikova, J.; Vinsova, J.; Jampilek, J. Investigating spectrum of biological activity of 4- and 5-chloro-2-hydroxy-*N*-[2-(arylamino)-1-alkyl-2-oxoethyl]benz-amides. *Molecules* **2011**, *16*, 2414–2430. [[CrossRef](#)] [[PubMed](#)]
77. Gonec, T.; Kos, J.; Zadrazilova, I.; Pesko, M.; Govender, R.; Chambel, B.; Pereira, D.; Kollar, P.; Imramovsky, A.; O'Mahony, J.; et al. Antibacterial and herbicidal activity of ring-substituted 2-hydroxynaphthalene-1-carboxanilides. *Molecules* **2013**, *18*, 9397–9419. [[CrossRef](#)] [[PubMed](#)]
78. Gonec, T.; Kos, J.; Zadrazilova, I.; Pesko, M.; Keltosova, S.; Tengler, J.; Bobal, P.; Kollar, P.; Cizek, A.; Kralova, K.; et al. Antimycobacterial and herbicidal activity of ring-substituted 1-hydroxynaphthalene-2-carboxanilides. *Bioorg. Med. Chem.* **2013**, *21*, 6531–6541. [[CrossRef](#)] [[PubMed](#)]
79. Morgan, J.A.; Tatar, J.F. Calculation of the residual sum of squares for all possible regressions. *Technometrics* **1972**, *14*, 317–325. [[CrossRef](#)]
80. Cheng, B.; Tong, H. On residual sums of squares in non-parametric autoregression. *Stoch. Process. Their Appl.* **1983**, *48*, 157–174. [[CrossRef](#)]
81. Kubinyi, H. QSAR: Hansch Analysis and Related Approaches. In *Methods and Principles in Medicinal Chemistry*; Mannhold, R., Krogsgaard-Larsen, P., Timmerman, H., Eds.; Wiley-VCh Verlag: Weinheim, Germany, 1993; Volume 1, pp. 22–56.
82. Weisberg, S. Multiple Regression. In *Applied Linear Regression*, 3rd ed.; Weisberg, S., Ed.; Wiley-Interscience (John Wiley and Sons): Hoboken, NJ, USA, 2005; pp. 47–68. [[CrossRef](#)]
83. Nakagawa, S.; Schielzeth, H. General and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods Ecol. Evol.* **2013**, *4*, 133–142. [[CrossRef](#)]
84. Mevik, B.H.; Cederkvist, H.R. Mean squared error of prediction (MSEP) estimates for principal component regression (PCR) and partial least squares regression (PLSR). *J. Chemom.* **2004**, *18*, 422–429. [[CrossRef](#)]
85. Ying, X.; Yang, L.; Zha, H. A fast algorithm for multidimensional ellipsoid-specific fitting by minimizing a new defined vector norm of residuals using semidefinite programming. *IEEE Trans. Pattern Anal. Mach. Intell.* **2012**, *34*, 1856–1863. [[CrossRef](#)] [[PubMed](#)]

Sample Availability: Samples of the compounds **8a–h** are available from the authors Tomáš Goněc and Ivan Malík.



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).