

Synthesis, Antimicrobial and Antioxidant Activities of Novel series of Cinnamamide Derivatives having Morpholine Moiety

Gangadhara Seelolla, Prasad Cheera and Venkateswarlu Ponneri*

Department of Chemistry, Sri Venkateswara University, Tirupati 517 502, AP, India

Abstract

Structurally morpholine containing Cinnamamide derivatives have been synthesized in quantitative yields by the reaction of different substituted cinnamic acids with morpholine. The structures of the newly synthesized compounds were confirmed by their IR, LC-MS, ¹H & ¹³C-NMR spectral data. The synthesized compounds were evaluated for their antimicrobial activity against bacterial species *Bacillus subtilis* and *Escherichia coli* etc and in vitro antioxidant activity by employing DPPH, hydrogen peroxide, and nitric oxide radical scavenging assays. Among these prepared compounds 4i and 4j showed significant antioxidant, antibacterial as well as antifungal activity and these were found to be the most potent activity molecules when compared with that of standard drugs.

Keywords: Morpholine; Cinnamic acid; Amide; Antioxidant property; Antimicrobial activity

Introduction

Cinnamic acid, which is ubiquitous in cinnamon oil and many other balsams, is a naturally occurring aromatic fatty acid of low toxicity with a long history of human exposure [1]. Commercially, it is widely applied in the perfume industry because of its floral odor. In recent years, cinnamic acid derivatives have attracted much attention due to their antioxidative [2], antitumor [3] and antimicrobial properties [4]. Though some of the research papers on cinnamic acid derivatives were reported as antitubercular agents [5-8]. Among these cinnamic acid derivatives such as cinnamamides were reported to have many different biological activities such as anticancer, antimutagenic, antioxidant and seed-germination inhibitory effects [9-11] and some of the cinnamamide derivatives exhibit a variety of biological activities, for example as shown in Figure 1, the genus piperaceae molecules (A) have been widely studied, due to the biological properties of secondary metabolites from these plants [12-14]. The natural piperidine derivative (B) showed good insecticidal activity against the fall armyworm, *Spodoptera frugiperda* [15]. Compounds (C) represent a novel series of potent small molecule inhibitors that not only have excellent in vitro profiles but also have activity *in vivo* [16]. N-(2-hydroxy ethyl) cinnamamide (D) was found to possess good anticonvulsant activity and also low toxicity [17]. In addition, in the course of screening compounds for pharmacological action, it was observed that α -phenyl-N,N-diethyl cinnamamide potentiated Nembutal hypnosis in rats [18].

On the other hand, among the family of heterocyclic compounds,




Figure 1. Examples of molecule containing cinnamamide derivatives.

the heterocycles with N, S and O atoms have attracted the attention of chemical research due to their wide spectrum of biological activities. N and O atoms containing Morpholine has its unique position in heterocyclic chemistry, and its derivatives are gaining considerable importance due to diverse biological activities such as antimalarial [19], antibacterial [20], antimicrobial [21,22], antidepressant [23], antiproliferative [24], hypocholesterolemic [25], antibiotic, antileukemic [26,27] etc.

Many efforts to find and develop novel and more potent antioxidants without side effects, and effective antimicrobial derivatives become increasingly important in the food and the cosmetic industries, and in medicine. Because of the constrained availability of natural cinnamic acid analogues, the establishment of their biological activities is still limited. Therefore, we synthesized a series of morpholine containing cinnamamide derivatives and evaluated their antimicrobial effects and antioxidant properties.

Experimental Section

Chemistry

Melting points were determined in open capillaries on a Mel-Temp apparatus and are uncorrected. All the reactions were monitored by thin layer chromatography (TLC) on precoated silica gel 60 F254 (mesh); spots were visualized with UV light. Merck silica gel (60-120 mesh) was used for column chromatography. The IR spectra were recorded on a Perkin-Elmer BX1 FTIR Spectrophotometer using KBr pellets and the wave numbers were given in cm^{-1} . ¹H NMR (400 MHz), and ¹³C NMR (100 MHz) spectra were recorded on a Bruker AMX 400 MHz NMR spectrometer in $\text{CDCl}_3/\text{DMSO}-d_6$ solution using TMS as an internal standard. All chemical shifts were reported in δ (ppm) using TMS as an internal standard. The mass spectra were recorded on Agilent 1100 LC/MSD instrument with method API-ES at 70 eV.

*Corresponding author: Venkateswarlu Ponneri, Department of Chemistry, Sri Venkateswara University, Tirupati 517 502, AP, India, Tel: 91-9393600444, E-mail: chem.ponneri@gmail.com

Received October 19, 2014; Accepted December 09, 2014; Published December 10, 2014

Citation: Seelolla G, Cheera P, Ponneri V (2014) Synthesis, Antimicrobial and Antioxidant Activities of Novel series of Cinnamamide Derivatives having Morpholine Moiety. Med chem 4: 778-783. doi:10.4172/2161-0444.1000229

Copyright: © 2014 Seelolla G, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

The microanalyses were performed on a Perkin Elmer 240C elemental analyzer. The antioxidant property was carried out by using Shimadzu UV-2450s spectrophotometer.

General Procedure for the Synthesis of trans cinnamides (4a-j)

In this reaction, a mixture of phenyl substituted cinnamic acid (4a-j) and SOCl_2 (which acted as both reactant and solvent) was stirred and refluxed at 80–90°C for 4 hrs. Remaining SOCl_2 was removed by distillation under reduced pressure. Thus, the intermediate acid chloride (2a-j) was added drop wise to 0°C cooled solution of morpholine (3) in dry pyridine to give product cinnamide (4a-j) with good yield as described in scheme 1.

(E)-1-morpholino-3-phenylprop-2-en-1-one (4a)

Color less solid; Yield 72 %; mp: 158-160°C; IR (KBr, cm^{-1}): 1660 (C=O), 1580 (C=C), 1170 (C-O); $^1\text{H-NMR}$ (DMSO-*d*6) δ : 7.48 (d, 2H, -ArH), 7.22 (t, 1H, -ArH), 6.83 (dd, 2H, -ArH), 6.72 (d, 1H, =CH), 7.11 (d, 1H, =CH), 2.50 (t, 2H, -CH₂), 2.14 (t, 2H, -CH₂); ^{13}C NMR (DMSO-*d*6) δ (ppm) : 158.4, 140.2, 135.3, 127.7, 127.3, 126.4, 115.8, 59.6, 44.8; MS: m/z 218 (M+H)⁺ for C₁₃H₁₅NO₂; Anal. calcd. for C₁₃H₁₅NO₂: C 71.87, H 6.96, N 6.45; Found: C 71.80, H 6.94, N 6.40%.

(E)-3-(2-chlorophenyl)-1-morpholinoprop-2-en-1-one (4b)

Pale yellowish brown solid in 70 %; mp: 188-190°C; IR(KBr, cm^{-1}): 1675 (C=O), 1550 (C=C), 1140 (C-O); $^1\text{H-NMR}$ (DMSO-*d*6) δ : 7.80 (d, 1H, -ArH), 7.57 (d, 1H, -ArH), 7.26 (t, 2H, -ArH), 7.39 (d, 1H, =CH), 7.99 (d, 1H, =CH), 3.50 (t, 2H, -CH₂), 3.72 (t, 2H, -CH₂); ^{13}C NMR

(DMSO-*d*6) δ (ppm) : 165.2, 138.9, 134.6, 133.6, 130.3, 130.1, 127.6, 126.9, 119.9, 66.8, 47.4; MS: m/z 252 (M+H)⁺ for C₁₃H₁₄ClNO₂; Anal. calcd. for C₁₃H₁₄ClNO₂: C 62.03, H 5.61, N 5.56; Found: C 61.94, H 5.59, N 5.51%.

(E)-3-(2-methoxyphenyl)-1-morpholinoprop-2-en-1-one (4c)


Color less solid; Yield 68 %; mp: 168-170°C; IR (KBr, cm^{-1}): 1680 (C=O), 1545 (C=C), 1130 (C-O); $^1\text{H-NMR}$ (DMSO-*d*6) δ : 7.33 (dd, 1H, -ArH), 6.97 (d, 1H, -ArH), 6.93 (dd, 1H, -ArH), 6.89 (d, 1H, -ArH), 7.91 (d, 1H, =CH), 7.46 (d, 1H, =CH), 3.87 (s, 3H, -OCH₃), 3.70 (m, 8H, -CH); ^{13}C NMR (DMSO-*d*6) δ (ppm) : 166.2, 158.2, 138.6, 130.7, 129.0, 124.2, 120.6, 111.6, 111.2, 66.8, 55.4, 53.3; MS: m/z 248 (M+H)⁺ for C₁₄H₁₇NO₃; Anal. calcd. for C₁₄H₁₇NO₃: C 68.00, H 6.93, N 5.66; Found: C 67.91, H 6.90, N 5.60%.

(E)-3-(3-methoxyphenyl)-1-morpholinoprop-2-en-1-one (4d)


Pale yellow solid in 70 %; mp: 169-171°C; IR(KBr, cm^{-1}): 1665 (C=O), 1530 (C=C), 1070 (C-O); $^1\text{H-NMR}$ (DMSO-*d*6) δ : 7.34 (d, 1H, -ArH), 7.32 (d, 1H, -ArH), 6.95 (dd, 1H, -ArH), 7.28 (d, 1H, -ArH), 7.49 (d, 1H, =CH), 7.24 (d, 1H, =CH), 3.81 (s, 3H, -OCH₃), 3.62 (m, 8H, -CH); ^{13}C NMR (DMSO-*d*6) δ (ppm) : 160.9, 158.4, 140.1, 136.7, 130.3, 127.7, 122.1, 115.8, 111.7, 107.6, 62.2, 55.5, 46.2; MS: m/z 248 (M+H)⁺ for C₁₄H₁₇NO₃; Anal. calcd. for C₁₄H₁₇NO₃: C 68.00, H 6.93, N 5.66; Found: C 67.91, H 6.90, N 5.60%.

(E)-3-(4-methoxyphenyl)-1-morpholinoprop-2-en-1-one (4e)

Color less solid in 71 %; mp: 166-169°C; IR(KBr, cm^{-1}): 1640 (C=O),



A: SOCl_2 , reflux, 80–90 °C, 4 hours B: Pyridine, reflux, 80–90 °C, 4 hours



Scheme 1:

1540 (C=C), 1090 (C-O); ¹H-NMR (DMSO-*d*₆) δ: 7.58 (d, 2H, -ArH), 6.95 (d, 2H, -ArH), 7.29 (d, 1H, =CH), 7.04 (d, 1H, =CH), 3.62 (s, 3H, -OCH₃), 3.43 (m, 8H, -CH); ¹³C NMR (DMSO-*d*₆) δ (ppm) : 162.9, 159.2, 140.1, 132.7, 128.3, 115.8, 111.7, 66.2, 55.8, 43.0; MS: m/z 248 (M+H)⁺ for C₁₄H₁₇NO₃; Anal. calcd. for C₁₄H₁₇NO₃: C 68.00, H 6.93, N 5.66; Found: C 67.91, H 6.90, N 5.60%.

(E)-3-(2,4-dimethoxyphenyl)-1-morpholinoprop-2-en-1-one (4f)

Color less solid in 70 %; mp: 174-176°C; IR(KBr, cm⁻¹): 1666 (C=O), 1536 (C=C), 1140 (C-O); ¹H-NMR (DMSO-*d*₆) δ: 7.78 (d, 1H, -ArH), 6.60 (d, 1H, -ArH), 6.55 (s, 1H, -ArH), 7.69 (d, 1H, =CH), 7.02 (d, 1H, =CH), 3.85 (s, 3H, -OCH₃), 3.59 (s, 3H, -OCH₃), 3.38 (m, 8H, -CH); ¹³C NMR (DMSO-*d*₆) δ (ppm) : 165.22, 161.98, 158.93, 136.3, 129.1, 116.4, 114.8, 105.8, 98.2, 66.2, 55.6, 55.5, 44.3; MS: m/z 278 (M+H)⁺ for C₁₅H₁₉NO₄; Anal. calcd. for C₁₅H₁₉NO₄: C 64.97, H 6.91, N 5.05; Found: C 64.90, H 6.88, N 5.01%.

(E)-3-(3,4-dimethoxyphenyl)-1-morpholinoprop-2-en-1-one (4g)

Color less solid in 68 %; mp: 176-179°C; IR(KBr, cm⁻¹): 1683 (C=O), 1526 (C=C), 1133 (C-O); ¹H-NMR (DMSO-*d*₆) δ: 7.35 (s, 1H, -ArH), 7.19 (dd, 1H, -ArH), 7.08 (d, 1H, -ArH), 7.44 (d, 1H, =CH), 6.95 (d, 1H, =CH), 3.82 (s, 3H, -OCH₃), 3.71 (s, 3H, -OCH₃), 3.60 (m, 8H, -CH); ¹³C NMR (DMSO-*d*₆) δ (ppm) : 160.2, 150.5, 149.5, 138.7, 128.3, 122.1, 115.8, 111.7, 107.6, 68.6, 55.6, 48.4; MS: m/z 278 (M+H)⁺ for C₁₅H₁₉NO₄; Anal. calcd. for C₁₅H₁₉NO₄: C 64.97, H 6.91, N 5.05; Found: C 64.90, H 6.88, N 5.01%.

(E)-3-(benzo[d][1,3]dioxol-5-yl)-1-morpholinoprop-2-en-1-one (4h)

Pale yellow solid in 71 %; mp: 179-181°C; IR(KBr, cm⁻¹): 1675 (C=O), 1580 (C=C), 1149 (C-O); ¹H-NMR (DMSO-*d*₆) δ: 7.41 (s, 1H, -ArH), 7.15 (dd, 1H, -ArH), 6.92 (d, 1H, -ArH), 7.45 (d, 1H, =CH), 7.12 (d, 1H, =CH), 6.02 (s, 2H, -OCH₂), 3.70 (m, 8H, -CH); ¹³C NMR (DMSO-*d*₆) δ (ppm) : 164.2, 150.5, 148.4, 139.5, 128.3, 127.3, 122.1, 118.8, 108.4, 104.6, 66.2, 49.2; MS: m/z 262 (M+H)⁺ for C₁₄H₁₅NO₄; Anal. calcd. for C₁₄H₁₅NO₄: C 64.36, H 5.79, N 5.36; Found: C 64.26, H 5.77, N 5.31%.

(E)-1-morpholino-3-(2,3,4-trimethoxyphenyl)prop-2-en-1-one (4i)

Color less solid in 68 %; mp: 196-199°C; IR(KBr, cm⁻¹): 1645 (C=O), 1560 (C=C), 1123 (C-O); ¹H-NMR (DMSO-*d*₆) δ: 7.20 (d, 1H, -ArH), 6.66 (d, 1H, -ArH), 7.79 (d, 1H, =CH), 6.87 (d, 1H, =CH), 3.87 (s, 9H, -OCH₃), 3.71 (m, 8H, -CH); ¹³C NMR (DMSO-*d*₆) δ (ppm) : 165.8, 154.6, 152.7, 142.1, 138.0, 123.0, 121.8, 115.5, 107.1, 66.4, 60.7, 60.4, 55.6, 48.2; MS: m/z 308 (M+H)⁺ for C₁₆H₂₁NO₅; Anal. calcd. for C₁₆H₂₁NO₅: C 62.53, H 6.89, N 4.56; Found: C 62.44, H 6.86, N 4.50%.

(E)-1-morpholino-3-(3,4,5-trimethoxyphenyl)prop-2-en-1-one (4j)

Color less solid in 68 %; mp: 176-179°C; IR(KBr, cm⁻¹): 1666 (C=O), 1525 (C=C), 1136 (C-O); ¹H-NMR (DMSO-*d*₆) δ: 7.04 (s, 2H, -ArH), 7.45 (d, 1H, =CH), 7.15 (d, 1H, =CH), 3.62 (s, 9H, -OCH₃); ¹³C NMR (DMSO-*d*₆) δ (ppm) : 162.0, 150.5, 138.7, 135.3, 130.7, 115.8, 107.6, 66.2, 60.6, 56.5, 44.2; MS: m/z 330 (M+Na)⁺ for C₁₆H₂₁NO₅; Anal. calcd. for C₁₆H₂₁NO₅: C 62.53, H 6.89, N 4.56; Found: C 62.44, H 6.86, N 4.50%.

Pharmacological Screening

Antioxidant screening (*in vitro*)

The compounds 4a-j were tested for antioxidant property by DPPH, NO and H₂O₂ methods.

DPPH radical scavenging activity

The hydrogen atom or electron donation ability of the compounds was measured from the bleaching of the purple colored methanol solution of 1,1-diphenyl-1-picrylhydrazyl (DPPH). The spectrophotometric assay uses the stable radical DPPH as a reagent. 1 ml of various concentrations of the test compounds (25, 50, 75 and 100 µg/mL) in methanol were added to 4 ml of 0.004% (w/v) methanol solution of DPPH. After a 30 min incubation period at room temperature, the absorbance was read against blank at 517 nm. The percent of inhibition (I %) of free radical production from DPPH was calculated by the following equation

$$\% \text{ of scavenging} = [(A \text{ control} - A \text{ sample}) / A \text{ blank}] \times 100 \rightarrow Eq. (1)$$

Where A control is the absorbance of the control reaction (containing all reagents except the test compound) and A sample is the absorbance of the test compound. Tests were carried at in triplicate.

Nitric oxide (NO) scavenging activity

Nitric oxide scavenging activity was measured by slightly modified methods of Green et al. and Marcocci et al. Nitric oxide radicals (NO) were generated from sodium nitroprusside. 1 mL of sodium nitroprusside (10 mM) and 1.5 mL of phosphate buffer saline (0.2 M, pH 7.4) were added to different concentrations (25, 50, 75 and 100 µg/mL) of the test compounds and incubated for 150 min at 25°C and 1 mL of the reaction mixture was treated with 1 mL of Griess reagent (1% sulfanilamide, 2% H₃PO₄ and 0.1% naphthyl ethylenediamine dihydrochloride). The absorbance of the chromophore was measured at 546 nm. Nitric oxide scavenging activity was calculated by using Equation (1).

Hydrogen peroxide (H₂O₂) scavenging activity

The H₂O₂ scavenging ability of the test compound was determined according to the method of Ruch *et al.* A solution of H₂O₂ (40 mM) was prepared in phosphate buffer (pH 7.4). 25, 50, 75 and 100 g/ml concentrations of the test compounds in 3.4 ml phosphate buffer were added to H₂O₂ solution (0.6 mL, 40 mM). The absorbance value of the reaction mixture was recorded at 230 nm. The percent of scavenging of H₂O₂ was calculated by using Eq. (1)

Results and Discussion

Chemistry

Target compounds (4a-j) were prepared as outlined in Scheme 1. In order to investigate how substituents on the phenyl ring in cinnamamides influence the activities, a mixture of substituted cinnamic acid (1a-j) and SOCl₂ (which acted as both reactant and solvent) was stirred and refluxed at 80-90°C for 4 hrs. Remaining SOCl₂ was removed by distillation under reduced pressure. Thus, the intermediate acid chloride (2a-j) was added drop wise to 0°C cooled solution of morpholine (3) in dry pyridine to give product cinnamide (4a-j) with good yield.

The structure of all newly synthesized morpholine containing cinnamide derivatives were determined by IR, NMR and Mass spectroscopic analysis and the results were in agreement with the proposed structures. In ¹H NMR spectra, the aromatic protons resonated at δ 6.55–7.80 ppm. The morpholine ring proton resonated with two triplets around δ 2.50 and 3.72. The conjugated double bond of cinnamic acid showed two doublet of which C–H proton near to aromatic ring (Ph–CH=HC–) resonated at δ 7.11–7.99 while C–H proton towards the carbonyl end (Ph–CH=CH–CO) resonated at δ 6.72–7.46. The synthesized compounds exhibited characteristic peak of

cinnamic acid double bond appeared at 1525–1580 cm^{-1} and carbonyl peak appeared at 1640–1683 cm^{-1} . ^{13}C NMR further confirmed the structures of the desired products and the $(\text{M}+\text{H})^+$ molecular ion peak of all the compounds were confirmed by their molecular weights.

Antimicrobial activity

All the synthesized compounds (4a-j) were screened for their antimicrobial activity determined by well plate method [28,29]. The antimicrobial values showed that most of the synthesized derivatives exhibited excellent activities against different strains of bacteria and moderate activity against fungi. Ciprofloxacin and Fluconazole were used as the standard drugs for antimicrobial and antifungal testing, respectively.

Antibacterial activity: All the synthesized compounds were screened against Gram +ve bacterial strains such as *Staphylococcus aureus* and *Bacillus subtilis* and Gram -ve bacterial strains such as *Klebsiella pneumoniae* and *Escherichia coli*. The results obtained as MIC is presented in Table 1. The Compounds 4f and 4i were showed good activity 12.5 and 6.12 $\mu\text{g}/\text{mL}$ respectively against the Gram positive bacterial strain *Staphylococcus aureus*, when compared with standard ciprofloxacin (12.5 $\mu\text{g}/\text{mL}$). When *Bacillus subtilis* was introduced as another Gram positive bacterial strain for conducting the antibacterial test, it was found that 4b and 4i both compounds were also exhibited good activity 12.5 $\mu\text{g}/\text{mL}$ equivalent to the standard ciprofloxacin. Some of the compounds 4b, 4c, 4e and 4j were screened moderate activity and rest of the compounds 4d and 4g does not show any activity against Gram +ve bacterial strains. The compounds 4b and 4i were showed MIC value 25 $\mu\text{g}/\text{mL}$ equivalent to that reported by the standard ciprofloxacin (25 $\mu\text{g}/\text{mL}$) against Gram negative bacterial strain *Klebsiella pneumoniae*. When *E.coli* was introduced as another gram-negative strain for conducting the antibacterial test, it was found that Compounds 4f and 4i were exhibited MIC value 12.5 $\mu\text{g}/\text{mL}$ equivalent to that reported by the standard ciprofloxacin (12.5 $\mu\text{g}/\text{mL}$) against Gram negative bacterial strain *Escherichia coli*. Some of the compounds 4c, 4e and 4j showed less MIC values and 4g compound does not show any activity against Gram negative bacterial strains. By comparing all these values, the compound which is having highest electron releasing substituent on benzene ring in cinnamide derivative was found potent activity against all the tested organisms (Table 1).

Antifungal activity: All the synthesized compounds 4a-j was also screened for their antifungal activity against two fungal strains such

as *Candida albicans* and *Aspergillus flavus*. Fluconazole was taken as a standard drug throughout the experiment. Compound 4b was found to be the excellent and potent antifungal activity with MIC 3.12 $\mu\text{g}/\text{mL}$ and 6.25 $\mu\text{g}/\text{mL}$ against *C. albicans* and *A. flavus* respectively, while standard drug Fluconazole with 6.25 $\mu\text{g}/\text{mL}$. Only two compounds 4i and 4j were also showed good activity with 3.12 $\mu\text{g}/\text{mL}$ and 6.25 $\mu\text{g}/\text{mL}$ against *Aspergillus flavus* and *Candida albicans* respectively and rest of the compounds 4c, 4a, 4d, 4f exhibit very poor activity compared with standard Fluconazole (Table 1).

Antioxidant screening (in vitro)

In the present study, the antioxidant activity of the synthesized compounds was assessed in vitro by the 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging assay, nitric oxide (NO) and hydrogen peroxide (H_2O_2) [30-34]. These methods were based on measuring the continual absorbance decrease of the methanolic solution of the DPPH at 517 nm, in the presence of antioxidant compound. The DPPH has an odd electron so it can accept an electron or hydrogen free radical. In the presence of antioxidant, this odd electron becomes paired due to H transfer from antioxidant and hence DPPH absorbance decreases. The ability of newly synthesized compounds (4a-j) act as hydrogen donors or free radical scavengers was tested by in vitro antioxidant assays involving DPPH radical, NO radical, H_2O_2 and the results were compared with that of standard antioxidant Ascorbic acid. In DPPH method, the newly synthesized compounds showed moderate to good antioxidant activity compared to the standard (Table 2). Among the newly synthesized compounds 4f, 4i and 4j were found to be good antioxidant activity compared with that of the standard Ascorbic acid (18.33 ± 0.04) with the IC_{50} values 19.44 ± 0.14 , 19.43 ± 0.08 and 19.09 ± 0.09 $\mu\text{g}/\text{mL}$ respectively. Some of the compounds 4c, 4e and 4g were exhibited moderate activity with the IC_{50} values 21.06 ± 1.23 , 21.44 ± 0.33 and 20.04 ± 1.18 $\mu\text{g}/\text{mL}$ respectively and the remaining compounds 4a and 4b were exhibited less activity and compounds 4d and 4h does not show any activity. In DPPH scavenging method only two compounds 4i and 4j showed good antioxidant activity because of these two compounds having more electron releasing methoxy groups on phenyl ring in cinnamide.

Further all the newly synthesized compounds were subjected to the NO radical scavenging activity. The IC_{50} value of NO is the concentration of sample required to inhibit 50% of the NO radicals. Except 4b and 4d all the tested compounds exhibited strong NO radical scavenging with moderate to good IC_{50} values. The greater NO radical scavenging

Compound	Minimum inhibitory concentration (MIC) ($\mu\text{g mL}^{-1}$)					
	Antibacterial activity				Antifungal activity	
	Gram +Ve bacteria		Gram -Ve bacteria			
	<i>S. aureus</i>	<i>B. subtilis</i>	<i>K. pneumoniae</i>	<i>E. coli</i>	<i>C. albicans</i>	<i>A. flavus</i>
4a	100	50	>100	100	50	100
4b	25	12.5	25	50	3.12	6.25
4c	25	50	50	100	50	100
4d	—	—	100	50	25	50
4e	25	50	50	25	100	50
4f	12.5	25	100	12.5	50	>100
4g	—	—	>100	50	—	—
4h	50	100	—	—	100	>100
4i	6.12	12.5	25	12.5	12.5	3.12
4j	25	25	50	25	6.25	12.5
Ciprofloxacin	12.5	12.5	25	12.5	—	—
Fluconazole	—	—	—	—	6.25	6.25

(—) No activity. Lower MIC values indicate higher the antimicrobial activity.

S. aureus: *Staphylococcus aureus*, *B. subtilis*: *Bacillus subtilis*, *K. pneumoniae*: *Klebsiella pneumoniae*, *E. coli*: *Escherichia coli*, *C. albicans*: *Candida albicans*.

Table 1: Antimicrobial activity (MIC profiles) of the synthesized compounds (4a-j).

activity of the tested compounds was showed by compounds 4i and 4j showed 18.33 ± 0.09 and 18.86 ± 0.97 respectively (Table 3).

The prepared compounds (4a-j) having various concentrations (25, 50, 75 and 100 $\mu\text{g/mL}$) were subjected to H_2O_2 radical scavenging activity. The findings of present study (Table 4) indicated that the most of the prepared compounds exhibited moderate to good radical scavenging ability. The good H_2O_2 scavenging effect was detected in compounds 4i and 4j with the values of IC_{50} 18.02 ± 0.09 and 18.33 ± 0.10 respectively compared with the standard Ascorbic acid (17.80 ± 0.08). Rest of the compounds 4e, 4f, and 4g were showed moderate

activity with IC_{50} values 20.76 ± 1.33 , 20.01 ± 0.20 and 19.43 ± 0.16 respectively and the compounds 4a, 4b, 4c, and 4d exhibited less activity. The reason would be the presence of electron releasing methoxy groups present on the phenyl ring.

Structure activity relationship (SAR) study was helped to reveal the effect of different substituent on the microbial strains, depending upon the different electronic environments developed on the aromatic ring by substituting the electron donating groups. It was observed that the compounds 4j (3,4,5- OCH_3), 4i (2,3,4- OCH_3), 4f (2,4- OCH_3) and 4b (2-Cl) with electronic donating groups exhibited good antimicrobial

Compound	Concentration ($\mu\text{g mL}^{-1}$)				
	25	50	75	100	IC_{50}
4a	56.21 ± 0.76	57.15 ± 0.62	59.75 ± 0.65	60.20 ± 0.94	22.23 ± 0.30
4b	54.32 ± 0.27	55.35 ± 0.93	57.25 ± 0.07	61.16 ± 0.97	23.01 ± 1.41
4c	59.35 ± 0.25	60.34 ± 0.18	62.46 ± 0.16	64.16 ± 0.72	21.06 ± 1.23
4d	-	-	-	-	-
4e	58.29 ± 0.64	62.24 ± 0.96	64.54 ± 1.08	68.72 ± 0.98	21.44 ± 0.33
4f	64.29 ± 0.27	65.26 ± 0.17	66.72 ± 0.23	67.14 ± 0.10	19.44 ± 0.14
4g	62.35 ± 0.10	62.66 ± 0.36	63.49 ± 0.89	69.14 ± 1.06	20.04 ± 1.18
4h	-	-	-	-	-
4i	64.32 ± 0.09	66.13 ± 0.20	67.14 ± 0.61	72.14 ± 0.02	19.43 ± 0.08
4j	65.45 ± 0.22	68.29 ± 0.92	70.39 ± 0.12	73.28 ± 0.90	19.09 ± 0.09
Ascorbic acid	68.04 ± 0.18	71.24 ± 0.15	72.43 ± 0.22	75.29 ± 0.03	18.37 ± 0.04
Blank	-	-	-	-	-

(-) showed no scavenging activity. Values were the means of three replicates \pm SD

Table 2: The *in vitro* antioxidant activity of (4a-j) in DPPH method

Compound	Concentration ($\mu\text{g mL}^{-1}$)				
	25	50	75	100	IC_{50}
4a	57.29 ± 0.84	60.16 ± 0.72	62.24 ± 0.63	63.42 ± 0.94	21.81 ± 1.30
4b	-	-	-	-	-
4c	63.40 ± 0.23	64.29 ± 0.19	65.19 ± 0.14	66.18 ± 0.78	19.71 ± 1.03
4d	-	-	-	-	-
4e	64.10 ± 0.64	65.20 ± 0.96	67.20 ± 1.08	69.14 ± 0.98	19.50 ± 0.64
4f	62.32 ± 0.07	66.28 ± 0.07	69.25 ± 0.03	70.24 ± 0.10	20.05 ± 0.02
4g	63.19 ± 0.10	65.26 ± 0.06	68.76 ± 0.89	71.24 ± 1.06	19.78 ± 0.18
4h	62.28 ± 0.28	63.16 ± 0.17	64.26 ± 0.44	69.14 ± 1.26	20.07 ± 1.03
4i	68.19 ± 0.06	69.14 ± 0.03	72.34 ± 0.06	76.14 ± 0.22	18.33 ± 0.09
4j	66.27 ± 0.30	68.16 ± 0.62	71.32 ± 0.12	74.29 ± 0.40	18.86 ± 0.97
Ascorbic acid	69.24 ± 0.23	72.16 ± 0.02	76.29 ± 0.21	78.31 ± 0.15	18.05 ± 0.05
Blank	-	-	-	-	-

(-) showed no scavenging activity. Values were the means of three replicates \pm SD

Table 3: The *in vitro* antioxidant activity (4a-j) in NO method.

Compound	Concentration ($\mu\text{g mL}^{-1}$)				
	25	50	75	100	IC_{50}
4a	52.16 ± 0.14	54.29 ± 0.48	55.19 ± 0.63	57.29 ± 0.72	23.96 ± 0.82
4b	53.24 ± 0.49	54.16 ± 0.51	56.28 ± 0.14	59.14 ± 0.33	23.47 ± 1.44
4c	59.28 ± 0.42	60.24 ± 0.69	62.32 ± 0.14	64.18 ± 0.78	21.08 ± 0.23
4d	51.20 ± 0.64	52.60 ± 0.44	54.39 ± 0.36	56.27 ± 0.94	24.41 ± 1.25
4e	60.20 ± 0.45	61.14 ± 0.96	62.22 ± 1.12	64.19 ± 0.38	20.76 ± 1.33
4f	62.44 ± 0.24	64.27 ± 0.15	65.29 ± 0.56	68.19 ± 0.19	20.01 ± 0.20
4g	64.33 ± 0.22	65.28 ± 0.15	67.19 ± 0.39	70.23 ± 1.14	19.43 ± 0.16
4h	-	-	-	-	-
4i	69.35 ± 0.06	72.39 ± 0.23	73.14 ± 0.16	75.22 ± 0.24	18.02 ± 0.09
4j	68.16 ± 0.20	70.19 ± 0.90	71.23 ± 0.22	74.28 ± 0.90	18.33 ± 0.10
Ascorbic acid	70.20 ± 0.12	72.14 ± 0.09	74.28 ± 0.14	76.14 ± 0.06	17.80 ± 0.08
Blank	-	-	-	-	-

(-) showed no scavenging activity. Values were the means of three replicates \pm SD

Table 4: The *in vitro* antioxidant activity of (4a-j) in H_2O_2 method

results, displaying moderate to good MIC values and good antioxidant properties with good IC₅₀ values.

Conclusion

In this work conclusion, a new series of trans cinnamamides were synthesized from starting materials namely morpholine and substituted cinnamic acids with good yield. All the title compounds were screened for their antimicrobial and antioxidant activity studies. The investigation of antimicrobial and antioxidant activity screening data reveals that, among the all the compounds only few of them are exhibiting moderate to good activity. Compounds 4i and 4j showed an excellent, almost equivalent to that of standard, rest of the compounds showed moderate to mild inhibition activity.

Acknowledgement

The authors are grateful to the Council of Scientific and Industrial Research (CSIR), New Delhi, for financial assistance under a Junior Research Fellow (JRF).

References

- Liu L, Hudgins WR, Shack S, Yin MQ, Samid D (1995) Cinnamic acid: a natural product with potential use in cancer intervention. *Int J Cancer* 62: 345-350.
- Natella F, Nardini M, Di Felice M, Scaccini C (1999) Benzoic and cinnamic acid derivatives as antioxidants: structure-activity relation. *J Agric Food Chem* 47: 1453-1459.
- De P, Baltas M, Bedos-Belval F (2011) Cinnamic acid derivatives as anticancer agents-a review. *Curr Med Chem* 18: 1672-1703.
- Narasimhan B, Belsare D, Pharande D, Mourya V, Dhake A (2004) Esters, amides and substituted derivatives of cinnamic acid: synthesis, antimicrobial activity and QSAR investigations. *Eur J Med Chem* 39: 827-834.
- Carvalho SA, da Silva EF, de Souza MV, Lourenço MC, Vicente FR (2008) Synthesis and antimycobacterial activity of cinnamide derivatives: a molecular hybridization approach. *Bioorg Med Chem Lett* 18: 538-541.
- Yoya GK, Bedos-Belval F, Constant P, Duran H, Daffa M, et al. (2009) Synthesis and evaluation of a novel series of pseudo-cinnamic derivatives as antituberculosis agents. *Bioorg Med Chem Lett* 19: 341-343.
- Kakwani MD, Suryavanshi P, Ray M, Rajan MG, Majee S, et al. (2011) Design, synthesis and antimycobacterial activity of cinnamide derivatives: a molecular hybridization approach. *Bioorg Med Chem Lett* 21: 1997-1999.
- De P, Koumba Yoya G, Constant P, Bedos-Belval F, Duran H, et al. (2011) Design, synthesis, and biological evaluation of new cinnamic derivatives as antituberculosis agents. *J Med Chem* 54: 1449-1461.
- Leslie BJ, Holaday CR, Nguyen T, Hergenrother PJ (2010) Phenylcinnamides as novel antimetabolic agents. *J Med Chem* 53: 3964-3972.
- Natella F, Nardini M, Di Felice M, Scaccini C (1999) Benzoic and cinnamic acid derivatives as antioxidants: structure-activity relation. *J Agric Food Chem* 47: 1453-1459.
- Cutillo F, D'Ambrosia B, DellaGreca M, Di Marino C, Golino A, et al. (2003) Cinnamic acid amides from *Chenopodium album*: effects on seeds germination and plant growth. *Phytochemistry* 64: 1381-1387.
- Kato MJ, Furlan M (2007) Chemistry and evolution of the Piperaceae. *Pure Appl Chem* 79: 529-538.
- Dyer LA, Dodson CD, Stireman JO 3rd, Tobler MA, Smilanich AM, et al. (2003) Synergistic effects of three Piper amides on generalist and specialist herbivores. *J Chem Ecol* 29: 2499-2514.
- Richards LA, Dyer LA, Smilanich AM, Dodson CD (2010) Synergistic effects of amides from two piper species on generalist and specialist herbivores. *J Chem Ecol* 36: 1105-1113.
- Castral TC, Matos AP, Monteiro JL, Araujo FM, Bondancia TM, et al. (2011) Synthesis of a combinatorial library of amides and its evaluation against the fall armyworm, *Spodoptera frugiperda*. *J Agric Food Chem* 59: 4822-4827.
- Slee DH, Romano SJ, Yu J, Nguyen TN, John JK, et al. (2001) Development of potent non-carbohydrate imidazole-based small molecule selectin inhibitors with antiinflammatory activity. *J Med Chem* 44: 2094-2107.
- Guan LP, Sui X, Deng XQ, Zhao DH, Qu YL, et al. (2011) N-palmitoylethanol amide derivatives: synthesis and studies on anticonvulsant and antidepressant activities. *Med Chem Res* 20: 601-606.
- Papa D, Schwenk E, Villani F, Klingsberg E (1950) The Analgesic Activity of N,N-Dialkyl Amides. *J Am Chem Soc* 72: 3885-3886.
- Singh N, Sijwali PS, Pandey KC, Rosenthal PJ (2006) Plasmodium falciparum: biochemical characterization of the cysteine protease falcipain-2'. *Exp Parasitol* 112: 187-192.
- Hirokawa Y, Kinoshita H, Tanaka T, Nakamura T, Fujimoto K, et al. (2009) Pleuromutilin derivatives having a purine ring. Part 2: Influence of the central spacer on the antibacterial activity against Gram-positive pathogens. *Bioorg Med Chem Lett* 19: 170-174.
- Fung HB, Kirschenbaum HL, Ojofeitimi BO (2001) Linezolid: an oxazolidinone antimicrobial agent. *Clin Ther* 23: 356-391.
- Yoshida K, Nakayama K, Yokomizo Y, Ohtsuka M, Takemura M, et al. (2006) MexAB-OprM specific efflux pump inhibitors in *Pseudomonas aeruginosa*. Part 6: Exploration of aromatic substituents. *Bioorg Med Chem* 14: 8506-8518.
- Takeuchi H, Nakato K, Hattori H, Sonoda R, Koshiya K, et al. (1997) Pharmacological studies on YM992, a novel antidepressant with selective serotonin re-uptake inhibitory and 5-HT_{2A} receptor antagonistic activity. *Eur J Pharmacol* 329: 27-35.
- Li HQ, Zhu TT, Yan T, Luo Y, Zhu HL (2009) Design, synthesis and structure-activity relationships of antiproliferative ,3-disubstituted urea derivatives. *Eur J Med Chem* 44: 453-459.
- Chrysselis MC, Rekkas EA, Kourounakis PN (2000) Hypocholesterolemic and hypolipidemic activity of some novel morpholine derivatives with antioxidant activity. *J Med Chem* 43: 609-612.
- Szulawska A, Arkusinska J, Czyz M (2007) Accumulation of gamma-globin mRNA and induction of irreversible erythroid differentiation after treatment of CML cell line K562 with new doxorubicin derivatives. *Biochem Pharmacol* 73: 175-184.
- Jakubowska J, Wasowska-Lukawska M, Czyz M (2008) STI571 and morpholine derivative of doxorubicin collaborate in inhibition of K562 cell proliferation by inducing differentiation and mitochondrial pathway of apoptosis. *Eur J Pharmacol* 596: 41-49.
- Koneman EW, Allen SD, Winn WC, et al. (1997) *Colour Atlas and Textbook of Diagnostic Microbiology*, Lippincott Raven Pub, Philadelphia, 86-856.
- NCCLS (2002) Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts Approved Standard, second ed. ISBN 1-56238- 469-4 NCCLS document M27-A2.
- Burits M, Bucar F (2000) Antioxidant activity of *Nigella sativa* essential oil. *Phytother Res* 14: 323-328.
- Cuendet M, Hostettmann K, Potterat O, Dyatmiko W (1997) Iridoid Glucosides with Free Radicals Scavenging properties from *Fagvaea blumei*. *Helv Chim Acta* 80: 1144-1152.
- Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS, et al. (1982) Analysis of nitrate, nitrite, and [15N]nitrate in biological fluids. *Anal Biochem* 126: 131-138.
- Marcocci L, Maguire JJ, Droy-Lefaux MT, Packer L (1994) The nitric oxide-scavenging properties of Ginkgo biloba extract EGb 761. *Biochem Biophys Res Commun* 201: 748-755.
- Ruch RJ, Cheng SJ, Klaunig JE (1989) Prevention of cytotoxicity and inhibition of intercellular communication by antioxidant catechins isolated from Chinese green tea. *Carcinogenesis* 10: 1003-1008.