



Synthesis, characterization and antibacterial activity of aspirin and paracetamol-metal complexes

Amudat LAWAL and Joshua A. OBALEYE*

Department of Chemistry, Faculty of Science, University of Ilorin, Ilorin, Nigeria

Received 27 December 2006

MS/No BKM/2006/040, © 2007 Nigerian Society for Experimental Biology. All rights reserved.

Abstract

Novel complexes of Co (II), Ni (II) and Fe (III) with aspirin and paracetamol have synthesized and characterized using infrared, electronic and Hnmr spectral, melting point and conductivity measurements. The two ligands have been found to act as bidentate chelating agents. Aspirin complexes coordinate through the carbonyl oxygen of the carboxyl and the ester groups, while paracetamol complexes coordinate through the oxygen of the hydroxyl and the amide groups. Antibacterial screening of the complexes against *Bacillus subtilis*, *Serratia species* and *Escherichia coli*, was also investigated. The metal complexes were found to have varied degree of inhibitory effect against the bacteria.

.Keywords: Aspirin, Paracetamol, Metal Complexes, Antibacterial Activity

*To whom correspondence may be addressed, **E-mail:** jobaleye@yahoo.com, amudatlawal@yahoo.com **Tel:** +234-8033582048

INTRODUCTION

Aspirin is a derivative of salicylic acid. It has analgesic, anti-inflammatory and antipyretic actions and inhibits prostaglandin synthetase¹ while paracetamol is a derivative of 4-aminophenol which also has analgesic and antipyretic action². Paracetamol is useful in the treatment of pain such as headache, toothache, rheumatism and neuralgia².

However, it is known that some drugs act via chelation or by inhibiting metalloenzymes but for most of the drugs that act as potential ligands, a lot of studies are being carried out to ascertain how metal binding influences the activities of the drugs³. Although, in 1987, Abu-El-Wafa *et al*⁴ reported the synthesis of Co(ii), Ni(ii) and Fe(ii) complexes of metformin(L), also an analgesic, on the basis of analytical data obtained, the prepared complexes were formulated as ML₂.

Metal complexes are gaining increasing importance in the design of drugs on coordination with a metal. This has led to lots of study on metal drug complexes.

Therefore, it was considered necessary to have some other analgesic drug-metal complexes, due to their chemotherapeutic properties. We hereby report the synthesis, characterization and antibacterial activity studies of novel transition metal complexes of aspirin and paracetamol.

MATERIALS AND METHODS

All the chemicals were reagent grade and used as commercially obtained (Aldrich, BDH) without further purification.

Aspirin and paracetamol were obtained from Rajrab Pharmaceutical company Ltd., Ilorin, Kwara State, Nigeria.

The melting points of the ligands and the complexes were carried out using Gallenkamp melting apparatus.

The thin layer chromatography of the complexes was run using solvent mixture of ethanol,

acetone and petroleum ether in the 7:2:1 mole ratio.

The molar conductivity of the ligands and the complexes in methanol at room temperature ($25 \pm 0.1^{\circ}\text{C}$) were made on WTW Conductometer Bridge with a cell constant of 0.82cm^{-1} . The IR spectra were recorded in solid state as KBr pellets using Buck-Scientific M 500 model infrared spectrophotometer from 4000cm^{-1} to 600cm^{-1}

The uv/visible spectra were made on Sp8 –400 uv/visible spectrophotometer using methanol as solvents. The metal content of the complexes were determined by using an SP9, atomic absorption spectrophotometer with PM 8251 simple-pen recorder.

The proton nmr spectra of the ligands and the complexes were obtained on Bruker AC 200/300 spectrometer using deuterated chloroform as solvents.

Evaluation of antibacterial activity

The antibacterial activity of the test compounds was assayed against three bacteria: *Bacillus Subtilis*, *Serratia Species* and *Escherichia coli*.

The antibacterial activity of the compounds was determined as reported by Abd E1-Wahab and E1-sarrag⁵.

The antibacterial activity was determined on the seeded nutrient agar on which 0.9cm diameter wells punched. Different concentrations (0.1% and 1.0%w/v) of sterile filtered solutions of the ligands and the complexes were made using methanol as solvents, 0.1 ml of each concentration was applied into the wells and incubated at 37°C for one to three days.

The antibacterial activity was estimated on the basis of the size of inhibition zone formed around the well of the seeded agar plates and the inhibition growth in percentage was determined on the basis of the average diameter of bacterial colony on the growth medium to their respective controls as in the equation

$$\% \text{ Inhibition} = \frac{A-B}{A} \times 100$$

Where A= Average diameter of bacterial growth on the control and B= Average diameter of bacterial growth on the test plate

Synthesis of Aspirin-metal complexes

The complexes were prepared by adding on aqueous solution of the hydrated metal chloride (0.01mol) to an ethanolic solution of the ligand, Aspirin (3.604g, 0.02mol). The solution was refluxed with constant stirring for 3 hours. The complexes were recovered from their solutions and followed by washing with ethanol and dried in a desiccator.

Synthesis of Paracetamol- metal complexes

The preparation was similar to that of Aspirin metal complexes. An aqueous solution of hydrated metal chloride (0.01mol) in ethanol (10ml) was added to an ethanolic solution of the ligand, paracetamol 3.032g, 0.02mol) the mixture was refluxed for 3 hours and left for

about three weeks before complexation. The precipitated complex was removed by filtration and washed with ethanol and dried in a desiccator.

RESULTS AND DISCUSSION

The results of the analytical data, spectroscopic and antibacterial studies are presented in Tables 1-5.

All the complexes were found to be stable. They are non-hygroscopic solids with low melting points. The molar conductance data for the metal complexes in methanol show non-electrolytic behavior in this solvent.

All the complexes synthesized were coloured. The coloured complexes obtained for iron (III) complexes (d^5) were likely due to charge transfer from ligand to metal and *vice versa*.

Table 1: Analytical data and some physical properties of the complexes

Compound	Colour	M.P ($^{\circ}$ C)	Yield %	M %	Molar Conductance $\Omega^{-1}\text{g}/\text{dm}^3$	Electronic Transition (nm)
Aspirin(asp)	Whitish	138-140	-	-	5.20×10^7	275,243
Co(Asp) $_2$ Cl $_2$	Brown	122-124	60.4	12.10 (12.02)	6.12×10^{-5}	302,262,233
Ni(Asp) $_2$ Cl $_2$	White	118-120	52.3	10.98 (10.27)	4.20×10^{-5}	296,257,241
Fe(Asp) $_2$ Cl $_3$	Black	123-126	51.2	9.30 (9.76)	1.52×10^{-5}	297,265,235
Paracetamol	Whitish	168-170	-	-	4.25×10^{-7}	243
Co(Par) $_2$ Cl $_2$	Cream	136-138	73.9	12.14 (11.76)	2.43×10^{-6}	245,220212
Ni(Par) $_2$ Cl $_2$	Light Pint	144-146	64.7.	13.59 (13.75)	1.94×10^{-6}	248,218,210
Fe(Par) $_2$ Cl $_3$	Black	128-130	54.3	10.29 (10.24)	1.65×10^{-6}	238

Table 2: Selected IR spectral assignment of aspirin (asp) and its metal complexes


Aspirin (cm $^{-1}$)	Co(Asp) $_2$ Cl $_2$ (cm $^{-1}$)	Ni (Asp) $_2$ Cl $_2$ (cm $^{-1}$)	Fe(Asp) $_2$ Cl $_3$ (cm $^{-1}$)	Tentative Assignment
3776.3m,b	3235.3w 3101.5w	3235.3 s,b 3101.5m	3764.2 s 3235w,b	O-H str
1757.8 s 1675.6 v,s	1745.7m, b 1666.6 s,b	1745.7 m,b 1666.6 v,s	1754.7m 1666.6 s	C =O of ester C =O of Carboxylic acid
1374.7 s 1308.0v .s	1301.8 v.s 1247.2s	1247.2 v.s	1393 s,b 1295.9 v, s	C-O str of Carboxylic acid
1095.2, b 1016.1 m, b	1156.0 v, s 1028.2 m	1156.Os	1156.0 s	C-O str of ester
-	669.6 s	663.3 s, b	663.4 4s	M-OH & M-O =C

Table 3: Selected IR spectral assignment of *par*/acetamol (*par*) and its metal complexes

Paracetamol (Par)	Co(Par) ₂ C1 ₂	Ni(Par) ₂ C1 ₂	Fe(Par) ₂ C1 ₂	Tentaive Assignment
3785.2m,b 3694.9 w	3794.6w, b 3320.3 s 3162.3 w,b 3707.5 w,b	3758.2 b 3429.8 w,b 3320.3 m 3162.3 w	3782.4 m,b 3691.2 w,b 3630.4 w,b	O-H & N-H str
1625 s, b	1612.0 s	1612.0 s	1617.9 m,b	C=O str
-	693.8 s	681.7 s,b	-	[M-OH] and [M-O=C]

Table 4: HNMR Spectral Assignment of Aspirin (Asp) and Its Metal Complexes

Compound	Hnmr Signal(ppm)	Assignment
Asp	8.11-8.09(IH, d)	a
	7.63-7.58(IH,t)	b
	7.36-7.24(H,t)	c
	7.13-7.10(IH,d)	d
	2.33(3H,S)	e
Co(Asp) ₂ C1 ₂	10.45(IH,d)	f
	8.13-8.10(IH,d)	a
	7.93-7.90(IH,d)	b
	7.66-758(IH,t)	c
	2.35(3H,S)	e
Ni(Asp) ₂ C1 ₂	8.13-8.08(IH,d)	a
	7.65-7.58(IH,t)	b
	7.38-7.24(IH,t)	c
	7.14-7.10(IH,d)	d
	2.33(3H,s)	e
Fe (Asp) ₂ C1 ₃	7.23	Solvent Peak (CDC1 ₃)
	1.53	Solvent peak(H ₂ O)



The selective infrared spectra assignment of free Aspirin and its complexes are given in Table 2. The assignments have been carried out based on comparison of the spectra data with those of similar compounds⁶.

The absorption band at 3776.3cm⁻¹ in the spectrum of free Aspirin has been attributed to the O-H group. These bands undergo hypsochromic

shift to 3235.3cm⁻¹ and 3101.5cm⁻¹ in the metal complexes. The shifting of these (O-H) stretching vibrational bands provides evidence that this group is one of the coordination sites of Aspirin. This is also supported by broad bands at 294.9cm⁻¹, 852cm⁻¹ and 803.3cm⁻¹ attributed to O-H bending⁶. The bands at 1460cm⁻¹ and 1417.5cm⁻¹ have been assigned to C=O of ester and carboxylic acid respectively; these bands

also undergo hypsochromic shift in the spectra of the complexes. The shifting of these (C=O) stretching bands provides evidence that this group is also one of the coordination sites of Aspirin.

The strong absorption bands at 1374.7cm^{-1} and 1308cm^{-1} have been attributed to C-O stretching vibration of the carboxylic acid while the medium bands at 1095.2cm^{-1} and 1016.1cm^{-1} and 1016.1cm^{-1} have been attributed to C-O stretching vibration of the ester⁶.

The strong absorption bands between the ranges of 651.3cm^{-1} on the spectra of the metal complexes which could not be traced to free Aspirin have been tentatively assigned to [M-OH] and [M-O=C] stretch bands of the metal complexes⁷.

The IR spectra assignment of Paracetamol (Par) and its metal complexes are presented in Table 3.

The spectrum of free paracetamol was compared with the spectral of its metal complexes. The absorption bands at 3785.2cm^{-1} and 3694.9cm^{-1} of the free paracetamol have been assigned to O-H and N-H stretching vibrations. These bands have been shifted in the spectra of the metal complexes due to coordination.

The strong absorption bands at 1625.0cm^{-1} in the spectrum of free paracetamol has been assigned to C=O stretching. These bands have been shifted in the spectra of the metal that this complexes. The shifting of C=O group provide evidence of coordination through this group⁷. The strong absorption bands at 693.8cm^{-1} and 681.7cm^{-1} in the spectra of the $\text{Co}(\text{Par})_2\text{Cl}_2$ and $\text{Ni}(\text{Par})_2\text{Cl}_2$ complexes which could not be traced to free paracetamol have been tentatively assigned to [M-O=C] and [M-OH] stretch bands of the metal complexes⁷.

The H NMR signals of aspirin and its metal complexes are presented in Table 4. The multiplicity and integral values of the proton signals support the structure of the ligand. The assignment of the protons is supported by highly reduced intensity of the proton signals after deuterium exchange.

The $\text{Fe}(\text{Asp})_2\text{Cl}_3$ complex was insoluble in CDCl_3 , therefore the signals obtained were mainly the solvent peaks⁸.

By considering all the above analytical and spectroscopic data and in the absence of X-ray crystallographic data. The structures tentatively proposed for the complexes are shown in Figures I and II.




Fig I: Proposed structure for $\text{M}(\text{Asp})_2\text{X}_n$




Fig II: Proposed structure for $\text{M}(\text{Par})_2\text{X}_n$
M = $\text{Co}(\text{II})$, $\text{Ni}(\text{II})$, $n = 2$
M = $\text{Fe}(\text{II})$, $n = 3$

The aspirin complexes, $\text{M}(\text{Asp})_2\text{X}_n$ have insignificant effect on *Serratia* and *E. coli* species at both concentrations (Table 5). $\text{Co}(\text{Asp})_2\text{Cl}_2$ have the greatest inhibitory effect against *Bacillus subtilis* while $\text{Ni}(\text{Asp})_2\text{Cl}_2$ have the lowest inhibitory effect. At 1.0% concentration, Aspirin metal complexes show higher activity than the original aspirin.

The paracetamol complexes, $\text{M}(\text{Par})_2\text{X}_n$ have insignificant effect on *Serratia* and *Bacillus subtilis* species at both concentration. $\text{Fe}(\text{Par})_2\text{Cl}_3$ have the greatest inhibitory effect against *E. coli* while $\text{Ni}(\text{Par})_2\text{Cl}_2$ have the lowest inhibitory effects at both concentration. The

Table 5: Result of antibacterial test

Compound	Concentration	Serratia SP	Bacillus SP	Escherichia Coli
Asp	0.1%	- - -	0 0 0	- - -
Co(Asp) ₂ Cl ₂		- - -	50 51 50	- - -
Ni (Asp) ₂ Cl ₂		- - -	32 32 34	- - -
Fe (Asp) ₂ Cl ₃		- - -	45 47 48	- - -
Asp	1.0%	- - -	28 28 30	- - -
Co(Asp) ₂ Cl ₂		- - -	90 94 96	- - -
Ni(Asp) ₂ Cl ₂		- - -	52 55 57	- - -
Fe(Asp) ₂ Cl ₃		- - -	85 82 80	- - -
Par	0.1%	- - -	- - -	0 0 0
Co(Par) ₂ Cl ₂		- - -	- - -	25 27 27
Ni(Par) ₂ Cl ₂		- - -	- - -	22 22 24
Fe(Par) ₂ Cl ₃		- - -	- - -	30 31 30
Par	1.0%	- - -	- - -	18 20 18
Co(Par) ₂ Cl ₂		- - -	- - -	55 58 60
Ni(Par) ₂ Cl ₂		- - -	- - -	50 52 55
Fe(Par) ₂ Cl ₃		- - -	- - -	60 62 64

paracetamol complexes also show higher activity than the ligand⁹.

Acknowledgement

The authors are grateful to Rajrab Pharmaceutical Company limited, Ilorin, Kwara State, Nigeria for the supply of the ligands.

REFERENCES

1. **British Pharmaceutical Codex (1977)** 11th ed. Pharmaceutical Press, London, Pp 48 – 54.
2. **Remington's Pharmaceutical Science (1970)** 18th ed., John Wiley Pub. Inc. New York.
3. **Behrens, N. B., Diaz, G. M. and Goodgame, D.M.L (1986)** Metal complexes of the antibiotics Nalidixic acid. *Inorg. Chim. Acta.* **125**:21-264.
4. **Abu-El-Wafa, S.M El-Rifs, M.A and Ahmed, F.H (1987)** Formation of metformin complexes with some transition metal ions and their biological activity. *Inorg. Chim. Acta.* **136**:127-131
5. **Abu-El-Wahab, Z. H. and El-Sarrag, M. R. (2004)** Derivative of phosphate schift base transition metal complexes: synthesis, studies and biological activity. *Spec. Acta* **60**:271-277.
6. **William, K. (1991)** Organic Spectroscopy, 3rd edition, Macmillan Education Ltd, London , 49-54, 60-75

7. **Nakamoto, K (1970)** Infrared Spectra of Inorganic and Coordination Compounds, 2nd edition, John Wiley, New York , 166-219
8. **Issa, A (2001)** Synthesis, Characterization and Antimicrobial Activity of Metronidazole, Aspirin, Paracetamol and Tetracycline Metal Complexes. M.Sc Thesis, Department of Chemistry, University of Ilorin , Pp 50-98.
9. **Obaleye, J.A Nde -aga,, J.B and Balogun , E.A (1997)** Some Antimalarial drug-metal complexes, "Synthesis, characterization and *in vivo* evaluation against malarial parasite. *Afr. J. Sci.* **1**:10-12